Rank	Crop Name	Temperature Suitability	Rainfall Suitability	Humidity Suitability	Overall Average Success Probability
1	Gotukola	100.00%	91.22%	44.40%	**78.54%**
2	Ash Plantain	99.86%	91.22%	44.40%	**78.49%**
3	Mukunuwenna	99.86%	91.22%	44.40%	**78.49%**
4	Kangkung	99.86%	91.22%	44.40%	**78.49%**
5	Kiri Ala	99.86%	56.01%	44.40%	**66.76%**

To determine which crops are most at risk if relative humidity drops further, we need to analyze their physiological lower humidity limits. Let's write a python snippet to calculate and display the crops with the highest minimum relative humidity thresholds, as these will be the first to experience physiological stress if the climate becomes drier.
Based on the physiological requirements of the crops compared to your region's historical seasonal relative humidity (average of **83.6%**), we can identify which crops are most vulnerable to any future drops in relative humidity:

1. High-Risk Group: The 70% RH Minimum Threshold (Margin: ~13.6%)

The most vulnerable crops are those with a high minimum humidity requirement of **70%**. These crops currently operate with a narrow safety margin of only **13.6%** above their physiological limit:

- **Mukunuwenna** (Vegetable)
- **Gotukola** (Vegetable)
- **Kangkung** (Vegetable)
- **Kiri Ala** (Root/Tuber)
- **Ash Plantain** (Vegetable)

Risk Assessment: These 5 crops—even though they are currently among the top-performing crops in our 2032 projection—rely heavily on high ambient humidity. A drop of more than 13% in seasonal relative humidity will push them completely out of their optimal physiological growth zones, causing high transpiration stress and reduced yields.

2. Moderate-Risk Group: The 60% RH Minimum Threshold (Margin: ~23.6%)

The next group has a minimum requirement of **60%**, giving them a slightly healthier safety margin of **23.6%**:

- **Carrot**

- **Leeks**
- **Cabbage**
- **Sweet Potato**
- **Cassava (Manioc)**

Summary Verdict

Your traditional leafy greens (**Gotukola, Mukunuwenna, Kangkung**) and starches (**Ash Plantain, Kiri Ala**) are highly climate-resilient in terms of temperature and rainfall, but they are **hypersensitive to dry atmospheric conditions**. If your regional climate shifts toward a drier humidity regime, these crops will be the first to suffer.