

Model-based approach for determining COVID-19 incidence for different testing intensities

Rasmus Kristoffer Pedersen,
Christian Berrig, Viggo Andreasen

PandemiX Center, Dept. Science and Environment,
Roskilde University, Denmark
Email: rakrpe@ruc.dk

ECMTB 2022, Heidelberg
22nd of September, 2022



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

- Different approaches to COVID-19 mitigation throughout the world

Introduction to the problem

- ▶ Different approaches to COVID-19 mitigation throughout the world
- ▶ The impact of differences in data-collection must be understood, also for future research.

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

Introduction to the problem

- ▶ Different approaches to COVID-19 mitigation throughout the world
- ▶ The impact of differences in data-collection must be understood, also for future research.
- ▶ In particular: For each reported case of COVID-19, how many unidentified cases?

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

Introduction to the problem

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

- ▶ Different approaches to COVID-19 mitigation throughout the world
- ▶ The impact of differences in data-collection must be understood, also for future research.
- ▶ In particular: For each reported case of COVID-19, how many unidentified cases?
- ▶ The role of testing: Confirmation of symptoms, required for various activities or entirely voluntary?

- ▶ Different approaches to COVID-19 mitigation throughout the world
- ▶ The impact of differences in data-collection must be understood, also for future research.
- ▶ In particular: For each reported case of COVID-19, how many unidentified cases?
- ▶ The role of testing: Confirmation of symptoms, required for various activities or entirely voluntary?
- ▶ **How do we compare case-counts between periods and places where testing activity was different?**

Introduction to the problem

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data


Relating to data


Discussion

- ▶ Different approaches to COVID-19 mitigation throughout the world
- ▶ The impact of differences in data-collection must be understood, also for future research.
- ▶ In particular: For each reported case of COVID-19, how many unidentified cases?
- ▶ The role of testing: Confirmation of symptoms, required for various activities or entirely voluntary?
- ▶ **How do we compare case-counts between periods and places where testing activity was different?**

Let's look at some data...

From November 1st, 2021 to May 1st, 2022





- We aim to determine the ratio between observed cases and the total number of COVID-19 cases.

- ▶ We aim to determine the ratio between observed cases and the total number of COVID-19 cases.
- ▶ This ratio can be used as a correction-factor for observed data.

- ▶ We aim to determine the ratio between observed cases and the total number of COVID-19 cases.
- ▶ This ratio can be used as a correction-factor for observed data.
- ▶ We extend the classic SIR-model to include voluntary testing that identifies pre- and asymptomatic cases.

- ▶ We aim to determine the ratio between observed cases and the total number of COVID-19 cases.
- ▶ This ratio can be used as a correction-factor for observed data.
- ▶ We extend the classic SIR-model to include voluntary testing that identifies pre- and asymptomatic cases.
- ▶ This allows us to investigate the relationship between the rate of voluntary testing and the appropriate correction-factor.

The model

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



The model

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



$$\dot{S} = -\beta S(P + A)$$

$$\dot{E}_1 = \beta S(P + A) - \gamma E_1$$

$$\dot{E}_2 = \gamma E_1 - \gamma E_2 - \tau E_2$$

$$\dot{P} = \gamma E_2 - \gamma P - \tau P$$

$$\dot{I} = \gamma \rho P - \nu I$$

$$\dot{A} = \gamma(1 - \rho)P - \nu A - \tau A$$

$$\dot{Q} = \tau(E_2 + P + A) - \nu Q$$

$$\dot{R}_p = \nu Q + \nu I$$

$$\dot{R}_n = \nu A$$

The model

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic
Our approach

Model presentation

Analysis

Model dynamics
Fraction identified
Fixed final size

Data and simulations

The data
Relating to data




Discussion

$$\begin{aligned}\dot{S} &= -\beta S(P + A) & \dot{A} &= \gamma(1 - \rho)P - \nu A - \tau A \\ \dot{E}_1 &= \beta S(P + A) - \gamma E_1 & \dot{Q} &= \tau(E_2 + P + A) - \nu Q \\ \dot{E}_2 &= \gamma E_1 - \gamma E_2 - \tau E_2 & \dot{R}_p &= \nu Q + \nu I \\ \dot{P} &= \gamma E_2 - \gamma P - \tau P & \dot{R}_n &= \nu A \\ \dot{I} &= \gamma \rho P - \nu I\end{aligned}$$

<i>Symbol</i>	<i>Description</i>	<i>Default value</i>
β	Infectivity	2/3
ν	Rate of recovery	1/3
γ	Rate of disease progression	1/3
ρ	Fraction of symptomatic cases	1/2
τ	Rate of testing	0 to 0.5

All rates units of day⁻¹. Approximate R_0 of 1.4 initially.

General dynamics like classic SIR-model.



Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and simulations

The data

Relating to data

Discussion

Analysis of fraction of cases identified

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

We consider the fraction of cases identified:

$$K = \frac{r_p}{r_p + r_n} \quad (1)$$

where $r_p = \lim_{t \rightarrow \infty} R_p(t)$ and $r_n = \lim_{t \rightarrow \infty} R_n(t)$.

We consider the fraction of cases identified:

$$K = \frac{r_p}{r_p + r_n} \quad (1)$$

where $r_p = \lim_{t \rightarrow \infty} R_p(t)$ and $r_n = \lim_{t \rightarrow \infty} R_n(t)$.

The correction-factor we want is K^{-1} .

We consider the fraction of cases identified:

$$K = \frac{r_p}{r_p + r_n} \quad (1)$$

where $r_p = \lim_{t \rightarrow \infty} R_p(t)$ and $r_n = \lim_{t \rightarrow \infty} R_n(t)$.

The correction-factor we want is K^{-1} .

Through the methods previously applied in (Andreasen, 2018), we are able to analytically determine:

$$K = 1 - \left(\frac{\nu}{\nu + \tau} \right) \left(1 - \frac{\tau}{\gamma + \tau} \right) \left(1 - \frac{\gamma\rho + \tau}{\gamma + \tau} \right) \quad (2)$$

Analysis of fraction of cases identified

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

We consider the fraction of cases identified:

$$K = \frac{r_p}{r_p + r_n} \quad (1)$$

where $r_p = \lim_{t \rightarrow \infty} R_p(t)$ and $r_n = \lim_{t \rightarrow \infty} R_n(t)$.

The correction-factor we want is K^{-1} .

Through the methods previously applied in (Andreasen, 2018), we are able to analytically determine:

$$K = 1 - \left(\frac{\nu}{\nu + \tau} \right) \left(1 - \frac{\tau}{\gamma + \tau} \right) \left(1 - \frac{\gamma\rho + \tau}{\gamma + \tau} \right) \quad (2)$$

Note that K is independent of β .

Analysis of fraction of cases identified

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

We consider the fraction of cases identified:

$$K = \frac{r_p}{r_p + r_n} \quad (1)$$

where $r_p = \lim_{t \rightarrow \infty} R_p(t)$ and $r_n = \lim_{t \rightarrow \infty} R_n(t)$.

The correction-factor we want is K^{-1} .

Through the methods previously applied in (Andreasen, 2018), we are able to analytically determine:

$$K = 1 - \left(\frac{\nu}{\nu + \tau} \right) \left(1 - \frac{\tau}{\gamma + \tau} \right) \left(1 - \frac{\gamma\rho + \tau}{\gamma + \tau} \right) \quad (2)$$

Note that K is independent of β .

(Andreasen, V. (2018). *Epidemics in Competition: Partial Cross-Immunity*. *Bulletin of Mathematical Biology*, 80(11), 2957-2977. <https://doi.org/10.1007/s11538-018-0495-2>)

Fixed final size and (τ, β)

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data


Relating to data

Discussion

Although the fraction of cases identified, K , is independent of β , the epidemic final size, i.e. $r_n + r_p$, is not.

Let us take a look at the final size as a function of τ and β .

Fixed final size and (τ, β)



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations



The data

Relating to data

Discussion



Fixed final size and (τ, β)



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)



(β chosen such that final size is fixed)

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)



(β chosen such that final size is fixed)

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Fixed final size and (τ, β)



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data


Discussion

The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Daily cases and serology from blood-donors



Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and simulations

The data

Relating to data

Discussion

The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

December 2021: Variant-sampling of 80 to 90% of cases.





The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Smoothing: 7-day running mean.



Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size



Data and
simulations

The data

Relating to data

Discussion

Weekly sums (Monday to Sunday)



The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and simulations

The data

Relating to data

Discussion

Sum of weekly tests



(Only PCR shown, Antigen-tests at similar magnitude)

The Danish data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and simulations

The data

Relating to data


Discussion

Sum of weekly tests



Scaled to tests per population per day (i.e. τ)

Relating to data



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Relating to data



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Relating to data



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and
simulations

The data

Relating to data

Discussion

Relating to data



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion

Relating to data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



Relating to data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



Relating to data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size


Data and simulations

The data

Relating to data

Discussion

τ from data, β : 0.48, initial infections: 7000



Relating to data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



Relating to data

Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified


Fixed final size

Data and simulations

The data

Relating to data

Discussion



- For comparing the impact of COVID-19 between countries, accurate estimates of final size are necessary, particular when evaluating mitigation strategies.

- ▶ For comparing the impact of COVID-19 between countries, accurate estimates of final size are necessary, particular when evaluating mitigation strategies.
- ▶ Using an extended SIR-model, we are able to estimate the fraction of COVID-19 cases identified in the Omicron wave of early 2022 in Denmark.

- ▶ For comparing the impact of COVID-19 between countries, accurate estimates of final size are necessary, particular when evaluating mitigation strategies.
- ▶ Using an extended SIR-model, we are able to estimate the fraction of COVID-19 cases identified in the Omicron wave of early 2022 in Denmark.
- ▶ The simple model allows for analytical results about the epidemic final size in addition to simulations.

- ▶ For comparing the impact of COVID-19 between countries, accurate estimates of final size are necessary, particular when evaluating mitigation strategies.
- ▶ Using an extended SIR-model, we are able to estimate the fraction of COVID-19 cases identified in the Omicron wave of early 2022 in Denmark.
- ▶ The simple model allows for analytical results about the epidemic final size in addition to simulations.
- ▶ Results suggest between 58% and 63% were identified, a little lower than official estimates suggesting two-thirds.

- ▶ For comparing the impact of COVID-19 between countries, accurate estimates of final size are necessary, particular when evaluating mitigation strategies.
- ▶ Using an extended SIR-model, we are able to estimate the fraction of COVID-19 cases identified in the Omicron wave of early 2022 in Denmark.
- ▶ The simple model allows for analytical results about the epidemic final size in addition to simulations.
- ▶ Results suggest between 58% and 63% were identified, a little lower than official estimates suggesting two-thirds.
- ▶ Future work consists of further analysis, parameter-fitting and application to other countries.

Thank you for your attention.

Any questions?



Feel free to also contact me
with questions or comments
later

Website:

rasmuspedersen.com

Email: rakrpe@ruc.dk



Determining
COVID incidence

Pedersen, Berrig &
Andreasen

Introduction

The problematic

Our approach

Model presentation

Analysis

Model dynamics

Fraction identified

Fixed final size

Data and
simulations

The data

Relating to data

Discussion