

PATIENT NAME : B/O DEEPA THAVAMANI**REF. DOCTOR : DR. D V SURESH****CODE/NAME & ADDRESS : C000113625**SKL DIAGNOSTICS CENTRE
NEAR NEURO FOUNDATION, MEYANOR MAIN 3
ROAD,
SALEM 636004
9443240665**ACCESSION NO : 0081VL022010****PATIENT ID : BODEF28092281****CLIENT PATIENT ID:****ABHA NO :****AGE/SEX : 3 Months Female****DRAWN : 27/12/2022 00:00:00****RECEIVED : 28/12/2022 16:57:23****REPORTED : 07/01/2023 10:39:41****Test Report Status Final****Results****Biological Reference Interval Units****ALKALINE PHOSPHATASE ISOENZYMES ELECTROPHORESIS**

ALKALINE PHOSPHATASE	1381 High	122 - 469	U/L
L1 (LIVER)	2.2	1 - 31	%
L2 (LIVER)	14.4 High	1 - 7	%
BONE	83	60 - 100	%
I1 (INTESTINE)	0.4	< 14	%
I2 (INTESTINE)	-	< 14	%
I3 (INTESTINE)	-	< 14	%
P1 (PLACENTAL)	-		
P2 (PLACENTAL)	-		%

Comments

The sample for Alkaline Phosphatase Isoenzymes Electrophoresis, was referred to an external laboratory (code no.20002) for performance of the test requested.

Interpretation(s)

ALKALINE PHOSPHATASE ISOENZYMES ELECTROPHORESIS-

Alkaline Phosphatase Isoenzymes Electrophoresis Interpretation :

Liver Isoenzymes : There are two liver isoenzymes, L1 and L2. L1 is increased in some non - malignant diseases (Such as cholestasis, cirrhosis, viral hepatitis and in various biliary and hepatic pathologies). It is also increased in malignancies with hepatic metastasis, in cancer of the lungs and digestive tract and in lymphoma. The L2 fraction exists in healthy subjects at low concentration (< 8 IU/L). An increase of L2 may occur in cholestasis and biliary diseases and in malignancies (e.g., breast, liver, lung, prostate, digestive tract) with liver metastasis.

Bone Isoenzyme : The bone fraction is interpreted with respect to age. Increase seen in :

- Breast cancer with bone or liver metastasis, Osteosarcoma and lymphoma with or without metastasis
- Rheumatismal diseases, hyperparathyroidism, Paget disease and rickets.

- Transient hyperphosphatasemia in infancy that is associated with a particular liver fraction .

- In such case the total ALP is highly elevated (> 2000 IU/L).

- When associated with an increase of other fraction (e.g. L1, L2) it may indicate malignancy of rectum, Lung and prostate (with bone and liver metastasis).

Intestinal Isoenzymes : They are absent in about 60% of normal subjects and when present are less than 14% .

Placental Isoenzyme :

Exists as P1 (90%) and P2 (10%). It is found during pregnancy & in some malignant diseases, e.g.:- ovarian cancers, sarcomas, pancreatic & stomach cancers. Increase can be also due to heavy smoking.

****End Of Report******Please visit www.srlworld.com for related Test Information for this accession***S.S. Wadalkar***Dr. Sneha Wadalkar, M.D**
(Reg.no.MMC2012/06/1868
Junior Biochemist

Page 1 Of 1



View Details



View Report

PERFORMED AT :SRL Ltd
PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL ESTATE, S.V. ROAD, GOREGAON (W)
MUMBAI, 400062
MAHARASHTRA, INDIA
Tel : 9111591115, Fax : 022 - 67801212
CIN - U74899PB1995PLC045956**Patient Ref. No. 81000002327386**