


# *The dark matter of the genome and blood pressure regulation – modeling non-coding genetic mechanisms in cellular models and rats*

Aron Geurts, PhD  
Department of Physiology  
Oct 6, 2024

## Acknowledgements



P01HL149620



Mingyu Liang, MD, PhD

Sid Rao, MD, PhD

Anne Kwitek, PhD

Andy Greene, PhD

Aron Geurts, PhD

Allen Cowley, PhD

Yong Liu, PhD

Pengyuan Liu, PhD

Curt Sigmund, PhD

Paul Auer, PhD

Mike Widlansky, MD

Hong Xue, PhD

Manoj Mishra, PhD

Ranjan Pandey, PhD

Atrayee Ray, PhD

Chun Yang, PhD

Joan Qiu, PhD

Monika Tutaj, PhD



Cary Stelloh

Terry Kurth

Mark Vanden Avond

Mike Grzybowski, MS

Jason Klotz, MBA

Bhavika Therani

Lishu He

Kristie Usa



## UNIVERSITY OF FLORIDA CoM

Abdel Alli, PhD

Niharika Bala, PhD



## SOUTH-SIDE CREW

Mindy Dwinell, PhD

Akiko Takizawa, PhD

Angela Lemke

Becky Schilling

Allison Zappa

Joe Herbst

Tristan Dooley

Lynn Lazcares

Christopher Bach

Emily Novak (past)

## RODENT MODEL RESOURCE

Shawn Kalloway (past)


Jamie Foeckler (past)

Tiffany Raatz

# Human BP SNPs -> rat orthology & human iPSC-derived cell types

Overall hypothesis: Non-coding, non-transcribed SNPs must exert their effects on phenotypes by modifying expression of gene(s) in some tissue(s) to modify one or more physiological mechanisms relevant to the trait.


•PMCID: [PMC11096100](#)




<1% missense or nonsense variants,  
>20% >10-kbp from nearest gene

•PMCID: [PMC703516](#)

7




•PMID: 15057822





Syntenic and sequence-level conservation

## rs1173771 (rs771)





[GWAS Catalog \(ebi.ac.uk\)](http://www.ebi.ac.uk/gwas/)





# GTex Portal

# We deleted the entire homologous non-coding region in SS using CRISPR-Cas9








Systolic  
blood  
pressure was  
reduced in  
 $\Delta$ LD male  
rats




# CRISPR-SpCas9 gene edited iPSC models reveal similar effects of the haplotype and single rs771 on Npr3



Deletion model



Haplotype  
reconstituted  
models



Single rs1173771  
edited model





# NPR3 encodes NPR-C, a multifunctional receptor

- In the vasculature, NPR-C binds C-type natriuretic peptide (CNP), mediates the effect of CNP on vascular function and structure, and plays an important role in preserving vascular homeostasis in vivo.
  - Moyes, et al. *J Clin Invest.* 2014; PMID: 25105365
  - Villar, et al. *Cardiovasc Res.* 2007; PMID: 17391657
- In the kidney, NPR-C facilitates the clearance of atrial, B-type, and C-type natriuretic peptides (ANP, BNP, CNP) from the circulation via endocytosis.
  - Maack, et al. *Science*, 1987; PMID: 2823385
  - Almeida, et al. *Am J Physiol.* 1989; PMID: 2537040




# $\Delta$ LD has tissue-specific effects on Npr3 expression and function

Hong Xue  
Abdel Alli

## Mesenteric Artery




## Kidney






# The BP elevating haplotype increases contact frequency between the LD region and the Npr3 promoter


- Region Capture Micro-C reveals coalescence of enhancers and promoters into nested microcompartments | Nature Genetics




**A.** iEC contact map (BP lowering haplotype)







**B.** iEC contact map (BP elevating haplotype)



**C.** iVSMC contact map (BP lowering haplotype)




**D.** iVSMC contact map (BP elevating haplotype)




# rs1882961 C->T (rs961)

riskAllele	pValue	pValueAnnotation	riskFrequency	orValue	beta	ci	Mapped Genes	traitName	Traits	bkgTraits	accessionId	locations	pubmedId	author
rs1882961-?	3.00E-13	-	NR	-	-	-	LINC02920,CYCSP42	Systolic blood pressure	systolic blood pressure	-	GCST007087	21:15184047	30595370	Kichaev G

[GWAS Catalog \(ebi.ac.uk\)](http://www.ebi.ac.uk/gwas/)



*No evidence of eQTL, 120-kb away from nearest gene*



Editing this conserved region in SS rats results in sex-divergent effects on blood pressure traits

Fig 1







Fig 2



# Effects on local genes (very preliminary data)

$\log_2FC$  KO/WT,  
FDR<0.05





# Overall conclusions

- The majority of loci (now >2,000) associated with blood pressure are non-coding and hundreds of functional SNP may be quite distant from the genes they regulate
- Human iPSCs can be engineered to harbor blood pressure elevating and lowering alleles and haplotypes then differentiated to study their effects on gene expression
- At least some human non-coding loci can be manipulated in animal models to provide functional evidence of their role in complex traits
- Unlike knockout models, each engineered SNP model will have unique tissue(s)-specific (and sex-specific!) effects on gene(s) expression and phenotypes, leading to complex hypotheses about their mechanisms

Thanks!  
Questions?



**maeby fünke**





@26percentbad

After all, aren't we all just fat rats, stuck in the manhole covers of life?




12:31 PM - 27 Feb 2019

Systolic blood pressure was reduced in  $\Delta$ LD male rats





Proposed mechanisms by which non-coding SNPs could influence genes and blood pressure





[Lift Genome Annotations \(ucsc.edu\)](http://lift.ucsc.edu)

Conservation in rats extends to a ~30.4-kbp region

