

# Knockout Rats via Embryo Microinjection of Zinc-Finger Nucleases


Aron M. Geurts,<sup>1,2\*</sup> Gregory J. Cost,<sup>3\*</sup> Yevgeniy Freyvert,<sup>3</sup> Bryan Zeitler,<sup>3</sup> Jeffrey C. Miller,<sup>3</sup> Vivian M. Choi,<sup>3</sup> Shirin S. Jenkins,<sup>3</sup> Adam Wood,<sup>4</sup> Xiaoxia Cui,<sup>4</sup> Xiangdong Meng,<sup>3</sup> Anna Vincent,<sup>3</sup> Stephen Lam,<sup>3</sup> Mięczysław Michalkiewicz,<sup>1,2</sup> Rebecca Schilling,<sup>1,2</sup> Jamie Foeckler,<sup>3</sup> Shawn Kalloway,<sup>3</sup> Hartmut Weiler,<sup>1,2</sup> Séverine Ménoret,<sup>5</sup> Ignacio Anegón,<sup>5</sup> Gregory D. Davis,<sup>4</sup> Lei Zhang,<sup>3</sup> Edward J. Rebar,<sup>3</sup> Philip D. Gregory,<sup>3</sup> Fyodor D. Urnov,<sup>3</sup> Howard J. Jacob,<sup>1,2,6†</sup> Roland Buelow<sup>7†</sup>

The laboratory rat is a well-established model for the genetic dissection of human disease-related traits (1) despite the fact that targeted modification of its genome is largely intractable. We investigated the application of

We delivered these ZFNs to 36 hemizygous GFP-transgenic (5) inbred SS (Dahl S; GFP ZFNs), 91 inbred FHH (Fawn-hooded hypertensive; Rab38 ZFNs), and 2793 outbred SD (Sprague Dawley; IgM ZFNs) embryos by pronuclear or

breeding to wild-type animals, one out of one GFP and three out of four IgM mutations were transmitted through the germline, one of which was subsequently bred to homozygosity (table S1 and fig. S4).

The high percentage of disrupted chromosomes demonstrates that ZFNs are active in early rat embryos from three strains, leading to both mono- and biallelic gene disruption. Although we observed no cleavage at predicted off-target sites, such events could be segregated away from the desired mutation by backcrossing to the parental strain. ZFN-driven gene disruption and germline transmission can be accomplished in 4 months' time, and ZFNs can be engineered against a broad range of sequences (6, 7); this strategy adds a valuable tool to an increasingly powerful rat genetic toolbox, opening up a range of new experiments and models of human disease.



PNI - pronuclear injection; IC - intracytoplasmic; \* - of transferred; † - of liveborn founders; ND - not determined

**Fig. 1.** ZFN-mediated gene disruption in rat embryos. (A) ZFNs containing five or six fingers were designed to target coding sequences of interest (gray lines) for site-specific cleavage. (B) Two of five pups born after microinjection of GFP-targeted ZFNs were devoid of GFP expression. (C) Polymerase chain reaction using GFP-specific primers revealed truncated but no wild-type sequence in each of the GFP negative pups compared with positive littermates. SS indicates Dahl S control DNA; NT indicates no template. (D) Table of injection data revealing successful mutagenesis of the three gene targets after multiple delivery methods and doses in three rat strains.

engineered zinc-finger nucleases [ZFNs (2)] for the elimination of specific rat gene functions and generation of knockout rats. ZFNs induce site-specific, double-strand DNA breaks that can be repaired by the error-prone nonhomologous end-joining DNA repair pathway to result in a targeted mutation (Fig. 1A). In the fruit fly and zebrafish, direct embryo injection of ZFN-encoding mRNA has been used to generate heritable knockout mutations at specific loci (2).

The design and validation of three sets of ZFN reagents that target the green fluorescent protein (GFP) gene and two endogenous rat genes, *Immunoglobulin M* (IgM) and *Rab38*, were performed as described (3) and are detailed in (4). To take advantage of the potential for greater specificity of action afforded by longer (and therefore rarer) targets, we used five- and six-finger ZFNs.

intracytoplasmic injection of ZFN-encoding DNA or mRNA at different concentrations (table S1). Screening 295 founder animals yielded 35 (12%) that harbored targeted mutations.

Full knockout of the GFP transgene was achieved because mutant animals lacked both GFP expression and wild-type GFP sequence (Fig. 1, B and C). Thirty-two IgM mutants and the single Rab38 mutant carried 25 to 100% disrupted target chromosomes (fig. S1). Sequence analysis of 18 founders revealed deletion alleles ranging from 3 to 187 base pairs; of note, one animal carried biallelic mutations in IgM (table S1). Furthermore, ZFN-mediated gene disruption demonstrated high fidelity for each target sequence because no ZFN-induced mutations were detected in target gene-disrupted animals at any of 20 predicted ZFN off-target sites (figs. S2 and S3). After

## References and Notes

1. T. J. Aitman *et al.*, *Nat. Genet.* **40**, 516 (2008).
2. D. Carroll, *Gene Ther.* **15**, 1463 (2008).
3. Y. Doyon *et al.*, *Nat. Biotechnol.* **26**, 702 (2008).
4. Materials and methods are available as supporting material on Science Online.
5. M. Michalkiewicz *et al.*, *Am. J. Physiol. Heart Circ. Physiol.* **293**, H881 (2007).
6. C. O. Pabo, E. Peisach, R. A. Grant, *Annu. Rev. Biochem.* **70**, 313 (2001).
7. A. Klug, *Proc. Jpn. Acad.* **81**, 87 (2005).
8. We thank R. Jaenisch, R. Hammer, P. Sullivan, and three anonymous referees for helpful suggestions; D. Smoller and E. Lanphier for support; E. Eastlund for the Rab38 ZFN mRNA; R. DeKeller and R. Amora for technical assistance; and Caliper Life Sciences, Incorporated for excellent service. Supported by NIH grants 5U01HL066579-08 and 5P01HL082798-03, a sponsored research agreement between the Medical College of Wisconsin and Sigma-Aldrich, and the American Physiological Society Fellowship in Physiological Genomics to A.M.G. The authors are filing patents based on the results reported in this paper.

## Supporting Online Material

www.sciencemag.org/cgi/content/full/325/5939/433/DC1  
Materials and Methods  
Figs. S1 to S5  
Tables S1 and S2  
References

18 February 2009; accepted 1 May 2009  
10.1126/science.1172447

<sup>1</sup>Human and Molecular Genetics Center, Medical College of Wisconsin, Milwaukee, WI 52336, USA. <sup>2</sup>Department of Physiology, Medical College of Wisconsin, Milwaukee, WI 52336, USA. <sup>3</sup>Sangamo BioSciences, Incorporated, Richmond, CA 94804, USA. <sup>4</sup>Sigma-Aldrich Biotechnology, St. Louis, MO 63103, USA. <sup>5</sup>INSERM, UMR 643, CHU, Nantes, Université de Nantes, 44322 Nantes, France. <sup>6</sup>Department of Pediatrics, Medical College of Wisconsin, Milwaukee, WI 52336, USA. <sup>7</sup>Open Monoclonal Technology, Incorporated, Palo Alto, CA 94303, USA.

\*These authors contributed equally to this work.

†To whom correspondence should be addressed. E-mail: jacob@mcw.edu (H.J.); rbuelow@omtinc.net (R.B.)