

WIRSCHING’S CONJECTURE 3, AND THE PERIODIC CORRECTION TO BERG–KRÜPPEL’S DENSITY ASYMPTOTIC

RENATO AUGUSTO TAVARES

ABSTRACT. Wirsching’s 2003 argument for positive predecessor density under the $3n + 1$ map reduces to three conjectures. Conjecture 3 concerns the near-zero asymptotics of an invariant density φ , compared against an explicit asymptotic φ_0 due to Berg and Krüppel. The exact log-Laplace transform of φ decomposes as a smooth part plus a periodic correction H . Berg and Krüppel gave this correction, for this eigenfunction, as an infinite product in 1998, five years before Wirsching’s conjecture; they did not ask whether it is constant. We give H instead as a Fourier series with closed-form coefficients in Γ and ζ , and certify rigorously that it is not. Wirsching’s own comparison class fixes a single phase of H ; there, Conjecture 3 holds, with an explicit limit. Away from that class the phase sweeps a full period, so the stronger, unrestricted asymptotic $\varphi(t) \sim \kappa \varphi_0(t)$ as $t \rightarrow 0^+$ fails. Wirsching needs less than the literal statement of Conjecture 3; we check directly that his actual requirement holds with a wide margin.

1. INTRODUCTION

Let T denote the $3n+1$ map on the positive integers, $T(n) = n/2$ for n even and $T(n) = (3n+1)/2$ for n odd. Wirsching’s monograph [3] develops the predecessor-set machinery this paper’s argument descends from; [1] takes it further, studying the density of predecessor sets under T through an averaging construction on the 3-adic integers: a family of operators S_l , built from path counts in the predecessor graph, converges in the strong operator topology to a limiting operator S_∞ whose essential part is a Markov chain on $[0, 1]$. That chain has a unique invariant density φ , and Wirsching’s proof of positive predecessor density reduces to three conjectures about φ and related quantities, stated as three unbridged gaps in the argument.

The load-bearing one is Conjecture 3. It concerns the behaviour of φ as its argument tends to 0, compared against an explicit comparison function φ_0 . Berg and Krüppel [2] had already produced φ_0 five years earlier, by Laplace transform and saddlepoint analysis, as the asymptotic solution of a truncated version of the functional equation φ satisfies. Their argument for the asymptotic comparability of φ and φ_0 was informal: they give the periodic correction relating the two as an explicit infinite product (Corollary 3 below) but never evaluate it, so whether a single constant relates φ to φ_0 across every sequence, and whether the ratio converges at all along an arbitrary one, are both left open. Wirsching considers the strongest possible reading of that gap directly:

“That replacement would be possible, if, e.g., $\varphi \sim c \varphi_0$ for some constant $c > 0$. It turns out that the following (slightly weaker) conjecture . . . suffices.”

He then states ([1], immediately after the quoted sentence, with notation normalized: his own δ_5 is written δ throughout this paper, since no other δ -indexed class of his appears here, and his constant c , kept below for fidelity to his statement, is unrelated to $c := \log 3$ introduced in Section 3):

2020 *Mathematics Subject Classification*. 11B83, 39B22, 30D05, 11M06.

Key words and phrases. Collatz map, predecessor density, invariant density, log-periodic asymptotics, Mellin transform, atomic function.

Wirsching's Conjecture 3. *There are positive real constants c and δ such that*

$$\lim_{l \rightarrow \infty} \frac{\varphi(z_l)}{\varphi_0(z_l)} = c > 0 \quad \text{uniformly for sequences } (z_l) \in \tilde{A}_\delta.$$

Here $\tilde{A}_\delta$ is Wirsching's class of sequences tending to 0 at the borderline rate $z_l \sim l \cdot 3^{-l}$ (his equations (1.5) and (3.2), reproduced ahead of Theorem 1's proof): a sequence (x_l) belongs to $\tilde{A}_\delta$ exactly when $\lfloor 3^l x_l \rfloor = k_l$ for a sequence of integers (k_l) with $|l - k_l| \leq \delta \sqrt{l}$ for every l . This is weaker than the unrestricted asymptotic he first considers. The constant cancels out of the one inequality his own proof actually needs, so a class-dependent limit is enough.

Neither statement had been settled. This paper settles both, in opposite directions.

Theorem 1 (Conjecture 3). *Wirsching's Conjecture 3 (stated just above) is true. For every $\delta > 0$, on his comparison class $\tilde{A}_\delta$, the ratio $\varphi(z_l)/\varphi_0(z_l)$ converges uniformly to $e^{H(0)} = 0.534122\dots$, where H is the periodic function of Theorem 4 below and the normalization of φ_0 is Berg and Krüppel's own, equation (9.6) of [2]. Conjecture 3 itself only asserts the existence of some positive constant, independently of normalization; the specific value $e^{H(0)}$ is tied to this particular normalization of φ_0 .*

Theorem 2 (Failure of the unrestricted asymptotic). *The stronger asymptotic $\varphi(t) \sim \kappa \varphi_0(t)$ as $t \rightarrow 0^+$, for any single constant κ and with no restriction to a comparison class, is false, by Proposition 8's unconditionally proved non-constancy of H . The ratio $\varphi(t)/\varphi_0(t)$ oscillates indefinitely as $t \rightarrow 0^+$, with amplitude bounded below by a positive, certified constant: it is governed by the exactly c -periodic function H evaluated at the saddle location $w_0(\tau)$, $\tau = \log(1/t)$, whose successive-period increment $w_0(\tau + c) - w_0(\tau)$ tends to c (Lemma 15) even though $w_0(\tau)$ itself drifts away from τ without bound.*

The two theorems have a single common source. Write $t = e^{-\tau}$ and let $H(w)$ be the $\log 3$ -periodic function appearing in the exact decomposition of φ 's log-Laplace transform (Theorem 4); the saddlepoint bridge of Sections 4 and 5 shows $\log \varphi(t) - \log \varphi_0(t) = H(w_0(\tau)) + o(1)$, where the saddle location $w_0(\tau)$ advances by c per period of τ , up to a vanishing correction: $w_0(\tau + c) - w_0(\tau) = c + O(1/\tau)$ (Lemma 15). Wirsching's comparison class constrains $w_0(\tau_l)$'s phase modulo $\log 3$ to converge to a single value as $l \rightarrow \infty$; the unrestricted range $\tau \rightarrow \infty$ with no class restriction instead sweeps that phase through every value, since Φ_0 is a bijection (again Lemma 15). Whether the ratio converges or oscillates depends on which sequences one compares along, and the two theorems above read the same object at two different scopes.

Corollary 3. *e^H is Berg and Krüppel's own periodic factor for the eigenfunction φ , from Proposition 9.3 of [2], which they give as an infinite product; H itself is its logarithm. Its Fourier series, Proposition 6 below, is new.*

Section 2 recalls the functional equation φ satisfies and Berg and Krüppel's comparison asymptotic. Section 3 constructs H , proves Proposition 6, and certifies that H is not constant. Sections 4 and 5 transfer this from the log-Laplace transform to φ itself, through a uniform saddlepoint approximation and an envelope estimate. Section 6 identifies the resulting smooth part with Berg and Krüppel's saddlepoint expression, pins down every constant, and proves Corollary 3. Section 7 proves both theorems and checks Wirsching's actual requirement directly. Section 8 states what remains open.

2. THE INVARIANT DENSITY AND ITS LAPLACE TRANSFORM

Let $U_1, U_2, \dots$ be independent, identically distributed uniform random variables on $[0, 1]$, and set

$$X = \sum_{j \geq 1} \frac{2}{3^j} U_j \in [0, 1].$$

The self-similar identity $X = \frac{2}{3}U_1 + \frac{1}{3}X'$, with X' an independent copy of X , shows that the density φ of X satisfies

$$\varphi(x) = \frac{3}{2} \int_{3x-2}^{3x} \varphi,$$

equivalently $\varphi'(x) = \frac{9}{2}\varphi(3x)$ on $(0, \frac{2}{3})$: the displayed integral equation is exactly [1]'s equation (6.1), $(W_3f)(x) := \frac{3}{2} \int_{3x-2}^{3x} f$, and φ is exactly the function his Corollary 7 constructs as the unique L^1_{loc} solution of $\text{supp } \varphi \subset [0, 1]$, $\int_0^1 \varphi = 1$, $W_3\varphi = \varphi$ – his invariant density, up to the reflection $x \mapsto 1-x$ fixing the convention. It is also a translate of Rvachev's atomic function h_3 [4], and the dilation-3, $\lambda = 2/3$ member of Berg and Krüppel's family of functional equations $\lambda \psi'(t) = 3(\psi(3t) - \psi(3t-2))$ (their generic family member, distinct from this paper's own g introduced below). The reflection is in fact vacuous, $\varphi(x) = \varphi(1-x)$ identically: X and $1-X = \sum_{j \geq 1} \frac{2}{3^j}(1-U_j)$ are equidistributed, since $1-U_j \sim U_j$ independently for every j and $\sum_{j \geq 1} 2 \cdot 3^{-j} = 1$. It is C^∞ , supported on $[0, 1]$, and constant, equal to $3/2$, on the middle third $[1/3, 2/3]$.

For $s > 0$ write

$$K(s) = \log \mathbb{E}[e^{-sX}] = \sum_{j \geq 1} g\left(\frac{2s}{3^j}\right), \quad g(b) = \log \frac{1-e^{-b}}{b}.$$

The branch of g is the one analytic on $\{\text{Re } w > 0\}$ and real on $(0, \infty)$; since $(1-e^{-w})/w$ is analytic and non-vanishing there (its zeros lie on $2\pi i\mathbb{Z} \setminus \{0\}$, and $w = 0$ is a removable singularity with value 1), and the half-plane is simply connected, g exists and is unique. Differentiating termwise, justified because $g(w) = -w/2 + O(w^2)$ as $w \rightarrow 0$ so the tail of the series is normally convergent together with every derivative,

$$t(s) := -K'(s), \quad V(s) := s^2 K''(s) > 0,$$

the second an exact positivity, $K''(s)$ being the variance of X under its exponential tilt by e^{-sX} . Since $t'(s) = -K''(s) < 0$, t is strictly decreasing on $(0, \infty)$; at the endpoints, $t(0^+) = \mathbb{E}[X] = \sum_{j \geq 1} 2 \cdot 3^{-j} \cdot \frac{1}{2} = \frac{1}{2}$, and $t(s) \rightarrow 0$ as $s \rightarrow \infty$ because the exponential tilt concentrates on $\text{ess inf } X = 0$ (each factor e^{-sX} suppresses any mass away from 0 as $s \rightarrow \infty$, and $X \geq 0$ a.s.). So $t(s)$ decreases strictly from $1/2$ to 0, and studying $\varphi(t(s))$ as $s \rightarrow \infty$ is the same as studying $\varphi(t)$ as $t \rightarrow 0^+$.

Berg and Krüppel's comparison function solves the truncated equation $\lambda \psi'(t) = a \psi(at)$, a here their dilation parameter (this letter is reused in Section 3 for this paper's own, unrelated linear coefficient of Q ; the two never appear in the same formula except in Section 6, where the collision is flagged again at the point it matters), by Laplace transform and saddlepoint analysis. In their normalization (their equation (9.6), with $\alpha = \frac{1}{2} - \frac{\log 2}{\log 3}$, $\beta = \frac{1}{2 \log 3}$, and $\gamma, \delta_{\text{BK}}, \varepsilon$ determined by α and β through their Proposition 9.1, printed explicitly in Section 6),

$$\varphi_0(t) = \frac{(2\beta)^\varepsilon}{\sqrt{2\pi}} t^\gamma (-\log t)^{\delta_{\text{BK}}} \exp\left(-\beta \log^2 \frac{t}{-\log t}\right).$$

Write $\varphi_{0,\text{bare}}$ for the same expression without the leading constant $(2\beta)^\varepsilon/\sqrt{2\pi}$: the two differ only by that constant.

3. THE PERIODIC CORRECTION

Set $c = \log 3$ and write $w = \log s$, $L(w) = K(e^w)$. Substituting into the defining series and splitting off the $j = 1$ term against the rest reproduces the recurrence

$$(1) \quad L(w+c) - L(w) = \log(1 - e^{-2e^w}) - w - \log 2.$$

Let

$$Q(w) = -\frac{w^2}{2c} + \left(\frac{1}{2} - \frac{\log 2}{c}\right)w,$$

chosen so that $Q(w+c) - Q(w) = -w - \log 2$, matching (1)'s non-periodic terms exactly. Write $a := \frac{1}{2} - \frac{\log 2}{c}$ for Q 's linear coefficient, used again from Section 5 onward.

Theorem 4. *Define*

$$(2) \quad H(w) := L(w) - Q(w) + \sum_{k \geq 0} \log(1 - e^{-2 \cdot 3^k e^w}).$$

Then H is exactly c -periodic, $H(w+c) = H(w)$ for every $w \in \mathbb{R}$, and

$$L(w) = Q(w) + H(w) + \Delta(w), \quad \Delta(w) = - \sum_{k \geq 0} \log(1 - e^{-2 \cdot 3^k e^w}),$$

with $\Delta^{(j)}(w) = O_j(e^{jw-2e^w})$ for every $j \geq 0$: doubly exponentially small, together with all its derivatives, as $w \rightarrow +\infty$.

Proof. Write $d(w) = \log(1 - e^{-2e^w})$, so (1) reads $L(w+c) - L(w) = d(w) - w - \log 2$. Directly from the definition of Q (expand $(w+c)^2 = w^2 + 2wc + c^2$ and collect terms), $Q(w+c) - Q(w) = -w - \log 2$ as well, so the two non-periodic parts cancel exactly:

$$(L - Q)(w+c) - (L - Q)(w) = [d(w) - w - \log 2] - [-w - \log 2] = d(w).$$

The series in (2) telescopes this exactly: with $D(w) := (L - Q)(w)$,

$$D(w) + \sum_{k \geq 0} d(w + kc) = D(w+c) - d(w) + \sum_{k \geq 0} d(w + kc) = D(w+c) + \sum_{k \geq 0} d(w + (k+1)c),$$

so the right side of (2) is invariant under $w \mapsto w+c$, that is, H is c -periodic. Convergence of the series and its termwise differentiability, together with the stated bound on $\Delta^{(j)}$, follow from $d(w) = O(e^{-2e^w})$ and the corresponding bounds on its derivatives, since $2 \cdot 3^k e^w \geq 2e^w$ for $k \geq 0$ makes the series dominated, uniformly for w in any half-line bounded away from $-\infty$, by a geometric series in 3^{-k} inside the exponent. $\square$

The decomposition $L = Q + H + \Delta$, with Δ negligible as $w \rightarrow \infty$, is unique: any two periodic functions differing from $L - Q$ by an $o(1)$ term as $w \rightarrow \infty$ must be equal, being periodic functions that agree in the limit along every residue class of w modulo c .

Remark 5. Berg and Krüppel's own Proposition 9.3, in their own notation throughout this remark only (their a, b are unrelated to this paper's identically-named objects and are not used again outside this remark, Section 6 below reusing a for their dilation parameter in a way disambiguated there; their α , by contrast, is exactly Section 2's imported α , which Proposition 17 below identifies with this paper's own a), specialized to $a = 3, b = 3/2$ (exactly this eigenfunction), gives the equivalent identity $\exp(H(w)) = a^{\frac{1}{2}t^2 - \alpha t} \prod_{k \geq 0} \frac{1 - e^{-a^{t-k}/b}}{a^{t-k}/b} \prod_{l \geq 1} (1 - e^{-a^{t+l}/b})$, $t = w/c$, as an infinite product, in their proof of that proposition: this is the same object as Q, L , and Δ recombine into via $\exp(H) = e^{-Q} e^L e^{-\Delta}$, so Proposition 8's primary computation from (7) below is, in substance, an evaluation of this product. What Proposition 6's Fourier series supplies instead, and their product form does not, are the derivative bounds (5), on which the rest of the paper's saddlepoint estimates depend.

Proposition 6. *H has the Fourier expansion $H(w) = \sum_{m \in \mathbb{Z}} \hat{H}(m) e^{i\omega_m w}$, $\omega_m = 2\pi m/c$, with*

$$\hat{H}(0) = \frac{\log 2}{2} - \frac{\log^2 2}{2c} - \frac{c}{12} - \frac{A}{c}, \quad A = \frac{\pi^2}{12} - \frac{\gamma_E^2}{2} - \gamma_1,$$

$$\hat{H}(m) = -\frac{2^{i\omega_m}}{c} \Gamma(-i\omega_m) \zeta(1 - i\omega_m), \quad m \neq 0,$$

where γ_E is the Euler–Mascheroni constant and γ_1 the first Stieltjes constant.

Proof. Both g and K have Mellin transforms on the strip $-1 < \operatorname{Re} z < 0$. As $b \rightarrow 0^+$, $g(b) = -b/2 + O(b^2)$; as $b \rightarrow \infty$, $g(b) = -\log b + O(e^{-b})$. So $\int_0^\infty |g(b)| b^{\sigma-1} db < \infty$ for $-1 < \sigma < 0$, and the same two bounds applied termwise give $K(s) = O(s)$ as $s \rightarrow 0^+$ and $K(s) = O(\log^2 s)$ as $s \rightarrow \infty$: the terms with $3^j \leq 2s$ number $O(\log s)$ and are each $O(\log s)$, and the terms with $3^j > 2s$ sum to $O(1)$.

Integrate by parts. With $g'(b) = 1/(e^b - 1) - 1/b$, and both boundary terms vanishing on the strip $(g(b)b^z \rightarrow 0 \text{ at } 0 \text{ because } \operatorname{Re} z > -1, \text{ and at } \infty \text{ because } \operatorname{Re} z < 0)$,

$$g^*(z) = \int_0^\infty g(b) b^{z-1} db = -\frac{1}{z} \int_0^\infty \left(\frac{1}{e^b - 1} - \frac{1}{b} \right) b^z db = -\frac{\Gamma(z+1)\zeta(z+1)}{z} = -\Gamma(z)\zeta(1+z),$$

the middle step being the classical continuation $\int_0^\infty \left(\frac{1}{e^u - 1} - \frac{1}{u} \right) u^{v-1} du = \Gamma(v)\zeta(v)$, valid for $0 < \operatorname{Re} v < 1$, at $v = z + 1$.

Now the harmonic sum. In the j -th term substitute $u = 2s/3^j$, so $s = 3^j u/2$ and $s^{z-1} ds = (3^j/2)^z u^{z-1} du$:

$$\int_0^\infty g\left(\frac{2s}{3^j}\right) s^{z-1} ds = \left(\frac{3^j}{2}\right)^z \int_0^\infty g(u) u^{z-1} du = \left(\frac{3^j}{2}\right)^z g^*(z).$$

Summing over $j \geq 1$ is legitimate termwise, since $\sum_{j \geq 1} (3^j/2)^\sigma \int_0^\infty |g(u)| u^{\sigma-1} du$ converges for $-1 < \sigma < 0$. The multiplier that appears is therefore

$$\sum_{j \geq 1} \left(\frac{3^j}{2}\right)^z = 2^{-z} \sum_{j \geq 1} 3^j = \frac{2^{-z} 3^z}{1 - 3^z} = \frac{(3/2)^z}{1 - 3^z},$$

a geometric series of ratio 3^z , of modulus $3^{\operatorname{Re} z} < 1$, convergent on $\operatorname{Re} z < 0$ and nowhere else. Hence

$$(3) \quad K^*(z) = \frac{(3/2)^z}{1 - 3^z} \cdot (-\Gamma(z)\zeta(1+z)) = \frac{(3/2)^z \Gamma(z) \zeta(1+z)}{3^z - 1}, \quad -1 < \operatorname{Re} z < 0.$$

The dilations $\mu_j = 2/3^j$ enter inverted, through μ_j^{-z} , which is where the 2^{-z} comes from, and with it the $2^{i\omega_m}$ of the statement.

In $\operatorname{Re} z > -1$ the right side of (3) is meromorphic, with a pole of order three at $z = 0$ (each of $\Gamma(z)$, $\zeta(1+z)$ and $(3^z - 1)^{-1}$ contributing one) and a simple pole at each remaining zero $z = i\omega_m$, $m \neq 0$, of $3^z - 1$. To the right of the imaginary axis it is analytic.

Fix $\sigma \in (-1, 0)$ and $\sigma' \in (0, 1)$, and shift the inversion contour in $K(s) = (2\pi i)^{-1} \int_{(\sigma)} K^*(z) s^{-z} dz$ from σ to σ' , using rectangles with horizontal sides at $\operatorname{Im} z = \pm T_M$, $T_M := \pi(2M+1)/c$. On those sides $3^z = 3^{\operatorname{Re} z} e^{i\pi(2M+1)} = -3^{\operatorname{Re} z}$, so $|3^z - 1| = 1 + 3^{\operatorname{Re} z} \geq 1$; Stirling makes $|\Gamma(z)|$ decay like $e^{-\pi T_M/2}$ times a power of T_M , uniformly for $\sigma \leq \operatorname{Re} z \leq \sigma'$, and $\zeta(1+z)$ grows at most polynomially there. The horizontal contributions therefore vanish as $M \rightarrow \infty$, the sum of residues converges absolutely, and on the right vertical line $|3^z - 1| \geq 3^{\sigma'} - 1 > 0$ makes $\int_{(\sigma')} |K^*(z) s^{-z}| |dz| = O(s^{-\sigma'})$. So

$$K(s) = -\operatorname{Res}_{z=0} [K^*(z) s^{-z}] - \sum_{\substack{m \neq 0 \\ z=i\omega_m}} \operatorname{Res} [K^*(z) s^{-z}] + O(s^{-\sigma'}).$$

Take the simple poles first. At $z = i\omega_m$ one has $3^{i\omega_m} = e^{2\pi i m} = 1$, so $(3^z - 1)' = 3^z \log 3$ equals c there and $(3/2)^{i\omega_m} = 3^{i\omega_m} 2^{-i\omega_m} = 2^{-i\omega_m}$. With $s = e^w$,

$$-\operatorname{Res}_{z=i\omega_m} [K^*(z) e^{-zw}] = -\frac{2^{-i\omega_m}}{c} \Gamma(i\omega_m) \zeta(1+i\omega_m) e^{-i\omega_m w},$$

and replacing m by $-m$ throughout the sum turns the whole family into $\hat{H}(m) e^{i\omega_m w}$ with $\hat{H}(m)$ exactly as stated.

At the origin, insert

$$\Gamma(z) = \frac{1}{z} - \gamma_E + \left(\frac{\gamma_E^2}{2} + \frac{\pi^2}{12} \right) z + O(z^2), \quad \zeta(1+z) = \frac{1}{z} + \gamma_E - \gamma_1 z + O(z^2).$$

The two $1/z$ terms cancel against each other and

$$\Gamma(z)\zeta(1+z) = \frac{1}{z^2} + A + O(z), \quad A = \frac{\pi^2}{12} - \frac{\gamma_E^2}{2} - \gamma_1,$$

which is where A comes from. Also $(3^z - 1)^{-1} = (cz)^{-1} - \frac{1}{2} + cz/12 + O(z^3)$, and $(3/2)^z e^{-zw} = e^{\theta z}$ with $\theta := \log(3/2) - w$. Multiplying the three expansions and reading off the coefficient of z^{-1} ,

$$\text{Res}_{z=0} [K^*(z)e^{-zw}] = \frac{1}{c} \cdot \frac{\theta^2}{2} - \frac{\theta}{2} + \frac{c}{12} + \frac{A}{c}.$$

Substitute $\theta = \log(3/2) - w$ and $\log(3/2) = c - \log 2$. The w -dependent part of $-\text{Res}$ is

$$-\frac{w^2}{2c} + \left(\frac{\log(3/2)}{c} - \frac{1}{2} \right) w = -\frac{w^2}{2c} + \left(\frac{1}{2} - \frac{\log 2}{c} \right) w = Q(w),$$

and the constant part is $\frac{1}{2} \log 2 - \frac{\log^2 2}{2c} - \frac{c}{12} - \frac{A}{c} = \hat{H}(0)$.

Collecting, with $s = e^w$,

$$L(w) = Q(w) + \tilde{H}(w) + O(e^{-\sigma'w}), \quad \tilde{H}(w) := \hat{H}(0) + \sum_{m \neq 0} \hat{H}(m) e^{i\omega_m w}.$$

That series converges absolutely and uniformly, because $|\Gamma(-i\omega_m)| = \sqrt{\pi/(\omega_m \sinh \pi\omega_m)}$ decays like $e^{-\pi\omega_m/2}$ while $\zeta(1 - i\omega_m)$ grows polynomially, so $\tilde{H}$ is continuous and exactly c -periodic. It is real valued: Γ and ζ are real on the real axis, so $\hat{H}(-m)$ is the complex conjugate of $\hat{H}(m)$.

The remainder $O(e^{-\sigma'w})$ is only ordinarily exponentially small, weaker than the doubly exponential Δ of Theorem 4, and it is enough. Theorem 4 gives $L - Q - H = \Delta = o(1)$ as $w \rightarrow +\infty$, so $\tilde{H} - H = O(e^{-\sigma'w}) + \Delta(w) \rightarrow 0$. Both are c -periodic, and a c -periodic function tending to 0 as $w \rightarrow +\infty$ vanishes identically, its value at any w being its value at $w + kc$ for every k . Hence $\tilde{H} = H$, and by uniform convergence the $\hat{H}(m)$ are the Fourier coefficients of H . $\square$

Remark 7. This is an instance of the de Bruijn–Mahler phenomenon: a base-change harmonic sum, Mellin transformed, picks up simple poles on a vertical line, and their residues assemble into an exactly periodic correction with Fourier coefficients in Γ and ζ . De Bruijn found the first case of this in Mahler’s partition asymptotic [5]; Erdős and Richmond name it on p. 448 of [6]; Kato and McLeod [7] and Derfel [8] established the log-squared decay rate of Section 2’s comparison asymptotic φ_0 as generic across this whole class of rescaled functional equations. Our contribution here is the closed form for the periodic correction itself, not the mechanism producing it.

The series for H' and H'' converge absolutely and uniformly, since

$$|\Gamma(-i\omega_m)| = \sqrt{\pi/(\omega_m \sinh \pi\omega_m)}$$

decays exponentially in m while $\zeta(1 - i\omega_m)$ grows only polynomially. Precisely, with $\alpha_H := 2\pi/c$, $q := e^{-\pi\alpha_H/2}$, $A_H := c^{-1} \sqrt{3\pi/\alpha_H}$ and $C_H := \frac{3}{2} + \alpha_H^{-1} + \frac{1}{2} \sqrt{1 + \alpha_H^{-2}}$, the bound $|\Gamma(-i\omega_m)| \leq \sqrt{3\pi/\omega_m} e^{-\pi\omega_m/2}$ (valid for $\omega_m \geq \alpha_H$, since $\sinh x \geq e^x/3$ there) and the Euler-Maclaurin bound $|\zeta(1 - i\omega_m)| \leq \log(\omega_m + 1) + C_H$ (valid for $\omega_m \geq \alpha_H$) together give the majorant

$$(4) \quad |\hat{H}(m)| \leq A_H q^m m^{-1/2} (\log(\alpha_H m + 1) + C_H), \quad m \geq 1.$$

Since $\sup_w |H^{(k)}(w)| \leq \sum_{m \neq 0} |\omega_m|^k |\hat{H}(m)| = 2 \sum_{m \geq 1} \omega_m^k |\hat{H}(m)|$, summing the first six terms exactly and bounding the tail $m > 6$ by (4) (the same closed-form geometric-times-polynomial tail

sum used in the proof of Proposition 8 below) gives explicit, certified bounds:

$$(5) \quad \sup_w |H'(w)| \leq 0.0011977472315550332, \quad \sup_w |H''(w)| \leq 0.0068518962896650951.$$

Proposition 8 (Certified non-constancy). *H is not constant. Rigorously,*

$$-0.000377190280943987 < H(0) - H(\log(3/2)) < -0.000377190280943985,$$

and consequently $\text{osc}(H) := \sup H - \inf H \geq 3.7719 \times 10^{-4}$; a grid search with a Lipschitz enclosure (below) gives the certified bound $\text{osc}(H) \in [4.1874494771 \times 10^{-4}, 4.1874620262 \times 10^{-4}]$.

Proof. Write $g(b) = -b/2 + \log S(b/2)$, $S(x) = \sinh(x)/x = \sum_{n \geq 0} x^{2n}/(2n+1)!$ (this is g 's defining formula rewritten around its removable singularity at 0, since $e^{-b/2}(e^{b/2} - e^{-b/2}) = 1 - e^{-b}$). Termwise domination of $6^n n! \leq (2n+1)!$, which holds at $n = 0$ and propagates because $(2n+3)(2n+2) \geq 6(n+1)$, gives $1 \leq S(x) \leq e^{x^2/6}$, that is

$$(6) \quad 0 \leq \log S(x) \leq \frac{x^2}{6}, \quad x \geq 0.$$

Since $\sum_{j \geq 1} 3^{-j} = \frac{1}{2}$, the $-b/2$ halves of the terms of L sum in closed form, and (2) becomes, with $s = e^w$,

$$(7) \quad H(w) = -\frac{s}{2} + \sum_{j \geq 1} \log S\left(\frac{s}{3^j}\right) - Q(w) + \sum_{k \geq 0} \log(1 - e^{-2 \cdot 3^k s}).$$

Truncate the first series at $j = R$ and the second at $k = M$. By (6) the first tail satisfies

$$0 \leq \sum_{j > R} \log S\left(\frac{s}{3^j}\right) \leq \frac{s^2}{6} \sum_{j > R} 9^{-j} = \frac{s^2 9^{-R}}{48},$$

and, since $0 \leq -\log(1-y) \leq y/(1-y)$ and $e^{-2 \cdot 3^k s} = (e^{-2 \cdot 3^{M+1} s})^{3^{k-M-1}}$ for $k > M$, the second tail satisfies

$$0 \leq -\sum_{k > M} \log(1 - e^{-2 \cdot 3^k s}) \leq \frac{y}{(1-y)^2} \leq 4y, \quad y := e^{-2 \cdot 3^{M+1} s} \leq \frac{1}{2}.$$

Take $R = 30$, $M = 4$. For $s \leq 3$ (which covers every evaluation point used below, including the oscillation grid) the first tail is at most 4.43×10^{-30} , and for $s \geq 1$ the second is at most $4e^{-486} < 10^{-210}$. The remaining 35 terms are elementary functions of s ; evaluated at 50 significant decimal digits, where the accumulated rounding stays below 10^{-45} , they give

$$H(0) = -0.62713093305152597830\dots, \quad H(\log \frac{3}{2}) = -0.62675374277058199247\dots,$$

whose difference is $-0.00037719028094398583\dots$, inside the stated interval; this first route is floating-point (50 significant digits, accumulated rounding bounded as stated) rather than interval arithmetic. Independently, and rigorously in the sense of interval arithmetic, the Fourier series of Proposition 6 gives the same difference in Arb ball arithmetic: the $m = 0$ term cancels, four modes suffice, and the majorant (4) gives $\sum_{m > 4} |\hat{H}(m)| \leq 9.05 \times 10^{-20}$; the discarded modes therefore move the difference by less than 3.7×10^{-19} . The two routes agreeing to 18 digits is itself strong evidence the floating-point route's own error estimate is not being violated by an unmodeled effect.

For the oscillation, let $w_i = ic/N$, $0 \leq i < N$, with $N = 2^{20}$, so $h := c/N = 1.04772 \times 10^{-6}$. Every real w lies within $h/2$ of some w_i modulo c , so (5) gives $|H(w) - H(w_i)| \leq M_1 h/2$ with $M_1 := 1.1977472315550332 \times 10^{-3}$, whence

$$\max_i H(w_i) \leq \sup H \leq \max_i H(w_i) + \frac{M_1 h}{2}, \quad \min_i H(w_i) - \frac{M_1 h}{2} \leq \inf H \leq \min_i H(w_i),$$

and therefore $\text{osc}(H) \in [D, D + M_1 h]$ for $D := \max_i H(w_i) - \min_i H(w_i)$. The lower bound $D \leq \text{osc}(H)$ (the direction Theorem 2 actually uses) follows from $\max_i H(w_i) \leq \sup H$ for the true values $H(w_i)$; since the working precision below keeps floating-point evaluation error many orders

below $M_1 h \approx 1.25 \times 10^{-9}$, this does not depend on which arithmetic evaluated the grid. The upper bound $D + M_1 h \geq \text{osc}(H)$, not used elsewhere in this paper, additionally requires the computed extrema not to be underestimates, which is not independently certified for the full sweep below (only the two located extremal points are re-confirmed in Arb ball arithmetic). Evaluating the N grid values from Proposition 6 truncated at $|m| \leq 10$ in floating point, where the same majorant bounds the discarded modes by 2.71×10^{-43} , locates the maximum at $i = 486746$ and the minimum at $i = 1011118$; re-evaluating H there to 40 digits, from (7) and from the Fourier series alike,

$$H(w_{486746}) = -0.6267174403736425786\dots, \quad H(w_{1011118}) = -0.6271361853213578093\dots$$

So $D = 4.187449477152 \times 10^{-4}$ and $M_1 h = 1.2549 \times 10^{-9}$, which is the stated enclosure. $\square$

4. A UNIFORM SADDLEPOINT APPROXIMATION

Define, for $s > 0$,

$$\varphi_{\text{sp}}(t(s)) := \frac{\exp(K(s) + st(s))}{\sqrt{2\pi K''(s)}}.$$

This section proves that $\varphi(t(s))/\varphi_{\text{sp}}(t(s)) \rightarrow 1$ as $s \rightarrow \infty$, uniformly over $\rho := 2s/3^N \in [1, 3)$ (defined below), which is what makes the two theorems of Section 7 depend on different scopes rather than on different mathematics.

For $n \geq 2$, define the derivative-normalized functions $\kappa_n(w) = (-1)^n w^n g^{(n)}(w)$ on $\{\text{Re } w > 0\}$; $\kappa_n(b)$, $b > 0$, is b^n times the n -th cumulant of the random variable with density $b e^{-bw}/(1 - e^{-b})$ on $[0, 1]$. Writing $x = w/2$,

$$\kappa_2(w) = 1 - \frac{x^2}{\sinh^2 x}, \quad \kappa_3(w) = 2 - \frac{2x^3 \cosh x}{\sinh^3 x}, \quad \kappa_4(w) = 6 + \frac{2x^4(1 - 3 \coth^2 x)}{\sinh^2 x},$$

and from the Bernoulli expansion $g(w) + w/2 = \sum_{k \geq 1} B_{2k} w^{2k}/(2k(2k)!)$ on $|w| < 2\pi$,

$$\kappa_n(w) = (-1)^n \sum_{k \geq \lceil n/2 \rceil} \frac{B_{2k} w^{2k}}{2k(2k - n)!}, \quad n \geq 2,$$

so that $\kappa_n(w) = O(w^2)$ as $w \rightarrow 0$ for every $n \geq 2$.

Write $s = \rho \cdot 3^N/2$ for the unique $N = \lfloor \log_3(2s) \rfloor \in \mathbb{Z}$ and $\rho \in [1, 3)$, and $b_j = 2s/3^j$. Then $\{b_j : j \geq 1\} = \{\rho \cdot 3^m : m \leq N - 1\}$; for $N \geq 1$ (the only range used from here on, corresponding to $s \geq 3/2$), exactly N of these, the smallest being $\rho \geq 1$, are ≥ 1 , and the rest form a geometric tail below 1. This single structural fact is the source of every phase-independent constant below.

Lemma 9 (Boundedness off the real axis). *For $w = b(1 + iu)$, $b > 0$, $|u| \leq 1$, with $e(r) = r^2/\sinh^2 r$ and $f(r) = r^3 \cosh r/\sinh^3 r$,*

$$|\kappa_2(w)| \leq 1 + 2e(b/2), \quad |\kappa_3(w)| \leq 2 + 2 \cdot 2^{3/2} f(b/2).$$

Both bounds are finite over the whole sector $|\text{Im } w| \leq \text{Re } w$; $M_2 := \sup |\kappa_2| \leq 3$, $M_3 := \sup |\kappa_3| \leq 2 + 2^{5/2} < 7.657$. In addition, for $|w| \leq 2$, $|\kappa_2(w)| \leq 0.114|w|^2$ and $|\kappa_3(w)| \leq 0.0119|w|^4$.

Proof. $|\sinh z|^2 = \sinh^2(\text{Re } z) + \sin^2(\text{Im } z) \geq \sinh^2(\text{Re } z)$ and $|\cosh z|^2 \leq \cosh^2(\text{Re } z)$; the first excludes the poles of $\coth$, since $\text{Re}(w/2) = b/2 > 0$ never approaches $\pi i k$. This gives the sector bounds directly. Monotonicity of e is immediate from the increasing power series of $\sinh(r)/r$. For f , set $u = 2r$; a direct computation gives $2r \sinh(r) \cosh(r) (\log f)'(r) = F(u)$ with $F(u) := 3 \sinh u - u \cosh u - 2u$, so $(\log f)'(r)$ has the sign of $F(2r)$. Since $F(0) = F'(0) = F''(0) = 0$ and $F'''(u) = -u \sinh u < 0$ for $u > 0$, integrating three times from 0 gives $F(u) < 0$ for all $u > 0$, hence $(\log f)'(r) < 0$ for $r > 0$ and f is strictly decreasing. (Lazarević's inequality, $(\sinh r/r)^3 > \cosh r$, separately gives $f < 1$.) The local bounds at $|w| \leq 2$ follow by applying the maximum-modulus principle to $\kappa_2(w)/w^2$ and $\kappa_3(w)/w^4$ (analytic on $|w| < 2\pi$ by the Bernoulli expansion): bound

each on $|w| = 2$ by the triangle inequality, using $|B_{2k}| = 2(2k)! \zeta(2k)/(2\pi)^{2k}$ and $\zeta(2k) \leq \zeta(2)$, respectively $\zeta(4)$. $\square$

Lemma 10 (Variance). *For all $\rho \in [1, 3)$, $N \geq 1$, with $V := s^2 K''(s) = \sum_j \kappa_2(b_j)$,*

$$N - A_V \leq V \leq N + 0.1283, \quad A_V := \sum_{m \geq 0} e(3^m/2) = 1.4269413069 \dots$$

Proof. Split $V = \sum_{m=0}^{N-1} \kappa_2(\rho 3^m) + \sum_{m < 0} \kappa_2(\rho 3^m)$, using $\kappa_2(b) = 1 - e(b/2) \in (0, 1)$. For the lower bound, drop the second (non-negative) sum and use $e(\rho 3^m/2) \leq e(3^m/2)$ (e decreasing, $\rho \geq 1$) in the first, giving $\sum_{m=0}^{N-1} \kappa_2(\rho 3^m) \geq N - A_V$. For the upper bound, the first sum is $< N$ term by term, and the second sum, where $\rho 3^m < 1$, is bounded by Lemma 9's local estimate: $\sum_{m < 0} \kappa_2(\rho 3^m) \leq 0.114 \rho^2 \sum_{m < 0} 9^m = 0.114 \rho^2/8 \leq 0.1283$ for $\rho < 3$. $\square$

Lemma 11 (Tail bound and existence of the density). *If $b_0 \geq 1$ and $b_k = 3^k b_0$, then*

$$\sum_{k \geq 0} \log \coth(b_k/2) \leq 3e^{-b_0}.$$

Consequently, with $\Lambda(y) := K(s(1+iy)) - K(s) + isy t(s)$,

$$|e^{\Lambda(y)}| \leq e^{3/e} (1+y^2)^{-N/2} \leq 3.0152 (1+y^2)^{-N/2}$$

for every real y . In particular X has a continuous density, the tilted characteristic function is integrable for $N \geq 3$, and Fourier inversion at $x = t(s)$ gives

$$(8) \quad R(s) := \frac{\varphi(t(s))}{\varphi_{\text{sp}}(t(s))} = \sqrt{\frac{V}{2\pi}} \int_{\mathbb{R}} e^{\Lambda(y)} dy,$$

a real quantity, since $\Lambda(-y) = \overline{\Lambda(y)}$.

Proof. $\log \coth(x/2) = 2 \operatorname{artanh}(e^{-x}) \leq 2e^{-x}/(1 - e^{-2x})$; summing over the geometric orbit $b_k = 3^k b_0$ and bounding the resulting series by its first few terms gives the stated constant. X 's density is continuous because the sum of the first two summands already has a continuous, compactly supported density and convolution preserves continuity. For Λ : $t(s) = -K'(s)$ cancels the linear term of the tilted transform exactly, and each remaining factor $e^{g(b_j(1+iy)) - g(b_j)}$ has modulus at most 1 (a tilted characteristic function), with modulus at most $\coth(b_j/2)/\sqrt{1+y^2}$ on the N indices where $b_j \geq 1$; the tail estimate applies to exactly that geometric set, with $b_0 = \rho \geq 1$. $\square$

Lemma 12 (Central expansion). *With $h(u) = g(b(1+iu))$, Taylor's theorem in integral form (the Lagrange form is invalid for a complex-valued function of a real variable) gives*

$$h(y) = h(0) + h'(0)y + h''(0)\frac{y^2}{2} + \frac{1}{2} \int_0^y (y-u)^2 h'''(u) du.$$

Consequently, writing $h_j(u) := g(b_j(1+iu))$ for each $j \geq 1$, for $\theta \in (0, 1]$,

$$S := \sum_j \sup_{|u| \leq \theta} |h_j'''(u)| \leq 2N + 10.559,$$

and hence $|\Lambda(y) + Vy^2/2| \leq B|y|^3$ for $|y| \leq \theta$, with $B := S/6 \leq (2N + 10.559)/6$.

Proof. Write $w_j := b_j(1+iu)$. The chain rule and the definition of κ_3 give $h_j'''(u) = i \kappa_3(w_j)/(1+iu)^3$, so

$$|h_j'''(u)| = \frac{|\kappa_3(w_j)|}{(1+u^2)^{3/2}}.$$

Split the index set as in Section 4. For the N indices with $b_j = \rho 3^m \geq 1$, $0 \leq m \leq N-1$, drop the denominator and use Lemma 9's sector bound, legitimate since $|u| \leq \theta \leq 1$:

$$|h_j'''(u)| \leq |\kappa_3(w_j)| \leq 2 + 2^{5/2} f(b_j/2) \leq 2 + 2^{5/2} f(3^m/2),$$

the last step by monotonicity of f and $\rho \geq 1$. Summing over $m \geq 0$ bounds this block by $2N + 2^{5/2}\Sigma$, $\Sigma := \sum_{m \geq 0} f(3^m/2)$. The first three terms of Σ are $0.99614738\dots$, $0.82241066\dots$, $0.04500508\dots$; for $r \geq 13$ one has $f(r) \leq 4.001 r^3 e^{-2r}$, so the remaining terms sum to less than 2×10^{-8} , and $2^{5/2}\Sigma < 10.541906$.

For the remaining indices $b_j = \rho 3^{-n}$, $n \geq 1$, one has $b_j < 1$ and $|w_j| = b_j \sqrt{1+u^2} \leq b_j \sqrt{2} < 2$, so Lemma 9's local bound applies and, keeping the denominator this time,

$$|h_j'''(u)| \leq \frac{0.0119 |w_j|^4}{(1+u^2)^{3/2}} = 0.0119 b_j^4 (1+u^2)^{1/2} \leq 0.0119 \sqrt{2} b_j^4.$$

Summing, $\sum_{n \geq 1} (\rho 3^{-n})^4 = \rho^4/80 < 81/80$, so this block is below $0.0119 \sqrt{2} \cdot 81/80 < 0.01704$. Together, $S < 2N + 10.541906 + 0.01704 < 2N + 10.559$.

For the remainder bound, $\sum_j h_j(y) = K(s(1+iy))$, $\sum_j h_j'(0) = isK'(s) = -ist(s)$ and $\sum_j h_j''(0) = -s^2 K''(s) = -V$, so summing the integral-form Taylor expansion over j gives

$$\Lambda(y) + \frac{Vy^2}{2} = \sum_j \frac{1}{2} \int_0^y (y-u)^2 h_j'''(u) du,$$

whose modulus is at most $S|y|^3/6 = B|y|^3$ for $|y| \leq \theta$. $\square$

Theorem 13 (Uniform saddlepoint asymptotic). *For $N \geq 19$, uniformly over $\rho \in [1, 3)$, $|R(s) - 1| \leq E(N) := e_1(N) + e_2(N) + e_3(N)$, with $\theta_N := N^{-1/3}$, $V_{\text{lo}} := N - A_V$, $V_{\text{up}} := N + 0.1283$, $B := (2N + 10.559)/6$,*

$$(9) \quad \begin{aligned} e_1(N) &:= \frac{4e^{B\theta_N^3} B}{\sqrt{2\pi} V_{\text{lo}}^{3/2}}, & e_2(N) &:= \frac{2}{\sqrt{2\pi}} \cdot \frac{e^{-\theta_N^2 V_{\text{lo}}/2}}{\theta_N \sqrt{V_{\text{lo}}}}, \\ e_3(N) &:= 2e^{3/e} \sqrt{\frac{V_{\text{up}}}{2\pi}} \cdot \frac{(1 + \theta_N^2)^{-(N-2)/2}}{\theta_N(N-2)}. \end{aligned}$$

$E(19) < 1$, and $\sqrt{N} E(N) \rightarrow \frac{4}{3} e^{1/3} (2\pi)^{-1/2} = 0.742358\dots$ as $N \rightarrow \infty$ (proved below, from e_1 alone: e_2 and e_3 vanish faster than any power of N). Numerically, E is also strictly decreasing and satisfies $E(N) \leq 4.18 N^{-1/2}$ throughout $19 \leq N \leq 5000$ (checked directly in the accompanying repository); neither fact, nor $E(19) < 1$ itself, is used below, where only $E(N) \rightarrow 0$ as $N \rightarrow \infty$ is needed. Consequently $\varphi(t) = \varphi_{\text{sp}}(t) (1 + o(1))$ as $t \rightarrow 0^+$, with no restriction on the sequence of t 's.

Proof. Split (8) at $|y| = \theta_N$. On $|y| \leq \theta_N$, Lemma 12 bounds the cubic remainder $|\Lambda(y) + Vy^2/2|$ by $B\theta_N^3$, which stays bounded (not $o(1)$): this is exactly why e_1 carries the factor $e^{B\theta_N^3}$ rather than vanishing) as $N \rightarrow \infty$, since $B = O(N)$ and $\theta_N^3 = N^{-1}$. Write $C(y) := e^{\Lambda(y)} - e^{-Vy^2/2}$ and factor the Gaussian out before estimating, $e^{\Lambda(y)} = e^{-Vy^2/2} e^{z(y)}$ with $z(y) := \Lambda(y) + Vy^2/2$, so that $C(y) = e^{-Vy^2/2} (e^{z(y)} - 1)$. Since $|z(y)| \leq B|y|^3 \leq B\theta_N^3$ on $|y| \leq \theta_N$, the elementary $|e^z - 1| \leq |z|e^{|z|}$ gives

$$|C(y)| \leq e^{-Vy^2/2} B|y|^3 e^{B\theta_N^3} \quad (|y| \leq \theta_N),$$

the Gaussian factor retained. Integrating that inequality, $\int_{\mathbb{R}} e^{-Vy^2/2} |y|^3 dy = 4/V^2$, so

$$\sqrt{V/2\pi} \int_{|y| \leq \theta_N} |C(y)| dy \leq \frac{4Be^{B\theta_N^3}}{\sqrt{2\pi} V^{3/2}},$$

which is decreasing in V ; using $V \geq V_{\text{lo}}$ from Lemma 10 gives the central Taylor-remainder contribution e_1 . Since $\sqrt{V/2\pi} \int_{\mathbb{R}} e^{-Vy^2/2} dy = 1$, the same splitting leaves exactly two further terms. The truncated Gaussian tail $\sqrt{V/2\pi} \cdot 2 \int_{\theta_N}^{\infty} e^{-Vy^2/2} dy$, substituting $v = y\sqrt{V}$ to get $(2/\sqrt{2\pi}) \int_c^{\infty} e^{-v^2/2} dv$, $c = \theta_N \sqrt{V}$, and bounding via $\int_c^{\infty} e^{-v^2/2} dv \leq e^{-c^2/2}/c$ at $V = V_{\text{lo}}$, gives e_2 . On $|y| > \theta_N$, Lemma 11's bound $|e^{\Lambda(y)}| \leq e^{3/e} (1+y^2)^{-N/2}$ integrates, via $\int_a^{\infty} (1+y^2)^{-N/2} dy \leq$

$(1 + a^2)^{-(N-2)/2}/(a(N-2))$ at $a = \theta_N$ (using $V \leq V_{\text{up}}$ for the leading $\sqrt{V/2\pi}$ factor in (8)), to give e_3 ; this needs $\theta_N \gg N^{-1/2}$, satisfied since $\frac{1}{3} < \frac{1}{2}$. Every constant entering E (via A_V , Lemma 10, and the $3.0152 \approx e^{3/e}$ of Lemma 11) is a supremum over $b > 0$ or a sum over one complete geometric orbit bounded using $\rho \geq 1$, hence independent of ρ . Directly evaluating (9) gives $E(18) = 1.00170\dots \geq 1$ and $E(19) = 0.95673\dots < 1$, confirming that $N \geq 19$ is where the stated bound $E(N) < 1$ first becomes non-vacuous.

For the asymptotic rate, e_1, e_2, e_3 separate on very different scales as $N \rightarrow \infty$. Since $\theta_N^3 = 1/N$, $B\theta_N^3 \rightarrow 1/3$ and $B/N \rightarrow 1/3$, $V_{\text{lo}}/N, V_{\text{up}}/N \rightarrow 1$, so

$$\sqrt{N} e_1(N) = \frac{4e^{B\theta_N^3}}{\sqrt{2\pi}} \cdot \frac{\sqrt{N} B}{V_{\text{lo}}^{3/2}} \rightarrow \frac{4e^{1/3}}{\sqrt{2\pi}} \cdot \frac{1}{3} = \frac{4}{3} e^{1/3} (2\pi)^{-1/2}.$$

For e_2 and e_3 , $\theta_N^2 V_{\text{lo}} \sim N^{1/3} \rightarrow \infty$, so both carry a factor $e^{-N^{1/3}/2}(1 + o(1))$ (for e_3 , via $(1 + \theta_N^2)^{-(N-2)/2} = \exp(-\frac{N-2}{2}\theta_N^2(1 + o(1)))$), against polynomial prefactors of order $N^{-1/6}$; a factor decaying like $e^{-N^{1/3}/2}$ beats every power of N , so $N^k e_2(N), N^k e_3(N) \rightarrow 0$ for every k . Hence $\sqrt{N} E(N) = \sqrt{N} e_1(N) + o(1) \rightarrow \frac{4}{3} e^{1/3} (2\pi)^{-1/2}$, and in particular $E(N) \rightarrow 0$. Since $t(s)$ ranges bijectively over $(0, 1/2)$, this is exactly the statement $\varphi(t) = \varphi_{\text{sp}}(t)(1 + o(1))$ as $t \rightarrow 0^+$. $\square$

Remark 14. The rate is not sharp, and a sharper one is plausible but not established here. Carrying the central expansion to fourth order, using that the odd cubic term integrates to zero against the Gaussian core, and using the limits $V_3 := \sum_j \kappa_3(b_j) = 2N + O(1)$, $V_4 := \sum_j \kappa_4(b_j) = 6N + O(1)$ (the third and fourth cumulant sums, by the same argument as Lemma 10) suggests the Edgeworth correction $R(s) = 1 - \frac{1}{12N} + O(N^{-3/2})$, matching the leading term of the classical Stirling correction for a $\text{Gamma}(N, 1)$ law: after tilting, the $b_j \geq 1$ summands behave asymptotically like independent $\text{Exp}(1)$ variables, so X under its tilt is asymptotically $\text{Gamma}(N, 1)$ -distributed, and ρ enters only through $O(1)$ shifts of the cumulants, at order N^{-2} . We do not need this sharper rate and do not prove the remainder term here; Theorem 13's qualitative $o(1)$ is all that Section 7 uses.

5. THE ENVELOPE LEMMA

Theorem 13 compares φ to φ_{sp} , a quantity built from the full transform $K = Q + H + \Delta$. It remains to compare φ_{sp} to a smooth auxiliary system built from Q alone, so that the periodic part H can be extracted explicitly. Write $t = e^{-\tau}$, $g_\tau(w) = L(w) + e^{w-\tau}$, $g_{0,\tau}(w) = Q(w) + e^{w-\tau}$. The true and smooth systems each have their own analogue of $-K'$: write $B_{\text{tr}}(w) := e^{wt}(e^w) = -L'(w)$ and $B_{\text{sm}}(w) := -Q'(w) = w/c - a$.

Lemma 15 (True and smooth saddles). *Let $\Phi(w) = w - \log B_{\text{tr}}(w)$ and $\Phi_0(w) = w - \log B_{\text{sm}}(w)$, and set $\tau_0 := 1 + ca + \log c = 0.9502067913\dots$, so $\tau_0 > \log 2$.*

For $\tau > \log 2$, g_τ has a unique global minimizer $w^(\tau)$, at which $g_\tau''(w^*) = V(r^*) > 0$, $r^* = e^{w^*}$; it is characterized by $\tau = \Phi(w^*)$, and Φ is a strictly increasing bijection $\mathbb{R} \rightarrow (\log 2, \infty)$. For $\tau \leq \log 2$ there is no critical point at all. For $\tau > \tau_0$, $g_{0,\tau}$ has a unique local minimizer $w_0(\tau)$, characterized by $\tau = \Phi_0(w_0)$ together with $B_0 := B_{\text{sm}}(w_0) = e^{w_0-\tau} > 1/c$; Φ_0 restricted to $(ca+1, \infty)$ is a strictly increasing bijection onto (τ_0, ∞) , and B_0 increases from $1/c$ to ∞ along it. For $\tau \leq \tau_0$ there is no such minimizer.*

Once $B_0 \geq 5$ (equivalently $\tau \geq 3.73978\dots$),

$$|w^* - w_0| \leq \frac{0.00652}{B_0}, \quad 0 \leq g_\tau(w_0) - g_\tau(w^*) \leq \frac{5.77 \times 10^{-5}}{B_0}.$$

Moreover $w_0(\tau + c) - w_0(\tau) = c + O(1/\tau)$ as $\tau \rightarrow \infty$.

Proof. Substituting $s = e^w$ and $t = e^{-\tau}$ turns g_τ into $K(s) + st$, strictly convex in $s > 0$ because $K'' > 0$, with derivative $K'(s) + t$ vanishing exactly where $t = t(s)$. As $t(\cdot)$ decreases strictly from

$1/2$ to 0 , that happens for exactly one s when $t \in (0, 1/2)$, that is when $\tau > \log 2$, and for no s when $t \geq 1/2$. Since $w \mapsto e^w$ is a bijection $\mathbb{R} \rightarrow (0, \infty)$, $w^* = \log s$ is the unique global minimizer of g_τ , and Φ inverts $\tau \mapsto w^*$. Differentiating twice, $g_\tau''(w) = e^w(K'(e^w) + t) + e^{2w}K''(e^w)$, whose first term vanishes at w^* , leaving $g_\tau''(w^*) = V(r^*) > 0$.

For the smooth system, $g_{0,\tau}'(w) = e^{w-\tau} - B_{\text{sm}}(w)$ and $g_{0,\tau}''(w) = e^{w-\tau} - 1/c$, so a critical point is a local minimizer exactly when $B_0 > 1/c$. Also $\Phi_0'(w) = 1 - 1/(c B_{\text{sm}}(w))$, positive exactly where $B_{\text{sm}}(w) > 1/c$, that is on $w > ca + 1$; there Φ_0 increases from $\Phi_0(ca + 1) = ca + 1 + \log c = \tau_0$ to ∞ . This gives the stated existence, uniqueness and range.

Now take $B_0 \geq 5$. The decomposition $L = Q + H + \Delta$ evaluates the true gradient at the smooth saddle exactly: $Q'(w_0) = -B_0 = -e^{w_0-\tau}$, so the leading terms cancel and

$$(10) \quad g_\tau'(w_0) = L'(w_0) + e^{w_0-\tau} = H'(w_0) + \Delta'(w_0), \quad |g_\tau'(w_0)| \leq \varepsilon_1 := \sup |H'| + \sup_{[w_0-1, w_0+1]} |\Delta'|.$$

On $[w_0 - 1, w_0 + 1]$ write $g_\tau''(w) = e^{w-\tau} - 1/c + H''(w) + \Delta''(w)$. Here $e^{w-\tau} = B_0 e^{w-w_0}$ ranges over $[B_0/e, B_0e]$, so g_τ'' admits no bound independent of τ ; it is of exact order B_0 ,

$$(11) \quad \frac{B_0}{e} - \eta_0 \leq g_\tau''(w) \leq B_0e, \quad \eta_0 := \frac{1}{c} + \sup |H''| + \sup_{[w_0-1, w_0+1]} |\Delta''|,$$

the upper bound because $-1/c + \sup |H''| + \sup |\Delta''| < 0$. Since $w_0 = c(B_0 + a)$, $B_0 \geq 5$ forces $w_0 \geq 5.3492$ and so $w \geq 4.3492$ on the interval, where $|\Delta'|, |\Delta''| \leq 10^{-60}$: with $b := 2e^w \geq 2e^{4.3492} > 154$ and $x_k := 3^k b$, differentiating $\Delta(w) = -\sum_{k \geq 0} \log(1 - e^{-x_k})$ termwise (justified as in the proof of Theorem 4) gives $\Delta'(w) = -\sum_{k \geq 0} x_k/(e^{x_k} - 1)$ and, applying $|d/dx[x/(e^x - 1)]| \leq 8xe^{-x}$ for $x \geq 1$ (elementary, since $e^x - 1 \geq e^x/2$ there) together with $dx_k/dw = x_k$, $\Delta''(w) = -\sum_{k \geq 0} x_k \cdot \frac{d}{dx} \left[\frac{x}{e^x - 1} \right]_{x=x_k}$, so $|\Delta''(w)| \leq \sum_{k \geq 0} 8x_k^2 e^{-x_k}$; bounding both sums by their dominant $k = 0$ term via $x_k - b \geq 2kb$ (from $3^k - 1 \geq 2k$) gives $|\Delta'(w)| \leq 4be^{-b}$ and $|\Delta''(w)| \leq 16b^2 e^{-b}$, both well under 10^{-60} at $b > 154$. With (5) this makes $\varepsilon_1 \leq 0.0011978$ and $\eta_0 \leq 0.9170912$, whence $2e\eta_0 = 4.9859 \dots \leq 5 \leq B_0$, that is $\eta_0 \leq B_0/(2e)$, and (11) improves to $g_\tau'' \geq B_0/(2e) > 0$ throughout the interval.

Put $h_0 := 2e\varepsilon_1/B_0 \leq 2e \cdot 0.0011978/5 < 0.0014$. For $h \in (h_0, 1]$ the mean value theorem applied on $[w_0 - 1, w_0 + 1]$ gives $g_\tau'(w_0 + h) \geq g_\tau'(w_0) + hB_0/(2e) \geq -\varepsilon_1 + hB_0/(2e) > 0$ and, symmetrically, $g_\tau'(w_0 - h) \leq \varepsilon_1 - hB_0/(2e) < 0$. As w^* is the only zero of g_τ' , it lies in $(w_0 - h, w_0 + h)$ for every such h , so $|w^* - w_0| \leq h_0 = 2e\varepsilon_1/B_0 \leq 0.00652/B_0$.

Finally $g_\tau(w_0) - g_\tau(w^*) \geq 0$ because w^* is the global minimizer, and Taylor's theorem about w^* , where $g_\tau'(w^*) = 0$, bounds it by $\frac{1}{2}(\sup_{[w_0-1, w_0+1]} g_\tau'')(w_0 - w^*)^2 \leq \frac{1}{2}B_0e h_0^2 = 2e^3 \varepsilon_1^2/B_0 \leq 5.77 \times 10^{-5}/B_0$.

For the phase drift, $w_0 = c(B_0 + a)$ and $\tau = w_0 - \log B_0$ give $dw_0/d\tau = (1 - 1/(cB_0))^{-1} = 1 + O(1/B_0)$ along the curve $\tau \mapsto w_0(\tau)$; since $B_0 \rightarrow \infty$ as $\tau \rightarrow \infty$ (via $B_0 \sim \tau/c$), integrating over one period gives $w_0(\tau + c) - w_0(\tau) = \int_\tau^{\tau+c} (1 + O(1/B_0(u))) du = c + O(1/\tau)$. $\square$

Proposition 16 (Envelope estimate). *With $S(\tau) := \log \varphi_{\text{sp}}(t) = g_\tau(w^*) + w^* - \frac{1}{2} \log(2\pi V(r^*))$ and $P(\tau) := Q(w_0) + B_0 + w_0 - \frac{1}{2} \log(2\pi(B_0 - \frac{1}{c}))$,*

$$|S(\tau) - P(\tau) - H(w_0(\tau))| = O(1/\tau) \quad (\tau \rightarrow \infty),$$

uniformly in the phase of τ modulo c .

Proof. Start from the identity. Since $g_{0,\tau}(w_0) = Q(w_0) + e^{w_0-\tau} = Q(w_0) + B_0$, the definition of P reads $P(\tau) = g_{0,\tau}(w_0) + w_0 - \frac{1}{2} \log(2\pi(B_0 - \frac{1}{c}))$, so

$$S(\tau) - P(\tau) = [g_\tau(w^*) - g_{0,\tau}(w_0)] + (w^* - w_0) - \frac{1}{2} \log \frac{V(r^*)}{B_0 - \frac{1}{c}}.$$

Theorem 4 gives $L = Q + H + \Delta$, hence $g_\tau = g_{0,\tau} + H + \Delta$ pointwise, and in particular $g_\tau(w^*) = g_{0,\tau}(w^*) + H(w^*) + \Delta(w^*)$. Substituting and subtracting $H(w_0)$,

$$(12) \quad \begin{aligned} S(\tau) - P(\tau) - H(w_0) = & \underbrace{[g_{0,\tau}(w^*) - g_{0,\tau}(w_0)]}_{T_1} + \underbrace{[H(w^*) - H(w_0)]}_{T_2} + \underbrace{(w^* - w_0)}_{T_3} \\ & - \underbrace{\frac{1}{2} \log \frac{V(r^*)}{B_0 - \frac{1}{c}}}_{T_4} + \underbrace{\Delta(w^*)}_{T_5}. \end{aligned}$$

This is exact for every τ with $B_0 \geq 5$. The five terms are bounded in turn.

T_1 is the envelope gap of the smooth system. Taylor about w_0 , where $g'_{0,\tau}$ vanishes, with $g''_{0,\tau}(w) = e^{w-\tau} - 1/c \leq B_0 e$ on $[w_0 - 1, w_0 + 1]$ and $|w^* - w_0| \leq 0.00652/B_0$ from Lemma 15, gives $0 \leq T_1 \leq \frac{1}{2} B_0 e (0.00652/B_0)^2 < 5.78 \times 10^{-5}/B_0$.

T_2 is the periodic-part difference: $|T_2| \leq \sup |H'| \cdot |w^* - w_0| < 7.82 \times 10^{-6}/B_0$ by (5).

T_3 is the saddle-location shift itself, $|T_3| \leq 0.00652/B_0$, again from Lemma 15.

T_4 compares the two normalizations. By $V(r^*) = g''_\tau(w^*)$ and the expression for g''_τ used in Lemma 15,

$$V(r^*) - \left(B_0 - \frac{1}{c}\right) = B_0(e^{w^*-w_0} - 1) + H''(w^*) + \Delta''(w^*),$$

whose modulus is at most $0.00652e^{0.00652/5} + \sup |H''| + 10^{-60} < 0.0135$, since $B_0|e^{w^*-w_0} - 1| \leq B_0|w^* - w_0|e^{|w^*-w_0|}$. Dividing by $B_0 - 1/c$ and taking $\frac{1}{2} \log(1 + \cdot)$ gives $|T_4| \leq 0.0068/(B_0 - 1/c)$.

T_5 is the remainder of Theorem 4, evaluated at the true saddle rather than dropped. Here $B_0 \geq 5$ forces $w_0 \geq 5.3492$ and $w^* \geq w_0 - 0.0014$, so $e^{w^*} \geq 210.1$ and $|T_5| \leq \sum_{k \geq 0} 2e^{-2 \cdot 3^k \cdot 210.1} < 10^{-182}$.

Each bound is $O(1/B_0)$ (for T_4 , via $B_0 - 1/c \geq B_0 - 1$, comparable to B_0 once $B_0 \geq 5$). Finally $\tau = w_0 - \log B_0$ and $w_0 = c(B_0 + a)$ give $\tau = cB_0 + ca - \log B_0$, so $B_0/\tau \rightarrow 1/c$, and $O(1/B_0) = O(1/\tau)$ with constants free of the phase of τ . $\square$

6. THE BERG-KRÜPPEL IDENTITY

Proposition 17. Write $f(p) = \alpha \log p - \beta \log^2 p$, so that Berg and Krüppel's transform $G_0(p) = p^\alpha \exp(-\beta \log^2 p)$ of their equation (9.3) is $e^{f(p)}$ and their equation (9.5) reads

$$g_0(t) = \frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \exp[pt + f(p)] dp, \quad \sigma > 0.$$

Take α, β from their equation (9.4) at $a = 3$, $\lambda = 2/3$. Then $f(p) = Q(\log p)$ identically; the exact saddle of that integral, $t + f'(x) = 0$ with $t = e^{-\tau}$, is $x^\sharp = e^{w_0(\tau)}$; and

$$P(\tau) = x^\sharp t + f(x^\sharp) - \frac{1}{2} \log(2\pi f''(x^\sharp)),$$

the saddlepoint expression invoked in the proof of their Proposition 9.1, evaluated at that exact saddle. In the variable $Y := cB_0$, with $r = -ca - \log c$, the saddle equation is $Y - \log Y = \tau + r$.

Proof. Their truncated equation (9.2) is $\lambda \psi'(t) = a \psi(at)$ for $a > 1$, $\lambda > 0$; Laplace transforming (the boundary term $\lambda \psi(0)$ from $\mathcal{L}[\psi'](p) = pG(p) - \psi(0)$ vanishes, since the solutions of interest are supported on $[0, \infty)$ with $\psi(0) = 0$) gives $\lambda p G(p) = G(p/a)$, whose general solution (9.3) is G_0 times an arbitrary 1-periodic function of $\log p / \log a$, with

$$\alpha = -\frac{1}{2} - \frac{\log \lambda}{\log a}, \quad \beta = \frac{1}{2 \log a}$$

their equation (9.4). Section 2 placed φ in that family at $a = 3$, $\lambda = 2/3$: their untruncated (9.1) reads $\frac{2}{3}\varphi'(t) = 3(\varphi(3t) - \varphi(3t - 2))$, which on $(0, \frac{2}{3})$ is $\varphi'(t) = \frac{9}{2}\varphi(3t)$. At those values

$$\alpha = -\frac{1}{2} - \frac{\log(2/3)}{\log 3} = \frac{1}{2} - \frac{\log 2}{\log 3} = a, \quad \beta = \frac{1}{2\log 3} = \frac{1}{2c},$$

where the a on the right is this paper's linear coefficient of Q ; their a , the dilation factor 3, is a different quantity and is not written again below. Hence, writing $y := \log p$,

$$f(p) = \alpha y - \beta y^2 = ay - \frac{y^2}{2c} = Q(y),$$

the third of the three coincidences asserted in the proposition above. The other two are the chain rule:

$$-p f'(p) = -Q'(y) = B_{\text{sm}}(y) = 2\beta y - \alpha, \quad p^2 f''(p) = Q''(y) - Q'(y) = B_{\text{sm}}(y) - \frac{1}{c} = 2\beta y - 2\beta - \alpha.$$

Berg and Krüppel's own displayed formula for $f''(p)$, on p. 179 of [2], in the proof of their Proposition 9.1 (the unnumbered display right after "In view of"), reads $p^2 f''(p) = 2\beta \ln p + 2\beta - \alpha$, with $+2\beta$ where direct differentiation of f gives -2β : $f''(p) = \frac{1}{p^2}(2\beta \ln p - 2\beta - \alpha)$, since $f'(p) = \frac{1}{p}(\alpha - 2\beta \ln p)$ differentiates to $-\frac{2\beta}{p^2} - \frac{\alpha - 2\beta \ln p}{p^2}$. Their Proposition 9.1 is unaffected, the term being of lower order there than $2\beta \log p$, but the exact expression used here is not.

The saddle condition $t + f'(x) = 0$ at $x = e^y$ now reads $e^{-\tau} = B_{\text{sm}}(y)e^{-y}$, that is $\tau = y - \log B_{\text{sm}}(y) = \Phi_0(y)$, whose unique solution on the branch $B_{\text{sm}} > 1/c$ is $y = w_0(\tau)$ by Lemma 15. So $x^\sharp = e^{w_0}$, and there $x^\sharp t = e^{w_0 - \tau} = B_0$, $f(x^\sharp) = Q(w_0)$, $(x^\sharp)^2 f''(x^\sharp) = B_0 - 1/c > 0$. Substituting,

$$\begin{aligned} x^\sharp t + f(x^\sharp) - \frac{1}{2} \log(2\pi f''(x^\sharp)) &= B_0 + Q(w_0) - \frac{1}{2} \log \frac{2\pi(B_0 - 1/c)}{(x^\sharp)^2} \\ &= Q(w_0) + B_0 + w_0 - \frac{1}{2} \log \left(2\pi(B_0 - \frac{1}{c}) \right), \end{aligned}$$

which is $P(\tau)$. Finally $w_0 = c(B_0 + a)$ and $\tau = w_0 - \log B_0$ give $\tau = cB_0 + ca - \log B_0 = Y - \log Y + ca + \log c = Y - \log Y - r$. $\square$

Series-reverting $Y - \log Y = \tau + r$ to second order, $Y = \tau + \log \tau + r + (\log \tau + r)/\tau + O(\tau^{-2} \log^2 \tau)$, and substituting into P gives

$$\begin{aligned} P(\tau) &= -\frac{(\tau + \log \tau)^2}{2c} + \left(1 + \frac{1}{c} - \frac{r}{c}\right)\tau + \left(\frac{1}{2} - \frac{r}{c}\right)\log \tau + C_P + O\left(\frac{\log^2 \tau}{\tau}\right), \\ C_P &= -\frac{r^2}{2c} + r + ca + \frac{ca^2}{2} - \frac{\log 2\pi}{2} + \frac{\log c}{2} = -0.9576743183133982760669122497855855\dots \end{aligned}$$

Berg and Krüppel's own equations following (9.6) give, in their notation,

$$\delta_{\text{BK}} = \frac{1}{2} + \alpha - 2\beta \log(2\beta), \quad \gamma = -2\beta - \delta_{\text{BK}} - \frac{1}{2}, \quad \varepsilon = \frac{1}{2} + \alpha - \beta \log(2\beta),$$

with $\alpha = a$, $\beta = 1/(2c)$ as identified above. Substituting and simplifying (using $r = -ca - \log c$) gives, matching the τ , $\log \tau$, and constant coefficients of P term for term,

$$\gamma = -\left(1 + \frac{1}{c} - \frac{r}{c}\right) = -1.8649154949\dots, \quad \delta_{\text{BK}} = \frac{1}{2} - \frac{r}{c} = 0.4546762683\dots,$$

$$\varepsilon = a + \frac{1}{2} + \frac{\log c}{2c} = 0.4118732574\dots,$$

so that, in particular, $C_P = \varepsilon \log(2\beta) - \frac{1}{2} \log(2\pi)$ holds identically (verified by substituting ε 's expression directly into C_P 's), giving

$$C_P = \varepsilon \log(2\beta) - \frac{1}{2} \log(2\pi) = \log \frac{(2\beta)^\varepsilon}{\sqrt{2\pi}}.$$

Hence $P(\tau) - \log \varphi_{0,\text{bare}}(e^{-\tau}) \rightarrow C_P$ and, using the prefactor-included normalization, $P(\tau) - \log \varphi_0(e^{-\tau}) \rightarrow 0$ with rate $O(\tau^{-1} \log^2 \tau)$: the τ , $\log \tau$, and constant coefficients of P reproduce γ , δ_{BK} , ε exactly, so this convergence is a direct consequence of the identity of Proposition 17 together with the series reversion above, not an appeal to Berg and Krüppel's own asymptotic analysis of φ_0 .

Proof of Corollary 3. By Proposition 16 and 17, $S(\tau) = \log \varphi_0(e^{-\tau}) + H(w_0(\tau)) + O(\tau^{-1} \log^2 \tau)$, i.e. $\varphi_{\text{sp}} = \varphi_0 \cdot e^{H(w_0(\tau))}(1+o(1))$; by Theorem 13, $\varphi = \varphi_{\text{sp}}(1+o(1))$ as well, so $\varphi = \varphi_0 \cdot e^{H(w_0(\tau))}(1+o(1))$: the periodic factor multiplying Berg and Krüppel's asymptotic solution, in equation (9.7) for this eigenfunction, is e^H , matching the product formula their Proposition 9.3 supplies for it, as noted after Theorem 4. Proposition 6's Fourier series for H is the new, closed form; Proposition 8's certificate is what neither their product formula nor their closing remark (that they only *expect* φ bounded and bounded away from zero relative to φ_0) settles. $\square$

7. PROOFS OF THE MAIN THEOREMS

Proof of Theorem 1. Wirsching's class $\tilde{A}_\delta$ (his equations (1.5) and (3.2)) consists of sequences (x_l) with $\lfloor 3^l x_l \rfloor = k_l$ satisfying $|l - k_l| \leq \delta \sqrt{l}$ for every l ; hence $\lambda_l := 3^l x_l / l \rightarrow 1$ uniformly at rate $O(\delta / \sqrt{l})$ on the class. Writing $z_l = \lambda_l l 3^{-l}$, $\tau_l = -\log z_l = lc - \log l - \log \lambda_l$, the saddle location satisfies $w_0(\tau_l) = lc - \log \lambda_l + O(1/l)$ by Lemma 15's defining equation, so $w_0(\tau_l) \bmod c \rightarrow 0$ uniformly on the class, at rate $O(1/\sqrt{l})$. By Theorem 13, Proposition 16, Proposition 17 (which gives $P(\tau) - \log \varphi_0(e^{-\tau}) \rightarrow 0$, Section 6), and continuity of H (indeed Lipschitz, by (5)),

$$\log \varphi(z_l) - \log \varphi_0(z_l) = H(w_0(\tau_l)) + o(1) \rightarrow H(0)$$

uniformly on $\tilde{A}_\delta$, i.e. $\varphi(z_l)/\varphi_0(z_l) \rightarrow e^{H(0)}$ uniformly. This is exactly Conjecture 3, with the constant given explicitly. $\square$

Proof of Theorem 2. As $t = e^{-\tau}$ ranges over $(0, 1/2)$ with no restriction, $\tau \rightarrow \infty$. By Lemma 15, $w_0 = \Phi_0^{-1}$ is continuous and strictly increasing and unbounded on (τ_0, ∞) , so $w_0(\tau) \bmod c$ sweeps every value in $[0, c)$ infinitely often as $\tau \rightarrow \infty$ (a continuous, strictly increasing, unbounded function hits every residue class modulo c infinitely often), so by the same combination of results used above, $\log \varphi(t) - \log \varphi_0(t) = H(w_0(\tau)) + o(1)$ takes values arbitrarily close to both $\sup H$ and $\inf H$ infinitely often. By Proposition 8, $\text{osc}(H) > 0$ rigorously, so $\varphi(t)/\varphi_0(t)$ does not converge, and $\limsup / \liminf \geq e^{\text{osc}(H)} \geq 1.0004188$. $\square$

Proposition 18 (Wirsching's actual requirement). *For $(x_l) \in \tilde{A}_\delta$, write $x_l^+ := x_l + 3^{-(l+1)}$ (his equation (7.14)); since $x_l \sim l \cdot 3^{-l}$, x_l^+ is asymptotic to x_l and plays the same role as this paper's z_l). Wirsching's own use of Conjecture 3, via $\varphi'(x) = \frac{9}{2}\varphi(3x)$ on $(0, \frac{2}{3})$ and his own calculation, his equation (7.13), that $\lim_l (2/3^{l+1})\varphi'_0(x_l)/\varphi_0(x_l) = 2/3$ uniformly on the class, reduces to*

$$\limsup_l \Lambda_l < \frac{3}{2}, \quad \Lambda_l := \frac{\varphi(3x_l^+)/\varphi_0(3x_l^+)}{\varphi(x_l^+)/\varphi_0(x_l^+)}.$$

This holds with a wide margin: $\Lambda_l \rightarrow 1$.

Proof. First, the reduction itself. Wirsching's actual requirement is his inequality (7.5),

$$\limsup_l \frac{2}{3^{l+1}} \cdot \frac{\varphi'(x_l^+)}{\varphi(x_l^+)} < 1;$$

by $\varphi' = \frac{9}{2}\varphi(3\cdot)$ this is $\limsup_l (9/3^{l+1})\varphi(3x_l^+)/\varphi(x_l^+) < 1$. By definition of Λ_l , $\varphi(3x_l^+)/\varphi(x_l^+) = \Lambda_l \cdot \varphi_0(3x_l^+)/\varphi_0(x_l^+)$, and Berg and Krüppel's φ_0 is an asymptotic solution of the same truncated equation (their equation (7.12) in Wirsching's notation: $\varphi'_0(t)/\varphi_0(3t) \rightarrow 9/2$ as $t \rightarrow 0$), so $\varphi_0(3x_l^+)/\varphi_0(x_l^+) = \frac{2}{9}\varphi'_0(x_l^+)/\varphi_0(x_l^+)(1+o(1))$. To see that (7.13) transfers from x_l to x_l^+ with

an $o(1)$ correction, write $v = \log t$; the explicit form of φ_0 above gives $\varphi'_0(t)/\varphi_0(t) = B(v)/t$ with $B(v) = \gamma + \delta_{\text{BK}}/v - 2\beta(v - \log(-v))(1 - 1/v)$, and, as $v \rightarrow -\infty$, $B(v) \sim -2\beta v$ while $B'(v) \rightarrow -2\beta$, so $(\log B)'(v) = B'(v)/B(v) \rightarrow 0$ and $(\log[\varphi'_0/\varphi_0])'(v) = (\log B)'(v) - 1 \rightarrow -1$, bounded. Since $x_l^+/x_l = 1 + 3^{-(l+1)}/x_l \rightarrow 1$ (as $x_l \sim l \cdot 3^{-l}$, in fact $3^{-(l+1)}/x_l = O(1/l)$), $v(x_l^+) - v(x_l) = \log(x_l^+/x_l) = O(1/l)$, and the mean value theorem gives $\log[\varphi'_0/\varphi_0(x_l^+)] - \log[\varphi'_0/\varphi_0(x_l)] = O(1/l)$, i.e. $\varphi'_0(x_l^+)/\varphi_0(x_l^+) = \varphi'_0(x_l)/\varphi_0(x_l) \cdot (1 + O(1/l))$. Substituting and using his equation (7.13), $\lim_l (2/3^{l+1})\varphi'_0(x_l)/\varphi_0(x_l) = 2/3$ uniformly on the class,

$$\frac{9}{3^{l+1}} \cdot \frac{\varphi(3x_l^+)}{\varphi(x_l^+)} = \Lambda_l \cdot \frac{2}{3^{l+1}} \cdot \frac{\varphi'_0(x_l^+)}{\varphi_0(x_l^+)} (1 + o(1)) \rightarrow \frac{2}{3} \Lambda_l \cdot (1 + o(1)),$$

so the requirement $\limsup_l (9/3^{l+1})\varphi(3x_l^+)/\varphi(x_l^+) < 1$ is exactly $\limsup_l \Lambda_l < 3/2$.

For the value of Λ_l : the phase shift under $t \mapsto 3t$, i.e. $\tau \mapsto \tau - c$, is $w_0(\tau - c) - w_0(\tau) + c = O(1/\tau)$ by Lemma 15's defining equation, independently of the phase of τ . By Theorem 13, Proposition 16, and Proposition 17 together (the same chain used in the proof of Theorem 1), $\log \varphi(t) - \log \varphi_0(t) = H(w_0(\tau)) + o(1)$ uniformly in phase, so

$$\log \Lambda_l = H(w_0(\tau_l - c)) - H(w_0(\tau_l)) + o(1) = O(1/\tau_l) + o(1) \rightarrow 0,$$

using that H is Lipschitz ((5)) and the phase shift above is $O(1/\tau_l)$; hence $\Lambda_l \rightarrow 1$, well inside $\limsup \Lambda_l < 3/2$. $\square$

8. DISCUSSION

These results settle Conjecture 3 alone. Wirsching's positive-predecessor-density argument reduces to three conjectures; Conjectures 1 and 2, concerning respectively a pointwise-versus-average transfer for the Elka functions and a uniformity-near-a-singularity statement for the limiting operator S_∞ , are untouched here, and the full theorem is not established by this paper.

Wirsching stated the weaker Conjecture 3 instead of the stronger $\varphi \sim \kappa\varphi_0$ he considers first. Theorem 2 shows why: the stronger statement fails by a mechanism with a certified, positive amplitude, not by some correctable technical looseness. Every phase $\theta \in [0, c)$ swept by H admits some sequence along which the ratio converges to $e^{H(\theta)}$: take $\tau_l := \Phi_0(\theta + lc)$, $z_l := e^{-\tau_l}$, so $w_0(\tau_l) = \theta + lc$ exactly, and apply Theorem 13, Proposition 16, and Proposition 17 as in the proof of Theorem 1. What is special about phase 0 is only that it is the one Wirsching's own class $\tilde{A}_\delta$ happens to fix. That phase sits close to H 's minimum: $H(0) - \min H = 5.25 \times 10^{-6}$ against a total oscillation of 4.19×10^{-4} . We do not know whether this is coincidental.

The rate is not established beyond $o(1)$. The sharper Edgeworth correction of Remark 14, and a correspondingly sharper form of the envelope and Berg–Krüppel comparisons, would give an explicit polynomial rate throughout; neither theorem needs it.

9. CODE AND DATA AVAILABILITY

Every certified or numerically-checked claim in this paper is backed by a runnable script in the accompanying repository, <https://github.com/faculdade/wirsching-conjecture3-proof>, organized by section with a README per script stating what it verifies and how to run it. This includes the interval-arithmetic certificates behind Proposition 8 and the bounds (5), the constant chain behind Theorem 13, and the identity of Proposition 17.

Acknowledgments. AI assistance was used for textual review and translation.

REFERENCES

- [1] G. J. Wirsching, *On the problem of positive predecessor density in $3N + 1$ dynamics*, Discrete Contin. Dyn. Syst. **9** (2003), no. 3, 771–787.
- [2] L. Berg and M. Krüppel, *On the solution of an integral-functional equation with a parameter*, Z. Anal. Anwendungen **17** (1998), no. 1, 159–181.
- [3] G. J. Wirsching, *The Dynamical System Generated by the $3n + 1$ Function*, Lecture Notes in Mathematics, vol. 1681, Springer, Berlin, 1998.
- [4] V. A. Rvachev, *Compactly supported solutions of functional-differential equations and their applications*, Russian Math. Surveys **45** (1990), no. 1, 87–120.
- [5] N. G. de Bruijn, *On Mahler's partition problem*, Nederl. Akad. Wetensch., Proc. **51** (1948), 659–669 (also Indag. Math. **10**, 210–220).
- [6] P. Erdős and L. B. Richmond, *Concerning periodicity in the asymptotic behaviour of partition functions*, J. Austral. Math. Soc. Ser. A **21** (1976), 447–456.
- [7] T. Kato and J. B. McLeod, *The functional-differential equation $y'(x) = ay(\lambda x) + by(x)$* , Bull. Amer. Math. Soc. **77** (1971), no. 6, 891–937.
- [8] G. Derfel, *Functional-differential and functional equations with rescaling*, in Operator Theory: Advances and Applications, vol. 80, Birkhäuser, Basel, 1995, 100–111.

UNIVERSIDADE FEDERAL DE GOIÁS

Email address: `dr.renatotavares@gmail.com`