

SDS 385: Exercises 1 - Preliminaries

August 23, 2016

Professor James Scott

Spencer Woody

Problem 1

(A)

$$\hat{\beta} = \arg \min_{\beta \in \mathbb{R}^p} \sum_{i=1}^N \frac{w_i}{2} (y_i - x_i^T \beta)^2 \quad (1)$$

$$= \arg \min_{\beta \in \mathbb{R}^p} \frac{1}{2} (Y - X\beta)^T W (Y - X\beta) \quad (2)$$

$$\frac{1}{2} (Y - X\beta)^T W (Y - X\beta) = \frac{1}{2} (Y^T - \beta^T X^T) W (Y - X\beta) \quad (3)$$

$$= \frac{1}{2} (Y^T W - \beta^T X^T W) (Y - X\beta) \quad (4)$$

$$= \frac{1}{2} (Y^T W Y - \beta^T X^T W Y - Y^T W X \beta + \beta^T X^T W X \beta) \quad (5)$$

$$= \frac{1}{2} (Y^T W Y - 2(X\beta)^T W Y + \beta^T X^T W X \beta) \quad (6)$$

$$= \frac{1}{2} Y^T W Y - (X\beta)^T W Y + \frac{1}{2} \beta^T X^T W X \beta, \quad (7)$$

because

$$\beta^T X^T W Y = (X\beta)^T W Y, \quad (8)$$

and

$$Y^T W X \beta = (Y^T W X \beta)^T \because Y^T W X \beta \in \mathbb{R}^1 \quad (9)$$

$$(Y^T W X \beta)^T = (W X \beta)^T Y = (X\beta)^T W^T Y = (X\beta)^T W Y. \quad (10)$$

We want to minimize the objective function from Eqn. (7), so we take the gradient with respect to β and set it equal to zero. For each of the three terms, their are respective gradients with respect to β are

(i)

$$\frac{\partial}{\partial \beta} \frac{1}{2} Y^T W Y = 0 \quad (11)$$

(ii)

$$\frac{\partial}{\partial \beta} - (X\beta)^T W Y = -X^T W Y \quad (12)$$

(iii)

$$\frac{\partial}{\partial \beta} \frac{1}{2} \beta^T X^T W X \beta = \frac{1}{2} \beta^T (X^T W X + (X^T W X)^T) \quad (13)$$

$$= X^T W X \beta. \quad (14)$$

Summing these terms and equaling them to zero yields

$$X^T W X \beta - X^T W Y = 0 \therefore \quad (15)$$

$$(X^T W X) \hat{\beta} = X^T W Y \quad (16)$$

(B) The brute force method of solving Eqn. (16) is the *inversion method*, i.e.

$$\hat{\beta} = (X^T W X)^{-1} X^T W y. \quad (17)$$

However, this method is computationally expensive. Therefore I propose an alternative methods to solving this matrix equation using the Cholesky decomposition. **Cholesky Decomposition**
Let

$$C = X^T W X, \quad D = X^T W y \quad (18)$$

so

$$C \hat{\beta} = D. \quad (19)$$

We decompose matrix C into a product of a lower-triangular matrix and an upper-triangular matrix, such that $U = L^T$ so

$$C = LU = LL^T \therefore \quad (20)$$

$$LL^T \hat{\beta} = D. \quad (21)$$

Furthermore we define matrix $A = L^T \hat{\beta}$. Thus we are left with two matrix equations to solve.

$$LA = D \quad (22)$$

$$L^T \hat{\beta} = A \quad (23)$$

This method will be much less computationally intensive than the inversion method because of the fact that the two left-matrices L and $U = L^T$ are triangular. We still must invert L and L^T but this is simpler than taking an inverse of a more complicated matrix $X^T W X$. This is similar to an LU decomposition, with the exception that we necessarily have two triangular matrices that are transposes of one another. Therefore, this method gains a computational advantage over LU decomposition from symmetric exploitation.

(C) Code for implementing this method is shown in the appendix to this paper.

(D)

Problem 2

(A) We have $y_i \sim \text{Binomial}(m_i, w_i)$, where

$$w_i = \frac{1}{1 + \exp(-x_i^T \beta)}, \quad 1 - w_i = \frac{\exp(-x_i^T \beta)}{1 + \exp(-x_i^T \beta)}, \quad (24)$$

so the negative log likelihood is

$$\ell(\beta) = -\log \left\{ \prod_{i=1}^N p(y_i | \beta) \right\} \quad (25)$$

$$= -\log \left\{ \prod_{i=1}^N \binom{m_i}{y_i} (w_i)^{y_i} (1 - w_i)^{m_i - y_i} \right\} \quad (26)$$

$$= -\left\{ \sum_{i=1}^N \left(\log \binom{m_i}{y_i} + y_i \log(w_i) + (m_i - y_i) \log(1 - w_i) \right) \right\} \quad (27)$$

$$= -\left\{ \sum_{i=1}^N \left(\log \binom{m_i}{y_i} + y_i \log \left(\frac{1}{1 + \exp(-x_i^T \beta)} \right) + (m_i - y_i) \log \left(\frac{\exp(-x_i^T \beta)}{1 + \exp(-x_i^T \beta)} \right) \right) \right\} \quad (28)$$

$$= -\left\{ \sum_{i=1}^N \left(\log \binom{m_i}{y_i} - y_i \log(1 + \exp(-x_i^T \beta)) - (m_i - y_i) x_i^T \beta - m_i \log(1 + \exp(-x_i^T \beta)) + y_i \log(1 + \exp(-x_i^T \beta)) \right) \right\} \quad (29)$$

$$= -\left\{ \sum_{i=1}^N \left(\log \binom{m_i}{y_i} - (m_i - y_i) x_i^T \beta - m_i \log(1 + \exp(-x_i^T \beta)) \right) \right\} \quad (30)$$

$$= \sum_{i=1}^N \left((m_i - y_i) x_i^T \beta + m_i \log(1 + \exp(-x_i^T \beta)) - \log \binom{m_i}{y_i} \right) \quad (31)$$

$$(32)$$

The gradient for this expression is,

$$\nabla \ell(\beta) = \sum_{i=1}^N \left((m_i - y_i) x_i - m_i \frac{\exp(-x_i^T \beta)}{1 + \exp(-x_i^T \beta)} x_i \right) \quad (33)$$


$$= \sum_{i=1}^N ((m_i - y_i) x_i - m_i (1 - w_i) x_i) \quad (34)$$

$$= \sum_{i=1}^N (m_i w_i - y_i) x_i \quad (35)$$

(B) Code for implementing the gradient descent method is shown in the appendix. Note that we normalize the values in the X matrix and add a column of 1's to make an intercept term. We start by having an initial arbitrary guess for β , which we define as β_0 . Then we use an iterative process to converge upon the true value of β based on the calculated gradient of the log likelihood at $\hat{\beta}_t$ and an arbitrary step size, α as follows

$$\hat{\beta}_{t+1} = \hat{\beta}_t - \alpha \times \nabla \ell(\hat{\beta}_t) \quad (36)$$

We use an initial guess of $\beta_0 = 0$, a step size of $\alpha = 0.025$, and 50,000 iterations and reach convergence in optimizing the log likelihood, as shown in the trace plot below. Our final estimations of β are reported below along with estimations from R's native `glm` function. The two sets of estimates are in close agreement with one another.



	Grad descent	R: glm
$\hat{\beta}_1$	0.48553	0.48702
$\hat{\beta}_2$	-7.14618	-7.22185
$\hat{\beta}_3$	1.65481	1.65476
$\hat{\beta}_4$	-1.80713	-1.73763
$\hat{\beta}_5$	13.99290	14.00485
$\hat{\beta}_6$	1.07426	1.07495
$\hat{\beta}_7$	-0.07319	-0.07723
$\hat{\beta}_8$	0.67573	0.67512
$\hat{\beta}_9$	2.59383	2.59287
$\hat{\beta}_{10}$	0.44615	0.44626
$\hat{\beta}_{11}$	-0.48276	-0.48248

Table 1: Comparison of results from gradient descent and glm

- (C) We need to calculate the Hessian matrix of the log likelihood function, $\nabla^2(\ell(\beta))$. The Hessian will be a $P \times P$ matrix, with the element in row i and column j being¹

$$\frac{\partial^2}{\partial \beta_i \partial \beta_j} \ell(\beta) = \frac{\partial}{\partial \beta_i} \left(\frac{\partial}{\partial \beta_j} \ell(\beta) \right) \quad (37)$$

$$= \frac{\partial}{\partial \beta_i} \left(\frac{\partial}{\partial \beta_j} \sum_{k=1}^N (\dots) \right) \quad (38)$$

$$= \frac{\partial}{\partial \beta_i} \left(\sum_{k=1}^N (m_k w_k - y_k) x_{kj} \right) \quad (39)$$

$$= \sum_{k=1}^N x_{ki} x_{kj} m_k w_k (1 - w_k) \quad (40)$$

Note:

$$\frac{\partial}{\partial \beta_i} w_k = x_{ki} \frac{\exp(-x_k^T \beta)}{(1 + \exp(-x_k^T \beta))^2} \quad (41)$$

$$= x_{ki} w_k (1 - w_k) \quad (42)$$

This matrix is equivalent to $X^T W X$ where $W = \text{diag}(m_1 w_1 (1 - w_1), \dots, m_N w_N (1 - w_N))$

- (D) Now we use Newton's to estimate β . This is also an iterative process, though now we need far fewer iterations to achieve convergence because we are taking the curvature of our objective function ($\ell(\beta)$) into account. In fact, we only use 10 iterations and achieve estimates $\hat{\beta}$ which are *exactly* in line with estimates from `glm`.

Newton's Method:

$$\hat{\beta}_{t+1} = \hat{\beta}_t - (\nabla^2 \ell(\hat{\beta}_t))^{-1} \nabla \ell(\hat{\beta}_t) \quad (43)$$

	N.'s method	R: <code>glm</code>
$\hat{\beta}_1$	0.48702	0.48702
$\hat{\beta}_2$	-7.22185	-7.22185
$\hat{\beta}_3$	1.65476	1.65476
$\hat{\beta}_4$	-1.73763	-1.73763
$\hat{\beta}_5$	14.00485	14.00485
$\hat{\beta}_6$	1.07495	1.07495
$\hat{\beta}_7$	-0.07723	-0.07723
$\hat{\beta}_8$	0.67512	0.67512
$\hat{\beta}_9$	2.59287	2.59287
$\hat{\beta}_{10}$	0.44626	0.44626
$\hat{\beta}_{11}$	-0.48248	-0.48248

Table 2: Comparison of results from Newton's method and `glm`

- (E) Gradient descent: many iterations; Newton's method: must invert a matrix

¹Notice the reindexing shown below for summations.

```
#####
##### Created by Spencer Woody on 24 Aug 2016 #####
#####

5 library(Matrix)
  library(microbenchmark)

  ### No. 1 pt C

10 # Set N, P, X, W, and y

  N <- 2000
  P <- 500

15 X <- matrix(rnorm(N * P), nrow = N)
  y <- matrix(rnorm(N), nrow = N)
  W <- diag(rep(1, N))

  # Inversion method

20 Inv.method <- function(X.Inv, W.Inv, y.Inv) {
  XtWX <- (t(X.Inv)*diag(W.Inv)) %*% X.Inv
  XtWY <- (t(X.Inv)*diag(W.Inv)) %*% y.Inv
  bhat.Inv <- solve(XtWX) %*% XtWY
25   return(bhat.Inv)
}

  Cho.decomp <- function(X.Cho, W.Cho, y.Cho) {
  D.Cho <- (t(X.Cho)*diag(W.Cho)) %*% y.Cho
30   C.Cho <- (t(X.Cho)*diag(W.Cho)) %*% X.Cho

  U.Cho <- chol(C.Cho)
  L.Cho <- t(U.Cho)

35   u <- forwardsolve(L.Cho, D.Cho)
  bhat.Cho <- backsolve(U.Cho, u)

  return(bhat.Cho)
}

40 microbenchmark(
  Inv.method(X, W, y),
  Cho.decomp(X, W, y),
  times=5)

45

  ### No. 1 pt D

  N <- 2000
50 P <- 500

  X <- matrix(rnorm(N * P), nrow = N)
  mask <- matrix(rbinom(N * P, 1, 0.05), nrow = N)
```

```
X <- mask * X

55 Inv.methodSPARSE <- function(X.Inv, W.Inv, y.Inv) {
  X <- Matrix(X, sparse = TRUE)
  XtWX <- (t(X.Inv)*diag(W.Inv)) %*% X.Inv
  XtWY <- (t(X.Inv)*diag(W.Inv)) %*% y.Inv
60  bhat.Inv = Matrix::solve(XtWX, XtWY, sparse = TRUE)
  return(bhat.Inv)
}

Inv.methodSPARSE2 <- function(X.Inv, W.Inv, y.Inv) {
65  XtWX <- (t(X.Inv)*diag(W.Inv)) %*% X.Inv
  XtWY <- (t(X.Inv)*diag(W.Inv)) %*% y.Inv
  bhat.Inv = solve(XtWX, XtWY)
  return(bhat.Inv)
}

70 microbenchmark(
  Inv.methodSPARSE(X, W, y),
  Inv.methodSPARSE2(X, W, y),
  Cho.decomp(X, W, y),
75  times=5)

microbenchmark(
  solve(XtWX, XtWY),
80  Matrix::solve(XtWX, XtWY, sparse = TRUE),
  solve(XtWX) %*% XtWY,
  times = 5
)

85
# END
```



```
#####  
##### Created by Spencer Woody on 24 Aug 2016 #####  
#####  
5 wdbc <- read.csv("wdbc.csv", header = FALSE)  
  
X <- as.matrix(wdbc[, 3:12])  
X <- scale(X)  
X <- cbind(rep(1, nrow(X)), X)  
10  
y <- wdbc[, 2]  
y <- y == "M"  
beta <- as.matrix(rep(0, ncol(X)))  
mi <- 1  
15  
comp.wi <- function (X, beta) {  
  wi <- 1 / (1 + exp(-X %*% beta))  
  return(wi)  
}  
20  
loglik <- function(beta, y, X, mi) {  
  loglik <- apply((mi - y) * (X %*% beta) + mi*log(1 + exp(-X %*% beta)), 2, sum)  
  return(loglik)  
}  
25  
grad.loglik <- function(beta, y, X, mi){  
  grad <- array(NA, dim = length(beta))  
  wi <- comp.wi(X, beta)  
  grad <- apply(X*as.numeric(mi * wi - y), 2, sum)  
30  return(grad)  
}  
  
stepfactor <- 0.025  
n.steps <- 50000  
35 log.lik <- NULL  
  
for (step in 1:n.steps) {  
  log.lik[step] <- loglik(beta, y, X, mi)  
  beta <- beta - stepfactor * grad.loglik(beta, y, X, mi)  
40 }  
  
png("beta_trace1.png")  
plot(log.lik,  
      main = "Trace Plot for log likelihood(beta)",  
45      xlab = "Step",  
      ylab = "log likelihood(beta)")  
dev.off()  
  
mymodel <- glm(y ~ X[, c(-1)], family = "binomial")  
50 summary(mymodel)  
  
print(beta)
```

```

# Newton's method
55 beta.N <- as.matrix(rep(0, ncol(X)))

n.steps <- 10
log.lik2 <- NULL
60
for (step in 1:n.steps) {
  log.lik2[step] <- loglik(beta, y, X, mi)
  w.i <- as.numeric(comp.wi(X, beta.N))
  W <- diag(w.i*(1-w.i))
65 Hessian <- t(X) %*% W %*% X
  beta.N <- beta.N - solve(Hessian) %*% grad.loglik(beta.N, y, X, mi)
}

round(as.matrix(coef(mymodel)) - beta.N, 8)
70
# RESULTS
#
# FROM glm
# Coefficients:
75 #
#           Estimate Std. Error z value Pr(>|z|)
# (Intercept)   0.48702    0.56432   0.863   0.3881
# X[, c(-1)]V3  -7.22185   13.09494  -0.551   0.5813
# X[, c(-1)]V4   1.65476    0.27758   5.961 2.5e-09 ***
# X[, c(-1)]V5  -1.73763   12.27499  -0.142   0.8874
80 # X[, c(-1)]V6  14.00485    5.89090   2.377   0.0174 *
# X[, c(-1)]V7   1.07495    0.44942   2.392   0.0168 *
# X[, c(-1)]V8  -0.07723    1.07434  -0.072   0.9427
# X[, c(-1)]V9   0.67512    0.64733   1.043   0.2970
# X[, c(-1)]V10  2.59287    1.10701   2.342   0.0192 *
85 # X[, c(-1)]V11  0.44626    0.29143   1.531   0.1257
# X[, c(-1)]V12 -0.48248    0.60406  -0.799   0.4244
# ----
#
# FROM GRADIENT DESCENT
90 # > print(beta)
#           [,1]
# [1,]  0.48553491
# [2,] -7.14617798
# [3,]  1.65480926
95 # [4,] -1.80712669
# [5,] 13.99289624
# [6,]  1.07426088
# [7,] -0.07318783
# [8,]  0.67573353
100 # [9,]  2.59382838
# [10,] 0.44615349
# [11,] -0.48275720

# FROM NEWTON'S METHOD
105 # > round(beta.N, 5)
#           [,1]

```

```
110 #      0.48702
    # V3  -7.22185
    # V4   1.65476
    # V5  -1.73763
    # V6  14.00485
    # V7   1.07495
    # V8  -0.07723
    # V9   0.67512
115 # V10  2.59287
    # V11  0.44626
    # V12 -0.48248

120 #END
```