POLLUTION MAP OF INDIAN CITIES

A Geospatial Analysis of Air Quality Using Python

Capstone-II Mid-Semester Project Report

Submitted by the Student of

Hybrid UG Program in Computer Science & Data Analytics

Group - 121

S.No.	Student Name	Roll Number
1	Ritu Ranjan	2312RES533
2	Ratnesh Ranjan	2312RES506
3	Shishir Ranjan	2312RES600
4	Roopesh Ranjan	2312RES542
5	Priyansh Ranjan	2312RES468

INDIAN INSTITUTE OF TECHNOLOGY PATNA

BIHTA - 801106, INDIA

30 March 2026

ABSTRACT

Air pollution has become a critical environmental and public health crisis in India, with millions of citizens exposed to hazardous air quality levels daily. Understanding the spatial distribution of pollution across different cities is essential for effective policy formulation, resource allocation, and public awareness. This project presents a comprehensive geospatial visualization of air pollution levels across major Indian cities using Python's powerful geospatial libraries.

The primary objective of this work is to create an informative map that visually represents PM2.5 concentrations across 15 major Indian cities, enabling quick identification of highly affected areas and facilitating spatial pattern analysis. The methodology employs GeoPandas for geospatial data handling, Shapely for geometric operations, and Matplotlib for visualization. The India boundary is constructed using polygon coordinates, while city-specific pollution data is processed using Pandas.

Key outcomes of this project include:

1. A detailed geospatial map of India with cities marked using distinct markers and sizes corresponding to their pollution severity.

2. Comprehensive statistical analysis including maximum, minimum, and mean PM2.5 values.

3. Bar charts and pie charts illustrating pollution category distribution.

4. Map elements such as legend, scale bar, north arrow, and statistical information box for enhanced readability.

The results reveal significant spatial variation in pollution levels, with northern cities in the Indo-Gangetic Plain showing substantially higher $PM_{2.5}$ concentrations compared to coastal and southern regions. Delhi recorded the highest pollution level at 310 $\mu g/m^3$, while Chennai showed the lowest at 45 $\mu g/m^3$. The project successfully demonstrates the effectiveness of geospatial data visualization in environmental analysis and provides a reproducible framework for similar studies.

Summary of the Project

This project creates a geospatial visualization of air pollution levels across major Indian cities using Python. The work combines geographic data with pollution statistics from a CSV file to produce an informative map for identifying highly affected areas and analyzing environmental patterns.

Methodology: The implementation follows a systematic approach:

(1) loading and validating pollution data from `pollution_data.csv`,

(2) creating India's outline using polygon coordinates,

(3) merging geographic and pollution datasets,

(4) generating the base map with India's boundary,

(5) plotting cities with pollution-based markers,

(6) adding essential map elements, and

(7) creating supplementary visualizations.

Key Results: The maximum $PM_{2.5}$ concentration is 310 $\mu g/m^3$ (Delhi), minimum is 45 $\mu g/m^3$ (Chennai), and mean is 113.5 $\mu g/m^3$. Category-wise distribution shows 6.7% cities in "Very Unhealthy", 13.3% in "Unhealthy", 13.3% in "Unhealthy for Sensitive Groups", 53.3% in "Moderate", and 13.4% with "Good" air quality. Northern cities exhibit significantly higher pollution levels than coastal and southern regions.

Conclusion: This project highlights the potential of geospatial data visualization for analyzing environmental conditions. It provides a scalable and reproducible framework that can be further refined and expanded to include predictive models for air pollution. By leveraging historical datasets, the system can be adapted to estimate future pollution levels and trends, thereby aiding in informed decision-making and long-term environmental planning.

Work Done by Each Member

1. **Shishir Ranjan (2312RES600)** - Data Collection and Preprocessing: Gathered pollution data for major Indian cities, created CSV structure, performed data cleaning, validation, and implemented coordinate verification.

2. **Priyansh Ranjan (2312RES468)** - Geospatial Data Processing: Downloaded and processed shapefiles, created India outline polygon using Shapely, and implemented coordinate reference systems (CRS).

3. **Roopesh Ranjan (2312RES542)** - Map Visualization: Implemented the main map plotting using Matplotlib, created marker-based differentiation for pollution levels, and added map elements including legend, scale bar, north arrow, and grid lines.

4. **Ritu Ranjan (2312RES533)** - Statistical Analysis: Performed comprehensive data analysis using Pandas and NumPy, calculated statistical measures (mean, median, standard deviation), created bar charts and pie charts for pollution distribution visualization, and generated statistical summary reports.

5. **Ratnesh Ranjan (2312RES506)** - System Integration, Documentation, and Testing: Integrated geospatial data with visualization module, added city labels with annotations, prepared complete project documentation in LaTeX, tested code functionality across different environments, debugged implementation issues, created final report with all sections, and ensured code reproducibility.

My Work: Ratnesh Ranjan

Roll Number: 2312RES506

Overview of Contribution

My role was multifaceted, acting as the integrator and quality assurance lead for the project. While my teammates focused on individual components, I was responsible for stitching these components together into a cohesive and functional whole. This involved system integration, ensuring code reliability through testing, and creating comprehensive documentation to make our work reproducible and understandable.

System Integration

A major part of my work was to integrate the geospatial data and visualization modules created by my teammates. This involved:

a. **City Labels with Annotations:** I enhanced the main map by adding clear city labels. To prevent labels from overlapping with markers, I implemented leader lines (arrows) that connect the city name to its corresponding point on the map, a detail that significantly improves readability.

b. **Data-Visualization Pipeline:** I wrote the glue code that took the `india_gdf` (from Priyansh) and the `pollution_df` (from Shishir) and fed them into the plotting functions (from Roopesh) to generate the final map. I ensured that all components were compatible and that the workflow from data loading to final output was seamless.

c. **Final Report Compilation:** I compiled all individual contributions—the text, the code, the figures, and the analysis—into a single, cohesive LaTeX document, ensuring a consistent format and narrative flow across all chapters.

Documentation

To ensure the project's longevity and usability, I prepared comprehensive documentation. This included:

1. **LaTeX Documentation:** I structured the final report in LaTeX, managing chapters, sections, figures, tables, and cross-references to create a professional, academic-grade document.

2. **Code Comments:** While my teammates commented their own code, I reviewed and standardized the comments to ensure clarity and consistency across all modules.

3. **Installation Instructions:** I created a clear and concise installation guide (Appendix B), listing the required libraries (`geopandas`, `pandas`, `matplotlib`, `numpy`, `shapely`) and the necessary commands (Listing B.1).

4. **Complete Code Listing:** I prepared a complete, well-formatted listing of the entire codebase (Appendix A), making it easy for others to view and run the project in its entirety.

Testing and Debugging

Ensuring the code worked correctly and reproducibly was a critical responsibility. I performed:

A) **Cross-Environment Testing:** I ran the complete code on different operating systems and Python environments to ensure there were no hidden dependencies or platform-specific issues.

B) **Debugging Implementation Issues:** I helped resolve integration bugs, such as coordinate mismatches, legend formatting errors, and file path issues that arose when combining the different modules.

C) **Reproducibility Assurance:** I verified that the entire project—from running the code to generating the final report—could be reproduced from scratch using the provided data and instructions. This ensures the integrity and scientific validity of our work.

Contribution to Project Outcomes

My work was essential for delivering a complete and professional project:

1. I ensured that the final map (Figure 5.1) was not just a plot, but a complete figure with clear, annotated city labels.

2. I produced the comprehensive final report, ensuring all contributions were properly documented and presented.

3. My testing and documentation guarantee that this project is not a one-off exercise but a reproducible framework that can be used for future work, as mentioned in the "Future Work" section of the report.

Contents

Summary of the Project **2**

Work Done by Each Member **3**

My Work: Ratnesh Ranjan **4**

1 Introduction **11**
1.1 Background and Motivation 11
1.2 Problem Statement 11
1.3 Project Objectives 11
1.4 Scope and Limitations 12
1.5 Report Organization 12

2 Literature Review **13**
2.1 Geospatial Data Visualization 13
2.1.1 Python Geospatial Libraries 13
2.2 Air Pollution Studies in India 13
2.3 Related Work 14

3 Methodology **15**
3.1 Data Collection 15
3.1.1 Pollution Data 15
3.1.2 Geospatial Data 16
3.2 Data Preprocessing 16
3.2.1 Pollution Categorization 16
3.3 Technical Approach 17
3.3.1 Software Stack 17
3.3.2 System Architecture 17
3.4 Implementation Strategy 17

4 Implementation **18**
4.1 Code Structure 18
4.1.1 Data Loading and Preparation 18
4.1.2 India Outline Creation 19
4.1.3 Map Visualization 19
4.1.4 Statistical Analysis 20
4.2 Key Implementation Decisions 20

4.2.1 Marker-Based Differentiation 21
4.2.2 Map Elements 21

5 Results and Analysis **22**
5.1 Generated Maps 22
5.2 Key Findings 23
5.2.1 Pollution Distribution 23
5.2.2 Statistical Summary 23
5.2.3 City Rankings 23
5.2.4 Category Distribution 24
5.3 Discussion 24
5.3.1 Geographic Patterns 24
5.3.2 Interpretation 24

6 Conclusion and Future Work **26**
6.1 Conclusion 26
6.2 Future Work 26
6.2.1 Data Improvements 26
6.2.2 Technical Enhancements 26
6.2.3 Application Extensions 27
6.3 Concluding Remarks 27

References/Bibliography **27**

A Complete Code Listing **29**

B Installation Instructions **30**

List of Figures

2.1 Statistical Analysis of Pollution Distribution 14

5.1 India Pollution Map 22

5.2 Statistical Analysis: Bar Chart and Pie Chart 24

List of Tables

2.1 Comparison of Python Geospatial Libraries 13

3.1 Complete Pollution Dataset from pollution_data.csv 16
3.2 Pollution Categories and AQI Standards 16

5.1 Statistical Summary of PM2.5 Levels 23
5.2 Top 5 Most Polluted Cities 23

List of Listings

4.1 Data Loading and Preparation 18
4.2 Creating India Boundary 19
4.3 Main Map Generation 19
4.4 Statistical Analysis 20
B.1 Installation Commands 30

1. Introduction

1.1 Background and Motivation

Air pollution has emerged as one of the most pressing environmental challenges of the 21st century, particularly in developing nations like India. According to the World Health Organization (WHO), India is home to several of the world's most polluted cities, with particulate matter (PM2.5) levels often exceeding safe limits by significant margins. The health implications are severe, ranging from respiratory diseases to cardiovascular problems, and even premature death.

The primary pollutants affecting Indian cities include: (1) PM2.5 (fine particulate matter with diameter less than 2.5 micrometers), (2) PM10 (coarse particulate matter with diameter less than 10 micrometers), (3) NO2 (nitrogen dioxide from vehicle emissions), and (4) SO2 (sulfur dioxide from industrial processes).

Understanding the spatial distribution of pollution is crucial for: (i) identifying highly affected areas for targeted intervention, (ii) analyzing environmental patterns and pollution sources, (iii) planning urban development and infrastructure, and (iv) raising public awareness about air quality issues.

1.2 Problem Statement

Air pollution levels vary significantly across different cities in India. Displaying this information on a map makes it easier to identify highly affected areas and analyze environmental patterns. Tabular pollution data alone cannot reveal spatial insights effectively. Therefore, the question addressed in this project is: **How can we create a map that visually displays air pollution levels across major Indian cities using geospatial data in Python?**

1.3 Project Objectives

The primary objectives of this project are:

1. Load and process geospatial data of India using GeoPandas.
2. Prepare and clean air pollution data for major Indian cities from `pollution_data.csv`.
3. Merge geographic and pollution datasets for spatial analysis.
4. Create a visual map with cities marked based on pollution levels.

5. Generate supporting statistical visualizations for comprehensive analysis.

6. Save the final output as high-quality image files for reporting.

1.4 Scope and Limitations

The scope of this project includes: (1) 15 major Indian cities with available pollution data in the provided CSV file, (2) PM2.5 as the primary air quality indicator, (3) geospatial visualization using Python libraries, and (4) professional presentation suitable for academic reporting.

Limitations of the current implementation include: (a) limited number of cities due to data availability, (b) static representation without real-time updates, (c) simplified India boundary representation, and (d) lack of temporal analysis capabilities.

1.5 Report Organization

This report is organized into the following chapters: Chapter 2 (Literature Review) discusses existing work in geospatial pollution mapping. Chapter 3 (Methodology) details the approach, tools, and techniques used. Chapter 4 (Implementation) describes the code structure and implementation details. Chapter 5 (Results and Analysis) presents the generated maps and statistical findings. Chapter 6 (Conclusion and Future Work) summarizes achievements and discusses future directions.

2. Literature Review

2.1 Geospatial Data Visualization

Geospatial data visualization has evolved significantly over the past decade. Traditional approaches relied on Geographic Information Systems (GIS) such as ArcGIS and QGIS. However, the rise of Python as a data science language has led to the development of powerful geospatial libraries like GeoPandas, Shapely, and Folium [1].

2.1.1 Python Geospatial Libraries

Table 2.1: Comparison of Python Geospatial Libraries

Library	**Purpose**	**Key Features**	**Use Case**
GeoPandas	Spatial data manipulation	Extends pandas for geospatial data	This project
Shapely	Geometric operations	Creating and manipulating geometric objects	Polygon creation
Matplotlib	Visualization	Plotting and rendering maps	Map visualization
Folium	Interactive maps	Leaflet.js integration	Interactive web maps
Cartopy	Map projections	Advanced cartographic projections	Geographic projections

2.2 Air Pollution Studies in India

Several studies have examined air pollution patterns in Indian cities. The Central Pollution Control Board (CPCB) maintains a network of monitoring stations across the country, providing valuable data for analysis [2]. Research by Ghude et al. (2016) demonstrated the seasonal variation in PM2.5 levels across the Indo-Gangetic Plain [3].



Figure 2.1: Statistical Analysis of Pollution Distribution

2.3 Related Work

Previous work in this domain includes: (1) Urban Heat Island Analysis combining temperature and pollution data, (2) Source Apportionment identifying pollution sources using receptor models, (3) Health Impact Assessment correlating pollution levels with health outcomes, and (4) Predictive Modeling using machine learning for pollution forecasting.

3. Methodology

3.1 Data Collection

3.1.1 Pollution Data

The pollution data for this project was provided through `pollution_data.csv`, which includes: (1) 15 major Indian cities with their pollution levels, (2) geographic coordinates (latitude and longitude), and (3) state information for regional analysis.

The dataset includes the following cities with their PM2.5 levels:

- Delhi (310 μg/m^3)
- Mumbai (98 μg/m^3)
- Chennai (45 μg/m^3)
- Kolkata (112 μg/m^3)
- Bangalore (68 μg/m^3)
- Hyderabad (72 μg/m^3)
- Ahmedabad (95 μg/m^3)
- Pune (85 μg/m^3)
- Jaipur (110 μg/m^3)
- Lucknow (140 μg/m^3)
- Patna (180 μg/m^3)
- Bhopal (88 μg/m^3)
- Chandigarh (55 μg/m^3)
- Guwahati (65 μg/m^3)
- Bhubaneshwar (58 μg/m^3)

Table 3.1: Complete Pollution Dataset from pollution_data.csv

City	State	Latitude	Longitude	PM2.5 (μg/m^3)
Delhi	Delhi	28.61	77.20	310
Mumbai	Maharashtra	19.07	72.87	98
Chennai	Tamil Nadu	13.08	80.27	45
Kolkata	West Bengal	22.57	88.36	112
Bangalore	Karnataka	12.97	77.59	68
Hyderabad	Telangana	17.38	78.48	72
Ahmedabad	Gujarat	23.02	72.57	95
Pune	Maharashtra	18.52	73.85	85
Jaipur	Rajasthan	26.91	75.78	110
Lucknow	Uttar Pradesh	26.84	80.94	140
Patna	Bihar	25.59	85.13	180
Bhopal	Madhya Pradesh	23.25	77.41	88
Chandigarh	Punjab	30.73	76.78	55
Guwahati	Assam	26.14	91.73	65
Bhubaneshwar	Odisha	20.29	85.82	58

3.1.2 Geospatial Data

The India boundary data was created using polygon coordinates approximating the country's outline. The coordinates were derived from geographic data sources and represent the major boundaries of India.

3.2 Data Preprocessing

The preprocessing steps include: (1) data cleaning (removing duplicates and handling missing values), (2) coordinate validation (ensuring all coordinates are within India's boundaries), and (3) categorization (classifying pollution levels into meaningful categories).

3.2.1 Pollution Categorization

PM2.5 levels were categorized according to the Air Quality Index (AQI) standards:

Table 3.2: Pollution Categories and AQI Standards

PM2.5 Range (μg/m^3)	Category	Health Impact
0-50	Good	Minimal impact
51-100	Moderate	Acceptable for sensitive groups
101-150	Unhealthy for Sensitive Groups	May affect sensitive individuals
151-200	Unhealthy	May affect general population
Greater than 200	Very Unhealthy	Health alert for all

3.3 Technical Approach

3.3.1 Software Stack

The project utilizes the following Python libraries: (1) GeoPandas for geospatial data handling, (2) Pandas for data manipulation and analysis, (3) Matplotlib for visualization and plotting, (4) Shapely for geometric operations, and (5) Numpy for numerical computations.

3.3.2 System Architecture

The system architecture consists of four main components: (A) Data Layer (CSV files containing pollution data), (B) Processing Layer (data cleaning and merging operations), (C) Visualization Layer (map generation and rendering), and (D) Output Layer (image file generation and storage).

3.4 Implementation Strategy

The implementation follows a modular approach: (1) load and validate input data, (2) create India outline using Shapely polygons, (3) merge geographic and pollution data, (4) generate base map with India boundary, (5) plot cities with pollution-based markers, (6) add map elements (legend, scale, north arrow), (7) create statistical visualizations, and (8) save outputs for reporting.

4. Implementation

4.1 Code Structure

The implementation is organized into logical sections for clarity and maintainability. Below is the complete code with detailed comments:

4.1.1 Data Loading and Preparation

```python
import geopandas as gpd
import pandas as pd
import matplotlib.pyplot as plt
import matplotlib.patches as mpatches
import numpy as np
import os
from shapely.geometry import Point, Polygon

# Load pollution data from CSV file
if not os.path.exists('pollution_data.csv'):
    # Create sample data with 15 major Indian cities
    sample_data = {
        'City': ['Delhi', 'Mumbai', 'Chennai', 'Kolkata', 'Bangalore',
                 'Hyderabad', 'Ahmedabad', 'Pune', 'Jaipur', 'Lucknow',
                 'Patna', 'Bhopal', 'Chandigarh', 'Guwahati', 'Bhubaneshwar'],
        'State': ['Delhi', 'Maharashtra', 'Tamil Nadu', 'West Bengal', 'Karnataka',
                  'Telangana', 'Gujarat', 'Maharashtra', 'Rajasthan', 'Uttar Pradesh',
                  'Bihar', 'Madhya Pradesh', 'Punjab', 'Assam', 'Odisha'],
        'Latitude': [28.61, 19.07, 13.08, 22.57, 12.97,
                     17.38, 23.02, 18.52, 26.91, 26.84,
                     25.59, 23.25, 30.73, 26.14, 20.29],
        'Longitude': [77.20, 72.87, 80.27, 88.36, 77.59,
                      78.48, 72.57, 73.85, 75.78, 80.94,
                      85.13, 77.41, 76.78, 91.73, 85.82],
        'PM2.5': [310, 98, 45, 112, 68,
                  72, 95, 85, 110, 140,
                  180, 88, 55, 65, 58]
    }
    pollution_df = pd.DataFrame(sample_data)
    pollution_df.to_csv('pollution_data.csv', index=False)
```

```
else:
    pollution_df = pd.read_csv('pollution_data.csv')
```

Listing 4.1: Data Loading and Preparation

4.1.2 India Outline Creation

```
# Create India outline using polygon coordinates
india_coords = [
    (68.5, 8.0), (70.0, 8.5), (72.0, 10.0), (75.0, 12.0),
    (78.0, 12.5), (80.0, 14.0), (82.0, 16.0), (85.0, 18.0),
    (88.0, 20.0), (90.0, 22.0), (92.0, 24.0), (95.0, 26.0),
    (97.0, 27.5), (97.0, 28.5), (96.5, 30.0), (95.0, 31.0),
    (92.0, 32.5), (90.0, 34.0), (88.0, 35.0), (85.0, 36.0),
    (82.0, 36.5), (80.0, 37.0), (78.0, 36.5), (75.0, 36.0),
    (73.0, 35.0), (71.0, 33.0), (69.0, 30.0), (68.0, 28.0),
    (67.5, 25.0), (67.0, 22.0), (67.5, 20.0), (68.0, 18.0),
    (68.0, 15.0), (68.2, 12.0), (68.5, 8.0)
]

india_polygon = Polygon(india_coords)
india_gdf = gpd.GeoDataFrame(geometry=[india_polygon], crs="EPSG:4326")
```

Listing 4.2: Creating India Boundary

4.1.3 Map Visualization

```
# Create figure and axis
fig, ax = plt.subplots(1, 1, figsize=(16, 12), facecolor='white')

# Plot India outline
india_gdf.boundary.plot(ax=ax, color='black', linewidth=2, zorder=2)

# Plot cities with pollution-based markers
for idx, row in pollution_df.iterrows():
    if row['PM2.5'] <= 50:
        marker = 'o'       # circle for good
        size = 50
        color = 'darkgray'
    elif row['PM2.5'] <= 100:
        marker = 's'       # square for moderate
        size = 80
        color = 'gray'
    elif row['PM2.5'] <= 150:
        marker = '^'       # triangle for unhealthy for sensitive
        size = 120
```

```
        color = 'dimgray'
    elif row['PM2.5'] <= 200:
        marker = 'D'       # diamond for unhealthy
        size = 160
        color = 'darkslategray'
    else:
        marker = '*'       # star for very unhealthy
        size = 200
        color = 'black'

    ax.scatter(row['Longitude'], row['Latitude'],
               marker=marker, s=size, c=color,
               edgecolors='black', linewidth=1.5, zorder=5)
```

Listing 4.3: Main Map Generation

4.1.4 Statistical Analysis

```
# Calculate statistics
max_pollution = pollution_df['PM2.5'].max()
min_pollution = pollution_df['PM2.5'].min()
avg_pollution = pollution_df['PM2.5'].mean()

# Categorize pollution levels
def categorize_pollution(pm25):
    if pm25 <= 50:
        return 'Good'
    elif pm25 <= 100:
        return 'Moderate'
    elif pm25 <= 150:
        return 'Unhealthy for Sensitive Groups'
    elif pm25 <= 200:
        return 'Unhealthy'
    else:
        return 'Very Unhealthy'

pollution_df['Category'] = pollution_df['PM2.5'].apply(
    categorize_pollution)
```

Listing 4.4: Statistical Analysis

4.2 Key Implementation Decisions

The following decisions were made during implementation:

4.2.1 Marker-Based Differentiation

Different marker shapes and sizes were used to represent pollution levels: (1) Circle (o) for Good quality (0-50 μg/m^3), (2) Square (s) for Moderate quality (51-100 μg/m^3), (3) Triangle ($^{)}forUnhealthyforsensitive(101-150\mu$g/m^3), (4) Diamond (D) for Unhealthy (151-200 μg/m^3), and (5) Star (*) for Very unhealthy (greater than 200 μg/m^3).

4.2.2 Map Elements

The map includes several essential elements: (1) scale bar with approximate distance calculation, (2) north arrow for orientation, (3) grid lines for coordinate reference, (4) statistical information box, and (5) city labels with arrows for clarity.

5. Results and Analysis

5.1 Generated Maps

The project successfully generated a comprehensive pollution map of India showing PM2.5 levels across major cities.



Figure 5.1: India Pollution Map

5.2 Key Findings

5.2.1 Pollution Distribution

Analysis of the data reveals several important patterns: (1) cities in the Indo-Gangetic Plain show consistently higher pollution levels, (2) coastal cities like Mumbai and Chennai have relatively lower PM2.5 concentrations, (3) the capital region (Delhi) exhibits the highest pollution level at 310 $\mu g/m^3$, and (4) significant variation exists across different regions.

5.2.2 Statistical Summary

Table 5.1: Statistical Summary of PM2.5 Levels

Statistic	**Value ($\mu g/m^3$)**
Maximum	310
Minimum	45
Mean	113.5
Median	88.0
Standard Deviation	71.2

5.2.3 City Rankings

Table 5.2: Top 5 Most Polluted Cities

Rank	**City**	**PM2.5 ($\mu g/m^3$)**
1	Delhi	310
2	Patna	180
3	Lucknow	140
4	Kolkata	112
5	Jaipur	110

5.2.4 Category Distribution



Figure 5.2: Statistical Analysis: Bar Chart and Pie Chart

The analysis shows that: (1) 6.7% of cities fall under the "Very Unhealthy" category (Delhi only), (2) 6.7% fall under "Unhealthy" category (Patna only), (3) 20.0% are "Unhealthy for Sensitive Groups" (Lucknow, Kolkata, Jaipur), (4) 53.3% are in "Moderate" category (Mumbai, Ahmedabad, Pune, Hyderabad, Bangalore, Bhopal, Guwahati, Bhubaneshwar, Chandigarh), and (5) 6.7% show "Good" air quality (Chennai only).

5.3 Discussion

5.3.1 Geographic Patterns

The results clearly demonstrate the influence of geography on pollution levels. The Indo-Gangetic Plain shows elevated pollution due to agricultural burning practices, industrial concentration, geographic bowl effect trapping pollutants, and high population density. Coastal areas show better air quality due to sea breeze dispersion, less industrial concentration, and better infrastructure.

5.3.2 Interpretation

The mapping approach successfully achieves its objective of providing spatial insights that would be difficult to obtain from tabular data alone. The visual representation clearly shows the concentration of high pollution in northern India, the relatively cleaner

air in southern coastal regions, the urban-rural pollution gradient, and the impact of industrial corridors.

6. Conclusion and Future Work

6.1 Conclusion

This project successfully demonstrates the creation of a geospatial map for visualizing air pollution levels across major Indian cities. The following objectives were achieved: (1) data integration (successfully merged geographic and pollution data from `pollution_data.csv`), (2) visualization (created a comprehensive map with pollution-based markers), (3) analysis (generated statistical insights from the pollution data), and (4) documentation (produced a complete report with findings).

The project contributes to the field of environmental data analysis by: (i) demonstrating Python's capabilities for geospatial analysis, (ii) providing a reproducible methodology for pollution mapping, (iii) creating visual tools for public awareness and policy-making, and (iv) establishing a foundation for more advanced studies.

6.2 Future Work

Several enhancements could be made to extend this project:

6.2.1 Data Improvements

A. **Real-time Data Integration:** Connect to live API feeds from pollution monitoring stations.

B. **Expanded City Coverage:** Include tier-2 and tier-3 cities for comprehensive analysis.

C. **Multiple Pollutants:** Add PM10, NO2, SO2, and other pollutants for multidimensional analysis.

D. **Temporal Analysis:** Include seasonal and annual trends.

6.2.2 Technical Enhancements

Interactive Maps: Implement using Folium or Plotly for interactive web-based maps.

Advanced Geospatial Analysis: Incorporate spatial interpolation (Kriging, IDW) for continuous surfaces.

Machine Learning: Develop predictive models for pollution forecasting.

Mobile Application: Create a mobile app for real-time pollution information.

6.2.3 Application Extensions

- ⋆ **Health Impact Assessment:** Correlate pollution levels with health data.
- ⋆ **Source Apportionment:** Identify pollution sources using advanced algorithms.
- ⋆ **Policy Impact Analysis:** Evaluate effectiveness of pollution control measures.
- ⋆ **Public Awareness Campaigns:** Develop accessible visualization tools for public use.

6.3 Concluding Remarks

The project demonstrates the power of geospatial data visualization in understanding and communicating environmental challenges. As air pollution continues to be a critical issue in India, tools like these become increasingly important for policymakers, researchers, and the public. The modular and well-documented implementation provides a solid foundation for future work in this important domain.

Bibliography

[1] GeoPandas Development Team. (2020). *GeoPandas: Python tools for geographic data*. Available online at: https://geopandas.org/

[2] Central Pollution Control Board. (2023). *National Air Quality Monitoring Programme*. Ministry of Environment, Forest and Climate Change, Government of India. Available at: https://cpcb.nic.in/

[3] Ghude, S. D., et al. (2016). *Seasonal variation of PM2.5 over the Indo-Gangetic Plain*. Atmospheric Environment, 145, 210-223.

[4] World Health Organization. (2021). *WHO global air quality guidelines*. WHO Press. Available at: https://www.who.int/publications/i/item/9789240034228

[5] Sharma, S., et al. (2020). *Effect of restricted emissions during COVID-19 on air quality in India*. Science of The Total Environment, 728, 138878.

[6] Chowdhury, S., et al. (2019). *Indian annual ambient air quality standard is achievable by completely mitigating emissions from household sources*. Proceedings of the National Academy of Sciences, 116(22), 10711-10716.

[7] Balakrishnan, K., et al. (2019). *The impact of air pollution on deaths, disease burden, and life expectancy across the states of India*. The Lancet Planetary Health, 3(1), e26-e39.

[8] Hunter, J. D. (2007). *Matplotlib: A 2D graphics environment*. Computing in Science & Engineering, 9(3), 90-95. Available at: https://matplotlib.org/

[9] McKinney, W. (2010). *Data structures for statistical computing in Python*. Proceedings of the 9th Python in Science Conference, 51-56. Available at: https://pandas.pydata.org/

[10] Gillies, S., et al. (2021). *Shapely: Manipulation and analysis of geometric objects*. Available at: https://shapely.readthedocs.io/

[11] World Air Quality Index Project. (2024). *Real-time Air Quality Index (AQI) data for cities worldwide*. Available at: https://aqicn.org/

[12] Ministry of Environment, Forest and Climate Change. (2022). *National Clean Air Programme (NCAP)*. Government of India. Available at: https://moef.gov.in/

A. Complete Code Listing

The complete code for the project is available in the project repository. Key functions and their purposes are documented in Chapter 4, with code listings provided in Listings 4.1, 4.2, 4.3, and 4.4.

Project Repository Available at: https://github.com/ratneshranjan484/Pollution-Map-of-Indian-Cities

B. Installation Instructions

To run this project, install the required dependencies:

```
pip install geopandas pandas matplotlib numpy shapely
```

Listing B.1: Installation Commands

System Requirements

- Python 3.7 or higher.
- Minimum 4GB RAM.
- 500MB free disk space.
- Operating System: Windows, Linux, or macOS.