



A MTBLS79 cow versus sheep
Effects estimated separately by batch after joint correction
Pearson effect correlation
Spearman effect correlation
Sign concordance
Raw
ComBat
QC−RLSC
TIGER
SERRF
WiNN (QC−tuned)
WiNN (inferred segments)
WiNN
0.0
0.5
1.0
Fold bootstrap mean and 95% percentile interval
Held−out batch
1
2
3
4
5
6
7
8
Intervals describe individual folds after joint correction.



B WaveICA biological groups
Effects estimated separately by batch after joint correction
Pearson effect correlation
Spearman effect correlation
Sign concordance
Raw
ComBat
QC−RLSC
TIGER
SERRF
WiNN (QC−tuned)
WiNN (inferred segments)
WiNN
0.00
0.25
0.0
0.2
0.0
0.3
0.6
Fold bootstrap mean and 95% percentile interval
Held−out batch 1 2 3
Intervals describe individual folds after joint correction.

C Sacurine age

Effects estimated separately by batch after joint correction




Intervals describe individual folds after joint correction.


D Sacurine BMI
Effects estimated separately by batch after joint correction
Pearson effect correlation
Spearman effect correlation
Sign concordance
Raw
ComBat
QC−RLSC
TIGER
SERRF
WiNN (QC−tuned)
WiNN (inferred segments)
WiNN
0.00
0.25
0.00
0.25
0.0
0.4
0.8
Fold bootstrap mean and 95% percentile interval
Held−out batch
ne1
ne2
Intervals describe individual folds after joint correction.

E Sacurine gender

Effects estimated separately by batch after joint correction



Intervals describe individual folds after joint correction.