

Gleason's Theorem and Quantum Logic

Changjae Lee

November 23, 2024

Contents

1	Preliminaries	3
1.1	Elements of Operator Algebras	3
1.2	Quantum Logic	4
2	Gleason's Theorem	5
2.1	Gleason's Theorem	5

Introduction

In 1932, John von Neumann noticed that the projection operators of Hilbert space could be viewed as quantum mechanical propositions about observables [8]. The principles governing these quantum propositions were then called quantum logic by von Neumann and Birkhoff [2]. Von Neumann also managed to derive the Born rule in his textbook [8]. However, his assumptions were regarded to not be well-motivated by John Bell [1], [3].

By the late 1940s, George Mackey was wondering whether the Born rule was the only possible rule for calculating probabilities in a theory that represented measurements as orthonormal bases on a Hilbert space. In his investigation of mathematical foundation of quantum mechanics, Mackey had proposed the following problem [7, p.50-51][4, p.129]: Determine all measures on the closed subspaces of a Hilbert space. Gleason provided the answer to this problem in the same year [5], which is later called Gleason's theorem. Gleason succeeded in describing all σ -additive probability measures on the logical algebra of all closed subspaces of a separable Hilbert space and in showing that, except for the obvious two-dimensional counterexamples, all probability measures can be identified with normal states in the sense of von Neumann approach. Gleason's achievement confirmed von Neumann's original insight and put the calculus of Hilbert space quantum mechanics on natural physical grounds [6, p.87-88]. Gleason's theorem is of particular importance for the field of quantum logic and its attempt to find a minimal set of mathematical axioms for quantum theory.

After a considerable effort the Gleason's theorem was established in the early 90's for finitely additive vector measures on the projection lattices of von Neumann algebras. It has turned out that the lattice homomorphisms on nonabelian von Neumann algebras are σ -additive, or that finitely additive measures on projection lattices whose kernels are lattice ideal enjoy many continuity properties [6, p.3-4].

The present paper is intended to serve as an introduction to topic of the relation between Gleason's theorem and the theory of quantum logic. Most of the contents were referred to chapters 1, 2, 3 and 5 of [4] and chapters 1, 2, 3, and 9 of [6].

Chapter 1

Preliminaries

1.1 Elements of Operator Algebras

A C^* -algebra A is a Banach algebra over the field \mathbb{C} together with a map $*$: $A \rightarrow A$, $a \mapsto a^*$ which following properties:

1. $a^{**} = a$ for all $a \in A$,
2. $(a + b)^* = a^* + b^*$ for all $a, b \in A$,
3. $(ab)^* = b^*a^*$ for all $a, b \in A$,
4. $(\lambda a)^* = \bar{\lambda}a^*$ for all $\lambda \in \mathbb{C}$ and $a \in A$,
5. $\|aa^*\| = \|a\| \|a^*\|$ for all $a \in A$.

A bounded linear map $\pi : A \rightarrow B$ between C^* -algebras A and B is called a $*$ -homomorphism if

1. $\pi(ab) = \pi(a)\pi(b)$
2. $\pi(a^*) = \pi(a)^*$

for all $a, b \in A$.

Let A be a C^* -algebra. A *state* in C^* -algebra is a positive functional ρ which is $\|\rho\| = 1$. We shall denote the set of all states on A by $S(A)$ which is called the *state space* of A .

Definition 1. Let $\xi \in H$. A *vector functional* ω_ξ is defined by

$$\omega_\xi(a) = (a\xi, \xi) \quad \text{for all } a \in A.$$

Moreover, if ξ is a unit vector and A contains the identity operator, then ω_ξ is a state that is called the *vector state*.

A $*$ -representation of a C^* -algebra A on a Hilbert space H is a ring homomorphism $\pi : A \rightarrow B(H)$ such that

1. $\pi(a^*) = \pi(a)^*$ for all $a \in A$,
2. π is nondegenerate.

Definition 2 (GNS construction). Let φ be a positive functional on the C^* -algebra A . Then we can define a semidefinite inner product

$$\langle x, y \rangle = \varphi(y^* x) \quad \text{for all } x, y \in A$$

We say that φ is *faithful* if $\varphi(x^* x) = 0$ implies $x = 0$. The product is inner product if and only if φ is faithful.

Theorem 1 (Bicommutant theorem). *A $*$ -subalgebra M of $B(H)$ that contains the unit of $B(H)$ if M is a*

1.2 Quantum Logic

Chapter 2

Gleason's Theorem

2.1 Gleason's Theorem

Theorem 2. *Let H be a Hilbert space with $\dim H \geq 3$. Then any bounded completely additive measure μ on the projection lattice $P(H)$ extends uniquely to a normal functional on the algebra $B(H)$ of all bounded operators action on H .*

Bibliography

- [1] J. S. Bell. On the problem of hidden variables in quantum mechanics. *Rev. Modern Phys.*, 38:447–452, 1966.
- [2] G. Birkhoff and J. von Neumann. The logic of quantum mechanics. *Ann. of Math. (2)*, 37(4):823–843, 1936.
- [3] J. Bub. Von Neumann’s ‘no hidden variables’ proof: a re-appraisal. *Found. Phys.*, 40(9-10):1333–1340, 2010.
- [4] A. Dvurečenskij. *Gleason’s theorem and its applications*, volume 60 of *Mathematics and its Applications (East European Series)*. Kluwer Academic Publishers Group, Dordrecht; Ister Science Press, Bratislava, 1993.
- [5] A. M. Gleason. Measures on the closed subspaces of a Hilbert space. *J. Math. Mech.*, 6:885–893, 1957.
- [6] J. Hamhalter. *Quantum measure theory*, volume 134 of *Fundamental Theories of Physics*. Kluwer Academic Publishers Group, Dordrecht, 2003.
- [7] G. W. Mackey. Quantum mechanics and Hilbert space. *Amer. Math. Monthly*, 64(8):45–57, 1957.
- [8] J. von Neumann. *Mathematical Foundations of Quantum Mechanics*. Princeton University Press, Princeton, 2018.