

LATEX PRACTICE

RAUL VILLALOBOS

(6)

$$m_1 = 0.650 \text{ kg}$$

$$T_{i1} = 352 + 273 = 625 \text{ K}$$

$$m_2 = 0.650 \text{ kg}$$

$$T_{i2} = 704 + 273 = 977 \text{ K}$$

$$C_{fe} = 450.0 \frac{\text{J}}{\text{kg K}}$$

a. Find T_f in °C and K

$$Q = mC\Delta T$$

$$0 = Q_1 + Q_2 = m_1 C_{fe} (T_f - T_{i1}) + m_2 C_{fe} (T_f - T_{i2})$$

$$m_1 C_{fe} T_{i1} + m_2 C_{fe} T_{i2} = m_1 C_{fe} T_f + m_2 C_{fe} T_f$$

$$m_1 C_{fe} T_{i1} + m_2 C_{fe} T_{i2} = T_f (m_1 C_{fe} + m_2 C_{fe})$$

$$T_f = \frac{m_1 C_{fe} T_{i1} + m_2 C_{fe} T_{i2}}{m_1 C_{fe} + m_2 C_{fe}}$$

$$T_f = \frac{T_{i1} + T_{i2}}{2} = \frac{(625 + 977) \text{ K}}{2} = 801 \text{ K} = 528 \text{ °C}$$

b. Find ΔS

$$\Delta S_{tot} = \Delta S_1 + \Delta S_2 = m_1 C_{fe} \ln \left(\frac{T_f}{T_{i1}} \right) + m_2 C_{fe} \ln \left(\frac{T_f}{T_{i2}} \right)$$

$$\Delta S_{tot} = mC \left(\ln \left(\frac{T_f}{T_{i1}} \right) + \ln \left(\frac{T_f}{T_{i2}} \right) \right) = mC \ln \left(\frac{T_f}{T_{i1}} \times \frac{T_f}{T_{i2}} \right)$$

$$\Delta S_{tot} = mC \ln \left(\frac{T_f^2}{T_{i1} \times T_{i2}} \right) = (0.650 \text{ kg}) \left(450.0 \frac{\text{J}}{\text{kg K}} \right) \ln \left(\frac{(801 \text{ K})^2}{625 \text{ K} \times 977 \text{ K}} \right)$$

$$\Delta S_{tot} = 14.47 \frac{\text{J}}{\text{K}}$$

c. Find ΔS_{env}

$$\Delta S_{env} = 0 \frac{\text{J}}{\text{K}}$$

d. Find ΔS_{uni}

$$\Delta S_{uni} = \Delta S_{sys} + \Delta S_{env} = 14.47 \frac{\text{J}}{\text{K}} + 0 \frac{\text{J}}{\text{K}} = 14.47 \frac{\text{J}}{\text{K}}$$

The process is irreversible.