



Disentangling the Components of the Milky Way

Inferring the Structure of the Milky Way in Phase-Space Using Gaussian Mixture
Modelling with Extreme Deconvolution

A REPORT PRESENTED

BY

RAUNAQ RAI

Departments

Department of Physics (Cavendish Laboratory)
Institute of Astronomy

Degree

MPhil in Data Intensive Science

Supervision

Dr Anke Arentsen



St Edmund's College
University of Cambridge

29th June 2025

Contents

| | | |
|----------|---|----------|
| 1 | Introduction | 2 |
| 1.1 | Components of the Milky Way | 2 |
| 1.2 | Metallicity as a Cosmic Clock | 2 |
| 1.3 | Λ CDM : hierarchical growth and a lopsided halo | 3 |
| 1.4 | Accretion versus <i>in-situ</i> disc formation | 3 |
| 1.5 | Origins of very-metal-poor disc candidates | 4 |
| 1.6 | This Work | 4 |
| 2 | Data | 4 |
| 3 | Methodology | 4 |
| 3.1 | Extreme Deconvolution | 4 |
| 4 | Results | 5 |
| 5 | Extension direction | 5 |
| 6 | Conclusion | 5 |

1 Introduction

The Milky Way Galaxy, host to our solar system, is a spiral galaxy with a centre located approximately 150 000 trillion miles (or 25 000 lightyears) from Earth. Its formation history is complex and remains an active area of research. Being embedded within the Milky Way means we can study it in greater detail than any external galaxy, testing models of galaxy formation with high-precision observational data. One of the central aims of Galactic Archaeology is to reconstruct the Milky Way’s assembly by examining the chemical compositions and dynamical properties of its stars.

In this project, we replicate and extend the analysis of [Zhang et al. \[2024\]](#), who investigated a very metal-poor disc component in the Milky Way. Very metal poor stars, formed from an interstellar medium unpolluted by earlier generations of supernovae, are among the oldest relics in the Galaxy. Discovering them on disc-like orbits would challenge the conventional view that the disc formed later from already enriched gas [[Bland-Hawthorn and Gerhard, 2016](#)], implying instead an earlier onset of disc assembly. Using Gaia DR3, the original study applied a Gaussian Mixture Model with Extreme Deconvolution to the velocity distributions of stars across metallicity bins, probing whether a coherent disc signal persists down to the lowest metallicities.

1.1 Components of the Milky Way

The Milky Way is commonly decomposed into four stellar components: a *thin disc*, a *thick disc*, a central *bulge/bar*, and a roughly spherical *halo* [[Bland-Hawthorn and Gerhard, 2016](#), [Helmi, 2020](#)]. The thin disc dominates the census, containing $\sim 90\%$ of all stars and most of the interstellar gas. Ongoing star formation is concentrated in the “molecular–gas ring” at Galactocentric radii $R \simeq 4\text{--}8$ kpc, where young ($\lesssim 1$ Gyr), metal-rich stars trace nearly circular, co-rotating orbits with low velocity dispersion ($\sigma_\phi \simeq 20$ km s $^{-1}$). Above the mid-plane lies the thick disc: an older ($\gtrsim 8\text{--}10$ Gyr), moderately metal-poor population with $[\text{Fe}/\text{H}] \sim -0.6$ to -1.0 , a scale height of $z_{\text{scale}} \approx 1$ kpc, and hotter kinematics ($\sigma_z \simeq 40$ km s $^{-1}$) while still retaining net prograde rotation. Inside $R \lesssim 2$ kpc, the central bulge—partly bar-shaped—hosts both old, metal-rich stars and a younger, actively forming component; stellar motions there combine bar-driven streaming with high random velocities ($\sigma \sim 100$ km s $^{-1}$). Encasing all of these is the stellar halo, which contributes only a few per cent of the total stellar mass yet harbours the Galaxy’s oldest, most metal-poor stars ($[\text{Fe}/\text{H}] \lesssim -1.5$) on highly eccentric or even retrograde orbits. Its low density, rich substructure, and extreme kinematics reveal an origin in the hierarchical accretion and tidal disruption of dwarf galaxies and globular clusters. Together, the spatial distribution, chemistry, and dynamics of these four components encode the Milky Way’s star-formation history and its sequence of merger events.

1.2 Metallicity as a Cosmic Clock

Precise ages for individual old stars are notoriously difficult to measure, so their chemical composition—most commonly the iron-to-hydrogen ratio, $[\text{Fe}/\text{H}]$ —is often used as a surrogate clock. Very metal-poor (VMP) stars must have formed before successive generations of Type II and Type Ia supernovae had substantially enriched the interstellar medium, and therefore exhibit low $[\text{Fe}/\text{H}]$ values. Metallicity is inferred spectroscopically from the equivalent widths of metal absorption lines such as Fe I and the Ca II K line; after

correcting for effective temperature and surface gravity, their relative strengths yield elemental abundances to a precision of $\sim 0.1\text{--}0.2$ dex. Large surveys—including APOGEE, GALAH, LAMOST, and the Gaia XP spectra—now provide such measurements for millions of stars, enabling empirical age–metallicity relations that link chemistry to stellar chronometry [e.g. Nordström et al., 2004, Haywood et al., 2013, Leung and Bovy, 2019, Anders et al., 2023]. These studies consistently show that stars with $[\text{Fe}/\text{H}] \lesssim -1$ are typically older than ~ 10 Gyr, making low-metallicity populations valuable probes of the Milky Way’s earliest disc-building epochs.

1.3 Λ CDM: hierarchical growth and a lopsided halo

In the concordance Λ CDM cosmology, structure grows *bottom-up*: small dark-matter haloes collapse first and subsequently merge to build larger systems. Cosmological N -body simulations show that the differential sub-halo mass function follows $\text{d}n/\text{d}M \propto M^{-1.9}$, so a Milky-Way-mass halo is expected to experience $\mathcal{O}(10^2)$ low-mass mergers ($M_{\text{sub}} \lesssim 10^9 M_{\odot}$) and a handful of *massive* events ($M_{\text{sub}} \gtrsim 10^{10} M_{\odot}$) over a Hubble time [Bullock and Johnston, 2005, Cooper et al., 2010]. Yet only a tiny fraction of these haloes ever become luminous. Because star-formation efficiency declines sharply below $M_{\text{vir}} \sim 10^{11} M_{\odot}$ —a consequence of re-ionisation and stellar feedback—the galaxy stellar-mass–halo-mass (SMHM) relation is steep at the low-mass end [Purcell et al., 2007, Bullock and Boylan-Kolchin, 2017]. As a result, *one or two* relatively massive dwarfs deposit the bulk of the stellar material in the halo, while hundreds of low-mass sub-haloes remain dark.

Once accreted, dynamical friction drags the most massive satellites deep into the Galactic potential, their orbits radialise, and their debris is dispersed throughout the *inner* halo. The disrupted stars inherit coherent signatures—high radial anisotropy, distinctive angular momenta, and chemically narrow sequences—that survive to the present [e.g. Helmi and Tim de Zeeuw, 2000]. Consequently, the stellar halo is not a smooth spheroid but a palimpsest of the Galaxy’s merger history, with the inner halo overwhelmingly shaped by a few dominant progenitors (e.g. Gaia–Sausage/Enceladus), and the outer halo supplied by many low-mass accretions.

1.4 Accretion versus *in-situ* disc formation

Chemical and kinematic evidence confirms that the metal-poor halo is primarily accreted. The debris of the Gaia–Sausage/Enceladus (GSE) event, for instance, is traced by stars with $-2 < [\text{Fe}/\text{H}] < -1$ and extreme orbital anisotropy ($\beta \gtrsim 0.8$; Belokurov et al. 2018, Helmi et al. 2018). At $[\text{Fe}/\text{H}] \lesssim -2$ an even broader mix of minor mergers emerges, erasing any global rotation signal [Lancaster et al., 2019, Bird et al., 2021].

Against this backdrop, a number of studies have uncovered stars in the range $-2 < [\text{Fe}/\text{H}] < -1$ whose velocities resemble a *disc*: modest eccentricities and net prograde rotation [Norris et al., 1985, Chiba and Beers, 2000, Carollo et al., 2019, An and Beers, 2020]. Gaia has pushed this frontier to $[\text{Fe}/\text{H}] < -2$ [Sestito et al., 2019, Venn et al., 2020, Cordoni et al., 2020, Mardini et al., 2022]. Whether these objects represent (i) an *in-situ* metal-poor disc or (ii) the spun-up debris of earlier mergers remains hotly debated.

1.5 Origins of very-metal-poor disc candidates

Three broad formation scenarios have been proposed:

1. **Early *in-situ* disc.** Stars form in a nascent, gas-rich disc before $z \sim 4$, and later migrate outward or are dynamically heated; such stars would share the chemistry of the proto-Galaxy.
2. **Proto-galactic building blocks.** VMP stars originate in several massive, gas-rich satellites accreted at high redshift; their debris is dragged into the disc plane as the gaseous disc settles [e.g. [Sestito et al., 2020](#)].
3. **Late, minor prograde mergers.** Low-mass satellites on aligned orbits are assimilated after the disc forms, depositing a thin layer of metal-poor stars that retain disc-like kinematics [[Santistevan et al., 2021](#)].

State-of-the-art cosmological simulations generally reproduce scenario 2, finding that early mergers dominate the VMP budget while a coherent disc does not appear until $z \lesssim 2$ [[Gurvich et al., 2023](#)].

Observationally, [Belokurov and Kravtsov \[2022\]](#) identified *Aurora*—a kinematically hot, weakly rotating population with $-2 \lesssim [\text{Fe}/\text{H}] \lesssim -1.3$ —arguing against an extremely early disc. Follow-up work shows *Aurora* to be centrally concentrated [[Rix et al., 2022](#), [Arentsen et al., 2020a,b](#)], consistent with heated debris rather than a long-lived thin disc. Furthermore, secular processes such as bar–halo resonances can impart a modest prograde bias to halo stars, mimicking a disc signal [[Dillamore et al., 2023](#)].

Unravelling these possibilities demands six-dimensional phase-space information and precision abundances—the focus of the present study.

1.6 This Work

We revisit the existence of a VMP disc using Gaia DR3, combining 6D phase-space and spectroscopic $[\text{Fe}/\text{H}]$ and $[\alpha/\text{Fe}]$. By fitting Gaussian Mixture Models with Extreme Deconvolution to 3D velocity distributions in narrow metallicity bins, we test for a rotating disc component at $[\text{Fe}/\text{H}] < -2.0$ and explore its implications for the timing and mechanisms of disc formation.

2 Data

3 Methodology

Details of your methodology.

3.1 Extreme Deconvolution

Discussion of XD.

4 Results

What you found.

5 Extension direction

6 Conclusion

Summary of your findings.

References

- D. An and T. C. Beers. A blueprint for the milky way’s stellar populations: The power of large photometric and astrometric surveys. *The Astrophysical Journal*, 897(1):39, July 2020. doi: 10.3847/1538-4357/ab8d39. URL <https://ui.adsabs.harvard.edu/abs/2020ApJ...897...39A>.
- F. Anders, P. Gispert, B. Ratcliffe, C. Chiappini, I. Minchev, S. Nepal, A. B. A. Queiroz, J. A. S. Amarante, T. Antoja, G. Casali, L. Casamiquela, A. Khalatyan, A. Miglio, H. Perottoni, and M. Schultheis. Spectroscopic age estimates for apogee red-giant stars: Precise spatial and kinematic trends with age in the galactic disc. *Astronomy and astrophysics*, 678:A158, Oct. 2023. ISSN 1432-0746. doi: 10.1051/0004-6361/202346666. URL <http://dx.doi.org/10.1051/0004-6361/202346666>.
- A. Arentsen, E. Starkenburg, N. F. Martin, D. S. Aguado, D. B. Zucker, C. Allende Prieto, V. Hill, K. A. Venn, R. G. Carlberg, J. I. González Hernández, L. I. Mashonkina, J. F. Navarro, R. Sánchez-Janssen, M. Schultheis, G. F. Thomas, K. Youakim, G. F. Lewis, J. D. Simpson, Z. Wan, R. E. Cohen, D. Geisler, and J. E. O’Connell. The pristine inner galaxy survey (pigs) ii: Uncovering the most metal-poor populations in the inner milky way. *Monthly Notices of the Royal Astronomical Society*, 496(4): 4964–4978, July 2020a. ISSN 1365-2966. doi: 10.1093/mnras/staa1661. URL <http://dx.doi.org/10.1093/mnras/staa1661>.
- A. Arentsen, E. Starkenburg, N. F. Martin, V. Hill, R. Ibata, A. Kunder, M. Schultheis, K. A. Venn, D. B. Zucker, D. Aguado, R. Carlberg, J. I. González Hernández, C. Lardo, N. Longeard, K. Malhan, J. F. Navarro, R. Sánchez-Janssen, F. Sestito, G. Thomas, K. Youakim, G. F. Lewis, J. D. Simpson, and Z. Wan. The pristine inner galaxy survey (pigs) i: Tracing the kinematics of metal-poor stars in the galactic bulge. *Monthly Notices of the Royal Astronomical Society*, 491(1):L11–L16, January 2020b. doi: 10.1093/mnrasl/slz156. URL <https://ui.adsabs.harvard.edu/abs/2020MNRAS.491L..11A>.
- V. Belokurov and A. Kravtsov. From dawn till disc: Milky way’s turbulent youth revealed by the apogee+gaia data. *Monthly Notices of the Royal Astronomical Society*, 514(1): 689–714, May 2022. ISSN 1365-2966. doi: 10.1093/mnras/stac1267. URL <http://dx.doi.org/10.1093/mnras/stac1267>.
- V. Belokurov, D. Erkal, N. W. Evans, S. E. Koposov, and A. J. Deason. Co-formation of the disc and the stellar halo. *Monthly Notices of the Royal Astronomical Society*,

- 478(1):611–619, June 2018. ISSN 1365-2966. doi: 10.1093/mnras/sty982. URL <http://dx.doi.org/10.1093/mnras/sty982>.
- S. A. Bird, X.-X. Xue, C. Liu, J. Shen, C. Flynn, C. Yang, G. Zhao, and H.-J. Tian. Constraints on the assembly history of the milky way’s smooth, diffuse stellar halo from the metallicity-dependent, radially dominated velocity anisotropy profiles probed with k giants and bhb stars using lamost, sdss/segue, and gaia. *The Astrophysical Journal*, 919(1):66, Sept. 2021. ISSN 1538-4357. doi: 10.3847/1538-4357/abfa9e. URL <http://dx.doi.org/10.3847/1538-4357/abfa9e>.
- J. Bland-Hawthorn and O. Gerhard. The galaxy in context: Structural, kinematic, and integrated properties. *Annual Review of Astronomy and Astrophysics*, 54(1):529–596, Sept. 2016. ISSN 1545-4282. doi: 10.1146/annurev-astro-081915-023441. URL <http://dx.doi.org/10.1146/annurev-astro-081915-023441>.
- J. S. Bullock and M. Boylan-Kolchin. Small-scale challenges to the lambda cdm paradigm. *Annual Review of Astronomy and Astrophysics*, 55(1):343–387, Aug. 2017. ISSN 1545-4282. doi: 10.1146/annurev-astro-091916-055313. URL <http://dx.doi.org/10.1146/annurev-astro-091916-055313>.
- J. S. Bullock and K. V. Johnston. Tracing galaxy formation with stellar halos. i. methods. *The Astrophysical Journal*, 635(2):931–949, Dec. 2005. ISSN 1538-4357. doi: 10.1086/497422. URL <http://dx.doi.org/10.1086/497422>.
- D. Carollo, M. Chiba, M. Ishigaki, K. Freeman, T. C. Beers, Y. S. Lee, P. Tissera, C. Battistini, and F. Primas. Evidence for the third stellar population in the milky way’s disk. *The Astrophysical Journal*, 887(1):22, December 2019. doi: 10.3847/1538-4357/ab517c. URL <https://ui.adsabs.harvard.edu/abs/2019ApJ...887...22C>.
- M. Chiba and T. C. Beers. Kinematics of metal-poor stars in the galaxy. iii. formation of the stellar halo and thick disk as revealed from a large sample of nonkinematically selected stars. *The Astronomical Journal*, 119(6):2843–2865, June 2000. ISSN 0004-6256. doi: 10.1086/301409. URL <http://dx.doi.org/10.1086/301409>.
- A. P. Cooper, S. Cole, C. S. Frenk, S. D. M. White, J. Helly, A. J. Benson, G. De Lucia, A. Helmi, A. Jenkins, J. F. Navarro, V. Springel, and J. Wang. Galactic stellar haloes in the cdm model: Galactic stellar haloes in the cdm model. *Monthly Notices of the Royal Astronomical Society*, 406(2):744–766, May 2010. ISSN 0035-8711. doi: 10.1111/j.1365-2966.2010.16740.x. URL <http://dx.doi.org/10.1111/j.1365-2966.2010.16740.x>.
- G. Cordoni, G. S. Da Costa, D. Yong, A. D. Mackey, A. F. Marino, S. Monty, T. Nordlander, J. E. Norris, M. Asplund, M. S. Bessell, A. R. Casey, A. Frebel, K. Lind, S. J. Murphy, B. P. Schmidt, X. D. Gao, T. Xylakis-Dornbusch, A. M. Amarsi, and A. P. Milone. Exploring the galaxy’s halo and very metal-weak thick disc with skymapper and gaia dr2. *Monthly Notices of the Royal Astronomical Society*, 503(2):2539–2561, Nov. 2020. ISSN 1365-2966. doi: 10.1093/mnras/staa3417. URL <http://dx.doi.org/10.1093/mnras/staa3417>.
- A. M. Dillamore, V. Belokurov, N. W. Evans, and E. Y. Davies. Stellar halo substructure generated by bar resonances. *Monthly Notices of the Royal Astronomical Society*, 524

- (3):3596–3608, July 2023. ISSN 1365-2966. doi: 10.1093/mnras/stad2136. URL <http://dx.doi.org/10.1093/mnras/stad2136>.
- A. B. Gurvich, J. Stern, C.-A. Faucher-Giguère, P. F. Hopkins, A. Wetzel, J. Moreno, C. C. Hayward, A. J. Richings, and Z. Hafen. Rapid disc settling and the transition from bursty to steady star formation in milky way-mass galaxies. *Monthly Notices of the Royal Astronomical Society*, 519(2):2598–2614, February 2023. doi: 10.1093/mnras/stac3712. URL <https://ui.adsabs.harvard.edu/abs/2023MNRAS.519.2598G>.
- M. Haywood, P. Di Matteo, M. D. Lehnert, D. Katz, and A. Gómez. The age structure of stellar populations in the solar vicinity: Clues of a two-phase formation history of the milky way disk. *Astronomy and Astrophysics*, 560:A109, Dec. 2013. ISSN 1432-0746. doi: 10.1051/0004-6361/201321397. URL <http://dx.doi.org/10.1051/0004-6361/201321397>.
- A. Helmi. Streams, substructures, and the early history of the milky way. *Annual Review of Astronomy and Astrophysics*, 58(1):205–256, Aug. 2020. ISSN 1545-4282. doi: 10.1146/annurev-astro-032620-021917. URL <http://dx.doi.org/10.1146/annurev-astro-032620-021917>.
- A. Helmi and P. Tim de Zeeuw. Mapping the substructure in the galactic halo with the next generation of astrometric satellites. *Monthly Notices of the Royal Astronomical Society*, 319(3):657–665, 2000.
- A. Helmi, C. Babusiaux, H. H. Koppelman, D. Massari, J. Veljanoski, and A. G. A. Brown. The merger that led to the formation of the milky way’s inner stellar halo and thick disk. *Nature*, 563(7729):85–88, Oct. 2018. ISSN 1476-4687. doi: 10.1038/s41586-018-0625-x. URL <http://dx.doi.org/10.1038/s41586-018-0625-x>.
- L. Lancaster, S. E. Koposov, V. Belokurov, N. W. Evans, and A. J. Deason. The halo’s ancient metal-rich progenitor revealed with bhb stars. *Monthly Notices of the Royal Astronomical Society*, 486(1):378–389, Mar. 2019. ISSN 1365-2966. doi: 10.1093/mnras/stz853. URL <http://dx.doi.org/10.1093/mnras/stz853>.
- H. W. Leung and J. Bovy. Deep learning of multi-element abundances from high-resolution spectroscopic data. *Monthly Notices of the Royal Astronomical Society*, 483(3):3255–3277, 2019.
- M. K. Mardini, A. Frebel, A. Chiti, Y. Meiron, K. V. Brauer, and X. Ou. The atari disk, a metal-poor stellar population in the disk system of the milky way. *The Astrophysical Journal*, 936(1):78, Sept. 2022. ISSN 1538-4357. doi: 10.3847/1538-4357/ac8102. URL <http://dx.doi.org/10.3847/1538-4357/ac8102>.
- B. Nordström, M. Mayor, J. Andersen, J. Holmberg, F. Pont, B. R. Jørgensen, E. H. Olsen, S. Udry, and N. Mowlavi. The geneva-copenhagen survey of the solar neighbourhood: Ages, metallicities, and kinematic properties of $\sim 14\,000$ f and g dwarfs. *Astronomy and Astrophysics*, 418(3):989–1019, Apr. 2004. ISSN 1432-0746. doi: 10.1051/0004-6361:20035959. URL <http://dx.doi.org/10.1051/0004-6361:20035959>.
- J. Norris, M. S. Bessell, and A. J. Pickles. Population studies. i. the bidelman-macconnell “weak-metal” stars. *The Astrophysical Journal Supplement Series*, 58:463–492, July

1985. doi: 10.1086/191049. URL <https://ui.adsabs.harvard.edu/abs/1985ApJS...58..463N>.
- C. W. Purcell, J. S. Bullock, and A. R. Zentner. Shredded galaxies as the source of diffuse intrahalo light on varying scales. *The Astrophysical Journal*, 666(1):20–33, Sept. 2007. ISSN 1538-4357. doi: 10.1086/519787. URL <http://dx.doi.org/10.1086/519787>.
- H.-W. Rix, V. Chandra, R. Andrae, A. M. Price-Whelan, D. H. Weinberg, C. Conroy, M. Fouesneau, D. W. Hogg, F. De Angeli, R. P. Naidu, M. Xiang, and D. Ruz-Mieres. The poor old heart of the milky way. *The Astrophysical Journal*, 941(1):45, Dec. 2022. ISSN 1538-4357. doi: 10.3847/1538-4357/ac9e01. URL <http://dx.doi.org/10.3847/1538-4357/ac9e01>.
- I. B. Santistevan, A. Wetzel, R. E. Sanderson, K. El-Badry, J. Samuel, and C.-A. Faucher-Giguère. The origin of metal-poor stars on prograde disc orbits in fire simulations of milky way-mass galaxies. *Monthly Notices of the Royal Astronomical Society*, 505(1):921–938, July 2021. doi: 10.1093/mnras/stab1345. URL <https://ui.adsabs.harvard.edu/abs/2021MNRAS.505..921S>.
- F. Sestito, N. Martin, and E. Starkenburg. Tracing the formation of the milky way through ultra metal-poor stars. In *The Gaia Universe*, page 47, Apr. 2019. doi: 10.5281/zenodo.3236051.
- F. Sestito, T. Buck, E. Starkenburg, N. F. Martin, J. F. Navarro, K. A. Venn, A. Obreja, P. Jablonka, and A. V. Macciò. Exploring the origin of low-metallicity stars in milky-way-like galaxies with the nihao-uhd simulations. *Monthly Notices of the Royal Astronomical Society*, 500(3):3750–3762, Nov. 2020. ISSN 1365-2966. doi: 10.1093/mnras/staa3479. URL <http://dx.doi.org/10.1093/mnras/staa3479>.
- K. A. Venn, C. L. KIELTY, F. Sestito, E. Starkenburg, N. Martin, D. S. Aguado, A. Arentsen, P. Bonifacio, E. Caffau, V. Hill, P. Jablonka, C. Lardo, L. Mashonkina, J. F. Navarro, C. Sneden, G. Thomas, K. Youakim, J. I. González-Hernández, R. S. Janssen, R. Carlberg, and K. Malhan. The pristine survey - ix. cfht espadons spectroscopic analysis of 115 bright metal-poor candidate stars. *Monthly Notices of the Royal Astronomical Society*, 492(3):3241–3262, Mar. 2020. doi: 10.1093/mnras/stz3546.
- H. Zhang, A. Arden-Arentsen, and V. Belokurov. On the existence of a very metal-poor disc in the milky way, 2024. URL <https://arxiv.org/abs/2311.09294>.