

# DISENTANGLING THE COMPONENTS OF THE MILKY WAY

INFERRING THE STRUCTURE OF THE MILKY WAY IN PHASE-SPACE USING GAUSSIAN MIXTURE MODELLING WITH EXTREME DECONVOLUTION

Raunaq Singh Rai — MPhil Data Intensive Science — University of Cambridge

## Motivation and Scientific Justification

A central question in Galactic Archaeology is *when* the Milky Way’s disc first settled. Standard models suggest that the disc formed relatively late, after the interstellar medium had been enriched by multiple generations of stars. This implies a lack of stars with disc-like kinematics at very low metallicities ( $[\text{Fe}/\text{H}] \lesssim -1.5$ ). Since metallicity generally increases over cosmic time, it serves as a rough proxy for stellar age—lower-metallicity stars are typically older, having formed when the Galaxy was less chemically evolved. Thus, identifying a coherent, metal-poor disc population would challenge the “late-disc” paradigm and force a reassessment of the balance between in-situ star formation and accreted growth.

Zhang et al. (2024) [1] addressed this question by applying Extreme-Deconvolution Gaussian Mixture Modelling (XD-GMM) to Gaia DR3 red giants, and found no evidence for a cold, rotation-supported disc at very low metallicity. However, their analysis did not account for  $\alpha$ -abundance, which can be used as tracer of star formation timescales.

We reproduce their metallicity-binned XD-GMM using the same bright RGB catalogue, and extend the work by dividing stars into high and low  $\alpha$  sequences following Viswanathan et al. (2024) [2]. Within each  $\alpha$  branch, we perform XD-GMM fits in successive metallicity bins, selecting the optimal number of components via the Bayesian Information Criterion. By inspecting the weights and kinematic properties of these components, we assess whether disc-like signatures at low metallicity are genuine or attributable to contamination, noise, or accreted substructure.

## Methodology

We use a cleaned sample of red giant branch (RGB) stars from Gaia DR3, with metallicities and  $\alpha$ -abundances from Andrae et al. Catalogue [3] and the Li et al. Catalogue [4], and distances from the Bailer-Jones et al. Catalogue [5]. As analysis largely depends on the accuracies of metallicities regions of high extinction, where XP spectra is known to be bias, are excluded with the sacrifice of losing a large proportion of RGB stars from subsequent analysis..

The velocity distribution ( $v_R, v_\phi, v_z$ ) is modelled using Extreme-Deconvolution Gaussian Mixture Modelling (XD-GMM) [6][7], which accounts for observational uncertainties. We bin stars by metallicity and use the Bayesian Information Criterion (BIC) to determine the

number of Gaussian components per bin. This allows us to identify structure without over fitting and introducing too many gaussians.

We extend the original method by splitting the sample into high- and low- $\alpha$  sequences [2], fitting separate XD-GMMs to each. By inspecting the component weights and kinematic properties, we quantify the emergence of rotational support and assess the presence of disc-like populations across both chemical tracks.

## Key Findings




Figure 1: XD-GMM decomposition of red giant kinematics across metallicity bins.

Below  $[\text{M}/\text{H}] \sim -2$ , no disc-like component is detected. The kinematics are fully explained by two broad halo Gaussians: one stationary and one mildly prograde likely associated with the Aurora population [8]. In the intermediate metallicity bin, two additional highly radial, non-rotating components emerge, linked to the Gaia-Sausage/Enceladus (GS/E) merger [9]. These represent remnants of a major head-on collision between the Milky Way and a massive dwarf galaxy, which deposited stars on radial orbits and contributed significantly to the inner halo’s anisotropic structure.

Disc-like rotation only emerges above  $[\text{M}/\text{H}] \sim -1.6$ : a distinct prograde component with relatively low dispersion, consistent with an early thick disc is observable. This component strengthens with increasing metallicity, supporting a gradual build-up of ordered rotation as the Galaxy chemically enriched.

As shown in Figure 2, stars in the VMP and IMP bins have negligible rotational support, with  $v_\phi/\sigma_\phi < 1$ . It is generally accepted that a disc-like population should have  $v_\phi/\sigma_\phi \gtrsim 3$ , confirming the absence of a cold disc for  $[\text{M}/\text{H}] < -1.6$ . From Figure 2, we can conclude that the onset of disc-like rotation occurs above  $[\text{M}/\text{H}] \sim -1.6$ .




Figure 2: Rotational support ( $v_\phi/\sigma_\phi$ ) for each Gaussian component across metallicity bins. The size of the points indicates the weight of each component.

Following the methodology proposed by Viswanathan et al. [2], we separated the sample into high- and low- $\alpha$  sequences. This  $\alpha$ -abundance reflects star formation timescales: high- $\alpha$  stars typically formed rapidly in intense, early starbursts, and are thus older, while low- $\alpha$  stars formed more gradually over extended periods, often in lower-density environments.

Figure 3 shows how these two populations differ in their kinematic evolution. In the high- $\alpha$  branch, we observe a smooth increase in mean azimuthal velocity and a concurrent decrease in dispersion with increasing metallicity, indicating a gradual spin-up from a dynamically hot halo to a thick disc. In contrast, the low- $\alpha$  population undergoes a sharper transition: rotation and cooling only emerge above  $[M/H] \sim -1.3$ , consistent with a younger, thin-disc origin.




Figure 3: Median  $v_\phi$  and dispersion vs. metallicity, split by  $\alpha$ -sequence.

We repeated the GMM decomposition separately on each  $\alpha$  sequence (Figure 4). In the high- $\alpha$  population, a disc-like component emerges as early as  $[M/H] \sim -1.6$ , while in the low- $\alpha$  branch, such a component is only detected above  $[M/H] \sim -1.3$ . This supports the interpretation that the high- $\alpha$  thick disc formed earlier and more gradually, whereas the low- $\alpha$  thin disc emerged later in the Galaxy’s evolution.




Figure 4: XD-GMM decomposition across  $\alpha$ -sequences and metallicity bins. Top: high  $\alpha$ . Bottom: low  $\alpha$ .

Interestingly, we detect GS/E-like radial components even in the high- $\alpha$  regime. Since GS/E stars are expected to be  $\alpha$ -poor due to their extended star formation histories, their presence here likely reflects misclassification at low signal-to-noise rather than genuinely  $\alpha$ -rich accreted progenitors. This highlights the importance of reliable chemical separation in metal-poor regimes, where contamination can obscure interpretations of Galactic structure.

## References

- [1] Hanyuan Zhang, Anke Arden-Arentsen, and Vasily Belokurov. On the existence of a very metal-poor disc in the milky way, 2024.
- [2] Akshara Viswanathan, Danny Horta, Adrian M. Price-Whelan, and Else Starkenburg. A slow spin to win – the gradual evolution of the proto-galaxy to the old disc, 2024.
- [3] René Andrae, Hans-Walter Rix, and Vedant Chandra. Robust data-driven metallicities for 175 million stars from gaia xp spectra. *The Astrophysical Journal Supplement Series*, 267(1):8, jul 2023.
- [4] Jiadong Li, Kaze W. K. Wong, David W. Hogg, Hans-Walter Rix, and Vedant Chandra. Aspgap: Augmented stellar parameters and abundances for 23 million rgb stars from gaia xp low-resolution spectra, 2023.
- [5] C. A. L. Bailer-Jones, J. Rybizki, M. Fouesneau, M. Demleitner, and R. Andrae. VizieR Online Data Catalog: Distances to 1.47 billion stars in Gaia EDR3 (Bailer-Jones+, 2021). VizieR On-

- line Data Catalog: I/352. Originally published in: 2021AJ....161..147B, February 2021.
- [6] Jo Bovy, David W. Hogg, and Sam T. Roweis. Extreme deconvolution: Inferring complete distribution functions from noisy, heterogeneous and incomplete observations. *The Annals of Applied Statistics*, 5(2B), June 2011.
  - [7] Peter Melchior and Andy Goulding. Filling the gaps: Gaussian mixture models from noisy, truncated or incomplete samples. *Astronomy and Computing*, 25, 11 2016.
  - [8] Vasily Belokurov and Andrey Kravtsov. From dawn till disc: Milky way’s turbulent youth revealed by the apogee+gaia data. *Monthly Notices of the Royal Astronomical Society*, 514(1):689–714, May 2022.
  - [9] V Belokurov, D Erkal, N W Evans, S E Koposov, and A J Deason. Co-formation of the disc and the stellar halo. *Monthly Notices of the Royal Astronomical Society*, 478(1):611–619, June 2018.