

\$1.60

*linear
integral
equations*
*william
vernon
lovitt*

LINEAR INTEGRAL EQUATIONS by W. V. Lovitt

This unique book is the only English language introduction to linear integral equations. It considers both the general theory of linear equations with a single integration and their application to differential equations, calculus of variations, and special areas in mathematical physics. It develops in detail the Fredholm and Hilbert-Schmidt theories, and applies them to the solution of Dirichlet's problem and Neumann's problem, and similar topics. Applications are extensive, treatment is extremely clear, and demonstrations are full and detailed.

PARTIAL CONTENTS. I. Linear integral equations, linear differential equations. II. Solution of integral equations of second kind by successive substitutions. III. Solution of Fredholm's equation expressed as a ratio of two integral series in lambda. IV. Applications of Fredholm's theory. Free vibrations of an elastic string. Constrained vibrations of an elastic string. Auxiliary theorems on harmonic functions. Logarithmic potential of a double layer. Fredholm's solution of Dirichlet's problem, logarithmic potential of a simple layer, Fredholm's solution of Neumann's problem. V. Hilbert-Schmidt theory of integral equations with symmetric kernels. VI. Applications of the Hilbert-Schmidt theory. Boundary problems for ordinary linear differential equations, problems of the calculus of variations, vibration problems, flow of heat in a bar.

"The only textbook on linear integral equations in the English language . . . conception and exposition are clear and easily understandable and include all fundamental questions on the subject. . . . One of its merits is that nearly half of the volume is devoted to applications . . . an excellent introduction, can be warmly recommended," **JOURNAL OF THE ROYAL NAVAL SCIENTIFIC SERVICE.** Complete, unabridged. Index. 27 figures. ix + 253pp. 5 $\frac{3}{8}$ x 8.

S176 Paperbound \$1.60

THIS DOVER EDITION IS DESIGNED FOR YEARS OF USE

THE PAPER is chemically the same quality as you would find in books priced \$5.00 or more. It does not discolor or become brittle with age. Not artificially bulked, either; this edition is an unabridged full-length book, but is still easy to handle.

THE BINDING: The pages in this book are SEWN in signatures, in the method traditionally used for the best books. These books open flat for easy reading and reference. Pages do not drop out, the binding does not crack and split (as in the case with many paperbooks held together with glue).

THE TYPE IS LEGIBLE: Margins are ample and allow for cloth rebinding.

LINEAR

INTEGRAL EQUATIONS

BY

WILLIAM VERNON LOVITT, PH.D.

Professor of Mathematics, Colorado College

FIRST EDITION

1950

DOVER PUBLICATIONS INC.
NEW YORK

**COPYRIGHT, 1924, BY THE
McGRAW-HILL BOOK COMPANY, INC.**

DOVER REPRINTED EDITION

1950

PRINTED IN THE UNITED STATES OF AMERICA

PREFACE

For many years the subject of functional equations has held a prominent place in the attention of mathematicians. In more recent years this attention has been directed to a particular kind of functional equation, an integral equation, wherein the unknown function occurs under the integral sign. The study of this kind of equation is sometimes referred to as the inversion of a definite integral.

In the present volume I have tried to present in a readable and systematic manner the general theory of linear integral equations with some of its applications. The applications given are to differential equations, calculus of variations, and some problems in mathematical physics. The applications to mathematical physics herein given are to Neumann's problem, Dirichlet's problem, and certain vibration problems which lead to differential equations with boundary conditions. The attempt has been made to present the subject matter in such a way as to make the volume available as a text on this subject in Colleges and Universities.

The reader who so desires can omit the chapters on the applications. The remaining chapters on the general theory are an entity in themselves.

The discussion has been confined to those equations which are linear and in which a single integration occurs. The limits of the present volume forbid any adequate treatment of integral equations in several independent variables; systems of integral equations; integral equations of higher order; integro-differential equations; singular integral equations; integral equations with special or discontinuous kernels.

I desire here to express my thanks to Prof. Oscar Bolza (now of Freiburg University, formerly of the University of Chicago) for his permission to make use of my notes on his lectures on integral equations delivered during the

summer of 1913 at the University of Chicago. A bound volume of these notes in my handwriting has resided these past ten years on the shelves of the University of Chicago mathematical library and has been available to many students during this time. A number of copies of these notes are in circulation in this country at present.

The following books have been available and have been found to be of value in the preparation of this volume:

M. BOCHER: An Introduction to the Study of Integral Equations. No. 10, Cambridge Tracts, 1909. University Press. Cambridge.

É. GOURSAT: Cours D'Analyse Mathématique. Tome III. Chaps. 30, 31, 32, 33. Paris, Gauthier-Villars, 1923.

HEYWOOD-FRÉCHET: L'Équation de Fredholm et ses applications à la Physique Mathématique. Paris, Hermann et Fils, 1912.

KNESER: Die Integralgleichungen und ihre Anwendungen in der Math. Physik. Braunschweig. Vieweg et Sohn, 1922.

G. KOWALEWSKI: Einführung in die Determinanten Theorie. 18 and 19 cap. (p. 455–505). Verlag von Veit et C., 1909.

LALESKO: Introduction à la Théorie des Équations Intégrales. Paris. Hermann et Fils, 1912.

Volterra: Leçons sur les Équations Intégrales et les Équations Intégro-Differentielles. Paris, Gauthier-Villars, 1913.

For those who desire a bibliography on this subject we refer the reader to a short bibliography in the work by Heywood-Fréchet and to a more extensive bibliography in the work by Lalesko.

W. V. LOVITT.

COLORADO SPRINGS, COLO., June, 1924.

CONTENTS

CHAPTER I INTRODUCTORY

	PAGE
PREFACE	III
ART.	
1. Linear Integral Equation of the First Kind	1
2. Abel's Problem	2
3. Linear Integral Equation of the Second Kind	3
4. Relation between Linear Differential Equation and Volterra's Integral Equation	4
5. Non-linear Equations	5
6. Singular Equations	6
7. Types of Solutions	6
Exercises.	

CHAPTER II

SOLUTION OF INTEGRAL EQUATION OF SECOND KIND BY SUCCESSIVE SUBSTITUTIONS

8. Solution by Successive Substitutions	9
9. Volterra's Equation	13
10. Successive Approximations	15
11. Iterated Functions	17
12. Reciprocal Functions	17
13. Volterra's Solution of Fredholm's Equation	19
14. Discontinuous Solutions	20
Exercises.	

CHAPTER III

SOLUTION OF FREDHOLM'S EQUATION EXPRESSED AS RATIO OF TWO INTEGRAL SERIES IN λ

15. Fredholm's Equation as Limit of a Finite System of Linear Equations	23
16. Hadamard's Theorem	28

ART.		PAGE
17. Convergence Proof		32
18. Fredholm's Two Fundamental Relations		34
19. Fredholm's Solution of the Integral Equation when $D(\lambda) \neq 0$		39
20. Solution of the Homogeneous Equation when $D(\lambda) = 0$, $D'(\lambda) \neq 0$		42
21. Solution of the Homogeneous Integral Equation when $D(\lambda) = 0$		46
22. Characteristic Constant. Fundamental Functions		56
23. The Associated Homogeneous Integral Equation		56
24. The Non-homogeneous Integral Equation when $D(\lambda) = 0$		62
25. Kernels of the Form $\sum a_i(x)b_i(y)$		68
Exercises.		

CHAPTER IV

APPLICATIONS OF THE FREDHOLM THEORY

I. Free Vibrations of an Elastic String

26. The Differential Equations of the Problem	73
27. Reduction to a One-dimensional Boundary Problem	74
28. Solution of the Boundary Problem	74
29. Construction of Green's Function	76
30. Equivalence between the Boundary Problem and a Linear Integral Equation	78

II. Constrained Vibrations of an Elastic String

31. The Differential Equations of the Problem	81
32. Equivalence between the Boundary Problem and a Linear Integral Equation	82
33. Remarks on Solution of the Boundary Problem	83

III. Auxiliary Theorems on Harmonic Functions

34. Harmonic Functions	85
35. Definitions about Curves	85
36. Green's Theorem	86
37. The Analogue of Theorem IX for the Exterior Region	89
38. Generalization of the Preceding	90

IV. Logarithmic Potential of a Double Layer

39. Definition	92
40. Properties of $w(x, y)$ at Points not on C	93

CONTENTS

VII

ART.		PAGE
41. Behavior of $w(x, y)$ on C		95
42. Behavior of $\frac{\partial w}{\partial n}$ on the Boundary C and at Infinity.		98
43. Case where w_i or w_s Vanish along C		99

V. Fredholm's Solution of Dirichlet's Problem

44. Dirichlet's Problem		100
45. Reduction to an Integral Equation.		101
46. Solution of the Integral Equation		104
47. Index of $\lambda = 1$ for $K(s_0, s)$		105

VI. Logarithmic Potential of a Simple Layer

48. Definition.		107
49. Properties of $v(x, y)$		107

VII. Fredholm's Solution of Neumann's Problem

50. Neumann's Problem		110
51. Reduction to an Integral Equation.		110
52. Solution of the Integral Equation		112

CHAPTER V

HILBERT-SCHMIDT THEORY OF INTEGRAL EQUATIONS WITH
SYMMETRIC KERNELSSOLUTION EXPRESSED IN TERMS OF A SET OF FUNDAMENTAL
FUNCTIONS

I. Existence of at Least One Characteristic Constant

53. Introductory Remarks		116
54. Power Series for $\frac{D'(\lambda)}{D(\lambda)}$		118
55. Plan of Kneser's Proof		121
56. Lemmas on Iterations of a Symmetric Kernel		123
57. Schwarz's Inequality.		125
58. Application of Schwarz's Inequality		126

II. Orthogonality

59. Orthogonality Theorem.		128
60. Reality of the Characteristic Constants.		129

	PAGE
ART.	
61. Complete Normalized Orthogonal System of Fundamental Functions	131
III. Expansion of an Arbitrary Function According to the Fundamental Functions of a Complete Normalized Orthogonal System	
62. (a) Problem of the Vibrating String Resumed	138
(b) Determination of the Coefficients in the General Problem	140
IV. Expansion of the Kernel According to the Fundamental Functions of a Complete Normalized Orthogonal System	
63. (a) Determination of the Coefficients	141
(b) The Bilinear Formula for the Case of a Finite Number of Fundamental Functions	142
(c) The Bilinear Formula for Kernels Having an Infinite Number of Characteristic Constants.	144
64. The Complete Normalized Orthogonal System for the Iterated Kernel $K_n(x, t)$	146

V. Auxiliary Theorems

65. Bessel's Inequality.	149
66. Proof of the Bilinear Formula for the Iterated Kernel $K_n(x, t)$ for $n \geq 4$	152
67. An Auxiliary Theorem of Schmidt.	155
68. VI. Expansion of an Arbitrary Function According to the Complete Normalized Orthogonal System of Fundamental Functions of a Symmetric Kernel.	158

VII. Solution of the Integral Equation

69. Schmidt's Solution of the Non-homogeneous Integral Equation when λ is not a Characteristic Constant.	162
70. Schmidt's Solution of the Non-homogeneous Integral Equation when λ is a Characteristic Constant.	166
71. Remarks on Obtaining a Solution	168
Exercises.	

CHAPTER VI

APPLICATIONS OF THE HILBERT-SCHMIDT THEORY

I. Boundary Problems for Ordinary Linear Differential Equations	
72. Introductory Remarks	170
73. Construction of Green's Function	173

ART.		PAGE
74.	Equivalence between the Boundary Problem and a Homogeneous Linear Integral Equation	176
75.	Special Case $g(x) \equiv 1$	181
76.	Miscellaneous Remarks.	187
 II. Applications to some Problems of the Calculus of Variations		
77.	Some Auxiliary Theorems of the Calculus of Variations	192
78.	Dirichlet's Problem	196
79.	Applications to the Second Variation.	204
80.	Connection with Jacobi's Condition	207
 III. Vibration Problems		
81.	Vibrating String.	209
82.	Vibrations of a Rope.	215
83.	The Rotating Rope	222
 IV. Applications of the Hilbert-Schmidt Theory to the Flow of Heat in a Bar		
84.	The Partial Differential Equations of the Problem.	229
85.	Application to an Example	235
86.	General Theory of the Exceptional Case	240
87.	Flow of Heat in a Ring.	243
88.	Stationary Flow of Heat Produced by an Interior Source.	246
89.	Direct Computation of the Characteristic Constants and Fundamental Functions	247
 INDEX		251

LINEAR INTEGRAL EQUATIONS

CHAPTER I INTRODUCTORY

1. Linear Integral Equation of the First Kind.—An equation of the form

$$(1) \quad \int_a^b K(x, t)u(t)dt = f(x)$$

is said to be a **linear integral equation of the first kind**. The functions $K(x, t)$ and $f(x)$ and the limits a and b are known. It is proposed so to determine the unknown function u that (1) is satisfied for all values of x in the closed interval $a \leq x \leq b$. $K(x, t)$ is called the *kernel* of this equation.

Instead of equation (1), we have often to deal with equations of exactly the same form in which the upper limit of integration is the variable x . Such an equation is seen to be a special case of (1) in which the kernel $K(x, t)$ vanishes when $t > x$, since it then makes no difference whether x or b is used as the upper limit of integration.

The characteristic feature of this equation is that the unknown function u occurs under a definite integral. Hence equation (1) is called an *integral* equation and, since u occurs linearly, equation (1) is called a *linear* integral equation.

2. Abel's Problem.—As an illustration of the way in which integral equations arise, we give here a statement of Abel's problem.




FIG. 1.

Given a smooth curve situated in a vertical plane. A particle starts from rest at any point P . Let us find, under the action of gravity, the time T of descent to the lowest point O . Choose O as the origin of coordinates, the x -axis vertically upward, and the y -axis horizontal. Let the coordinates of P be (x, y) , of Q be (ξ, η) , and s the arc OQ .

The velocity of the particle at Q is

$$\frac{ds}{dt} = -\sqrt{2g(x - \xi)}.$$

Hence

$$t = - \int_P^Q \frac{ds}{\sqrt{2g(x - \xi)}}.$$

The whole time of descent is, then,

$$T = \int_0^P \frac{ds}{\sqrt{2g(x - \xi)}}.$$

If the shape of the curve is given, then s can be expressed in terms of ξ and hence ds can be expressed in terms of ξ . Let

$$ds = u(\xi)d\xi.$$

Then

$$T = \int_0^x \frac{u(\xi)d\xi}{\sqrt{2g(x - \xi)}}.$$

Abel set himself the problem¹ of finding that curve for which the time T of descent is a given function of x , say $f(x)$.

¹ For a solution of this problem, see BÔCHER, "Integral Equations," p. 8, Cambridge University Press, 1909.

Our problem, then, is to find the unknown function u from the equation

$$f(x) = \int_0^x \frac{1}{\sqrt{2g(x - \xi)}} u(\xi) d\xi.$$

This is a linear integral equation of the first kind for the determination of u .

3. Linear Integral Equation of the Second Kind.—An equation of the form

$$(2) \quad u(x) = f(x) + \int_a^b K(x, t) u(t) dt$$

is said to be a **linear integral equation of the second kind**.

$K(x, t)$ is called the kernel of this equation. The functions $K(x, t)$ and $f(x)$ and the limits a and b are known. The function u is unknown.

The equation

$$u(x) = f(x) + \int_a^x K(x, t) u(t) dt$$

is known as Volterra's linear integral equation of the second kind.

If $f(x) \equiv 0$, then

$$u(x) = \int_a^b K(x, t) u(t) dt.$$

This equation is said to be a homogeneous linear integral equation of the second kind.

Sometimes, in order to facilitate the discussion, a parameter λ is introduced, thus

$$u(x) = f(x) + \lambda \int_a^b K(x, t) u(t) dt.$$

This equation is said to be a linear integral equation of the second kind with a parameter.

Linear integral equations of the first and second kinds are special cases of the linear integral equation of the third kind:

$$\Psi(x)u(x) = f(x) + \int_a^b K(x, t)u(t)dt.$$

Equation (1) is obtained if $\Psi(x) \equiv 0$.

Equation (2) is obtained if $\Psi(x) \equiv 1$.

4. Relation between Linear Differential Equations and Volterra's Integral Equation.—Consider the equation

$$(3) \quad \frac{d^n y}{dx^n} + a_1(x) \frac{d^{n-1}y}{dx^{n-1}} + \dots + a_n(x)y = \varphi(x),$$

where the origin is a regular point for the $a_i(x)$.

Let us make the transformation

$$\frac{d^n y}{dx^n} = u(x).$$

Then

$$(4) \quad \left\{ \begin{array}{l} \frac{d^{n-1}y}{dx^{n-1}} = \int_0^x u(x)dx + C_1 \\ \dots \\ y = \int_0^x u(x)dx^n + C_1 \frac{x^{n-1}}{n-1} + C_2 \frac{x^{n-2}}{n-2} + \dots + C_n, \end{array} \right.$$

where $\int_0^x u(x)dx^n$ stands for a multiple integral of order n .

Equations (4) transform (3) into

$$(5) \quad u(x) + a_1(x) \int_0^x u(x)dx + \dots + a_n(x) \int_0^x u(x)dx^n = \varphi(x) + \sum_1^n C_i \alpha_i(x),$$

where

$$\alpha_i(x) = a_i(x) + \frac{x}{1} a_{i+1}(x) + \dots + a_n(x) \frac{x^{n-i}}{n-i}.$$

If we now put

$$\varphi(x) + \sum_1^n C_i \alpha_i(x) = f(x),$$

and make use of the well-known formula

$$\int_a^x u(t) dt^n = \int_a^x \frac{(x-t)^{n-1}}{|n-1|} u(t) dt,$$

equation (5) becomes

$$u(x) + \int_a^x \left[a_1(x) + a_2(x)(x-t) + \dots + a_n(x) \frac{(x-t)^{n-1}}{|n-1|} \right] u(t) dt = f(x),$$

which is a Volterra integral equation of the second kind.

In order that the right-hand member of (5) have a definite value it is necessary that the coefficients C_i have definite values. Then, inversely, the solution of the Volterra's equation (5) is equivalent to the solution of Cauchy's problem for the linear differential equation (3). The uniqueness of the solution of Volterra's equation follows from the fact that Cauchy's problem admits for a regular point one and only one solution.¹

5. Non-linear Equations.—This work will be confined to a discussion of linear integral equations. It is desirable, however, at this point to call the reader's attention to some integral equations which are non-linear.

The unknown function may appear in the equation to a power n greater than 1, for example;

$$u(x) = f(x) + \lambda \int_a^b K(x,t) u^n(t) dt.$$

The unknown function may appear in a more general way, as indicated by the following equation:

$$u(x) = f(x) + \lambda \int_a^b \varphi[x, t, u(t)] dt.$$

¹ For further discussion consult LALESKO, T., "Théorie Des Équations Intégrales," pp. 12ff, Herman and Fils, Paris, 1912.

In particular, the differential equation

$$\frac{du}{dx} = \varphi(x, u)$$

can be put in the integral form

$$u(x) = C + \int_0^x \varphi[t, u(t)]dt.$$

Still other general types of non-linear integral equations have been considered. Studies have also been made of systems of integral equations both linear and non-linear. Some study has been made of integral equations in more than one variable, for example;

$$u(x, y) = f(x, y) + \lambda \int_a^b \int_c^d K(x, y; t_1, t_2)u(t_1, t_2)dt_1dt_2.$$

6. Singular Equations.—An integral equation is said to be singular when either one or both of the limits of integration become infinite, for example;

$$u(x) = f(x) + \lambda \int_0^\infty \sin (xt)u(t)dt.$$

An integral equation is also said to be singular if the kernel becomes infinite for one or more points of the interval under discussion, for example;

$$f(x) = \int_0^x \frac{H(x, t)}{(x - t)^\alpha} u(t)dt \quad (0 < \alpha < 1).$$

Abel's problem, as stated in §2, is of this character. Abel set himself the problem of solving the more general equation

$$f(x) = \int_a^x \frac{u(t)dt}{(x - t)^\alpha} \quad (0 < \alpha < 1).$$

7. Types of Solutions.—By the use of distinct methods, the solution of a linear integral equation of the second kind with a parameter λ has been obtained in three different forms:

1. The first method, that of successive substitutions, due to Neumann, Liouville, and Volterra, gives us $u(x)$ as an integral series in λ , the coefficients of the various powers of λ being functions of x . The series converges for values of λ less in absolute value than a certain fixed number.

2. The second method, due to Fredholm, gives $u(x)$ as the ratio of two integral series in λ . Each series has an *infinite* radius of convergence. In the numerator the coefficients of the various powers of λ are functions of x . The denominator is independent of x . For those values of λ for which the denominator vanishes, there is, in general, no solution, but the method gives the solution in those exceptional cases in which a solution does exist. The solution is obtained by regarding the integral equation as the limiting form of a system of n linear algebraic equations in n variables as n becomes infinite.

3. The third method, developed by Hilbert and Schmidt, gives $u(x)$ in terms of a set of *fundamental functions*. The functions are, in the ordinary case, the solutions of the corresponding homogeneous equation

$$u(x) = \lambda \int_a^b K(x, t)u(t)dt.$$

In general, this equation has but one solution:

$$u(x) \equiv 0.$$

But there exists a set of numbers,

$$\lambda_1, \lambda_2, \dots, \lambda_n, \dots;$$

called *characteristic constants* or *fundamental numbers*, for each of which this equation has a finite solution:

$$u_1(x), u_2(x), \dots, u_n(x), \dots$$

These are the fundamental functions. The solution then is obtained in the form

$$u(x) = \sum C_n u_n(x),$$

where the C_n are arbitrary constants.

EXERCISES

Form the integral equations corresponding to the following differential equations with the given initial conditions:

1. $\frac{d^2y}{dx^2} + y = 0, x = 0, y = 0, y' = 1, y'' = 0.$

$$Ans. u(x) = x + \int_0^x (t - x)u(t)dt.$$

2. $\frac{d^2y}{dx^2} - 5\frac{dy}{dx} + 6y = 0, x = y = 0, y' = -1, y'' = -5.$

$$Ans. u(x) = 29 + 6x + \int_0^x (6x - 6t + 5)u(t)dt.$$

3. $\frac{d^2y}{dx^2} + y = \cos x, x = y = 0, y' = 1, y'' = 2.$

$$Ans. u(x) = \cos x - x - 2 + \int_0^x (t - x)u(t)dt.$$

4. $\frac{dy}{dx} - y = 0, x = 0, y = y' = 1.$

$$Ans. u(x) = 1 + \int_0^x u(t)dt.$$

5. $\frac{d^3y}{dx^3} - 3\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 5y = 0.$

$$Ans. u(x) = 1 - 2x - 4x^2 + \int_0^x [3 + 6(x - t) - 4(x - t)^2]u(t)dt.$$

CHAPTER II

SOLUTION OF INTEGRAL EQUATION OF SECOND KIND BY SUCCESSIVE SUBSTITUTIONS

8. Solution by Successive Substitutions.—We proceed now to a solution of the linear integral equation of the second kind with a parameter. We take up first the case where both limits of integration are fixed (*Fredholm's equation*). We assume that

$$(1) \quad a) \quad u(x) = f(x) + \lambda \int_a^b K(x,t)u(t)dt, \quad (a, b, \text{ constants}).$$

- b) $K(x, t) \neq 0$, is real and continuous in the rectangle R , for which $a \leq x \leq b$ and $a \leq t \leq b$.
- c) $f(x) \neq 0$, is real and continuous in the interval I , for which $a \leq x \leq b$.
- d) λ , constant.

We see at once that if there exists a continuous solution $u(x)$ of (1) and $K(x, t)$ is continuous, then $f(x)$ must be continuous. Hence the inclusion of condition (c) above.

Substitute in the second member of (1), in place of $u(t)$, its value as given by the equation itself. We find

$$\begin{aligned} u(x) &= f(x) + \lambda \int_a^b K(x,t) \left[f(t) + \lambda \int_a^b K(t,t_1)u(t_1)dt_1 \right] dt \\ &= f(x) + \lambda \int_a^b K(x,t)f(t)dt \\ &\quad + \lambda^2 \int_a^b K(x,t) \int_a^b K(t,t_1)u(t_1)dt_1 dt. \end{aligned}$$

Here again we substitute for $u(t_1)$ its value as given by (1).

We get

$$\begin{aligned}
 u(x) &= f(x) + \lambda \int_a^b K(x, t)f(t)dt \\
 &\quad + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1) \left[f(t_1) \right. \\
 &\quad \quad \quad \left. + \lambda \int_a^b K(t_1, t_2)u(t_2)dt_2 \right] dt_1 dt \\
 &= f(x) + \lambda \int_a^b K(x, t)f(t)dt \\
 &\quad + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1)f(t_1)dt_1 dt \\
 &\quad + \lambda^3 \int_a^b K(x, t) \int_a^b K(t, t_1) \int_a^b K(t_1, t_2)u(t_2)dt_2 dt_1 dt.
 \end{aligned}$$

Proceeding in this way we obtain

$$\begin{aligned}
 (2) \quad u(x) &= f(x) + \lambda \int_a^b K(x, t)f(t)dt \\
 &\quad + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1)f(t_1)dt_1 dt + \dots \\
 &\quad + \lambda^n \int_a^b K(x, t) \int_a^b K(t, t_1) \dots \\
 &\quad \quad \quad \int_a^b K(t_{n-2}, t_{n-1})f(t_{n-1})dt_{n-1} \dots dt_1 dt + R_{n+1}(x),
 \end{aligned}$$

where

$$R_{n+1}(x) = \lambda^{n+1} \int_a^b K(x, t) \int_a^b K(t, t_1) \dots \int_a^b K(t_{n-1}, t_n) \\
 u(t_n)dt_n \dots dt_1 dt.$$

This leads us to the consideration of the following infinite series:

$$\begin{aligned}
 (3) \quad f(x) &+ \lambda \int_a^b K(x, t)f(t)dt \\
 &\quad + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1)f(t_1)dt_1 dt + \dots
 \end{aligned}$$

Under our hypotheses *b*) and *c*), each term of this series is continuous in *I*. This series then represents a continuous function in *I*, provided it converges uniformly in *I*.

Since $K(x, t)$ and $f(x)$ are continuous in *R* and *I* respectively, $|K|$ has a maximum value *M* in *R* and $|f(x)|$ has a maximum value *N* in *I*:

$$\begin{aligned}|K(x, t)| &\leq M \quad \text{in } R \\ |f(x)| &\leq N \quad \text{in } I.\end{aligned}$$

$$\text{Put } S_n(x) = \lambda^n \int_a^b K(x, t) \int_a^b K(t, t_1) \dots \int_a^b K(t_{n-2}, t_{n-1}) f(t_{n-1}) dt_{n-1} \dots dt_1 dt.$$

$$\text{Then } |S_n(x)| \leq |\lambda|^n NM^n(b - a)^n.$$

The series of which this is a general term converges only when

$$|\lambda|M(b - a) < 1.$$

Thus we see that the series (3) converges absolutely and uniformly when

$$|\lambda| < \frac{1}{M(b - a)}.$$

If (1) has a continuous solution, it must be expressed by (2). If $u(x)$ is continuous in *I*, its absolute value has a maximum value *U*. Then

$$|R_{n+1}(x)| < |\lambda^{n+1}| UM^{n+1}(b - a)^{n+1}.$$

$$\text{If } |\lambda|M(b - a) < 1, \text{ then } \lim_{n \rightarrow \infty} R_{n+1}(x) = 0.$$

Thus we see that the function $u(x)$ satisfying (2) is the continuous function given by the series (3).

We can verify by direct substitution that the function $u(x)$ defined by (3) satisfies (1) or, what amounts to the same thing, place the series given by (3) equal to $u(x)$,

multiply both sides by $\lambda K(x, t)$ and integrate term by term,¹ as we have a right to do. We obtain

$$\begin{aligned} \lambda \int_a^b K(x, t) u(t) dt &= \lambda \int_a^b K(x, t) \left[f(t) + \lambda \int_a^b K(t, t_1) f(t_1) dt_1 \right. \\ &\quad \left. + \dots \right] dt \\ &= \lambda \int_a^b K(x, t) f(t) dt \\ &\quad + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1) f(t_1) dt_1 dt \\ &\quad + \dots \\ &= u(x) - f(x). \end{aligned}$$

Thus we obtain the following:

Theorem I.—If

- a) $u(x) = f(x) + \lambda \int_a^b K(x, t) u(t) dt$ (a, b , constants).
- b) $K(x, t)$ is real and continuous in a rectangle R ,
for which $a \leq x \leq b, a \leq t \leq b$.
 $|K(x, t)| \leq M$ in $R, K(x, t) \neq 0$.
- c) $f(x) \neq 0$, is real and continuous in $I: a \leq x \leq b$.
- d) λ constant, $|\lambda| < \frac{1}{M(b-a)}$,

then the equation (1) has one and only one continuous solution in I and this solution is given by the absolutely and uniformly convergent series (3).

The equation

$$(4) \quad u(x) = f(x) + \int_a^b K(x, t) u(t) dt$$

is a special case of the equation (1), for which $\lambda = 1$. The discussion just made holds without change after putting $\lambda = 1$.

¹ GOURSAT-HEDRICK, "Mathematical Analysis, vol. 1, §174, Ginn & Co.

Equations (1) and (4) may have a continuous solution, even though the hypothesis d),

$$|\lambda|M(b-a) < 1,$$

is not fulfilled. The truth of this statement is shown by the following example:

$$u(x) = \frac{x}{2} - \frac{1}{3} + \int_a^1 (x+t)u(t)dt,$$

which has the continuous solution $u(x) = x$, while

$$|\lambda|M(b-a) = 2 \not< 1.$$

9. Volterra's Equation.—The equation

$$(5) \quad u(x) = f(x) + \lambda \int_a^x K(x, t)u(t)dt$$

is known as *Volterra's equation*.

Let us substitute successively for $u(t)$ its value as given by (5). We find

$$(6) \quad u(x) = f(x) + \lambda \int_a^x K(x, t)f(t)dt \\ + \lambda^2 \int_a^x K(x, t) \int_a^t K(t, t_1)f(t_1)dt_1 dt + \dots \\ + \lambda^n \int_a^x K(x, t) \int_a^t K(t, t_1) \dots \\ \int_a^{t_{n-2}} K(t_{n-2}, t_{n-1})f(t_{n-1})dt_{n-1} \dots dt_1 dt + R_{n+1}(x),$$

where

$$R_{n+1}(x) = \lambda^{n+1} \int_a^x K(x, t) \int_a^t K(t, t_1) \dots \\ \int_a^{t_{n-1}} K(t_{n-1}, t_n) u(t_n)dt_n \dots dt_1 dt.$$

We consider the infinite series.

$$(6') \quad u(x) = f(x) + \lambda \int_a^x K(x, t)f(t)dt \\ + \lambda^2 \int_a^x K(x, t) \int_a^t K(t, t_1)f(t_1)dt_1 dt + \dots$$

The general term $V_n(x)$ of this series may be written

$$V_n(x) = \lambda^n \int_a^x K(x, t) \int_a^t K(t, t_1) \dots \int_a^{t_{n-2}} K(t_{n-2}, t_{n-1}) f(t_{n-1}) dt_{n-1} \dots dt_1 dt.$$

Then, since $|K(x, t)| \leq M$ in R and $|f(t)| \leq N$ in I , we have

$$|V_n(x)| \leq |\lambda^n| N M^n \frac{(x-a)^n}{n!} \leq |\lambda^n| N \frac{[M(b-a)]^n}{n!}, (a \leq x \leq b).$$

The series, for which the positive constant $|\lambda^n| N \frac{[M(b-a)]^n}{n!}$

is the general expression for the n th term, is convergent for all values of λ , N , M , $(b-a)$. Hence the series (6') is absolutely and uniformly convergent.

If (5) has a continuous solution, it must be expressed by (6'). If $u(x)$ is continuous in I , its absolute value has a maximum value U . Then

$$|R_{n+1}(x)| \leq |\lambda^{n+1}| U M^{n+1} \frac{(x-a)^{n+1}}{|n+1|} \leq |\lambda^{n+1}| U \frac{[M(b-a)]^{n+1}}{|n+1|}, \\ (a \leq x \leq b).$$

Whence

$$\lim_{n \rightarrow \infty} R_{n+1}(x) = 0.$$

Thus we see that the function $u(x)$, satisfying (6), is the continuous function given by the series (6'). As before, we can show that the expression for $u(x)$ given by (6') satisfies (5). Hence we have the following:

Theorem II.—If

$$(5) \quad a) \quad u(x) = f(x) + \lambda \int_a^x K(x, t) u(t) dt \quad (a, \text{ constant}).$$

- b) $K(x, t)$ is real and continuous in the rectangle R , for which $a \leq x \leq b$, $a \leq t \leq b$.
- c) $|K(x, t)| \leq M$ in R , $K(x, t) \neq 0$.
- d) $f(x) \neq 0$, is real and continuous in I : $a \leq x \leq b$.
- d) λ , constant.

then the equation (5) has one and only one continuous solution $u(x)$ in I , and this solution is given by the absolutely and uniformly convergent series (6').

The results of this article hold without change for the equation

$$u(x) = f(x) + \int_a^x K(x, t)u(t)dt$$

by putting throughout the discussion $\lambda = 1$.

10. Successive Approximations.—We would like to point out that the method of solution by successive approximations differs from that of successive substitutions.

Under the method of successive approximations we select any real function $u_o(x)$ continuous in I . Substitute in the right-hand member of

$$(1) \quad u(x) = f(x) + \lambda \int_a^b K(x, t) u(t) dt,$$

in place of $u(t)$, the function $u_o(t)$. We find

$$u_1(x) = f(x) + \lambda \int_a^b K(x, t) u_o(t) dt.$$

The function $u_1(x)$ so determined is real and continuous in I . Continue in like manner by replacing u_o by u_1 , and so on. We obtain a series of functions

$$u_o(x), u_1(x), u_2(x), \dots, u_n(x), \dots$$

which satisfy the equations,

$$(7) \quad \left\{ \begin{array}{l} u_2(x) = f(x) + \lambda \int_a^b K(x, t) u_1(t) dt \\ \dots \dots \dots \dots \dots \dots \\ u_{n-1}(x) = f(x) + \lambda \int_a^b K(x, t) u_{n-2}(t) dt \\ u_n(x) = f(x) + \int_a^b K(x, t) u_{n-1}(t) dt \\ \dots \dots \dots \dots \dots \dots \end{array} \right.$$

From these equations we find

$$(8) \quad u_n(x) = f(x) + \lambda \int_a^b K(x, t)f(t)dt + \lambda^2 \int_a^b K(x, t) \int_a^b K(t, t_1)f(t_1)dt_1 dt + \dots + \lambda^{n-1} \int_a^b K(x, t) \int_a^b K(t, t_1) \dots \int_a^b K(t_{n-3}, t_{n-2})f(t_{n-2})dt_{n-2} \dots dt_1 dt + R_n$$

where

$$R_n = \lambda^n \int_a^b K(x, t) \int_a^b K(t, t_1) \dots \int_a^b K(t_{n-2}, t_{n-1}) u_o(t_{n-1}) dt_{n-1} \dots dt_1 dt$$

$u_o(x)$ is real and continuous in I and so has a maximum value U in I . Then it is easy to see that

$$|R_n| \leq |\lambda^n| UM^n(b-a)^n.$$

If, then, $|\lambda|M(b-a) < 1$, we have

$$\lim_{n \rightarrow \infty} R_n = 0.$$

Thus, as n increases, the series of functions $u_n(x)$ approach a limit function which is given by the series in the right member of (8). We identify this series with the right member of (6'). Thus

$$\lim_{n \rightarrow \infty} u_n(x) \equiv u(x).$$

By this process at each step a new function $u_n(x)$ appears dependent upon the choice of $u_o(x)$. We notice, however, that the limit $u(x)$ is independent of the choice of $u_o(x)$.

We can now make an independent proof of the uniqueness of the solution. Suppose there was another solution $v(x)$. Choose $u_o(x) \equiv v(x)$. It is then clear that each $u_n(x)$ will be identical with $v(x)$ and hence the limit will be

$v(x)$. But we have just seen that the limit is independent of the choice of $u_o(x)$. Therefore,

$$v(x) \equiv u(x).$$

A similar discussion can be carried through without further difficulty for the Volterra equation.

11. Iterated Functions.—Place

$$(9) \quad \begin{cases} K_1(x, t) = K(x, t) \\ K_i(x, t) = \int_a^b K(x, s) K_{i-1}(s, t) ds. \end{cases}$$

The functions $K_1, K_2, \dots, K_n, \dots$ so formed are called *iterated functions*.

By successive applications of (9) it is evident that

$$(10) \quad K_i(x, t) = \int_a^b \dots \int_a^b K(x, s_1) K(s_1, s_2) \dots K(s_{i-1}, t) ds_{i-1} \dots ds_1.$$

From (10) $K_n(x, s)$ is an $(n - 1)$ -fold integral and $K_p(s, t)$ is a $(p - 1)$ -fold integral. Whence we see that $\int_a^b K_n(x, s) K_p(s, t) ds$ is an $(n + p - 1)$ -fold integral, which, by some simple changes in the order of integration, is seen to be identical with $K_{n+p}(x, t)$. Hence

$$(11) \quad K_{n+p}(x, t) = \int_a^b K_n(x, s) K_p(s, t) ds.$$

12. Reciprocal Functions.—Let

$$(12) \quad -k(x, t) = K_1(x, t) + K_2(x, t) + \dots + K_n(x, t) + \dots$$

It is easy to show that, when $K(x, t)$ is real and continuous in R , the infinite series for $k(x, t)$ is absolutely and uniformly convergent if $M(b - a) < 1$. Consequently, $k(x, t)$ is real

and continuous in R . On account of the first of equations (9) and equation (11), we have

$$\begin{aligned} -k(x, t) - K(x, t) &= K_2(x, t) + K_3(x, t) + \dots \\ &\quad + K_n(x, t) + \dots \\ &= \int_a^b K_1(x, s) K_1(s, t) ds + \dots \\ &\quad + \int_a^b K_1(x, s) K_{n-1}(s, t) ds + \dots \\ &= \int_a^b K_1(x, s) K_1(s, t) ds + \dots \\ &\quad + \int_a^b K_{n-1}(x, s) K_1(s, t) ds + \dots \end{aligned}$$

These equations may be written

$$\begin{aligned} -k(x, t) - K(x, t) &= \int_a^b K_1(x, s) \left[K_1(s, t) + \dots \right. \\ &\quad \left. + K_{n-1}(s, t) + \dots \right] ds \\ &= \int_a^b \left[K_1(x, s) + \dots + K_{n-1}(x, s) + \dots \right] K_1(s, t) ds. \end{aligned}$$

If we now make use of (12), we obtain the following characteristic formula:

$$(13) \quad K(x, t) + k(x, t) = \int_a^b K(x, s) k(s, t) ds \\ = \int_a^b k(x, s) K(s, t) ds.$$

Two functions $K(x, t)$ and $k(x, t)$ are said to be *reciprocal* if they are both real and continuous in R and if they satisfy the condition (13). A function $k(x, t)$ reciprocal to $K(x, t)$ will exist, provided the series in (12) converges uniformly. But we have seen that this series converges uniformly when $M(b - a) < 1$, where M is the maximum of $|K(x, t)|$ in R . Thus, we have the

Theorem III.—*If $K(x, t)$ is real and continuous in R , there exists a reciprocal function $k(x, t)$ given by (12) provided that*

$$M(b - a) < 1$$

where M is the maximum of $|K(x, t)|$ in R .

13. Volterra's Solution of Fredholm's Equation.—Volterra has shown how to find a solution of

$$(4) \quad u(x) = f(x) + \int_a^b K(x, t)u(t)dt$$

whenever the reciprocal function $k(x, t)$ of $K(x, t)$ is known. If (4) has a continuous solution $u(x)$, then

$$u(t) = f(t) + \int_a^b K(t, t_1)u(t_1)dt_1.$$

Multiplying by $k(x, t)$ and integrating, we find

$$\begin{aligned} \int_a^b k(x, t)u(t)dt &= \int_a^b k(x, t)f(t)dt \\ &\quad + \int_a^b \int_a^b k(x, t)K(t, t_1)u(t_1)dt_1 dt \\ &= \int_a^b k(x, t)f(t)dt \\ &\quad + \int_a^b \left[K(x, t_1) + k(x, t_1) \right] u(t_1)dt_1, \end{aligned}$$

which reduces to

$$(14) \quad 0 = \int_a^b k(x, t)f(t)dt + \int_a^b K(x, t_1)u(t_1)dt_1.$$

But from (4) we have

$$\int_a^b K(x, t_1)u(t_1)dt_1 = u(x) - f(x).$$

Therefore, (14) may be written

$$(15) \quad u(x) = f(x) - \int_a^b k(x, t)f(t)dt.$$

If (4) has a continuous solution, it is given by this formula and it is unique.

To see that the expression for $u(x)$ given by (15) is, indeed, a solution, we write (15) in the form

$$f(x) = u(x) + \int_a^b k(x, t)f(t)dt.$$

This is an integral equation for the determination of $f(x)$. The function reciprocal to $k(x, t)$ is $K(x, t)$. By what we have just proved, if this equation has a continuous solution, it is unique and is given by

$$f(x) = u(x) - \int_a^b K(x, t)u(t)dt.$$

But this is the equation (1) from which we started. Thus we see that (4) is satisfied by the value of $u(x)$ given by (15). Thus we have the following

Theorem IV.—If

a) $K(x, t)$ is real and continuous in R . $K(x, t) \neq 0$.

b) $f(x)$ is real and continuous in I . $f(x) \neq 0$.

c) A function $k(x, t)$ reciprocal to $K(x, t)$ exists, then the equation (4) has one and only one continuous solution in I and this solution is given by (15).

The same reasoning applied to (13), considered as an integral equation for the determination of $k(x, t)$, shows that, if a continuous reciprocal function exists, it is unique.

14. Discontinuous Solutions.—We have shown the existence, under proper assumptions, of a unique continuous solution for a linear integral equation of the second kind. This integral equation may have also, in addition, discontinuous solutions. To show this we exhibit the special equation¹

$$u(x) = \int_0^x t^{x-t}u(t)dt,$$

which has one and only one continuous solution, namely $u(x) \equiv 0$. We can show, by direct substitution, that this

¹ BÔCHER, "Integral Equations," p. 17, Cambridge Press, 1909.

equation has also an infinite number of discontinuous solutions given by

$$u(x) = Cx^{x-1},$$

where C is an arbitrary constant not zero.

EXERCISES

Solve the following linear integral equations:

1. a) $u(x) = x + \int_0^x (t - x)u(t)dt.$ *Ans.* $u(x) = \sin x.$
- b) $u(x) = 1 + \int_0^x (t - x)u(t)dt.$ *Ans.* $u(x) = \cos x.$
2. $u(x) = \frac{5x}{6} + \frac{1}{2} \int_0^1 xt.u(t)dt.$ *Ans.* $u(x) = x.$
3. $u(x) = \frac{5x}{6} - \frac{1}{9} + \frac{1}{3} \int_0^1 (t + x)u(t)dt.$ *Ans.* $u(x) = x.$
4. $u(x) = 1 + \int_0^x u(t)dt.$ *Ans.* $u(x) = e^x.$
5. $u(x) = e^x - \frac{e}{2} + \frac{1}{2} + \frac{1}{2} \int_0^1 u(t)dt.$ *Ans.* $u(x) = e^x.$
6. $u(x) = \sin x - \frac{x}{4} + \frac{1}{4} \int_0^{\frac{\pi}{2}} tx u(t)dt.$ *Ans.* $u(x) = \sin x.$
7. $u(x) = x + \int_0^{\frac{3\pi}{2}} u(t)dt.$ *Ans.* $u(x) = x + \text{constant.}$
8. $u(x) = 1 - 2x - 4x^2 + \int_0^x [3 + 6(x - t) - 4(x - t)^2]u(t)dt.$ *Ans.* $u(x) = e^x.$
9. $u(x) = \frac{3}{2}e^x - \frac{xe^x}{2} - \frac{1}{2} + \frac{1}{2} \int_0^1 t u(t)dt.$ *Ans.* $u(x) = e^x.$
10. $u(x) = 29 + 6x + \int_0^x (6x - 6t + 5)u(t)dt.$ *Ans.* $u(x) = e^{2x} - e^{3x}.$
11. $u(x) = \cos x - x - 2 + \int_0^x (t - x)u(t)dt.$ *Ans.* $u(x) = \sin x + x \sin x.$

- 12.** Using the method of successive approximations find five successive approximations in the solution of Exercises 1, 2, 3, 4, 5 after choosing $u_0(x) \equiv 0$.
- 13.** Show that

$$u(x) = A + Bx + \int_0^x [C + D(x-t)]u(t)dt$$

where A, B, C, D are arbitrary constants, has for solution

$$u(x) = K_1 e^{m_1 x} + K_2 e^{m_2 x},$$

where K_1, K_2, m_1, m_2 depend upon A, B, C, D .

- 14.** Show that

$$u(x) = f(x) + \lambda \int_a^b u(t) \left[\sum_1^p \alpha_q(t) \beta_q(t) \right] dt$$

has the solution

$$u(x) = f(x) + \lambda \sum_1^p A_q \alpha_q(x),$$

where the A_q are constants determined by the equation

$$\sum_1^p r A_r \left[\lambda \int_a^b \alpha_r(t) \beta_q(t) dt \right] - A_q = \int_a^b \beta_q(t) f(t) dt \quad (q = 1, \dots, p).$$

[Heywood-Frèchet].

CHAPTER III

**SOLUTION OF FREDHOLM'S EQUATION EXPRESSED AS RATIO OF TWO INTEGRAL SERIES
IN λ**

15. Fredholm's Equation As Limit of a Finite System of Linear Equations.—The solution given in the previous chapter for the equation

$$(1) \quad u(x) = f(x) + \lambda \int_a^b K(x,t) u(t) dt$$

has the disadvantage of holding only for restricted values of λ . It is desirable to have, if possible, a solution which holds for all values of λ . Such a solution was given by Fredholm in the form

$$u(x) = \frac{\beta_0(x) + \beta_1(x)\lambda + \dots}{\alpha_0 + \alpha_1\lambda + \dots},$$

the numerator and denominator being permanently converging power series in λ .

a) The System of Linear Equations Replacing the Integral Equation.—Before stating explicitly and proving Fredholm's result, we give an outline of the reasoning which led him to his discovery.

Divide the interval (ab) into n equal parts and call the points of division t_1, t_2, \dots, t_{n-1} . Then

$$(2) \quad t_0 = a, t_1 = a + h, t_2 = a + 2h, \dots, t_n = a + nh,$$

$$h = \frac{b - a}{n}.$$

Replace the definite integral in (1) by the sum, corresponding to the points of division (2), of which it is the limit. We obtain the approximate equation

$$u(x) - \lambda h [K(x, t_1)u(t_1) + K(x, t_2)u(t_2) + \dots + K(x, t_n)u(t_n)] = f(x).$$

Since this equation holds for every value of x , it must be satisfied for $x = t_1, t_2, \dots, t_n$. We thus obtain the following system of n linear equations for the determination of the n unknowns $u(t_1), u(t_2), \dots, u(t_n)$:

$$u(t_1) - \lambda h [K(t_1, t_1)u(t_1) + K(t_1, t_2)u(t_2) + \dots + K(t_1, t_n)u(t_n)] = f(t_1)$$

$$(3) \quad u(t_2) - \lambda h [K(t_2, t_1)u(t_1) + K(t_2, t_2)u(t_2) + \dots + K(t_2, t_n)u(t_n)] = f(t_2)$$

$$u(t_n) - \lambda h [K(t_n, t_1)u(t_1) + K(t_n, t_2)u(t_2) + \dots + K(t_n, t_n)u(t_n)] = f(t_n).$$

This system being solved with respect to $u(t_1), u(t_2), \dots, u(t_n)$, we can plot $u(t_i)$ ($i = 1, \dots, n$) as ordinates and by interpolation draw a curve $u(x)$, which we may expect




FIG. 2.

to be an approximation to the actual solution. We solve (3), making use of the abbreviations

$$f(t_i) = f_i \quad , \quad u(t_i) = u_i \quad , \quad K(t_i, t_j) = K_{ij}.$$

Denote by Δ the determinant of the coefficients of the u_i in (3). We have

$$\Delta = \begin{vmatrix} 1 - \lambda h K_{11} & -\lambda h K_{12} & \dots & -\lambda h K_{1n} \\ -\lambda h K_{21} & 1 - \lambda h K_{22} & \dots & -\lambda h K_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ -\lambda h K_{n1} & -\lambda h K_{n2} & \dots & 1 - \lambda h K_{nn} \end{vmatrix}$$

Denote by $\Delta_{\nu\mu}$ the first minor of the element in the ν th row and μ th column of Δ . Then, from Cramer's formula, we obtain, by solving (3) with respect to u_k ,

$$(4) \quad u_k = \frac{\sum_{i=1}^n i f_i \Delta_{ik}}{\Delta}, \text{ provided } \Delta \neq 0, K = 1, \dots, n.$$

b) *Limit of Δ .*—Expanding the determinant Δ , we obtain

$$\begin{aligned} \Delta = 1 - \lambda \sum_{i=1}^n K_{ii} h + \frac{\lambda^2}{2!} \sum_{i,j=1}^n h^2 \left| \begin{array}{cc} K_{ii} & K_{ij} \\ K_{ji} & K_{jj} \end{array} \right| + \dots \\ + (-1)^n h^n \lambda^n \left| \begin{array}{cccccc} K_{11} & \dots & K_{1n} \\ \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots \\ K_{n1} & \dots & K_{nn} \end{array} \right|. \end{aligned}$$

If we now let n increase indefinitely, we see that each term of this series has a definite limit. So that, at least formally,

$$\begin{aligned} \lim_{n \rightarrow \infty} \Delta = 1 - \lambda \int_a^b K(t, t) dt \\ + \frac{\lambda^2}{2!} \int_a^b \int_a^b \left| \begin{array}{cc} K(t_1, t_1) & K(t_1, t_2) \\ K(t_2, t_1) & K(t_2, t_2) \end{array} \right| dt_1 dt_2 \\ - \frac{\lambda^3}{3!} \int_a^b \int_a^b \int_a^b \left| \begin{array}{ccccc} K(t_1, t_1) & \dots & K(t_1, t_3) \\ \vdots & \ddots & \vdots \\ K(t_3, t_1) & \dots & K(t_3, t_3) \end{array} \right| dt_1 dt_2 dt_3 + \dots \end{aligned}$$

$$(5) \quad \equiv D(\lambda).$$

$D(\lambda)$ is called *Fredholm's determinant*, or the determinant of K .

c) *Limit of Δ_{ik} .*—The expression for $\Delta_{\mu\mu}$ is similar to that for Δ .

$$\Delta_{\mu\mu} = 1 - \lambda \sum_{i=1}^n K_{ii} h + \frac{\lambda^2}{2!} \sum_{i,j=1}^n h^2 \left| \begin{array}{cc} K_{ii} & K_{ij} \\ K_{ji} & K_{jj} \end{array} \right| + \dots,$$

where ' means omit $i = \mu$. Hence

$$(6) \quad \lim_{n \rightarrow \infty} \Delta_{\mu\mu} = D(\lambda).$$

Again, from rules for expansion of determinants,

$$\begin{aligned} \Delta_{\nu\mu} &= \lambda h \left\{ K_{\mu\nu} - \lambda \sum_{i=1}^n h \begin{vmatrix} K_{\mu\nu} & K_{\mu i} \\ K_{i\nu} & K_{ii} \end{vmatrix} \right. \\ &\quad \left. + \frac{\lambda^2}{2!} \sum_{i,j=1}^n h_2 \begin{vmatrix} K_{\mu\nu} & K_{\mu i} & K_{\mu j} \\ K_{i\nu} & K_{ii} & K_{ij} \\ K_{j\nu} & K_{ji} & K_{jj} \end{vmatrix} + \dots \right\}. \end{aligned}$$

$$\text{Put } h\mathfrak{D}_{\mu\nu} = \Delta_{\nu\mu}.$$

If, as n increases indefinitely, we let (t_μ, t_ν) vary in such a way that $\lim (t_\mu, t_\nu) = (x, y)$, we find, at least formally

$$\begin{aligned} (7) \quad \lim_{n \rightarrow \infty} \mathfrak{D}_{\mu\nu} &= \lambda K(x, y) - \lambda^2 \int_a^b \begin{vmatrix} K(x, y) & K(x, t) \\ K(t, y) & K(t, t) \end{vmatrix} dt \\ &\quad + \frac{\lambda^3}{2!} \int_a^b \int_a^b \begin{vmatrix} K(x, y) & K(x, t_1) & K(x, t_2) \\ K(t_1, y) & K(t_1, t_1) & K(t_1, t_2) \\ K(t_2, y) & K(t_2, t_1) & K(t_2, t_2) \end{vmatrix} dt_1 dt_2 + \dots \end{aligned}$$

$$(8) \quad \equiv D(x, y; \lambda).$$

This expression for $D(x, y; \lambda)$ is called *Fredholm's first minor*.

d) *Limit of u_k* .—We can now write (4) in the form

$$\begin{aligned} u_k &= f_k \frac{\Delta_{kk}}{\Delta} + \sum_{i=1}^n f_i \frac{\Delta_{ik}}{\Delta} \\ &= f_k \frac{\Delta_{kk}}{\Delta} + \sum_{i=1}^n f_i h D_{ki}, \end{aligned}$$

which in the passage to the limit as $n \rightarrow \infty$, becomes

$$u(t_k) = f(t_k) + \frac{1}{D(\lambda)} \int_a^b f(t) D(t_k, t; \lambda) dt.$$

But t_k is any point of division. Then we can replace t_k by x and write

$$(9) \quad u(x) = f(x) + \frac{1}{D(\lambda)} \int_a^b f(t) D(x, t; \lambda) dt.$$

This result has not been obtained by a rigorous mathematical procedure. However, we are inclined to believe that the above expression for $u(x)$ is a solution of (1). This belief is later shown to be correct.

e) *Fredholm's Two Fundamental Relations.*—We will now develop two relations which will be of use to us later in obtaining a solution of the integral equation (1). We recall the theorem, fundamental in the theory of determinants, that the sum of the products of the elements of any column by the corresponding minors of any other column is zero. This theorem applied to the determinant Δ gives

$$(1 - \lambda h K_{jj}) \Delta_{jk} - \lambda h K_{kj} \Delta_{kk} - \sum_{i=1}^n'' \lambda h K_{ij} \Delta_{ik} = 0,$$

where " means omit $i = j, k$. Making use of the relation $\Delta_{\nu\mu} = h \mathfrak{D}(t\mu, t\nu)$, we find

$$(1 - \lambda h K_{jj}) h \mathfrak{D}_{kj} - \lambda h K_{kj} \Delta_{kk} - \sum_{i=1}^n'' \lambda h^2 K_{ij} \mathfrak{D}_{ki} = 0.$$

Divide through by h , since $h \neq 0$. The passage to the limit, as $n \rightarrow \infty$, gives from the last equation by (6) and (7)

$$D(t_k, t_j; \lambda) - \lambda K(t_k, t_j) D(\lambda) - \lambda \int_a^b K(t, t_i) D(t_k, t; \lambda) dt = 0.$$

This last equation holds for any two points t_j, t_k on the interval (ab) . Let us put then $t_k = x, t_j = y$ and write

$$(10) \quad D(x, y; \lambda) - \lambda K(x, y) D(\lambda) = \lambda \int_a^b K(t, y) D(x, t; \lambda) dt.$$

This is called Fredholm's *first fundamental relation*.

Now apply the theorem: The sum of the products of the elements of any row by the corresponding minors of any other row is zero. This theorem applied to the determinant Δ gives

$$(1 - \lambda h K_{jj}) \Delta_{kj} - \lambda h K_{jk} \Delta_{kk} - \sum_{i=1}^n'' \lambda h K_{ji} \Delta_{ki} = 0.$$

Proceeding as before, we find

$$D(t_j, t_k; \lambda) - \lambda K(t_j, t_k) D(\lambda) - \lambda \int_a^b K(t_j, t) D(t, t_k; \lambda) dt = 0.$$

This equation holds for any two points t_j, t_k on (ab) . Let us put then $t_j = x, t_k = y$ and write

$$(11) \quad D(x, y; \lambda) - \lambda K(x, y) D(\lambda) - \lambda \int_a^b K(x, t) D(t, y; \lambda) dt = 0.$$

This is Fredholm's *second fundamental relation*.

16. Hadamard's Theorem.—We now proceed to establish rigorously the results of the preceding article. To this end we need a theorem due to Hadamard. To establish this theorem we make use of the following

Lemma.—If all of the elements a_{ik} of the determinant

$$A = \begin{vmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{vmatrix}$$

are real and satisfy the conditions

$$(12) \quad a_{r1}^2 + a_{r2}^2 + \dots + a_{rn}^2 = 1 \quad (r = 1, \dots, n),$$

then $|A| \leq 1$.




FIG. 3.

We give first two special cases of the lemma which have a geometric interpretation.

1) $n = 2$. The parallelogram $OP_1P_3P_2$ has the vertex O at the origin of a system of rectangular coordinates. The coordinates of P_1 and P_2 are as

indicated in the figure. The area A of $OP_1P_3P_2$ is given by

$$A = \begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \end{vmatrix}.$$

If $OP_1 = OP_2 = 1$, that is, if

$$x_1^2 + y_1^2 = 1 \text{ and } x_2^2 + y_2^2 = 1,$$

then it is geometrically evident that the greatest area is obtained when the figure is a rectangle, and then the area is 1. Hence, we have generally $|A| \leq 1$.

2) $n = 3$. The parallelopiped $OP_1P_2P_3$ has one vertex at the origin of a system of rectangular coordinates. The coordinates of P_1, P_2, P_3 are as indicated in the figure. The volume V of $OP_1P_2P_3$ is given by

$$V = \begin{vmatrix} x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \\ x_3 & y_3 & z_3 \end{vmatrix}.$$




FIG. 4.

If $OP_1 = OP_2 = OP_3 = 1$, that is, if

$x_1^2 + y_1^2 + z_1^2 = 1, x_2^2 + y_2^2 + z_2^2 = 1, x_3^2 + y_3^2 + z_3^2 = 1$, then it is evident geometrically that the volume is greatest when the figure is a rectangular parallelopiped, in which case the volume is 1. Hence, we have generally

$$|V| \leq 1.$$

Proof of Lemma.— $A(a_{11}, \dots, a_{nn})$ is a continuous function of its arguments a_{rs} in the region \mathfrak{A} defined by the equations (12). These conditions insure that $|a_{rs}| \leq 1$, and that the region \mathfrak{A} is bounded and closed. Hence A reaches a maximum and a minimum on the region \mathfrak{A} . The maximum and minimum which we are seeking are the so-called absolute maximum and minimum. But if a system of values furnishes the absolute maximum (minimum) for A , it furnishes also a relative maximum (minimum). Hence the ordinary methods of the differential calculus can be used for their determination.

Now if a function

$$f(x_1, x_2, \dots, x_n)$$

of n variables, connected by h distinct relations

$$\phi_1(x_1, \dots, x_n) = 0, \phi_2(x_1, \dots, x_n) = 0, \dots, \phi_h(x_1, \dots, x_n) = 0$$

has a maximum (minimum), then the n first partial derivatives of the auxiliary function

$$F = f + \lambda_1\phi_1 + \lambda_2\phi_2 + \dots + \lambda_n\phi_n,$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are constants, must vanish.¹

For our problem, $f = A$, $x_r = a_{rs}$

$$\phi_r = \sum_{s=1}^n a_{rs}^2 - 1 \quad (r = 1, \dots, n)$$

and the auxiliary function F becomes

$$F = A + \sum_{s=1}^n \frac{\lambda_s}{2} (a_{s1}^2 + a_{s2}^2 + \dots + a_{sn}^2 - 1).$$

For a maximum (minimum) by the theorem just stated, we must have

$$\frac{\partial F}{\partial a_{jk}} = \frac{\partial A}{\partial a_{jk}} + \lambda_j a_{jk} = 0, \text{ or}$$

$$(13) \quad A_{jk} + \lambda_j a_{jk} = 0, \quad (j, k = 1, 2, \dots, n),^2$$

where A_{jk} denotes the minor of a_{jk} in A . Multiply both sides of this equation by a_{jk} and sum with respect to k for $k = 1, 2, \dots, n$. We obtain

$$A + \lambda_j = 0, \quad \text{since } \sum_{k=1}^n a_{jk}^2 = 1$$

$$\text{or} \quad \lambda_j = -A \quad (j = 1, \dots, n).$$

Substitute this value of λ_j in equation (13). Then

$$A_{jk} = A a_{jk} \quad (jk = 1, \dots, n).$$

Whence the determinant

$$\begin{vmatrix} A_{11} & A_{12} & \dots & A_{1n} \\ A_{21} & A_{22} & \dots & A_{2n} \\ \vdots & \ddots & \ddots & \vdots \\ A_{n1} & A_{n2} & \dots & A_{nn} \end{vmatrix}$$

¹ GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §61, Ginn & Co.

² For differentiation of determinants, see BALTZER, "Die Determinanten," §3, 14.

adjoint to A , is equal to

$$\begin{vmatrix} Aa_{11} & Aa_{12} & \dots & Aa_{1n} \\ Aa_{21} & Aa_{22} & \dots & Aa_{2n} \\ \dots & \dots & \dots & \dots \\ Aa_{n1} & Aa_{n2} & \dots & Aa_{nn} \end{vmatrix}.$$

The first of these determinants is equal¹ to A^{n-1} . The second reduces to A^{n+1} . Hence

$$A^{n+1} = A^{n-1}$$

But the maximum and minimum of A must satisfy this equation. Therefore, the maximum of A is $+1$, the minimum is -1 , and

$$|A| \leq 1.$$

Hadamard's Theorem.—We are now in a position to prove a more general theorem due to Hadamard: *If the elements b_{ik} of the determinant*

$$B = \begin{vmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{n1} & b_{n2} & \dots & b_{nn} \end{vmatrix}$$

are real and satisfy the inequality

$$|b_{ik}| \leq M,$$

then $|B| \leq M^n \sqrt{n^n}$.

Proof.—Let

$$b_{i1}^2 + b_{i2}^2 + \dots + b_{in}^2 = s_i, \quad (i = 1, \dots, n)$$

Case I.—*Some one or more of the s_i vanish, say $s_k = 0$. Then $b_{ki} = 0$ ($i = 1, \dots, n$). Therefore $B = 0$, and the theorem is proved in this case.*

Case II.—*None of the s_i vanish. Then each s_i is positive.*

That is $s_1 > 0, s_2 > 0, \dots, s_n > 0$.

¹ BALTZER, *Loc. cit.*, §6, 1.

We now consider the determinant

$$\frac{B}{\sqrt{s_1 s_2 \dots s_n}} = \begin{vmatrix} \frac{b_{11}}{\sqrt{s_1}} & \dots & \frac{b_{1n}}{\sqrt{s_1}} \\ \dots & \dots & \dots \\ \frac{b_{n1}}{\sqrt{s_n}} & \dots & \frac{b_{nn}}{\sqrt{s_n}} \end{vmatrix}$$

for which $\frac{b_{i1}^2}{s_i} + \dots + \frac{b_{in}^2}{s_i} = 1$. ($i = 1, \dots, n$).

This determinant satisfies all of the conditions of our lemma. Hence

$$|B| \leq \sqrt{s_1 s_2 \dots s_n}.$$

But, since

$$|b_{ik}| \leq M,$$

we have from $s_i = b_{i1}^2 + \dots + b_{in}^2$

$$s_i \leq nM^2. \quad (i = 1, \dots, n)$$

Therefore,

$$|B| \leq M^n \sqrt{n^n}.$$

17. Convergence Proof.—With the help of Hadamard's theorem we can now prove the convergence of the series for $D(\lambda)$ and $D(x, y; \lambda)$.

a) *Convergence of $D(\lambda)$.*— $D(\lambda)$ is given by the series

$$(14) \quad D(\lambda) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^n}{n!} A_n, \text{ where}$$

$$(15) \quad A_n = \int_a^b \dots \int_a^b \begin{vmatrix} K(t_1 t_1) & \dots & K(t_1 t_n) \\ \dots & \dots & \dots \\ K(t_n t_1) & \dots & K(t_n t_n) \end{vmatrix} dt_1 \dots dt_n \quad (n > 0).$$

We have assumed in §8 that $|K(x, t)| \leq M$ in R . Thus the determinant in the expression for A_n satisfies all of the conditions of Hadamard's theorem, and hence

$$\begin{aligned} |A_n| &\leq \int_a^b \dots \int_a^b M^n \sqrt{n^n} dt_1 \dots dt_n \\ &= \sqrt{n^n} M^n (b - a)^n. \end{aligned}$$

Then

$$\left| (-1)^n \frac{\lambda^n}{n!} A_n \right| \leq M^n (b-a)^n |\lambda^n| \frac{\sqrt{n^n}}{n!} = C_n.$$

This identity defines C_n .

We now proceed to show that the series of which C_n is the general term converges. Applying the ratio test, we obtain

$$\frac{C_{n+1}}{C_n} = M(b-a)|\lambda| \sqrt{\left(1 + \frac{1}{n}\right)^n} \frac{1}{\sqrt{n+1}}.$$

This ratio has for its limit zero as n becomes infinite. Hence the series of which C_n is the general term converges. Consequently, the series for $D(\lambda)$ is absolutely and permanently convergent. We state this result in the following:

Theorem I.—*The series $D(\lambda)$ is an absolutely and permanently converging power series in λ .*

b) *Convergence of $D(x, y; \lambda)$.*—We have

$$(16) \quad D(x, y; \lambda) = \lambda K(x, y) + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} B_n(x, y), \text{ where}$$

$$(17) \quad B_n(x, y) = \int_a^b \dots \int_a^b \begin{vmatrix} K(x, y) & K(x, t_1) & \dots & K(x, t_n) \\ K(t_1, y) & K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(t_n, y) & K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

It is sometimes convenient to write

$$D(x, y; \lambda) = \sum_{n=0}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} B_n(x, y),$$

where we consider $B_0(x, y) = K(x, y)$ and $0! = 1$. The determinant in the expression for B_n satisfies all of the conditions of Hadamard's theorem and hence

$$|B_n| \leq \int_a^b \dots \int_a^b \sqrt{(n+1)^{n+1} M^{n+1}} dt_1 \dots dt_n = \sqrt{(n+1)^{n+1} M^{n+1}} (b-a)^n$$

Then

$$\left| (-1)^n \frac{\lambda^{n+1}}{n!} B_n \right| \leq M^{n+1} (b-a)^n |\lambda|^{n+1} \frac{\sqrt{(n+1)^{n+1}}}{n!} \equiv E_n.$$

This identity defines E_n . We now proceed to show that the series of which E_n is the general term converges. Applying the ratio test,

$$\frac{E_n}{E_{n-1}} = M(b-a)|\lambda| \sqrt{\left(1 + \frac{1}{n}\right)^n} \cdot \frac{1}{\sqrt{n+1}} \cdot \frac{n+1}{n}.$$

The ratio has for its limit zero as n becomes infinite. Hence the series of which E_n is the general term converges. Consequently, the series for $D(x, y; \lambda)$ is absolutely and permanently convergent in λ and, moreover, uniformly convergent in x and y on R . Hence the

Theorem II.—*The series $D(x, y; \lambda)$ converges absolutely and permanently in λ , and, moreover, uniformly¹ on R : $a \leq x \leq b$, $a \leq y \leq b$.*

18. Fredholm's Two Fundamental Relations.—We will now proceed to prove the two relations:

$$(18) \quad D(x, y; \lambda) - \lambda K(x, y)D(\lambda) = \lambda \int_a^b D(x, t; \lambda)K(t, y)dt$$

$$(19) \quad = \lambda \int_a^b K(x, t)D(t, y; \lambda) dt$$

which were previously heuristically obtained.

a) *Relation between the Coefficients $A_n(x, y)$ and $B_n(x, y)$.*— Substitute in (18), in place of $D(x, y; \lambda)$ and $D(\lambda)$, their series expressions. Then both sides of the equality in (18) become power series in λ . Hence, if (18) is true, the coefficients of corresponding powers of λ on the two sides must be equal. Conversely, if we can show that the coefficients of corresponding powers of λ on the two sides are equal, then (18) will be established. Making the substitutions, we obtain

¹ GOURSAT-HEDRICK, "Mathematical Analysis," §173, note.

$$\begin{aligned} \lambda K(x, y) + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} B_n(x, y) \\ - \lambda K(x, y) \left[1 + \sum_{n=1}^{\infty} (-1)^n A_n \frac{\lambda^n}{n!} \right] \\ = \lambda \int_a^b K(t, y) \left[\lambda K(x, t) + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} B_n(x, t) \right] dt. \end{aligned}$$

On the right-hand side, the series in the integrand is uniformly convergent and remains so when multiplied by $K(t, y)$. Therefore, we can integrate by terms and write

$$\lambda^2 \int_a^b K(x, t) K(t, y) dt + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+2}}{n!} \int_a^b B_n(x, t) K(t, y) dt.$$

If in this second integral we put $n' = n + 1$ and then drop the prime, we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} B_n(x, y) - K(x, y) \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^{n+1}}{n!} A_n \\ = \lambda^2 \int_a^b K(x, t) K(t, y) dt \\ + \sum_{n=1}^{\infty} (-1)^{n-1} \frac{\lambda^{n+1}}{(n-1)!} \int_a^b B_{n-1}(x, t) K(t, y) dt. \end{aligned}$$

Compare now the coefficients of corresponding powers of λ on the two sides. We obtain

$$(20) \quad B_n(x, y) = A_n K(x, y) - n \int_a^b B_{n-1}(x, t) K(t, y) dt.$$

If we establish the truth of this equality, then (18) will be shown to be true.

The relation (19) treated in exactly the same way leads to the relation

$$(21) \quad B_n(x, y) = A_n K(x, y) - n \int_a^b K(x, t) B_{n-1}(t, y) dt.$$

If we can establish the truth of this equality, then (19) will be shown to be true.

b) *Proof of (20) and (21).*—We prove (20) and also (21) by showing that the explicit expression for $B_n(x, y)$ can by proper expansion and change in notation be written in the form indicated.

The explicit expression for $B_n(x, y)$ is

$$B_n(x, y) = \int_a^b \dots \int_a^b \begin{vmatrix} K(x, y) & K(x, t_1) & \dots & K(x, t_n) \\ K(t_1, y) & K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(t_n, y) & K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

Develop the determinant which appears in the integrand in terms of the elements of the first column. We obtain

$$\begin{aligned} B_n(x, y) &= \int_a^b \dots \int_a^b K(x, y) \times \\ &\quad \begin{vmatrix} K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \ddots & \vdots \\ K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n \\ &+ \sum_{i=1}^n (-1)^i \int_a^b \dots \int_a^b K(t_i, y) \times \\ &\quad \begin{vmatrix} K(x, t_1) & \dots & K(x, t_n) \\ K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \ddots & \vdots \\ K(t_{i-1}, y) & \dots & K(t_{i-1}, t_n) \\ K(t_{i+1}, y) & \dots & K(t_{i+1}, t_n) \\ \vdots & \ddots & \vdots \\ K(t_n, y) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n. \end{aligned}$$

The first term on the right reduces to $K(x, y) \cdot A_n$, according to (15). In the terms which occur in the summation in place of

$$\text{write} \quad \begin{matrix} t_i, t_{i+1}, t_{i+2}, \dots, t_n \\ t, t_i, t_{i+1}, \dots, t_{n-1}, \end{matrix}$$

which means simply a change in the notation of the variables of integration in a multiple definite integral. We obtain

$$\sum_{i=1}^n (-1)^i \int_a^b \dots \int_a^b K(t, y) \times \begin{vmatrix} K(x, t_1) & \dots & K(x, t_{i-1}) & K(x, t_i) & \dots & K(x, t_{n-1}) \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ K(x, t_{n-1}) & \dots & \dots & \dots & \dots & K(t_{n-1}, t_{n-1}) \end{vmatrix} dt dt_1 \dots dt_{n-1},$$

which may be written by bringing t into the first column

$$\sum_{i=1}^n (-1)^{2i-1} \int_a^b \dots \int_a^b K(t, y) \times \begin{vmatrix} K(x, t) & K(x, t_1) & \dots & K(x, t_{n-1}) \\ K(t_1, t) & K(t_1, t_1) & \dots & K(t_1, t_{n-1}) \\ \dots & \dots & \dots & \dots \\ K(t_{n-1}, t) & K(t_{n-1}, t_1) & \dots & K(t_{n-1}, t_{n-1}) \end{vmatrix} dt dt_1 \dots dt_{n-1}.$$

This last expression shows that all of the n terms of the summation are equal. Furthermore, the integrations may be performed in any order. We then integrate first with respect to t_1, \dots, t_{n-1} . For these integrations, $K(t, y)$ may be regarded as a constant factor and may be taken before the $(n - 1)$ - fold integral, so that we obtain

$$-n \int_a^b K(t, y) \left\{ \int_a^b \dots \int_a^b \begin{vmatrix} K(x, t) & K(x, t_1) & \dots & \dots & K(x, t_{n-1}) \\ K(t_1, t) & K(t_1, t_1) & \dots & \dots & K(t_1, t_{n-1}) \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ K(t_{n-1}, t) & K(t_{n-1}, t_1) & \dots & \dots & K(t_{n-1}, t_{n-1}) \end{vmatrix} dt_1 \dots dt_{n-1} \right\} dt,$$

which, according to (17), may be written

$$-n \int_a^b B_{n-1}(x, t) K(t, y) dt.$$

Hence the development of $B_n(x, y)$ gives

$$B_n(x, y) = A_n \cdot K(x, y) - n \int_a^b B_{n-1}(x, t) K(t, y) dt,$$

which proves relation (20). Therefore (18), which is Fredholm's first relation, is true.

Take again the explicit expression for $B_n(x, y)$, but this time develop the determinant in the integrand in terms of the elements of the first row. Then, proceeding as before, we obtain for the development of $B_n(x, y)$

$$B_n(x, y) = A_n \cdot K(x, y) - n \int_a^b K(x, t) B_{n-1}(t, y) dt,$$

which proves the relation (21). Therefore, (19), which is Fredholm's second relation, is true. Thus we have established the following.

Theorem III.—*Between Fredholm's determinant $D(\lambda)$ and Fredholm's first minor $D(x, y; \lambda)$ the following double relation holds:*

$$(18) \quad D(x, y; \lambda) - \lambda K(x, y) D(\lambda) =$$

$$(19) \quad \begin{aligned} & \lambda \int_a^b D(x, t; \lambda) K(t, y) dt \\ & = \lambda \int_a^b K(x, t) D(t, y; \lambda) dt \end{aligned}$$

for all values of λ and for all values of x and y on R .

19. Fredholm's Solution of the Integral Equation When $D(\lambda) \neq 0$.—Fredholm's two fundamental relations, the equations (18) and (19), enable us now to obtain a solution of the integral equation

$$(1) \quad u(x) = f(x) + \lambda \int_a^b K(x, t)u(t)dt.$$

We obtain a hint as to the method of procedure from the method of solving the finite system

$$(3) \quad u_i - \lambda h \sum_{j=1}^n K_{ij} u_j = f_i \quad (i = 1, \dots, n).$$

To find u_k from this system we first multiply by Δ_{ik} and sum with respect to i . We obtain

$$\sum_{i=1}^n u_i \Delta_{ik} - \lambda h \sum_{i=1}^n \sum_{j=1}^n K_{ij} u_j \Delta_{ik} = \sum_{i=1}^n f_i \Delta_{ik}$$

whence

$$(4) \quad \Delta u_k = \sum_{i=1}^n f_i \Delta_{ik}.$$

Now

$$\Delta_{ik} = h \mathfrak{D}_{ki} \text{ by definition (see §15, c)}$$

and

$$\lim_{h \rightarrow 0} \sum_{i=1}^n h \mathfrak{D}_{ki} = \int_a^b D(x, t; \lambda) dt \text{ by (7).}$$

Let us now follow the analogy. Write (1) in the form

$$u(t) = f(t) + \lambda \int_a^b K(t, \xi)u(\xi)d\xi.$$

Multiply both sides of this equation, which we suppose is satisfied by a continuous function u , by $D(x, t; \lambda)$ and then integrate with respect to t from a to b . We obtain

$$(22) \quad \begin{aligned} \int_a^b D(x, t; \lambda)u(t)dt &= \int_a^b D(x, t; \lambda)f(t)dt \\ &+ \lambda \int_a^b \left[\int_a^b D(x, t; \lambda)K(t, \xi)u(\xi)d\xi \right] dt. \end{aligned}$$

The integrand of the double integral being continuous in x and ξ , we can interchange the order of integration¹ in the double integral and write it

$$\int_a^b u(\xi) \left[\lambda \int_a^b D(x, t; \lambda) K(t, \xi) dt \right] d\xi,$$

which, according to (18), becomes

$$\int_a^b \left[D(x, \xi; \lambda) - \lambda D(\lambda) K(x, \xi) \right] u(\xi) d\xi.$$

Hence (22) may be written

$$\begin{aligned} \int_a^b D(x, t; \lambda) u(t) dt &= \int_a^b D(x, t; \lambda) f(t) dt \\ &\quad + \int_a^b D(x, \xi; \lambda) u(\xi) d\xi - \lambda D(\lambda) \int_a^b K(x, \xi) u(\xi) d\xi, \end{aligned}$$

which, on account of (1), reduces to

$$O = \int_a^b D(x, t; \lambda) f(t) dt - D(\lambda) \left[u(x) - f(x) \right].$$

We solve now for $u(x)$, under the assumption $D(\lambda) \neq 0$. We obtain

$$(23) \quad u(x) = f(x) + \int_a^b \frac{D(x, t; \lambda) f(t)}{D(\lambda)} dt.$$

Hence, if u is a continuous function of x which satisfies (1), and if $D(\lambda) \neq 0$, then $u(x)$ is given by (23).

It remains for us to show that also, conversely, the expression for $u(x)$ given by (23) is a solution of (1). We do this by direct substitution. Substitute the value of $u(x)$ as given by (23) in (1). We obtain

$$\begin{aligned} f(x) + \int_a^b \frac{D(x, t; \lambda) f(t)}{D(\lambda)} dt &= f(x) + \lambda \int_a^b K(x, t) \times \\ &\quad \left\{ f(t) + \int_a^b \frac{D(t, \xi; \lambda) f(\xi)}{D(\lambda)} d\xi \right\} dt. \end{aligned}$$

¹ GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §123.

Break the last term up into two parts and in the double integral change the order of integration. We obtain

$$\int_a^b \frac{D(x, t; \lambda) f(t)}{D(\lambda)} dt = \lambda \int_a^b K(x, t) f(t) dt + \frac{1}{D(\lambda)} \int_a^b f(\xi) \left[\lambda \int_a^b K(x, t) D(t, \xi; \lambda) dt \right] d\xi,$$

which, according to (19), may be written

$$\int_a^b \frac{D(x, t; \lambda) f(t)}{D(\lambda)} dt = \lambda \int_a^b K(x, t) f(t) dt + \frac{1}{D(\lambda)} \int_a^b f(\xi) \left[D(x, \xi; \lambda) - \lambda K(x, \xi) D(\lambda) \right] d\xi.$$

But this last equation is seen to be an identity. Consequently, the expression for $u(x)$ given by (23) satisfies equation (1). Thus we have proved the following theorem, which is called *Fredholm's first fundamental theorem*:

Theorem IV.—If

- (a) $D(\lambda) \neq 0$.
- (b) $K(x, t)$ is continuous in R .
- (c) $f(x)$ is continuous in I .

then the equation

$$(1) \quad u(x) = f(x) + \lambda \int_a^b K(x, t) u(t) dt$$

has one and only one continuous solution given by

$$(23) \quad u(x) = f(x) + \int_a^b \frac{D(x, t; \lambda) f(t)}{D(\lambda)} dt$$

where $D(x, t; \lambda)$ and $D(\lambda)$ are absolutely and permanently convergent integral series in λ , and $D(x, t; \lambda)$ converges uniformly with respect to x and t on $R: a \leq x \leq b, a \leq y \leq b$.

We have at once, for the special case $f \equiv 0$, the following:

Corollary.—If $D(\lambda) \neq 0$, **then the homogeneous equation**

$$(24) \quad u(x) = \lambda \int_a^b K(x, t) u(t) dt$$

has one and only one continuous solution given by $u(x) \equiv 0$.

Let us point out the analogy with the finite system of linear equations

$$u_i - \lambda h \sum_{j=1}^n K_{ij} u_j = f_i \quad (i = 1, \dots, n)$$

with determinant Δ .

If $\Delta \neq 0$, then this system has one and only one solution.

If $f_i \equiv 0$, then the only solution is the trivial one

$$u_1 = \dots = u_n \equiv 0.$$

But the limit of Δ was $D(\lambda)$. Hence the results of Theorem IV and its corollary are exactly what was to be expected from the analogy with the finite system.

20. Solution of the Homogeneous Equation When $D(\lambda) = 0$, $D'(\lambda) \neq 0$.—The discussion up to this point has been made under the assumption $D(\lambda) \neq 0$. Let us now see what happens when $D(\lambda) = 0$, first with respect to the homogeneous equation (24).

Let λ_o be a value of λ for which

$$(25) \quad D(\lambda_o) = 0.$$

We now consider the solution of the homogeneous integral equation (24) for this particular value of λ :

$$(26) \quad u(x) = \lambda_o \int_a^b K(x, t)u(t)dt.$$

We obtain a solution of (26) by means of Fredholm's second fundamental relation (19), which is true for all values of λ and hence for $\lambda = \lambda_o$. With this value of λ , on account of (25), equation (19) becomes

$$D(x, y; \lambda_o) = \lambda_o \int_a^b K(x, t)D(t, y; \lambda_o)dt.$$

The equality holds for every value of y on the interval (ab) and, therefore, for $y = y_o$. Then

$$D(x, y_o; \lambda_o) = \lambda_o \int_a^b K(x, t)D(t, y_o; \lambda_o)dt.$$

But this is just the equation (26) with $u(x)$ replaced by $D(x, y_o; \lambda_o)$. Thus we see that $u(x) = D(x, y_o; \lambda_o)$ is a solution of (26). Moreover, this solution is continuous,¹ for $D(x, y; \lambda)$ is uniformly convergent in x and y and its terms are continuous. But $D(x, y_o; \lambda_o)$ may be identically zero in x , either on account of an unfortunate choice of y_o , in which case we could choose some other value for y_o , or because $D(x, y; \lambda_o) \equiv 0$ in x and y , in which case the above solution reduces to the trivial one $u \equiv 0$, no matter how we choose y_o . We have thus proved the following:

Theorem V.—*If $D(\lambda_o) = 0$ and $D(x, y; \lambda_o) \not\equiv 0$, then for a proper choice of y_o , $u(x) = D(x, y_o; \lambda_o)$ is a continuous solution of*

$$u(x) = \lambda_o \int_a^b K(x, t) u(t) dt$$

and $u(x) \not\equiv 0$.

In the theorem just stated, the condition $D(x, y; \lambda_o) \not\equiv 0$ may be replaced by the condition $D'(\lambda) \not\equiv 0$. To show this we prove the following

$$(27) \quad \int_a^b D(x, x; \lambda) dx = -\lambda D'(\lambda).$$

We prove (27) by making use of the series expressions for $D'(\lambda)$ and $D(x, y; \lambda)$. We have from (14)

$$D(\lambda) = 1 - \lambda A_1 + \frac{\lambda^2}{2!} A_2 - \frac{\lambda^3}{3!} A_3 + \dots$$

where the A_n are given by (15). Then

$$\begin{aligned} D'(\lambda) &= -A_1 + \lambda A_2 - \frac{\lambda^2}{2!} A_3 + \dots \\ &= -\sum_{n=0}^{\infty} (-1)^n \frac{\lambda^n}{n!} A_{n+1}. \end{aligned}$$

¹ GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §173.

In the expression for A_{n+1} :

$$A_{n+1} = \int_a^b \dots \int_a^b \begin{vmatrix} k(t_1, t_1)K(t_1, t_2) & \dots & K(t_1, t_{n+1}) \\ K(t_2, t_1)K(t_2, t_2) & \dots & K(t_2, t_{n+1}) \\ \vdots & \ddots & \vdots \\ K(t_{n+1}, t_1)K(t_{n+1}, t_2) & \dots & K(t_{n+1}, t_{n+1}) \end{vmatrix} dt_1 \dots dt_{n+1},$$

in place of $t_1, t_2, t_3, \dots, t_n, t_{n+1}$
put $x, t_1, t_2, \dots, t_{n-1}, t_n$.

Then

$$A_{n+1} = \int_a^b \dots \int_a^b \begin{vmatrix} K(x, x) K(x, t_1) & \dots & K(x, t_n) \\ K(t_1, x) K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \ddots & \vdots \\ K(t_n, x) K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dx dt_1 \dots dt_n.$$

In this multiple definite integral we change the order of integration¹ and integrate first with respect to $dt_1 \dots dt_n$, whence

$$A_{n+1} = \int_a^b \left\{ \int_a^b \dots \int_a^b \begin{vmatrix} K(x, x) K(x, t_1) & \dots & K(x, t_n) \\ K(t_1, x) K(t_1, t_1) & \dots & K(t_1, t_n) \\ \vdots & \ddots & \vdots \\ K(t_n, x) K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n \right\} dx,$$

which, on account of (17), becomes

$$A_{n+1} = \int_a^b B_n(x, x) dx.$$

¹ GOURSAT-HEDRICK "Mathematical Analysis," vol 1, §123; PIER-PONT, JAMES, "The Theory of Functions of Real Variables," vol. 1, §570.

Therefore,

$$D'(\lambda) = - \sum_{n=0}^{\infty} (-1)^n \frac{\lambda^n}{n!} \int_a^b B_n(x, x) dx.$$

But

$$\sum_{n=0}^{\infty} (-1)^n \frac{\lambda^n}{n!} B_n(x, x) dx$$

is a series uniformly convergent in x . We can then interchange the order of writing the summation and the integration¹ in the expression for $D'(\lambda)$ and write

$$D'(\lambda) = - \int_a^b \sum_{n=0}^{\infty} (-1)^n \frac{\lambda^n}{n!} B_n(x, x) dx.$$

Multiply both sides by $-\lambda$, and then by (16) we see that, indeed,

$$(27) \quad \int_a^b D(x, x; \lambda) dx = -\lambda D'(\lambda).$$

Let us suppose now that $D(\lambda_0) = 0$ and $D'(\lambda_0) \neq 0$, then certainly $\lambda_0 \neq 0$, since $D(0) = 1$. Hence, if we write (27) with $\lambda = \lambda_0$, the right-hand side of this equality is different from zero and so also the left-hand side, and, therefore, $D(x, x; \lambda_0) \neq 0$ in x and, consequently, $D(x, y; \lambda_0) \neq 0$ in x and y . Hence, indeed, in Theorem V the condition $D(x, y; \lambda_0) \neq 0$ may be replaced by $D'(\lambda_0) \neq 0$.

We remark further that if $u(x) = D(x, y_0; \lambda_0)$ is a solution of the homogeneous integral equation (26), then Cu (C , an arbitrary constant) is also a solution. Thus, there are an infinitude of solutions which differ only by a constant factor. We shall later show that there are no other solutions. This is in analogy with the finite system of linear equations

$$u_i - \lambda h \sum_{j=1}^n K_{ij} u_j = 0 \quad (i = 1, \dots, n)$$

with determinant Δ . If $\Delta = 0$, while not all of the first minors Δ_{ik} vanish, then these n equations determine uniquely the ratios of the u_1, \dots, u_n , that is, $u_j = C_j u_n$

¹ GOURSAT-HEDRICK, "Mathematical Analysis", vol. 1, §174.

($j = 1, \dots, n; C_n = 1$). Now, $\Delta = 0$ corresponds to $D(\lambda) = 0$, while *not all* Δ_{ik} vanish corresponds to $D(x, y; \lambda_o) \neq 0$.

21. Solution of the Homogeneous Integral Equation When $D(\lambda) = 0$.—It remains to consider the case

$$D(\lambda_o) = 0, D(x, y; \lambda_o) \equiv 0.$$

which corresponds in the linear system to the case where Δ and all its first minors are zero, and, where it becomes necessary, to consider the minors of higher order of Δ . Accordingly, we have to consider, in the treatment of the integral equation, the limits of these higher minors of Δ as h approaches zero.

In what follows let us use with Heywood-Fréchet ("L'Équation De Fredholm," page 53) the notation

$$K\begin{pmatrix} s_1, s_2, \dots, s_n \\ t_1, t_2, \dots, t_n \end{pmatrix} \equiv \begin{vmatrix} K(s_1, t_1) & \dots & K(s_1, t_n) \\ \dots & \dots & \dots \\ K(s_n, t_1) & \dots & K(s_n, t_n) \end{vmatrix}.$$

(a) *Definition of the pth Minor of $D(\lambda)$.*—Let

$$(28) \quad B_n\begin{pmatrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{pmatrix} = \int_a^b \dots \int_a^b K\begin{pmatrix} x_1, \dots, x_p, t_1, \dots, t_n \\ y_1, \dots, y_p, t_1, \dots, t_n \end{pmatrix} dt_1 \dots dt_n,$$

with

$$(29) \quad B_o\begin{pmatrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{pmatrix} = K\begin{pmatrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{pmatrix}.$$

Then the *pth minor of $D(\lambda)$* is defined by the infinite series

$$(30) \quad D\begin{pmatrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{pmatrix} \lambda = \sum_{n=0}^{\infty} (-1)^n \frac{\lambda^{n+p}}{n!} B_n\begin{pmatrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{pmatrix} \equiv D_p(x, y; \lambda),$$

which for $p = 1$ reduces to $D(x, y; \lambda)$.

By means of Hadamard's theorem we prove, exactly as before, the

Theorem VI.—*The infinite series for $D\left(\begin{matrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{matrix}, \lambda\right)$*

is absolutely and permanently convergent in λ , and uniformly convergent in $x_1, \dots, x_p, y_1, \dots, y_p$ for $a \leq x_\alpha \leq b$, $a \leq y_\beta \leq b$ ($\alpha, \beta = 1, \dots, p$).

Corollary.—*When two of the x 's become equal, say $x_r = x_s$, or when two of the y 's become equal, say $y_i = y_j$, then $D_p(x, y; \lambda)$ vanishes.* For then in the integrand of $B_n\left(\begin{matrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{matrix}\right)$ two rows (columns) become equal. Therefore,

$$B_n\left(\begin{matrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{matrix}\right) = 0, \text{ and hence } D_p(x, y; \lambda) = 0.$$

In like manner, if in $D_p(x, y; \lambda)$ two of the x 's or two of the y 's are interchanged, $D_p(x, y; \lambda)$ changes sign.

b) Generalization of the Two Fundamental Relations.—

Expand the determinant in $B_n\left(\begin{matrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{matrix}\right)$ according to the elements of the column y_β :

$$\begin{aligned} B_n\left(\begin{matrix} x_1, \dots, x_p \\ y_1, \dots, y_p \end{matrix}\right) &= \int_a^b \dots \int_a^b \left\{ \sum_{\alpha=1}^p (-1)^{\alpha+\beta} \times \right. \\ &K(x_\alpha, y_\beta) K\left(\begin{matrix} x_1, \dots, x_{\alpha-1}, x_{\alpha+1}, \dots, x_p, t_1, \dots, t_n \\ y_1, \dots, y_{\beta-1}, y_{\beta+1}, \dots, y_p, t_1, \dots, t_n \end{matrix}\right) \\ &+ \sum_{i=1}^n (-1)^{p+i+\beta} K(t_i, y_\beta) \times \\ &\left. K\left(\begin{matrix} x_1, \dots, x_p, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n \\ y_1, \dots, y_{\beta-1}, y_{\beta+1}, \dots, y_p, t_1, \dots, t_n \end{matrix}\right) \right\} dt_1 \dots dt_n. \end{aligned}$$

In the first sum, $K(x_\alpha, y_\beta)$ may be taken before the integral sign, and, according to (28), the sum becomes

$$\sum_{\alpha=1}^p (-1)^{\alpha+\beta} K(x_\alpha, y_\beta) B_n\left(\begin{matrix} x_1, \dots, x_{\alpha-1}, x_{\alpha+1}, \dots, x_p \\ y_1, \dots, y_{\beta-1}, y_{\beta+1}, \dots, y_p \end{matrix}\right).$$

In the second sum make a change of notation.

In place of $t_i, t_{i+1}, t_{i+2}, \dots, t_n$
write $t, t_i, t_{i+1}, \dots, t_{n-1}$

The i th term of the second sum becomes

$$(-1)^{p+i+\beta} K(t, y_\beta) \times \\ K \left(\begin{matrix} x_1, \dots, \dots, \dots, x_p, t_1, \dots, \\ y_1, \dots, y_{\beta-1}, y_{\beta+1}, \dots, y_p, t_1, \dots, \\ \dots, \dots, \dots, t_{n-1}, \\ , t_{i-1}, t, t_i, \dots, t_{n-1} \end{matrix} \right).$$

Bring the column t between the columns $y_{\beta-1}, y_{\beta+1}$ by $i + p - \beta - 1$ transpositions of columns. We get, then, for the i th term of the second sum

$$- K(t, y_\beta) \times \\ K \left(\begin{matrix} x_1, \dots, x_{\beta-1}, x_\beta, x_{\beta+1}, \dots, x_p, t_1, \dots, t_{n-1} \\ y_1, \dots, y_{\beta-1}, t, y_{\beta+1}, \dots, y_p, t_1, \dots, t_{n-1} \end{matrix} \right).$$

Hence, all terms in $\sum_{i=1}^n$ are equal and this sum may be written, if, moreover, we integrate first with respect to t_1, \dots, t_{n-1} ,

$$- n \int_a^b K(t, y_\beta) \left\{ \int_a^b \dots \int_a^b \right. \\ K \left(\begin{matrix} x_1, \dots, \dots, \dots, x_p, t_1, \dots, t_{n-1} \\ y_1, \dots, y_{\beta-1}, t, y_{\beta+1}, \dots, y_p, t_1, \dots, t_{n-1} \end{matrix} \right) \\ \left. dt_1 \dots dt_{n-1} \right\} dt,$$

which, according to (28), reduces to

$$- n \int_a^b K(t, y_\beta) \times \\ B_{n-1} \left(\begin{matrix} x_1, \dots, x_{\beta-1}, x_\beta, x_{\beta+1}, \dots, x_p \\ y_1, \dots, y_{\beta-1}, t, y_{\beta+1}, \dots, y_p \end{matrix} \right) dt.$$

Thus we arrive at the formula

$$(31) \quad B_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} =$$

$$\sum_{\alpha=1}^p (-1)^{\alpha+\beta} K(x_\alpha, y_\beta) B_n \begin{pmatrix} x_1, & \dots, & x_{\alpha-1}, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\beta-1}, & y_{\beta+1}, & \dots, & y_p \end{pmatrix}$$

$$- n \int_a^b K(t, y_\beta) B_{n-1} \begin{pmatrix} x_1, & \dots, & x_{\alpha-1}, & x_\alpha, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\beta-1}, & t, & y_{\beta+1}, & \dots, & y_p \end{pmatrix} dt.$$

In like manner, by expanding the integrand of $B_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix}$ according to the elements of the row x_α , we obtain

$$(32) \quad B_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} =$$

$$\sum_{\beta=1}^p (-1)^{\alpha+\beta} K(x_\alpha, y_\beta) B_n \begin{pmatrix} x_1, & \dots, & x_{\alpha-1}, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\beta-1}, & y_{\beta+1}, & \dots, & y_p \end{pmatrix}$$

$$- n \int_a^b K(x_\alpha, t) B_{n-1} \begin{pmatrix} x_1, & \dots, & x_{\alpha-1}, & t, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\beta-1}, & y_\beta, & y_{\beta+1}, & \dots, & y_p \end{pmatrix} dt.$$

If now we multiply both sides of (31) and (32) by $(-1)^n \frac{\lambda^{n+p}}{n!}$ and sum with respect to n from $n = 0$ to $n = \infty$, we obtain, on account of the definition given in (30), the following double relation, which is a *generalization of Fredholm's two fundamental relations* (10) and (11).

$$(33) \quad D \begin{pmatrix} x_1, & \dots, & x_p, \lambda \\ y_1, & \dots, & y_p \end{pmatrix} =$$

$$\sum_{\alpha=1}^p (-1)^{\alpha+\beta} \lambda K(x_\alpha, y_\beta) D \begin{pmatrix} x_1, & \dots, & x_{\alpha-1}, & x_{\alpha+1}, & \dots, & x_p, \lambda \\ y_1, & \dots, & y_{\beta-1}, & y_{\beta+1}, & \dots, & y_p \end{pmatrix}$$

$$+ \lambda \int_a^b K(t, y_\beta) D \begin{pmatrix} x_1, & \dots, & x_{\beta-1}, & x_\beta, & x_{\beta+1}, & \dots, & x_p, \lambda \\ y_1, & \dots, & y_{\beta-1}, & t, & y_{\beta+1}, & \dots, & y_p \end{pmatrix} dt$$

$$(34) = \sum_{\beta=1}^p (-1)^{\alpha+\beta} \lambda K(x_\alpha, y_\beta)$$

$$D\left(\begin{matrix} x_1, & \dots, & x_{\alpha-1}, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\beta-1}, & y_{\beta+1}, & \dots, & y_p \end{matrix} \lambda\right)$$

$$+ \lambda \int_a^b K(x_\alpha, t) D\left(\begin{matrix} x_1, & \dots, & x_{\alpha-1}, & t, & x_{\alpha+1}, & \dots, & x_p \\ y_1, & \dots, & y_{\alpha-1}, & y_\alpha, & y_{\alpha+1}, & \dots, & y_p \end{matrix} \lambda\right) dt.$$

c) Relation between $D^{(p)}(\lambda)$ and $D\left(\begin{matrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{matrix} \lambda\right)$.—

The relation (27) between $D'(\lambda)$ and $D(x, x; \lambda)$ generalizes as follows:

$$(35) \quad \int_a^b \dots \int_a^b D\left(\begin{matrix} x_1, & \dots, & x_p \\ x_1, & \dots, & x_p \end{matrix} \lambda\right) dx_1 \dots dx_p =$$

$$(-1)^p \lambda^p \frac{d^p D(\lambda)}{d\lambda^p}.$$

Proof.—From the series expression for $D(\lambda)$:

$$D(\lambda) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^n}{n!} A_n,$$

we get

$$D^{(p)}(\lambda) = \sum_{n=p}^{\infty} (-1)^n \frac{\lambda^{n-p}}{(n-p)!} A_n,$$

which, after the change in notation $n - p = n'$ and a final dropping of the prime ('), may be written

$$D^{(p)}(\lambda) = \sum_{n=0}^{\infty} (-1)^{n+p} \frac{\lambda^n}{n!} A_{n+p}.$$

But by (15)

$$A_{n+p} = \int_a^b \dots \int_a^b K\left(\begin{matrix} t_1, & \dots, & t_{n+p} \\ t_1, & \dots, & t_{n+p} \end{matrix}\right) dt_1 \dots dt_{n+p}.$$

If now

in place of $t_1, \dots, t_p, t_{p+1}, \dots, t_{p+n}$
we write $x_1, \dots, x_p, t_1, \dots, t_n$

and then change the order of integration, as we may, so that we integrate first with respect to t_1, \dots, t_n , we obtain

$$A_{n+p} = \int_a^b \dots \int_a^b \left\{ \int_a^b \dots \int_a^b K \begin{pmatrix} x_1, \dots, x_p, t_1, \dots, t_n \\ x_1, \dots, x_p, t_1, \dots, t_n \end{pmatrix} dt_1 \dots dt_n \right\} dx_1 \dots dx_p,$$

which, on account of (28), becomes

$$A_{n+p} = \int_a^b \dots \int_a^b B_n \begin{pmatrix} x_1, \dots, x_p \\ x_1, \dots, x_p \end{pmatrix} dx_1 \dots dx_p.$$

Multiply both sides of this equality by $(-1)^n \frac{\lambda^{n+p}}{n!}$ and sum with respect to n from $n = 0$ to $n = \infty$. We obtain

$$(-1)^p \lambda^p \frac{d^p D(\lambda)}{d\lambda^p} = \sum_{n=0}^{\infty} \int_a^b \dots \int_a^b (-1)^n \frac{\lambda^{n+p}}{n!} B_n \begin{pmatrix} x_1, \dots, x_p \\ x_1, \dots, x_p \end{pmatrix} dx_1 \dots dx_p.$$

It is permissible here to put the summation under the multiple integral sign. If we then make use of the equation of definition (30), we obtain

$$(-1)^p \lambda^p \frac{d^p D(\lambda)}{d\lambda^p} = \int_a^b \dots \int_a^b D \begin{pmatrix} x_1, \dots, x_p \\ x_1, \dots, x_p \end{pmatrix} \lambda dx_1 \dots dx_p,$$

which establishes the equality (35).

We make use of this result to prove that not all of the Fredholm minors vanish. Let λ_o be a root of $D(\lambda) = 0$. Then certainly $\lambda_o \neq 0$ for $D(0) = 1$. Furthermore, λ_o is a root of $D(\lambda)$ of finite multiplicity r ($r \geq 1$), defined by $D(\lambda_o) = 0, D'(\lambda_o) = 0, \dots, D^{(r-1)}(\lambda_o) = 0, D^{(r)}(\lambda_o) \neq 0$. The multiplicity r of the root must be finite, otherwise

$$D(\lambda) \equiv 0.$$

In (35) put $\lambda = \lambda_o$ and $p = r$, then the right member by our hypothesis does not vanish. Therefore, the left member does not vanish. Hence, since $\lambda_o \neq 0$,

$$D\left(\begin{matrix} x_1, & \dots, & x_r \\ x_1, & \dots, & x_r \end{matrix} \lambda_o\right) \not\equiv 0 \text{ in } x_1, \dots, x_r$$

and, consequently,

$$D\left(\begin{matrix} x_1, & \dots, & x_r \\ y_1, & \dots, & y_r \end{matrix} \lambda_o\right) \not\equiv 0 \text{ in } x_1, \dots, x_r, y_1, \dots, y_r.$$

Hence, in the series

$$D(\lambda_o) = 0, D\left(\begin{matrix} x \\ y \end{matrix} \lambda_o\right), D\left(\begin{matrix} x_1, & x_2 \\ y_1, & y_2 \end{matrix} \lambda_o\right), D\left(\begin{matrix} x_1, & x_2, & x_3 \\ y_1, & y_2, & y_3 \end{matrix} \lambda_o\right), \dots$$

we must come to a number $q \leq r$, called the *index of λ_o* , such that

$$D(\lambda_o) = 0, D(x, y; \lambda_o) \equiv 0, \dots,$$

$$D\left(\begin{matrix} x_1, & \dots, & x_{q-1} \\ y_1, & \dots, & y_{q-1} \end{matrix} \lambda_o\right) \equiv 0, D\left(\begin{matrix} x_1, & \dots, & x_q \\ y_1, & \dots, & y_q \end{matrix} \lambda_o\right) \not\equiv 0.$$

That is, there exists a particular set of values $x'_1, \dots, x'_q, y'_1, \dots, y'_q$ for the variables $x_1, \dots, x_q, y_1, \dots, y_q$ such that we have the following numerical inequality:

$$D\left(\begin{matrix} x'_1, & \dots, & x'_q \\ y'_1, & \dots, & y'_q \end{matrix} \lambda_o\right) \not\equiv 0.$$

Incidentally, we have proved the

Theorem VII.—*The index q of a root λ_o of $D(\lambda)$ is at most equal to the multiplicity r of λ_o : $q \leq r$.*

d) The q Independent Solutions of the Homogeneous Equation.—Let λ_o be a root of $D(\lambda) = 0$ of index q , so that

$$D(\lambda_o) = 0, D(x, y; \lambda_o) \equiv 0, \dots,$$

$$D\left(\begin{matrix} x_1, & \dots, & x_{q-1} \\ y_1, & \dots, & y_{q-1} \end{matrix} \lambda_o\right) \equiv 0$$

$$\text{but } D\left(\begin{matrix} x_1, & \dots, & x_q \\ y_1, & \dots, & y_q \end{matrix} \lambda_o\right) \not\equiv 0.$$

Write down the second of Fredholm's generalized fundamental relations (34) for $\lambda = \lambda_o$, $p = q$

$$\begin{aligned} x_1 &= x'_1, \dots, x_{\alpha-1} = x'_{\alpha-1}, x_\alpha = x, x_{\alpha+1} = x'_{\alpha+1}, \\ &\quad \dots, x_q = x'_q \\ y_1 &= y'_1, \dots, y_{\beta-1} = y'_{\beta-1}, y_\beta = y'_\beta, y_{\beta+1} = y'_{\beta+1}, \\ &\quad \dots, y_q = y'_q \end{aligned}$$

Then

$$\begin{aligned} D\left(\begin{matrix} x'_1, \dots, x'_{\alpha-1}, x, x'_{\alpha+1}, \dots, x'_q, \lambda_o \\ y'_1, \dots, y'_{\beta-1}, y'_\beta, y'_{\beta+1}, \dots, y'_q \end{matrix}\right) &= \\ \lambda_o \int_a^b K(x, t) D\left(\begin{matrix} x'_1, \dots, x'_{\alpha-1}, t, x'_{\alpha+1}, \dots, x'_q, \lambda_o \\ y'_1, \dots, y'_{\beta-1}, y'_\beta, y'_{\beta+1}, \dots, y'_q \end{matrix}\right) dt, \end{aligned}$$

since by hypothesis

$$D\left(\begin{matrix} x_1, \dots, x_{\alpha-1}, x_{\alpha+1}, \dots, x_q, \lambda_o \\ y_1, \dots, y_{\beta-1}, y_{\beta+1}, \dots, y_q \end{matrix}\right) = 0.$$

Hence, if we divide by $D\left(\begin{matrix} x'_1, \dots, x'_q, \lambda_o \\ y'_1, \dots, y'_q \end{matrix}\right)$

and put

$$(36) \quad D\left(\begin{matrix} x'_1, \dots, x'_{\alpha-1}, x, x'_{\alpha+1}, \dots, x'_q, \lambda_o \\ y'_1, \dots, y'_{\beta-1}, y'_\beta, y'_{\beta+1}, \dots, y'_q \end{matrix}\right) = \varphi_\alpha(x, \lambda_o) D\left(\begin{matrix} x'_1, \dots, x'_q, \lambda_o \\ y'_1, \dots, y'_q \end{matrix}\right)$$

we have

$$\varphi_\alpha(x, \lambda_o) = \lambda_o \int_a^b K(x, t) \varphi_\alpha(t, \lambda_o) dt,$$

which expresses that the q functions

$$\varphi_1(x, \lambda_o), \varphi_2(x, \lambda_o), \dots, \varphi_q(x, \lambda_o)$$

are solutions of the homogeneous equation (26). These solutions are continuous and

$$(37) \quad \varphi_\alpha(x_\beta, \lambda_o) = \begin{cases} 1, & \beta = \alpha \\ 0, & \beta \neq \alpha \end{cases}.$$

from (30) and (28). Furthermore, these functions are linearly independent. That is, if a relation of the form

$$C_1\varphi_1(x) + C_2\varphi_2(x) + \dots + C_q\varphi_q(x) \equiv 0$$

exists, where C_1, \dots, C_q are constants, then we must have $C_1 = C_2 = \dots = C_q = 0$. For, from (37), if $x = x_\alpha'$, then $C_\alpha = 0$.

From the homogeneity of (26) it follows that

$$(38) \quad u(x) = C_1\varphi_1(x) + \dots + C_q\varphi_q(x)$$

is again a solution of (26) for arbitrary values of C_1, \dots, C_q . We thus have a q -fold infinitude of solutions.

e) *Completeness Proof.*—It remains to show that every solution of (26) can be put in the form (38).

If $v(x)$ is any solution of (26), then

$$v(x) = \lambda_o \int_a^b K(x, t)v(t)dt.$$

Whence

$$0 \equiv \int_a^b H(x, t) \left\{ v(t) - \lambda_o \int_a^b K(t, s)v(s)ds \right\} dt,$$

where $H(x, t)$ is any continuous function. On subtracting the second equation from the first, we obtain

$$(39) \quad v(x) = \lambda_o \int_a^b N(x, t)v(t)dt$$

where

$$\lambda_o N(x, t) = \lambda_o K(x, t) - \left\{ H(x, t) - \lambda_o \int_a^b H(x, s)K(s, t)ds \right\}.$$

Now apply (33) with $p = q + 1$, $x_{q+1} = x$, $y_{q+1} = y$ and notice that a transposition of two x 's or two y 's changes the sign of D . Then

$$\begin{aligned} D\left(\begin{matrix} x, x_1, \dots, x_q & \lambda_o \\ y, y_1, \dots, y_q & \end{matrix}\right) &= \lambda K(x, y)D\left(\begin{matrix} x_1, \dots, x_q & \lambda \\ y_1, \dots, y_q & \end{matrix}\right) \\ &- \sum_{\alpha=1}^q \lambda K(x_\alpha, y)D\left(\begin{matrix} x_1, \dots, x_{\alpha-1}, x, x_{\alpha+1}, \dots, x_q & \lambda \\ y_1, \dots, y_{\beta-1}, y_\beta, y_{\beta+1}, \dots, y_q & \end{matrix}\right) \\ &+ \lambda \int_a^b K(t, y)D\left(\begin{matrix} x, x_1, \dots, x_q & \lambda \\ t, y_1, \dots, y_q & \end{matrix}\right) dt. \end{aligned}$$

Put $x_1 = x'_1, \dots, x_q = x'_q, y_1 = y'_1, \dots, y_q = y'_q$,
 $y = t, \lambda = \lambda_o$, divide by

$$D\left(\begin{matrix} x'_1, & \dots, & x'_q, \\ y'_1, & \dots, & y'_q, \\ & & \lambda_o \end{matrix}\right)$$

and put

$$(40) \quad H(x, y) = \frac{D\left(\begin{matrix} x, & x'_1, & \dots, & x'_q, \\ y, & y'_1, & \dots, & y'_q, \\ & & \lambda_o \end{matrix}\right)}{D\left(\begin{matrix} x'_1, & \dots, & x'_q, \\ y'_1, & \dots, & y'_q, \\ & & \lambda_o \end{matrix}\right)},$$

then we obtain

$$(41) \quad \sum_{\alpha=1}^q \lambda_o K(x'_\alpha, t) \varphi_\alpha(x) = \lambda_o K(x, t) - H(x, t) \\ + \lambda_o \int_a^b H(x, s) K(s, t) ds.$$

The equation (39) can now be written

$$v(x) = \lambda_o \sum_{\alpha=1}^q \varphi_\alpha(x) \int_a^b K(x'_\alpha, t) v(t) dt.$$

This shows that $v(x)$ can be written in the form (38) by taking for the constants C_α the values

$$C_\alpha = \lambda_o \int_a^b K(x'_\alpha, t) v(t) dt.$$

Thus we obtain *Fredholm's second fundamental theorem*:

Theorem VIII.—If $\lambda = \lambda_o$ is a root of $D(\lambda) = 0$ of order q , then the homogeneous integral equation

$$(26) \quad u(x) = \lambda_o \int_a^b K(x, t) u(t) dt$$

has q linearly independent solutions in terms of which every other solution is expressible linearly and homogeneously. Such a system of q independent solutions is given by

$$\varphi_\alpha(x) = \frac{D\left(\begin{matrix} x'_1, & \dots, & x'_{\alpha-1}, & x, & x'_{\alpha+1}, & \dots, & x'_q, \\ y'_1, & \dots, & y'_{\beta-1}, & y_\beta, & y'_{\beta+1}, & \dots, & y'_q, \\ & & & & & & \lambda_o \end{matrix}\right)}{D\left(\begin{matrix} x'_1, & \dots, & x'_q, \\ y'_1, & \dots, & y'_q, \\ & & \lambda_o \end{matrix}\right)} \quad (\alpha = 1, \dots, q).$$

22. Characteristic Constant. Fundamental Functions.

Definitions.—If $D(\lambda)$ is the Fredholm's determinant for the kernel $K(x, t)$ and $D(\lambda_o) = 0$, then λ_o is said to be a *characteristic constant* of the kernel $K(x, t)$. Further, if $\varphi(x)$ is continuous and not identically zero on the interval (ab) and

$$\varphi(x) = \lambda_o \int_a^b K(x, t)\varphi(t)dt,$$

then $\varphi(x)$ is called a *fundamental function* of the kernel $K(x, t)$, belonging to the characteristic constant λ_o .

$\varphi_1(x), \dots, \varphi_q(x)$ form a *complete system* of fundamental functions of the kernel $K(x, t)$, belonging to λ_o , if every other solution is expressible linearly in terms of these q solutions. Thus, if ψ_1, \dots, ψ_q are any other q solutions, we must have

$$\begin{aligned}\psi_1 &= C_{11}\varphi_1 + \dots + C_{1q}\varphi_q \\ &\vdots \\ \psi_q &= C_{q1}\varphi_1 + \dots + C_{qq}\varphi_q\end{aligned}$$

and if

$$\begin{vmatrix} C_{11} & \dots & C_{1q} \\ \vdots & \ddots & \vdots \\ C_{q1} & \dots & C_{qq} \end{vmatrix} \neq 0$$

ψ_1, \dots, ψ_q form again a complete system of fundamental functions belonging to λ_o .

23. The Associated Homogeneous Integral Equation.—Preparatory to the discussion of the non-homogeneous integral equation, for $D(\lambda) = 0$, we will discuss the homogeneous integral equation

$$(42) \quad v(x) = \lambda_o \int_a^b K(t, x)v(t)dt,$$

which is called the integral equation *associated* with the integral equation

$$(26) \quad u(x) = \lambda_o \int_a^b K(x, t)u(t)dt.$$

Notice that the kernel

$$\bar{K}(x, t) \equiv K(t, x)$$

of the associated equation is derived from the original kernel $K(x, t)$ by interchanging the arguments x and t . There exist important relations between the solutions of the two equations (42) and (26). To obtain them, we first compute the Fredholm determinant and Fredholm minors for the kernel $\bar{K}(x, t)$, which we indicate by the corresponding dashed notation.

a) *Fredholm's Determinant for the Associated Kernel.*—

$$D(\lambda) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^n}{n!} A_n$$

where

$$A_n = \int_a^b \dots \int_a^b \begin{vmatrix} K(t_1, t_1) & \dots & K(t_1, t_n) \\ \dots & \dots & \dots \\ K(t_n, t_1) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

Then

$$\bar{D}(\lambda) = 1 + \sum_{n=1}^{\infty} (-1)^n \frac{\lambda^n}{n!} \bar{A}_n$$

where

$$\bar{A}_n = \int_a^b \dots \int_a^b \begin{vmatrix} (\bar{K}t_1, t_1) & \dots & \bar{K}(t_1, t_n) \\ \dots & \dots & \dots \\ \bar{K}(t_n, t_1) & \dots & \bar{K}(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

In this expression for \bar{A}_n put $\bar{K}(x, t) = K(t, x)$. Then

$$\bar{A}_n = \int_a^b \dots \int_a^b \begin{vmatrix} K(t_1, t_1) & \dots & K(t_n, t_1) \\ \dots & \dots & \dots \\ \dots & \dots & \dots \\ K(t_1, t_n) & \dots & K(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

The determinant which appears in the integrand of \bar{A}_n is the same as that which appears in the integrand of the expression for A_n , with the exception that the rows and columns are interchanged. But this interchange leaves the value of the determinant unaltered. Therefore $\bar{A}_n = A_n$, and hence

$$(43) \quad \bar{D}(\lambda) \equiv D(\lambda).$$

Hence also we conclude that $K(t, x)$ and $K(x, t)$ have the same characteristic constants.

b) *Fredholm's Minors for the Associated Kernel.*—We have from (30)

$$D\begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \\ \lambda \end{pmatrix} = \sum_{n=0}^{\infty} (-1)^n \frac{\lambda^{n+p}}{n!} B_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix}$$

where the B_n are given by (28). Then

$$\bar{B}_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} = \int_a^b \dots \int_a^b \begin{vmatrix} \bar{K}(x_1, y_1) & \dots & \bar{K}(x_1, y_p) \bar{K}(x_1, t_1) & \dots & \bar{K}(x_1, t_n) \\ \dots & \dots & \dots & \dots & \dots \\ \bar{K}(x_p, y_1) & \dots & \bar{K}(x_p, y_p) \bar{K}(x_p, t_1) & \dots & \bar{K}(x_p, t_n) \\ \bar{K}(t_1, y_1) & \dots & \bar{K}(t_1, y_p) \bar{K}(t_1, t_1) & \dots & \bar{K}(t_1, t_n) \\ \dots & \dots & \dots & \dots & \dots \\ \bar{K}(t_n, y_1) & \dots & \bar{K}(t_n, y_p) \bar{K}(t_n, t_1) & \dots & \bar{K}(t_n, t_n) \end{vmatrix} dt_1 \dots dt_n.$$

But $\bar{K}(x, t) = K(t, x)$, then

$$\bar{B}_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} = \int_a^b \dots \int_a^b \left| \begin{array}{cccccc} K(y_1, x_1) & \dots & K(y_p, x_1) & K(t_1, x_1) & \dots & K(t_n, x_1) \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ K(y_1, x_p) & \dots & K(y_p, x_p) & K(t_1, x_p) & \dots & K(t_n, x_p) \\ K(y_1, t_1) & \dots & K(y_p, t_1) & K(t_1, t_1) & \dots & K(t_n, t_1) \\ \dots & \dots & \dots & \dots & \dots & \dots \\ K(y_1, t_n) & \dots & K(y_p, t_n) & K(t_1, t_n) & \dots & K(t_n, t_n) \end{array} \right| dt_1 \dots dt_n.$$

An interchange of rows and columns in the determinant of the integrand does not change the value of the determinant. Hence

$$\bar{B}_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} = \int_a^b \dots \int_a^b K \begin{pmatrix} y_1, & \dots, & y_p, & t_1, & \dots, & t_n \\ x_1, & \dots, & x_p, & t_1, & \dots, & t_n \end{pmatrix} dt_1 \dots dt_n.$$

Then, by means of the equation (28) defining B_n , we see that

$$\bar{B}_n \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} = B_n \begin{pmatrix} y_1, & \dots, & y_p \\ x_1, & \dots, & x_p \end{pmatrix}$$

and, therefore,

$$(44) \quad \bar{D} \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} \lambda = D \begin{pmatrix} y_1, & \dots, & y_p \\ x_1, & \dots, & x_p \end{pmatrix} \lambda.$$

Since λ_o is a characteristic constant of $K(x, t)$ of index q , we have

$$D \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix} \lambda_o = 0 \quad (p = 1, \dots, q - 1)$$

while

$$D \begin{pmatrix} x'_1, & \dots, & x'_q \\ y'_1, & \dots, & y'_q \end{pmatrix} \lambda_o \neq 0.$$

Hence, by (44)

$$\bar{D} \begin{pmatrix} x_1, & \dots, & x_p \\ y_1, & \dots, & y_p \end{pmatrix}_{\lambda_o} = D \begin{pmatrix} y_1, & \dots, & y_p \\ x_1, & \dots, & x_p \end{pmatrix}_{\lambda_o} \equiv 0$$

in $x_1, \dots, x_p, y_1, \dots, y_p$ for $p = 1, \dots, q - 1$.

Further, if we put

$$(45) \quad x_\alpha' = y_\alpha', \quad \bar{y}_\alpha' = x_\alpha',$$

we have

$$\begin{aligned} \bar{D} \begin{pmatrix} \bar{x}_1', & \dots, & \bar{x}_q' \\ \bar{y}_1', & \dots, & \bar{y}_q' \end{pmatrix}_{\lambda_o} &= D \begin{pmatrix} \bar{y}_1', & \dots, & \bar{y}_q' \\ \bar{x}_1', & \dots, & \bar{x}_q' \end{pmatrix}_{\lambda_o} \\ &= D \begin{pmatrix} x_1', & \dots, & x_q' \\ y_1', & \dots, & y_q' \end{pmatrix}_{\lambda_o} \neq 0. \end{aligned}$$

Hence, by definition, the index \bar{q} of λ_o as a characteristic constant of $K(t, x)$ is q . We state this result in the

Theorem IX.—If λ_o is a characteristic constant of $K(x, t)$ of index q , then λ_o is a characteristic constant of $K(t, x)$ of the same index:

$$\bar{q} = q.$$

c) *The Fundamental Functions of the Associated Equation.*

Apply, now, Theorem VIII to equation (42) and we find that it has q linearly independent solutions. A fundamental system of such solutions is given by

$$\begin{aligned} \bar{\varphi}_\alpha(x) &= \frac{\bar{D} \begin{pmatrix} \bar{x}_1', & \dots, & \bar{x}'_{\alpha-1}, & x, & \bar{x}'_{\alpha+1}, & \dots, & \bar{x}_q' \\ \bar{y}_1', & \dots, & \bar{y}'_{\alpha-1}, & \bar{y}_\alpha', & \bar{y}'_{\alpha+1}, & \dots, & \bar{y}_q' \end{pmatrix}_{\lambda_o}}{\bar{D} \begin{pmatrix} \bar{x}_1', & \dots, & \bar{x}_q' \\ \bar{y}_1', & \dots, & \bar{y}_q' \end{pmatrix}_{\lambda_o}} \\ &= \frac{D \begin{pmatrix} \bar{y}_1', & \dots, & \bar{y}'_{\alpha-1}, & \bar{y}_\alpha', & \bar{y}'_{\alpha+1}, & \dots, & \bar{y}_q' \\ \bar{x}_1', & \dots, & \bar{x}'_{\alpha-1}, & x, & \bar{x}'_{\alpha+1}, & \dots, & \bar{x}_q' \end{pmatrix}_{\lambda_o}}{D \begin{pmatrix} \bar{y}_1', & \dots, & \bar{y}_q' \\ \bar{x}_1', & \dots, & \bar{x}_q' \end{pmatrix}_{\lambda_o}}. \end{aligned}$$

If, now, we introduce the change of notation (45), we find

$$\bar{\varphi}_\alpha(x) = \frac{D(x_1', \dots, x'_{\alpha-1}, x_\alpha', x'_{\alpha+1}, \dots, x_q', \lambda_o)}{D(y_1', \dots, y'_{\alpha-1}, x, y'_{\alpha+1}, \dots, y_q', \lambda_o)} \\ (\varphi = 1, \dots, q).$$

The most general solution of (42) is now

$$v(x) = C_1 \bar{\varphi}_1(x) + C_2 \bar{\varphi}_2(x) + \dots + C_q \bar{\varphi}_q(x).$$

d) The Function $H(x, y)$ for the Associated Kernel.—From the definition of $H(x, y)$ given earlier, we have

$$\bar{H}(x, y) = \frac{\bar{D}(x, \bar{x}_1', \dots, \bar{x}_q', \lambda_o)}{\bar{D}(\bar{y}_1', \dots, \bar{y}_q', \lambda_o)}.$$

Apply (44) and make the change of notation (45). Then

$$(46) \quad \bar{H}(x, y) = \frac{D(y, y_1', \dots, y_q', \lambda_o)}{D(x, x_1', \dots, x_q', \lambda_o)} = H(y, x).$$

If, now, we take account of (46) and make the change of notation (45), then the relation (41) written for the kernel $\bar{K}(x, t) = K(t, x)$ becomes

$$(47) \quad \sum_{\alpha=1}^q \lambda_o K(t, y_\alpha') \bar{\varphi}_\alpha(x) = \lambda_o K(t, x) - H(t, x) \\ + \lambda_o \int_a^b H(s, x) K(t, s) ds.$$

e) The Orthogonality Theorem.—

Theorem X.—If λ_o and λ_1 are two distinct characteristic constants of $K(x, t)$, $\varphi_o(x)$ is a fundamental function of

$K(x, t)$ for λ_o , and $\bar{\varphi}_1(x)$ is a fundamental function of $\bar{K}(x, t)$ for λ_1 , that is

$$(48) \quad \varphi_o(x) = \lambda_o \int_a^b K(x, t) \varphi_o(t) dt$$

$$(49) \quad \bar{\varphi}_1(x) = \lambda_1 \int_a^b \bar{K}(x, t) \bar{\varphi}_1(t) dt = \lambda_1 \int_a^b K(t, x) \bar{\varphi}_1(t) dt$$

then

$$(50) \quad \int_a^b \varphi_o(x) \bar{\varphi}_1(x) = 0.$$

Proof.—From (48) and (49) we obtain

$$\begin{aligned} (\lambda_o - \lambda_1) \int_a^b \varphi_o(x) \bar{\varphi}_1(x) dx &= \\ &\quad \lambda_o \lambda_1 \int_a^b \int_a^b \varphi_o(x) K(t, x) \bar{\varphi}_1(t) dt dx \\ &\quad - \lambda_o \lambda_1 \int_a^b \int_a^b \bar{\varphi}_1(x) K(x, t) \varphi_o(t) dt dx. \end{aligned}$$

We see that the two integrals on the right are equal if in the last integral we write t and x in place of x and t . Then, since by hypothesis $\lambda_o \neq \lambda_1$, we must have (50).

Definition.—Two continuous functions $g(x)$, $h(x)$, for which

$$\int_a^b g(x) h(x) dx = 0,$$

are said to be *orthogonal* to each other.

Hence the above result may be stated as follows: $\varphi_o(x)$ and $\bar{\varphi}_1(x)$ are orthogonal to each other.

24. The Non-homogeneous Integral Equation When $D(\lambda) = 0$.—With the aid of the results established in the last article we can discuss completely the solution of the non-homogeneous integral equation

$$(51) \quad u(x) = f(x) + \lambda_o \int_a^b K(x, t) u(t) dt$$

when $D(\lambda_o) = 0$ and λ_o is of index q .

The finite system of linear equations

$$u_i - \lambda h \sum_{j=1}^n K_{ij} u_j = f_i \quad (i = 1, \dots, n),$$

of which the equation (51) may be considered as a limit, has, in general, for $\Delta = 0$, no finite solution. If, however, certain conditions on the f_i are satisfied, the system has an infinitude of solutions. In analogy we will find that (51), for $D(\lambda_0) = 0$, has, in general, no solutions. If, however, $f(x)$ satisfies certain conditions, then (51) has an infinitude of solutions.

a) *Necessary Conditions.*—To obtain these conditions $f(x)$ we assume that $u(x)$ is a continuous function of x satisfying (51). Multiply both sides of (51) by $\bar{\varphi}_\alpha(x)$, where

$$\bar{\varphi}_\alpha(x) = \lambda_0 \int_a^b K(t, x) \bar{\varphi}_\alpha(t) dt,$$

and integrate with respect to x from a to b . We obtain

$$(52) \quad \int_a^b f(x) \bar{\varphi}_\alpha(x) dx = \int_a^b u(x) \bar{\varphi}_\alpha(x) dx - \lambda_0 \int_a^b \bar{\varphi}_\alpha(x) \left\{ \int_a^b K(x, t) u(t) dt \right\} dx.$$

In the last integral on the right, $\bar{\varphi}_\alpha(x)$ is constant with respect to t and so can be placed under the second sign of integration. We may change the order of integration and then take $u(t)$ from under the sign of integration with respect to x . Thus the last term becomes

$$\int_a^b u(t) \left\{ \lambda_0 \int_a^b \bar{\varphi}_\alpha(x) K(x, t) dx \right\} dt = \int_a^b u(t) \bar{\varphi}_\alpha(t) dt.$$

Thus we see that the first and last terms on the right cancel and

$$(53) \quad \int_a^b f(x) \bar{\varphi}_\alpha(x) dx = 0. \quad (\alpha = 1, \dots, q)$$

Hence, in order that there may exist a continuous solution $u(x)$ of (51), $f(x)$ must satisfy the q conditions (53).

b) *Sufficiency Proof.*—Let us now show conversely that, if $f(x)$ satisfies the q conditions (53), then (51) does have a solution. By our hypothesis the q equations (53) are satisfied. Then

$$\sum_{\alpha=1}^q \lambda_\alpha K(x, y_\alpha') \int_a^b f(t) \bar{\varphi}_\alpha(t) dt = 0,$$

Now, $\lambda_\alpha K(x, y_\alpha')$ is independent of t and so may be placed under the sign of integration. It is permissible to change the order of performing the summation and the integration. We thus obtain

$$\int_a^b \left\{ \sum_{\alpha=1}^q \lambda_\alpha K(x, y_\alpha') \bar{\varphi}_\alpha(t) f(t) \right\} dt = 0,$$

which, on account of (47), becomes

$$(54) \quad 0 = \int_a^b \lambda_\alpha K(x, t) f(t) dt - \int_a^b H(x, t) f(t) dt \\ + \lambda_\alpha \int_a^b f(t) \left\{ \int_a^b H(s, t) K(x, s) ds \right\} dt.$$

The last term may be written

$$\lambda_\alpha \int_a^b \int_a^b f(t) H(s, t) K(x, s) ds dt.$$

Make now a change in notation. In place of t and s write s and t . We obtain

$$\lambda_\alpha \int_a^b \int_a^b f(s) H(t, s) K(x, t) dt ds.$$

In this definite double integral it is permissible to change the order of integration. We then obtain

$$\lambda_\alpha \int_a^b K(x, t) \left\{ \int_a^b H(t, s) f(s) ds \right\} dt.$$

After making these reductions in (54), combine the first and last terms and obtain

$$(55) \quad 0 = \lambda_o \int_a^b K(x, t) \left\{ f(t) + \int_a^b H(t, s)f(s)ds \right\} dt - \int_a^b H(x, t)f(t)dt$$

Now put

$$u_o(t) = f(t) + \int_a^b H(t, s)f(s)ds,$$

then

$$\int_a^b H(x, t)f(t)dt = u_o(x) - f(x).$$

Making use of these last two equations, (55) becomes

$$u_o(x) = f(x) + \lambda_o \int_a^b K(x, t)u_o(t)dt.$$

Thus we have proved that if (53) are satisfied then (51) has at least one solution, $u_o(x)$ given by

$$(56) \quad u_o(x) = f(x) + \int_a^b H(x, t)f(t)dt.$$

c) *Determination of All Solutions.*—Let us suppose that (51) has another continuous solution $u(x)$. Then $u(x) - u_o(x)$ is a solution of the homogeneous equation

$$(57) \quad v(x) = \lambda_o \int_a^b K(x, t)v(t)dt,$$

for, if we subtract the members of (56) from the corresponding members of (51), we obtain

$$(58) \quad u(x) - u_o(x) = \lambda_o \int_a^b K(x, t) [u(t) - u_o(t)] dt.$$

By Theorem VIII the most general solution of (57) is of the form

$$C_1\varphi_1(x) + C_2\varphi_2(x) + \dots + C_q\varphi_q(x),$$

where C_1, C_2, \dots, C_q are arbitrary constants.

Hence the most general solution of (58) is

$$u(x) - u_o(x) = C_1\varphi_1(x) + C_2\varphi_2(x) + \dots + C_q\varphi_q(x).$$

Therefore,

$$u(x) = f(x) + \int_a^b H(x, t)f(t)dt + C_1\varphi_1(x) + \dots + C_q\varphi_q(x)$$

is the complete solution of (51). We have thus proved *Fredholm's third fundamental theorem*:

Theorem XI.—*If λ_o is a characteristic constant of $K(x, t)$ of index q then*

$$(51) \quad u(x) = f(x) + \lambda_o \int_a^b K(x, t)u(t)dt$$

has, in general, no continuous solution. In order that a continuous solution exist it is necessary that

$$\int_a^b f(x)\bar{\varphi}_\alpha(x)dx = 0, \quad \alpha = 1, \dots, q$$

where the $\bar{\varphi}_\alpha(x)$ are a complete set of fundamental functions for the associated homogeneous equation

$$(42) \quad v(x) = \lambda_o \int_a^b K(t, x)v(t)dt.$$

If these conditions are satisfied, then there are a q -fold infinitude of solutions of (51) given by

$$u(x) = f(x) + \int_a^b H(x, t)f(t)dt + C_1\varphi_1(x) + \dots + C_q\varphi_q(x)$$

where C_1, \dots, C_q are arbitrary constants and where the $\varphi_\alpha(x)$ are a complete set of fundamental functions for

$$(26) \quad u(x) = \lambda_o \int_a^b K(x, t)u(t)dt \quad (\text{Theorem VIII})$$

and $H(x, t)$ is given by (40).

The following table exhibits the results of the solution of Fredholm's equation together with the analogy between the finite system and the integral equation.

$$u(x) = f(x) + \lambda \int_a^b K(x, t)u(t)dt$$

Case I: $D(\lambda) \neq 0$		Case II: $D(\lambda) = 0$, index q	
Non-homogeneous	Homogeneous	Non-homogeneous	Homogeneous
Unique solution $u(x) = f(x)$ + $\int_a^b \frac{D(x, t; \lambda)}{D(\lambda)} f(t)dt$	Unique solution $u \equiv 0$	In general, no solution Solutions exist only if f satisfies $\int_a^b f(x) \bar{\varphi}_\alpha(x) = 0$ Then ∞^q solutions	∞^q solutions: $\sum_{\alpha=1}^q C_\alpha \varphi_\alpha(x)$

$$u_i - \lambda h \sum_{j=1}^n K_{ij} u_j = f_i, \quad i = 1, \dots, n$$

Case I: $\Delta \neq 0$		Case II: $\Delta = 0$, index q	
Non-homogeneous	Homogeneous	Non-homogeneous	Homogeneous
Unique solution $u_k = \frac{\sum_{i=1}^n f_i \Delta_{ik}}{\Delta}$	Unique solution $u_k = 0$	In general, no solution ∞^q solutions if the f_i satisfy certain q relations of the form $C_{\alpha 1} f_1 + \dots + C_{\alpha n} f_n = 0,$ $\alpha = 1, \dots, q$	∞^q solutions if Δ is of rank r where $q = n - r$

25. Kernels of the Form $\sum a_i(x)b_i(y)$.—We give a brief discussion of the integral equation with a kernel of the form

$$K(x, y) = a_1(x)b_1(y) + \dots + a_n(x)b_n(y).$$

Fredholm's integral equation

$$u(x) = f(x) + \lambda \int_a^b K(x, t)u(t)dt$$

can now be written in the form

$$(59) \quad u(x) = f(x) + \lambda \left[a_1(x) \int_a^b b_1(t)u(t)dt + \dots + a_n(x) \int_a^b b_n(t)u(t)dt \right].$$

If, now, we put

$$(60) \quad \int_a^b b_i(t)u(t)dt = K_i \quad (i = 1, \dots, n),$$

we see that $u(x)$ is of the form

$$(61) \quad u(x) = f(x) + \lambda \left[a_1(x)K_1 + \dots + a_n(x)K_n \right].$$

In order to determine the constants K_i , let us substitute in (60) the value of u given by (61). We obtain the n equations

$$(62) \quad K_i - \lambda \left[\int_a^b a_1(t)b_i(t)K_1 dt + \dots + \int_a^b a_n(t)b_i(t)K_n dt \right] = \int_a^b b_i(t)f(t)dt \quad (i = 1, \dots, n).$$

Introducing the notation

$$\int_a^b a_k(t)b_i(t)dt = C_{ki}$$

the system (62) can be written in the form

$$(63) \quad K_i - \lambda \left[C_{1i}K_1 + \dots + C_{ni}K_n \right] = \int_a^b b_i(t)f(t)dt.$$

Equations (63) are a linear algebraic system of n non-homogeneous equations in the n unknowns K_1, \dots, K_n , with the determinant

$$D(\lambda) = \begin{vmatrix} 1 - \lambda C_{11} & -\lambda C_{12} & \dots & -\lambda C_{1n} \\ -\lambda C_{21} & 1 - \lambda C_{22} & \dots & -\lambda C_{2n} \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ -\lambda C_{n1} & -\lambda C_{n2} & \dots & 1 - \lambda C_{nn} \end{vmatrix}$$

From the theory of linear systems, we have at once the result:

a) If $D(\lambda) \neq 0$, the system (62) is satisfied by one and only one set of values of K_1, \dots, K_n and these values are given by Cramer's formulas. Therefore, Fredholm's equation (59) has one and only one solution, which is given by (61).

b) If $D(\lambda) = 0$ for $\lambda = \lambda_o$ (and this happens for n values of λ , real or complex), and one of the q th minors of $D(\lambda)$ is the first minor which does not vanish for $\lambda = \lambda_o$ (this q th minor is a determinant of order $n - q$), then the general solution of the homogeneous system (62) ($f(x) \equiv 0$) will be of the form¹

$$K_i = \alpha_1 m_{1i} + \alpha_2 m_{2i} + \dots + \alpha_q m_{qi} \quad (i = 1, \dots, n),$$

where $\alpha_1, \alpha_2, \dots, \alpha_q$ are arbitrary constants.

If we put the values of K_i so obtained in (61), we obtain

$$u(x) = \lambda_o \left[\alpha_1 u_1(x) + \alpha_2 u_2(x) + \dots + \alpha_q u_q(x) \right]$$

where the functions

$$u_r(x) = m_{r1} a_1(x) + m_{r2} a_2(x) + \dots + m_{rn} a_n(x) \quad (r = 1, \dots, q)$$

are linearly independent.

Thus we see that, under the circumstances specified, the homogeneous integral equation for $\lambda = \lambda_o$ has q linearly independent solutions.

¹ BÖCHER, "Introduction to Higher Algebra," §18.

The associated equation

$$(64) \quad \bar{u}(x) = f(x) + \lambda \left[b_1(x) \int_a^b a_1(t) \bar{u}(t) dt + \dots + b_n(x) \int_a^b a_n(t) \bar{u}(t) dt \right]$$

is obtained from (59) by interchanging the functions $a_i(x)$ and $b_i(x)$. The general term of the characteristic determinant of (59) being

$$\int_a^b a_k(t) b_i(t) dt = C_{ki},$$

the general term for the associated equation will be

$$\int_a^b b_k(t) a_i(t) dt = C_{ik}.$$

The characteristic determinants of these two equations are identical, since one can obtain one from the other by interchanging rows and columns. Therefore, the equation (59) and the associated equation (64) have exactly the same characteristic numbers and with the same index.

From the general theory we know that if λ_o is a root of $D(\lambda) = 0$ of index q , then, in order that the non-homogeneous equation (59) may have a solution, we must have

$$\int_a^b f(x) \bar{u}_i(x) dx = 0 \quad (i = 1, \dots, q).$$

EXERCISES

For the equation $u(x) = f(x) + \lambda \int_a^b K(x, t) u(t) dt$ compute $D(\lambda)$

and $D(x, y; \lambda)$ for the following kernels for the specified limits a and b :

Ans.

- | | |
|--|--|
| 1. $K(x, t) = 1, a = 0, b = 1.$ | $D(\lambda) = 1 - \lambda.$ |
| 2. $K(x, t) = -1, a = 0, b = 1.$ | $D(\lambda) = 1 + \lambda.$ |
| 3. $K(x, t) = \sin x, a = 0, b = \pi.$ | $D(\lambda) = 1 - 2\lambda.$ |
| 4. $K(x, t) = xt, a = 0, b = 10.$ | $D(\lambda) = 1 - \frac{1,000}{3}\lambda.$ |

5. $K(x, t) = t, a = 0, b = 10.$ $D(\lambda) = 1 - 50\lambda.$
 6. $K(x, t) = x, a = 4, b = 10.$ $D(\lambda) = 1 - 42\lambda.$
7. $K(x, t) = g(x), a = a, b = b.$ $D(\lambda) = 1 - \lambda \int_a^b g(t)dt.$
8. $K(x, t) = g(t), a = a, b = b.$ $D(\lambda) = 1 - \lambda \int_a^b g(t)dt.$
9. $K(x, t) = 2e^x \cdot e^t, a = 0, b = 1.$ $D(\lambda) = 1 - (e^2 - 1)\lambda.$
 10. $K(x, t) = x - t, a = 0, b = 1.$

Solve the following integral equations:

11. $u(x) = \sec^2 x + \lambda \int_0^1 u(t)dt.$

12. $u(x) = \sec x \cdot \tan x - \lambda \int_0^1 u(t)dt.$

13. $u(x) = \cos x + \lambda \int_0^\pi \sin x \cdot u(t)dt.$

14. $u(x) = e^x + \lambda \int_0^{10} xt \cdot u(t)dt.$

15. $u(x) = x^2 + \lambda \int_0^{10} t \cdot u(t)dt.$

16. $u(x) = \sin x + \lambda \int_4^{10} x \cdot u(t)dt.$

17. $u(x) = e^x + \lambda \int_0^1 2e^x e^t u(t)dt.$

Solve the following homogeneous integral equations:

18. $u(x) = \int_0^1 u(t)dt.$

19. $u(x) = \int_0^1 (-1)u(t)dt.$

20. $u(x) = \frac{1}{2} \int_0^\pi \sin x \cdot u(t)dt.$

21. $u(x) = \frac{3}{1,000} \int_0^{10} xt \cdot u(t)dt.$

22. $u(x) = \frac{1}{50} \int_0^{10} t \cdot u(t)dt.$

23. $u(x) = \frac{1}{42} \int_4^{10} x \cdot u(t) dt.$

24. $u(x) = \frac{1}{e^2 - 1} \int_0^1 2e^x e^t u(t) dt.$

Solve the following equation by the method of §25:

25. $u(x) = x^2 + \lambda \int_0^1 (1 + xt) u(t) dt.$

26. $u(x) = x + \lambda \int_0^\pi (1 + \sin x \sin t) u(t) dt.$

27. $u(x) = x + \lambda \int_0^1 (1 + x + t) u(t) dt.$

28. $u(x) = x + \lambda \int_0^1 (x - t) u(t) dt.$

29. $u(x) = x + \lambda \int_0^1 (x - t)^2 u(t) dt.$


30. Solve Exercises 11–17 inclusive by this method.

CHAPTER IV

APPLICATIONS OF THE FREDHOLM THEORY

I. FREE VIBRATIONS OF AN ELASTIC STRING

26. The Differential Equations of the Problem.—We consider an elastic string stretched between the two fixed points A and B . We pull it out of its position of equilibrium AQB into some other plane initial position, as ACB , and then release it. The string will describe transverse vibrations. Suppose that at time t the string occupies the position APB . Let $x = AQ$, $y = QP$ be the abscissa and ordinate of any one of its points P . Then y is a function of x and t . We suppose the cross-section of the string to be constant and infinitesimal compared with the length.



[FIG. 5.]

The string is of homogeneous density. The effect of gravity is to be neglected. Further, we take for simplicity $AB = 1$. It is then proved in the theory of elasticity that the motion of the string is given by the partial differential equation

$$(1) \quad \frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}, \quad (c = \text{constant})$$

with the initial conditions

$$(2) \quad y(0, t) = 0, \quad y(1, t) = 0,$$

$$(3) \quad y(x, 0) = g(x), \quad y_t(x, 0) = 0,$$

where $y = g(x)$ is the equation of the initial position ACB . Equations (2) are the analytic statement of the fact that the end points A and B remain fixed, while equations (3) state

that for $t = 0$ the string is in the initial position ACB and each particle starts its motion with an initial zero velocity.¹

27. Reduction to a One-dimensional Boundary Problem. Let us try to find a solution of (1) in the form

$$y = u(x)\varphi(t).$$

Substitution of this value of y in (1) gives

$$u(x)\frac{d^2\varphi}{dt^2} = c^2\varphi(t)\frac{d^2u}{dx^2},$$

which can be put in the form

$$\frac{\frac{d^2\varphi(t)}{dt^2}}{\varphi(t)} = c^2 \frac{\frac{d^2u(x)}{dx^2}}{u(x)}.$$

The right-hand side is independent of t , and the left-hand side is independent of x . Then either member is a constant, which we designate as $-\lambda c^2$. This gives us the two ordinary differential equations to solve:

$$\frac{d^2u}{dx^2} + \lambda u = 0, \quad \frac{d^2\varphi}{dt^2} + \lambda c^2\varphi = 0.$$

The initial conditions on u and φ are obtained from (2) and (3), and for u are as follows:

$$\begin{aligned} u(0)\varphi(t) &= 0 \quad \text{whence} \quad u(0) = 0 \quad \text{since } \varphi(t) \neq 0. \\ u(1)\varphi(t) &= 0 \quad u(1) = 0 \end{aligned}$$

We are thus led to the following *one-dimensional boundary problem*: to determine a function $u(x)$ which will satisfy the differential equation

$$(4) \quad \frac{d^2u}{dx^2} + \lambda u = 0$$

and the boundary conditions

$$u(0) = 0, \quad u(1) = 0.$$

28. Solution of the Boundary Problem.—We see at once that $u \equiv 0$ is a solution. This gives for the original prob-

¹ WEBER-RIEMANN, "Lehrbuch der Partielle Differentialgleichungen," §83.

lem of the vibrating string the trivial solution $y \equiv 0$, which means that the string remains at rest. Hereafter, when we refer to a solution of our boundary problem we shall mean a solution not identically zero. To obtain the solution we have three cases to consider.

Case I.— $\lambda > 0$. From the elementary theory of differential equations we know that the most general solution of (4) is

$$u(x) = A \cos \sqrt{\lambda}x + B \sin \sqrt{\lambda}x.$$

The conditions $u(0) = 0$, $u(1) = 0$ give us $A = 0$ and either

(1) $B = 0$, or (2) $\sin \sqrt{\lambda} = 0$.

(1) If $B = 0$, we obtain the trivial solution $u \equiv 0$.

(2) If $\sin \sqrt{\lambda} = 0$, then $\lambda = n^2\pi^2$ (n , an integer) and the solution is

$$u(x) = B \sin n\pi x$$

Case II.— $\lambda = 0$. The general solution of (4) is now

$$u = Ax + B.$$

But $u(0) = B = 0$, and $u(1) = A = 0$, so that we have again the trivial solution $u \equiv 0$.

Case III.— $\lambda < 0$. The general solution of (4) is now

$$u = Ae^{\sqrt{-\lambda}x} + Be^{-\sqrt{-\lambda}x}.$$

Applying the initial conditions, we obtain

$$u(0) = A + B = 0, \text{ whence } B = -A$$

$$u(1) = A(e^{\sqrt{-\lambda}} - e^{-\sqrt{-\lambda}}) = 0, \text{ whence } A = 0 (\lambda \neq 0).$$

Therefore, $A = B = 0$, and we have again the trivial solution $u \equiv 0$.

Thus we arrive at the

Theorem I.—*If $\lambda = n^2\pi^2$ (n , an integer), then the boundary problem*

$$(4) \quad \frac{d^2u}{dx^2} + \lambda u = 0, \quad u(0) = 0, \quad u(1) = 0$$

has an infinitude of solutions:

$$u = B \sin n\pi x.$$

If $\lambda \neq n^2\pi^2$, then the only solution is the trivial one $u \equiv 0$.

29. Construction of Green's Function.—We now propose to show that every solution of our boundary problem (4) satisfies at the same time a linear integral equation. We observe first that with the given boundary conditions the method of §4 cannot be used to determine an equivalent integral equation. In the present instance, in order to determine an equivalent integral equation, we first construct the *Green's function* belonging to the boundary problem.

The given boundary problem for $\lambda = 0$ has only the trivial solution $u \equiv 0$. This is true, however, only under the assumptions tacitly made throughout, namely, that u , together with its first and second derivatives, is continuous in the interval $[01]$. Let us use the notation $u^{c''}$ to denote this assumption.¹

Drop now the assumption $u^{c''}$ and allow the derivative u' to be discontinuous at an arbitrarily prescribed point ξ between 0 and 1, while u itself remains continuous. Accordingly, we propose to determine a function u satisfying the following conditions:

A) u^c in $[01]$.

B) $u^{c''}$ and $\frac{d^2u}{dx^2} = 0$ in $[0\xi]$.

$u^{c''}$ and $\frac{d^2u}{dx^2} = 0$ in $[\xi1]$.

C) $u(0) = 0, u(1) = 0$.

The solution then must be of the form

$$u = \begin{cases} \alpha_0 x + \beta_0 & \text{in } [0\xi] \\ \alpha_1 x + \beta_1 & \text{in } [\xi1]. \end{cases}$$

¹ Similarly, let us use the notation C to denote the class of all continuous functions and the notation $C^{(n)}$ to denote the class of all functions having continuous derivatives up to the order n inclusive.

From C) we find

$$u(0) = \beta_0 = 0; u(1) = \alpha_1 + \beta_1 = 0.$$

Thus the solution reduces to

$$u = \begin{cases} \alpha_0 x & [0\xi] \\ \beta_1(1 - x) & [\xi 1]. \end{cases}$$

The condition A) must be satisfied, whence

$$\alpha_0 \xi = \beta_1(1 - \xi).$$

Therefore $\alpha_0 = \rho(1 - \xi)$ and $\beta_1 = \rho \xi$.

The solution now takes the form

$$u = \begin{cases} \rho(1 - \xi)x & [0\xi] \\ \rho \xi(1 - x) & [\xi 1]. \end{cases}$$

Geometrically, this solution is represented by a broken line as in the adjoining figure.




FIG. 6.

For $x = \xi$, the derivative has a discontinuity measured by

$$u'(\xi - 0) - u'(\xi + 0) = \rho(1 - \xi) + \rho \xi = \rho.$$

We now impose the further condition that this discontinuity shall be $+1$. Then $\rho = 1$. The function u so obtained is called *Green's function* for the boundary problem (4). We use the notation $K(x, \xi)$ to represent this function. We have thus the following:

Theorem II.—*There exists one and only one function $K(x, \xi)$ which satisfies the conditions*

A) K^o in [01].

B) $K^{o''}$ and $\frac{d^2K}{dx^2} = 0$ in $[0\xi]$.

$K^{o''}$ and $\frac{d^2K}{dx^2} = 0$ in $[\xi 1]$.

C) $K(0) = 0, K(1) = 0$.

D) $\begin{cases} K' \\ x \end{cases} \begin{matrix} x = \xi - 0 \\ x = \xi + 0 \end{matrix} = +1, \quad 0 < \xi < 1.$

This function is given by the formula

$$(5) \quad K(x, \xi) = \begin{cases} (1 - \xi)x & \text{for } 0 \leq x \leq \xi \\ \xi(1 - x) & \text{for } \xi \leq x \leq 1. \end{cases}$$

Let us use the notation

$$K_0(x, \xi) = (1 - \xi)x$$

$$K_1(x, \xi) = \xi(1 - x).$$

The properties A) and D) of $K(x, \xi)$ may now be written

$$K_0(\xi, \xi) = K_1(\xi, \xi)$$

$$K_0'(\xi, \xi) - K_1'(\xi, \xi) = 1.$$

We can now prove the following:

Theorem III.—Green's function is symmetric in x and ξ :

$$K(x, \xi) = K(\xi, x).$$

Proof.—Let $0 \leq z_1 \leq z_2 \leq 1$.

$$\text{Then} \quad K(z_1, z_2) = K_0(z_1, z_2) = (1 - z_2)z_1$$

$$\text{while} \quad K(z_2, z_1) = K_1(z_1, z_2) = z_1(1 - z_2)$$

$$\text{whence} \quad K(z_1, z_2) = K(z_2, z_1).$$

30. Equivalence between the Boundary Problem and a Linear Integral Equation.—Take the equations

$$\frac{d^2u}{dx^2} = -\lambda u, \quad \frac{d^2K}{dx^2} = 0.$$

Multiply the first by $-K$ and the second by u and add. We obtain

$$uK'' - Ku'' = \lambda uK, \text{ or}$$

$$\frac{d}{dx}(uK' - Ku') = \lambda uK.$$

This equality holds in each of the two subintervals $[0\xi]$ and $[\xi 1]$. Integration over each of them gives

$$\begin{aligned} \left[uK' - Ku' \right]_0^{\xi-0} &= \lambda \int_0^{\xi-0} uKdx \\ \left[uK' - Ku' \right]_{\xi+0}^1 &= \lambda \int_{\xi+0}^1 uKdx. \end{aligned}$$

Both u and K vanish at 0 and 1. u , u' , and K are continuous functions of x over the whole interval $[01]$. Whence, adding the last two equations, we obtain

$$u(\xi) \left[K'(\xi - 0) - K'(\xi + 0) \right] = \lambda \int_0^1 K(x, \xi)u(x)dx.$$

But $K'(\xi - 0) - K'(\xi + 0) = 1$, and so

$$u(\xi) = \lambda \int_0^1 K(x, \xi)u(x)dx.$$

Now interchange x and ξ and remember that $K(x, \xi) = K(\xi, x)$. Then

$$u(x) = \lambda \int_0^1 K(x, \xi)u(\xi)d\xi.$$

This is a homogeneous linear integral equation of the second kind for the determination of $u(x)$. Every solution of the original boundary problem (4) satisfies this integral equation. Hence, we have the following:

Theorem IV.—*If $u(x)$ has continuous first and second derivatives, and satisfies the boundary problem*

$$(4) \quad \frac{d^2u}{dx^2} + \lambda u = 0, \quad u(0) = 0, \quad u(1) = 0,$$

then $u(x)$ is continuous and satisfies the homogeneous linear integral equation

$$u(x) = \lambda \int_0^1 K(x, \xi)u(\xi)d\xi,$$

where $K(x, \xi)$ is given by the formula (5).

Let us now prove the following converse

Theorem V.—*If $u(x)$ is continuous and satisfies the equation*

$$(6) \quad u(x) = \lambda \int_0^1 K(x, \xi)u(\xi)d\xi,$$

where $K(x, \xi)$ is given by the formula (5), then u has continuous first and second derivatives and satisfies the boundary problem

$$\frac{d^2u}{dx^2} + \lambda u = 0, \quad u(0) = 0, \quad u(1) = 0.$$

Proof.— x and ξ range from 0 to 1 and $K'(x, \xi)$ is discontinuous at $x = \xi$. Let us then write (6) in the form

$$u(x) = \lambda \int_0^x K_1(x, \xi)u(\xi)d\xi + \lambda \int_x^1 K_0(x, \xi)u(\xi)d\xi.$$

Now we may apply the general rule for the differentiation of a definite integral with respect to a parameter.¹ Then

$$\frac{du}{dx} = \lambda \int_0^x K_1'(x, \xi)u(\xi)d\xi + \lambda \int_x^1 K_0'(x, \xi)u(\xi)d\xi,$$

since $K_0(x, x) = K_1(x, x)$.

Moreover, since $K_1'(x, \xi)$ and $K_0'(x, \xi)$ are continuous in their respective intervals, we see that $\frac{du}{dx}$ is continuous.

A second differentiation gives

$$\begin{aligned} \frac{d^2u}{dx^2} &= \lambda \int_0^x K_1''(x, \xi)u(\xi)d\xi + \lambda \int_x^1 K_0''(x, \xi)u(\xi)d\xi \\ &\quad + \lambda K_1'(x, x)u(x) - \lambda K_0'(x, x)u(x) = -\lambda u(x), \end{aligned}$$

since by our hypothesis on K we have $K_1'' = K_0'' = 0$ and $K_1'(x, x) - K_0'(x, x) = -1$. Moreover, since $u(x)$ is continuous, we have u'' continuous. We have further

$$u(0) = \lambda \int_0^1 K(0, \xi)u(\xi)d\xi = 0,$$

¹ See GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §97.

since $K(0, \xi) = 0$, and

$$u(1) = \lambda \int_0^1 K(1, \xi)u(\xi)d\xi = 0,$$

since $K(1, \xi) = 0$.

We know the solution of the given boundary problem (4). This knowledge combined with Theorems IV and V gives the following:

Theorem VI.—*Only when $\lambda = n^2\pi^2(n, \text{ an integer})$ does the integral equation*

$$u(x) = \lambda \int_0^1 K(x, \xi)u(\xi)d\xi$$

have a solution not identically zero:

$$u(x) = B \sin n\pi x.$$

If we compare these results with the results obtained in the preceding chapter for the general homogeneous integral equation, we see that the characteristic constants for this particular problem are $\lambda = n^2\pi^2 \equiv \lambda_n$ and that they are of index $q = 1$. The fundamental function belonging to λ_n is

$$\varphi(x) = \sin n\pi x.$$

The kernel is a symmetric one, so that

$$K(x, \xi) = K(\xi, x),$$

and therefore the associated equation is identical with the original one and hence has the same solutions. The associated fundamental functions are, therefore,

$$\bar{\varphi}(x) = \sin n\pi x.$$

II. CONSTRAINED VIBRATIONS OF AN ELASTIC STRING

31. The Differential Equations of the Problem.—Let us suppose that an exterior force $\mu H(x, t)$ acts on each particle of mass μ in the y direction. Then it is known from the

mathematical theory¹ of vibrating strings that the equations of motion of the string are

$$(7) \quad \frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2} + H(x, t),$$

$$(8) \quad y(0, t) = 0, \quad y(1, t) = 0,$$

$$(9) \quad y(x, 0) = g(x), \quad y_t(x, 0) = 0.$$

Let us suppose now that $H(x, t)$ is harmonic, that is,

$$H(x, t) = C^2 r(x) \cos(\beta t + \gamma) \quad (c \neq 0).$$

Let us find, if possible, a solution of the form

$$y(x, t) = u(x) \cos(\beta t + \gamma).$$

Substitute this value of y in (7) and put $C^2 \lambda = \beta^2$. We find

$$(10) \quad \frac{d^2 u}{dx^2} + \lambda u + r(x) = 0,$$

while from (8) we derive the boundary conditions

$$(11) \quad u(0) = 0, \quad u(1) = 0.$$

32. Equivalence Between the Boundary Problem and a Linear Integral Equation.—Construct as before the Green's function $K(x, \xi)$. Then

$$\frac{d^2 K}{dx^2} = 0, \quad \frac{d^2 u}{dx^2} = -\lambda u - r.$$

Multiply the first of these by u , the second by $-K$ and add. We obtain

$$uK'' - Ku'' = \lambda uK + rK,$$

which may be written

$$\frac{d}{dx}(uK' - Ku') = \lambda uK + rK.$$

Proceed as before with the integration from 0 to ξ and from ξ to 1. We find

$$u(\xi) = \lambda \int_0^1 K(x, \xi) u(x) dx + \int_0^1 K(x, \xi) r(x) dx.$$

¹ WEBER, *Loc. cit.*, §83.

Interchange x and ξ . Then, on account of the symmetry of K , we have

$$u(x) = \lambda \int_0^1 K(x, \xi) u(\xi) d\xi + \int_0^1 K(x, \xi) r(\xi) d\xi.$$

If, now, we put

$$(12) \quad \int_0^1 K(x, \xi) r(\xi) d\xi = f(x)$$

we have

$$(13) \quad u(x) = f(x) + \lambda \int_0^1 K(x, \xi) u(\xi) d\xi,$$

which is a non-homogeneous linear integral equation of the second kind. It is satisfied by every solution of the boundary problem given by (10) and (11).

If we will proceed exactly as in the case of the homogeneous equation, we can now show conversely that if u is continuous and satisfies the equation (13), where $f(x)$ is given by (12), then u has continuous first and second derivatives and satisfies the differential equation (10) and the boundary conditions (11).

33. Remarks on Solution of the Boundary Problem.—Equations (6) and (13) have the same kernel, namely, the Green's function which we have constructed. Knowing that the characteristic constants for (6), and hence for (13), are $\lambda_n = n^2\pi^2$, we obtain from the general theory the following results for (13):

Case I.—If $\lambda \neq n^2\pi^2$, (13) has a unique solution.

Case II.—If $\lambda = n^2\pi^2$, there is, in general, no solution. A solution exists only when the condition

$$(14) \quad \int_0^1 f(x) \sin n\pi x dx = 0$$

is satisfied. This condition is what

$$\int_a^b f(x) \bar{\varphi}_n(x) dx = 0$$

becomes for this special problem. If (14) is satisfied, then (13) has ∞^1 solutions.

The boundary problem

$$\frac{d^2u}{dx^2} + \lambda u + r(x) = 0, \quad u(0) = 0, \quad u(1) = 0$$

was shown to be equivalent to the non-homogeneous integral equation (13) if

$$(15) \quad f(x) = \int_0^1 K(x, \xi) r(\xi) d\xi.$$

This enables us, from Cases I and II for the integral equation, to state that, when $\lambda \neq n^2\pi^2$, the boundary problem has a unique solution, and when $\lambda = n^2\pi^2$, there is, in general, no solution, but that when (14) is satisfied, there are ∞^1 solutions. If, now, in (14) we substitute for $f(x)$ its value as given by (15), this condition becomes

$$\int_0^1 \int_0^1 K(x, \xi) r(\xi) \sin n\pi x d\xi dx = 0.$$

Interchange the order of integration and remember that $\sin n\pi x$ is a solution of the homogeneous equation for $\lambda = \lambda_n$:

$$\sin n\pi x = \lambda_n \int_0^1 K(x, \xi) \sin n\pi \xi d\xi$$

$$\text{or} \quad \sin n\pi \xi = \lambda_n \int_0^1 K(x, \xi) \sin n\pi x dx$$

[since $K(x, \xi) = K(\xi, x)$], then the double integral becomes

$$\frac{1}{\lambda_n} \int_0^1 r(\xi) \sin n\pi \xi d\xi = 0.$$

That is, in order that the boundary problem (10) may have a solution, it is necessary and sufficient that $r(x)$ satisfies the equation

$$\int_0^1 r(x) \sin n\pi x dx = 0.$$

III. AUXILIARY THEOREMS ON HARMONIC FUNCTIONS

34. Harmonic Functions.—For the solution later of the two boundary problems of the potential theory known as *Dirichlet's problem* and *Neumann's problem* we shall need certain auxiliary theorems on *harmonic functions*, that is, functions of class C'' satisfying $\Delta u = 0$:

$$\Delta u \equiv \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0.$$

35. Definitions about Curves.—A curve C

$$C: \quad x = \varphi(t), \quad y = \psi(t), \quad t_0 \leq t \leq t_1$$

is said to be *continuous* if φ and ψ are continuous on $[t_0, t_1]$. We write this

$$C^c \sim \varphi^c \cdot \psi^c.$$

The symbol \sim is read *is equivalent to* or *implies and is implied by*. The symbol \cdot is read *and*. Further, we say that the curve C is of class c' if φ and ψ have continuous first derivatives and these first derivatives φ' and ψ' do not vanish simultaneously on $[t_0, t_1]$. We write this

$$C^{c'} \sim \varphi^{c'} \cdot \psi^{c'} (\varphi', \psi') \neq (0, 0).$$

Such a curve is sometimes called a *smooth curve*. A curve of class c' has a definite positive tangent at every point. The condition $(\varphi', \psi') \neq (0, 0)$ excludes singular points. Every arc has a definite finite length. Thus we are assured that for smooth curves we can choose as parameter t the length s of arc and write

$$C: \quad x = \xi(s), \quad y = \eta(s), \quad 0 \leq s \leq l$$

$$\xi'^2 + \eta'^2 = 1,$$

when l is the total length of arc.

In like manner, we define a curve of class C'' :

$$C^{c''} \sim \varphi^{c''} \psi^{c''} \cdot (\varphi', \psi') \neq (0, 0).$$

Curves of class c'' have at every point a definite curvature,¹ which varies continuously from point to point.

36. Green's Theorem.—Let C be a curve with the following properties:

- 1) It is closed.
- 2) It has no multiple points.
- 3) It is of class c'' . Hence we can represent it with the arc s as parameter:

$$x = \xi(s), \quad y = \eta(s), \quad 0 \leq s \leq l.$$

- 4) There exists a positive integer m such that every line parallel to the y or x axis meets the curve in at most m points.

According to Jordan's theorem² such a curve divides the plane into an interior and an exterior region. Let us denote by

I , the interior plus the boundary C ,
by E , the exterior plus the boundary C .

Then Green's theorem³ may be stated as follows:

Green's Theorem.—If $P(x, y)$ and $Q(x, y)$ are of class c' on I , then

$$(16) \quad \int_C \left[P(x, y)dx + Q(x, y)dy \right] = \int_I \int \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

where the line integral is taken in the positive sense around C .

¹ See GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §205, note.

² See OSGOOD, "Funktionentheorie," 2nd ed., p. 171.

³ See GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §126, for a proof of Green's theorem.

If we put $P = v \frac{\partial u}{\partial y}$, $Q = -v \frac{\partial u}{\partial x}$, then Green's theorem becomes: If v is of class c' and u of class c'' on I , then

$$(17) \quad \int_C v \left(\frac{\partial u}{\partial y} dx - \frac{\partial u}{\partial x} dy \right) = - \int_I \int v \Delta u dx dy - \int_I \int \left(\frac{\partial u}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial v}{\partial y} \right) dx dy.$$

The left-hand side can be simplified if we introduce the idea of the *directional derivative* of a function $f(x, y)$.

Let $f(x, y)$ be defined at every point of a region R of the xy -plane. The directional derivative is defined¹ as follows (see figure):

$$\frac{\partial f}{\partial p} = \lim_{h \rightarrow 0} \frac{f(x_1, y_1) - f(x, y)}{h},$$

if such limit exists.

If $f(x, y)$ is of class c' , then²

$$f(x_1, y_1) - f(x, y) = f_x(x, y)(x_1 - x) + f_y(x, y)(y_1 - y) + \alpha(x_1 - x) + \beta(y_1 - y),$$

where α and β approach 0 with $x_1 - x$ and $y_1 - y$.

Divide both members of this last equality by h and then let h approach zero. We obtain

$$(18) \quad \frac{\partial f(x, y)}{\partial p} = \frac{\partial f}{\partial x} \cos(px) + \frac{\partial f}{\partial y} \cos(py).$$



FIG. 8.

Let now n denote the direction of the inner normal to our curve C at a point (x, y) of C . Then, according to (18),

$$\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x} \cos(nx) + \frac{\partial u}{\partial y} \cos(ny).$$

¹ See Osgood, "Differential and Integral Calculus," rev. ed., p. 308, 1910.

² See Osgood, *Loc. cit.*, p. 292.




FIG. 7.

But

$$\cos(nx) = -\frac{dy}{ds}, \cos(ny) = \frac{dx}{ds},$$

then

$$\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial x} \frac{\partial y}{\partial s} + \frac{\partial u}{\partial y} \frac{\partial x}{\partial s}.$$

Thus, in Green's theorem, we may put

$$\int_{(C)} v \left(\frac{\partial u}{\partial y} \frac{dx}{ds} - \frac{\partial u}{\partial x} \frac{dy}{ds} \right) ds = \int_{(C)} v \frac{\partial u}{\partial n} ds.$$

We have then the following theorem:

Theorem VII.—*If u belongs to the class c'' and v to the class c' on I , then*

$$(19) \quad \int_{(C)} v \frac{\partial u}{\partial n} ds = \int_I \int v \Delta u dx dy - \int_I \int \left(\frac{\partial u}{\partial x} \frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \frac{\partial v}{\partial y} \right) dx dy.$$

We now apply this theorem to two special cases:

Case I.— $v = 1$, $\Delta u = 0$ on I . Then we have the theorem:

Theorem VIII.—*If u is harmonic on I , then*

$$(20) \quad \int_{(C)} \frac{\partial u}{\partial n} ds = 0.$$

Case II.— $v = u$, $\Delta u = 0$ on I . Then we have the

Theorem IX.—*If u is harmonic on I , then*

$$-\int_{(C)} u \frac{\partial u}{\partial n} ds = \int_I \int \left\{ \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right\} dx dy.$$

Corollary I.—*If u is harmonic on I and vanishes along the boundary of C , then $u \equiv 0$ on I .*

For, by these hypotheses,

$$= 0 \int \int \left\{ \left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right\} dx dy,$$

hence $\frac{\partial u}{\partial x} \equiv 0$, $\frac{\partial u}{\partial y} \equiv 0$ and, therefore, $u = \text{constant}$ on I .

But u is continuous and vanishes on the boundary C of I , hence $u \equiv 0$.

Corollary II.—If u is harmonic on I and $\frac{\partial u}{\partial n} = 0$ on the boundary C of I , then $u = \text{constant}$ on I .

Notice that in this case we cannot draw the conclusion $u \equiv 0$.

37. The Analogue of Theorem IX for the Exterior Region. Under the assumption that u is harmonic on E , it follows that u is harmonic in a region E_o exterior to C and interior to a circle S around the origin which includes C . From Theorem IX we would then have

$$(21) \quad \int \int \int \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy = - \int_C u \frac{\partial u}{\partial n_e} ds - \int_S u \frac{\partial u}{\partial n} ds,$$

where the normals must be drawn toward the interior of the region E_o . We now let the radius r of the circle S approach ∞ and examine the limits of the three terms in (21). The first of the single integrals remains unchanged. In order to see what happens to the second, make a transformation to polar coordinates. Then $u(x, y)$ becomes $U(r, \theta)$,

$ds = rd\theta$, and $\frac{\partial u}{\partial n} = -\frac{\partial u}{\partial r}$, since the




FIG. 9.

normal is opposite in direction to the radius r . Then the second single integral becomes

$$-\int_0^{2\pi} rU \frac{\partial U}{\partial r} d\theta.$$

Let us now add the assumption that

$$\lim_{r \rightarrow \infty} rU \frac{\partial U}{\partial r} = 0, \text{ uniformly with respect to } \theta.$$

Then $-\int_0^{2\pi} rU \frac{\partial U}{\partial r} d\theta$ approaches 0 as r approaches ∞ , and, accordingly, also the double integral on the left-hand side of (21) approaches a determinate limit denoted by $\iint_E \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy$. Hence we have the theorem:

Theorem X.—*If u is harmonic on E and*

$$(22) \quad \lim_{r \rightarrow \infty} rU \frac{\partial U}{\partial r} = 0, \text{ uniformly with respect to } \theta, \text{ then}$$

$$(23) \quad \iint_E \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy = - \int_C u \frac{\partial u}{\partial n_e} ds.$$

As before, we obtain the two corollaries:

Corollary I.—*If u is harmonic on E and (22) still holds, and, moreover, $u = 0$ on C , then $u \equiv 0$ on E .*

Corollary II.—*If u is harmonic on E and (22) still holds, and, moreover, $\frac{\partial u}{\partial n_e} = 0$ on C , then $u = \text{constant on } E$.*




FIG. 10.

38. Generalization of the Preceding.—

In the sequel we shall need a modification of the preceding theorems for the case where u is harmonic on the interior but not upon the boundary of C . Let us designate the interior without the boundary C by I' . Construct a closed curve C_ϵ .

$$C_\epsilon: \quad x = \xi(s, \epsilon), \quad y = \eta(s, \epsilon)$$

of the same character as C interior to C and such that

$$\lim_{\epsilon \rightarrow 0} C_\epsilon = C, \text{ uniformly as to } s.$$

Denote by I_ϵ the interior of C_ϵ plus the boundary C_ϵ . Theorem IX applies for this region and we have

$$\int \int_{I_\epsilon} \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy = - \int_{\bar{C}_\epsilon} u \frac{\partial u}{\partial n} ds.$$

Let us now impose the following three conditions (A):

- 1) $\lim_{\epsilon \rightarrow 0} u = u_i$, uniformly as to s .
- 2) $\lim_{\epsilon \rightarrow 0} \frac{\partial u}{\partial n} = \frac{\partial u_i}{\partial n}$, uniformly as to s .
- 3) $\left| \frac{\partial u}{\partial x} \right|, \left| \frac{\partial u}{\partial y} \right|$ are bounded on I' .

Then we obtain the following theorem:

Theorem XI.—*If u is harmonic on I' and satisfies the three conditions (A), then*

$$(24) \quad \int \int_I \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy = - \int_{\bar{C}} u_i \frac{\partial u_i}{\partial n} ds.$$

The following two corollaries follow as before.

Corollary I.—*If we add to the hypotheses of the theorem that $u_i = 0$ along C , then $u \equiv 0$ on I' .*

Corollary II.—*If we add to the hypotheses of the theorem that $\frac{\partial u_i}{\partial n} = 0$ along C , then $u = \text{constant}$ on I' .*

Make hypotheses similar to (A) for the exterior and call them (B). Then we get for the exterior minus the boundary C , which we designate by E' a corresponding theorem with two corollaries.

Theorem XII.—*If u is harmonic on E' and satisfies the conditions (B), then*

$$\int \int_E \left[\left(\frac{\partial u}{\partial x} \right)^2 + \left(\frac{\partial u}{\partial y} \right)^2 \right] dx dy = - \int_{\bar{C}} u_\epsilon \frac{\partial u_\epsilon}{\partial n_\epsilon} ds.$$

Corollary I.—If we add to the hypotheses of the theorem that $u_e = 0$ along C , then $u \equiv 0$ on E' .

Corollary II.—If we add to the hypotheses of the theorem that $\frac{\partial u_e}{\partial n_e} = 0$ on C , then $u = \text{constant}$ on E' .

IV. LOGARITHMIC POTENTIAL OF A DOUBLE LAYER

39. Definition.—We suppose that C

$$C: \quad x = \xi(s), \quad y = \eta(s), \quad s = 0, \dots, l$$

has the properties 1) . . . 4) of §36. Let (ξ, η) be a point on C , (x, y) a fixed point not on C , r their distance:

$$r = \sqrt{(x - \xi)^2 + (y - \eta)^2},$$

ν the inner normal to C at (ξ, η) , $\mu(s)$ a continuous function on C . Then the definite integral

$$(25) \quad w(x, y) = \int_0^l \mu(s) \frac{\partial \log \frac{1}{r}}{\partial \nu} ds$$

for physical reasons is called the *logarithmic potential of a double layer* of density $\mu(s)$ distributed over the curve C

We obtain as follows a more explicit form for $\frac{\partial}{\partial \nu} \log \frac{1}{r}$.

$$(26) \quad \frac{\partial}{\partial \nu} \log \frac{1}{r} = \cos(\nu\xi) \frac{\partial}{\partial \xi} \log \frac{1}{r} + \cos(\nu\eta) \frac{\partial}{\partial \eta} \log \frac{1}{r}$$

$$(27) \quad \begin{aligned} &= \frac{1}{r} \left[\frac{x - \xi}{r} \cos(\nu\xi) + \frac{y - \eta}{r} \cos(\nu\eta) \right] \\ &= \frac{1}{r} \left[\cos(r\xi) \cos(\nu\xi) + \cos(r\eta) \cos(\nu\eta) \right] \\ &= \frac{\cos(r\nu)}{r}. \end{aligned}$$




FIG. 11.

But from (27), since $\cos(\nu\xi) = -\eta'(s)$, $\cos(\nu\eta) = \xi'(s)$, we have

$$\frac{\partial}{\partial\nu} \log \frac{1}{r} = \frac{\left[y - \eta(s) \right] \xi'(s) - \left[x - \xi(s) \right] \eta'(s)}{r^2}$$

Thus $w(x, y)$ may be written in the form

$$w(x, y) = \int_0^l \mu(s) \frac{\cos(r\nu)}{r} ds,$$

or

$$(28) \quad w(x, y) = \int_0^l \mu(s) \frac{\left[y - \eta(s) \right] \xi'(s) - \left[x - \xi(s) \right] \eta'(s)}{\left[x - \xi(s) \right]^2 + \left[y - \eta(s) \right]^2} ds,$$

which shows explicitly the dependence of the integrand upon the parameters x and y .

40. Properties of $w(x, y)$ at Points Not on C.—Put

$$u(x, y) = \log \frac{1}{r}.$$

Then we have

$$\begin{aligned} \frac{\partial^2 u}{\partial x^2} &= -\frac{1}{r^2} + \frac{2(x - \xi)^2}{r^4} \\ \frac{\partial^2 u}{\partial y^2} &= -\frac{1}{r^2} + \frac{2(y - \eta)^2}{r^4}. \end{aligned}$$

Whence by adding we find $\Delta u = 0$. That is, $\log \frac{1}{r}$ is harmonic. Further

$$\frac{\partial^2}{\partial x^2} \left(\frac{\partial}{\partial \xi} \log \frac{1}{r} \right) = \frac{\partial}{\partial \xi} \left(\frac{\partial^2}{\partial x^2} \log \frac{1}{r} \right)$$

and

$$\frac{\partial^2}{\partial y^2} \left(\frac{\partial}{\partial \xi} \log \frac{1}{r} \right) = \frac{\partial}{\partial \xi} \left(\frac{\partial^2}{\partial y^2} \log \frac{1}{r} \right).$$

Whence, adding, we find

$$\Delta \left(\frac{\partial}{\partial \xi} \log \frac{1}{r} \right) = \frac{\partial}{\partial \xi} \left(\Delta \log \frac{1}{r} \right) = \frac{\partial}{\partial \xi} \Delta u = 0.$$

Therefore $\frac{\partial}{\partial \xi} \log \frac{1}{r}$ is harmonic. Similarly, we can show that $\frac{\partial}{\partial \eta} \log \frac{1}{r}$ is harmonic. Hence it follows from (26) that $\frac{\partial}{\partial \nu} \log \frac{1}{r}$ is harmonic. That is,

$$(29) \quad \Delta \left(\frac{\partial}{\partial \nu} \log \frac{1}{r} \right) = 0.$$

Let us now compute $\Delta w(x, y)$. We have

$$(30) \quad \begin{aligned} \frac{\partial^2}{\partial x^2} w(x, y) &= \frac{\partial^2}{\partial x^2} \int_0^l \mu(s) \frac{\partial}{\partial \nu} \log \frac{1}{r} ds \\ &= \int_0^{r^*} \mu(s) \frac{\partial^2}{\partial x^2} \left(\frac{\partial}{\partial \nu} \log \frac{1}{r} \right) ds, \end{aligned}$$

since the rules for the differentiation of a definite integral with respect to a parameter are applicable. Likewise, we have

$$(31) \quad \frac{\partial^2}{\partial y^2} w(x, y) = \int_0^l \mu(s) \frac{\partial^2}{\partial y^2} \left(\frac{\partial}{\partial \nu} \log \frac{1}{r} \right) ds.$$

Add (30) and (31) and take account of (29). We find

$$\Delta w(x, y) = \int_0^l \mu(s) \Delta \left(\frac{\partial}{\partial \nu} \log \frac{1}{r} \right) ds = 0.$$

From the explicit expression (28) we see that $w(x, y)$ is single-valued and continuous with all of its derivatives at every point (x, y) not on C . Thus we have proved the following theorem:

Theorem XIII.—The function

$$w(x, y) = \int_0^l \mu(s) \frac{\partial}{\partial \nu} \log \frac{1}{r} ds$$

is single-valued and continuous with all of its derivatives at every point (x, y) not on C , and in the same domain w is harmonic:

$$\Delta w(x, y) = 0.$$

41. Behavior of $w(x, y)$ on C .—In the preceding discussion (x, y) was supposed *not* to lie on C . Let now (x, y) coincide with a point (x_o, y_o) on C , corresponding to a value $s = s_o$, so that

$$x_o = \xi(s_o), \quad y_o = \eta(s_o).$$

Then the integrand becomes indeterminate. The indeterminate expression which we desire to investigate is

$$\frac{\cos(r_o v)}{r_o} = \frac{\left[\eta(s_o) - \eta(s) \right] \xi'(s) - \left[\xi(s_o) - \xi(s) \right] \eta'(s)}{\left[\xi(s_o) - \xi(s) \right]^2 + \left[\eta(s_o) - \eta(s) \right]^2}.$$

To evaluate, apply Taylor's remainder theorem to both numerator and denominator, stopping at the derivatives of the second order. A factor $(s - s_o)^2$ appears in both and cancels. The limit of what remains as $s \rightarrow s_o$ is

$$\frac{\xi' \eta'' - \xi'' \eta'}{2(\xi'^2 + \eta'^2)} = \frac{1}{2R_o},$$

where R_o is the radius of curvature of C at (x_o, y_o) . $w(x, y)$ is thus defined and has a determinate finite value at every point (x_o, y_o) of C :

$$w(x_o, y_o) = \int_0^l \mu(s) \frac{\left[\eta(s_o) - \eta(s) \right] \xi'(s) - \left[\xi(s_o) - \xi(s) \right] \eta'(s)}{\left[\xi(s_o) - \xi(s) \right]^2 + \left[\eta(s_o) - \eta(s) \right]^2} ds.$$

The existence of $w(x, y)$ on the boundary of C does not, however, imply that $w(x, y)$ remains continuous, as (x, y)

crosses the boundary C . On the contrary, $w(x, y)$ undergoes a discontinuity as (x, y) crosses C . Let us denote by P_i an interior point of C and by P_e an exterior point (x, y) in the vicinity of $P_o(x_o, y_o)$. If P_i approaches P_o , then $w(x, y)$ approaches a definite finite limit $w_i(x_o, y_o)$. If P_e approaches P_o ,

then $w(x, y)$ approaches a definite finite limit $w_e(x_o, y_o)$. Between these quantities and $w(x_o, y_o)$ defined above the following relations hold:

$$(32) \quad \begin{aligned} w_i(x_o, y_o) &= w(x_o, y_o) + \pi \cdot \mu(s_o) \\ w_e(x_o, y_o) &= w(x_o, y_o) - \pi \cdot \mu(s_o).^1 \end{aligned}$$

For future use we give here a proof for the special case $\mu(s) \equiv 1$. Let us use the notation

$$\xi = \xi + i\eta, \quad z = x + iy$$

and consider the integral H along C :

$$H_c = \int_C \frac{d\xi}{\xi - z}.$$

By Cauchy's first integral theorem

$$H_c = \begin{cases} 2\pi i, & xy \text{ an interior point of } C. \\ 0, & xy \text{ an exterior point of } C. \end{cases}$$

If we separate the real and the imaginary parts of the integral, we obtain

$$\begin{aligned} H_c &= \int \frac{(\xi - x)d\xi + (\eta - y)d\eta}{(x - \xi)^2 + (y - \eta)^2} \\ &\quad + i \int \frac{(y - \eta)d\xi - (x - \xi)d\eta}{(x - \xi)^2 + (y - \eta)^2} \equiv V_c + iW_c. \end{aligned}$$

¹ For a proof of these statements see HORN, "Einführung in die Theorie der Partiellen Differential-Gleichungen," §52.




FIG. 12.

Hence, by equation (28), $w(x, y)$ is the coefficient of i in the integral H_c . Thus we get

$$w(x, y) = \begin{cases} 2\pi, & xy \text{ an interior point of } C. \\ 0, & xy \text{ an exterior point of } C. \end{cases}$$

But if $w(x, y) = 2\pi$ constantly as x, y varies, then

$$\lim_{P_i \rightarrow P_o} w(x, y) = w_i(x_o, y_o) = 2\pi.$$

In like manner

$$\lim_{P_o \rightarrow P_e} w(x, y) = w_e(x_o, y_o) = 0.$$

We have now, in order to complete the proof, to compute $w(x_o, y_o)$ for a point (x_o, y_o) on C . Draw about P_o an arc α of a circle cutting C in Q and R . This arc will subtend an angle ω at P_o . Designate by C' the path C minus the arc QP_oR . Then $H_c + H_\alpha = 0$, since P_o is an exterior point for this path. But in the elements of the theory of functions of a complex variable it is shown that

$$H_\alpha = \int_\alpha \frac{d\xi}{\xi - z_o} = -i\omega,$$




FIG. 13.

whence

$$H_{C'} = i\omega.$$

Therefore,

$$W_{C'} = \omega.$$

Let the radius of the arc α be ρ . Then, as $\rho \rightarrow 0$,

$$W_{C'} \rightarrow W_C = w(x_o, y_o).$$

Now the integral W_C is convergent as shown above and $\omega \rightarrow \pi$. Therefore

$$w(x_o, y_o) = \pi.$$

The equations (32) are satisfied for these values of $w(x_o, y_o)$, $w_i(x_o, y_o)$, and $w_e(x_o, y_o)$ for the special case $\mu(s) \equiv 1$.

42. Behavior of $\frac{\partial w}{\partial n}$ on the Boundary C and at Infinity.—


At the point $P_o(x_o, y_o)$ draw the interior normal n_i and take on it a point $P_i(x, y)$. From Theorem

XIII we know that at P_i , $\frac{\partial w(x, y)}{\partial n_i}$ is definite and finite. Let the point P_i approach P_o along n_i and put

$$\lim_{P_i \rightarrow P_o} \frac{\partial w}{\partial n_i} = \frac{\partial w_i}{\partial n_i}, \text{ if the limit exists,}$$

$$\lim_{P_e \rightarrow P_o} \frac{\partial w}{\partial n_i} = \frac{\partial w_e}{\partial n_i}, \text{ if the limit exists,}$$

FIG. 14.



P_e designating a point on the prolongation of n_i beyond P_o . Then the following theorem is true.¹

Theorem XIV.—If one of the two limits $\frac{\partial w_i}{\partial n_i}$, $\frac{\partial w_e}{\partial n_i}$, exists, then also the other exists and

$$(33) \quad \frac{\partial w_i}{\partial n_i} = \frac{\partial w_e}{\partial n_i}.$$

If we denote by n_e the exterior normal to C at P_o , then

$$\frac{\partial}{\partial n_i} = -\frac{\partial}{\partial n_e}$$

from the definition of the directional derivative. Hence (33) may also be written

$$(34) \quad \frac{\partial w_i}{\partial n_i} = -\frac{\partial w_e}{\partial n_e}.$$

If we introduce polar coordinates r, θ , then $w(x, y)$ becomes $W(r, \theta)$. Then the following theorem is true:¹

¹ For proof, see HORN, *Loc. cit.*, §54.

Theorem XV.—

$$\lim_{r \rightarrow \infty} W(r, \theta) = 0, \text{ uniformly as to } \theta.$$

$$\lim_{r \rightarrow \infty} rW \frac{\partial W}{\partial r} = 0, \text{ uniformly as to } \theta.$$

43. Case Where w_i or w_e Vanish Along C.—We can now prove the following theorem:

Theorem XVI.—*If $w_i(x, y) \equiv 0$ along C , then*

$$\mu(s) \equiv 0.$$

Proof.—Theorem XIII assures us that $w(x, y)$ is harmonic on I' . By hypothesis $w_i(x, y) \equiv 0$ along C . Taking it for granted that the conditions (A) are always satisfied by w , it follows from Corollary I to Theorem XI that

$$w(x, y) \equiv 0 \text{ on } I'.$$

Hence it follows at once that $\frac{\partial w}{\partial n_i} \equiv 0$ for every interior point of C . Therefore, $\frac{\partial w_i}{\partial n_i}$ exists along C and vanishes there. Hence, also by Theorem XIV, $\frac{\partial w_e}{\partial n_e}$ exists along C and vanishes there. By Theorem XV, $\lim_{r \rightarrow \infty} rW \frac{\partial W}{\partial r} = 0$, uniformly as to θ . Hence, if we take it for granted that the conditions (B) on w are satisfied, then it follows from Corollary II to Theorem XII that w is constant on E' . We see from Theorem XV that this constant must be zero:

$$w(x, y) \equiv 0 \text{ on } E'.$$

Whence it follows that $w_e(x, y) = 0$ on C . If we now apply (32) we find

$$w_i(x_o, y_o) - w_e(x_o, y_o) = 2\pi\mu(s_o) = 0,$$

uniformly as to s_o . Therefore

$$\mu(s_o) \equiv 0, \text{ uniformly as to } s_o.$$

A similar theorem holds with respect to the limit w_e :

Theorem XVII.—*If $w_e(x, y) \equiv 0$ along C , then*

$$\mu(s) = \text{constant}.$$

Proof.—Theorem XIII assures us that w is harmonic on E' . By hypothesis $w_e(x, y) \equiv 0$ along C . Furthermore, $\lim_{r \rightarrow \infty} rW \frac{\partial W}{\partial r} = 0$, uniformly as to θ , according to Theorem XV. Taking it for granted that the conditions (B) of §38 are always satisfied by w , it follows from Corollary I to Theorem XII that $w(x, y) \equiv 0$ on E' . Hence it follows at once that $\frac{\partial w}{\partial n_e} \equiv 0$ for every point exterior to C . There-

fore $\frac{\partial w_e}{\partial n_e}$ exists along C and vanishes there. Hence also, by Theorem XIV, $\frac{\partial w_i}{\partial n_i}$ exists along C and vanishes there.

Taking it for granted that the conditions (A) are always satisfied by w , it follows from Corollary II to Theorem XI that

$$w(x, y) = \text{constant on } I'.$$

Whence it follows that $w_i(x, y) = \text{constant on } C$. If we now apply (32), we find

$$w_i(x_o, y_o) - w_e(x_o, y_o) = 2\pi\mu(s_o) = \text{constant},$$

uniformly as to s_o . Therefore,

$$\mu(s_o) = \text{constant, uniformly as to } s_o.$$

V. FREDHOLM'S SOLUTION OF DIRICHLET'S PROBLEM

44. Dirichlet's Problem.—We formulate Dirichlet's problem as follows:

Given.—1) A closed curve C

$$(35) \quad \begin{aligned} C : \quad x &= \xi(s), y = \eta(s), 0 \leq s \leq l \\ \xi(0) &= \xi(l), \eta(0) = \eta(l) \end{aligned}$$

with the properties 1) . . . 4) of §36.

2) A function $F(s)$, continuous on C for $0 \leq s \leq l$, and

$$(36) \quad F(0) = F(l)$$

Required.—A function $u(x, y)$ such that

- $\alpha)$ u is harmonic on I' , and
- $\beta)$ $u_i(x_o, y_o) = F(s_o)$, uniformly as to s_o ,

where $u_i(x_o, y_o)$ is the limit approached by $u(x, y)$ as the point (x, y) approaches from the interior a point (x_o, y_o) of parameter s_o on the boundary C .

45. Reduction to an Integral Equation. First Method.—The function $w(x, y)$:

$$w(x, y) = \int_0^l \mu(s) \frac{\partial}{\partial \nu} \log \frac{1}{r} ds$$

is harmonic on I' and thus satisfies the condition (α) for every choice of $\mu(s)$ for which $\mu(s)$ is continuous on $[0l]$ and

$$\mu(0) = \mu(l).$$

The function $w(x, y)$ will then furnish a solution of Dirichlet's problem if $\mu(s)$ can be so determined that the condition (β) is satisfied:

$$(37) \quad w_i(x_o, y_o) = F(s_o), \text{ uniformly as to } s_o.$$

From the first of equations (32) we have

$$w_i(x_o, y_o) = w(x_o, y_o) + \pi\mu(s_o).$$

The substitution of this value of w_i in (37) gives

$$(38) \quad w(x_o, y_o) + \pi\mu(s_o) = F(s_o).$$




FIG. 15.

If in this equality we substitute for $w(x_o, y_o)$ its explicit expression as given in §41, we obtain

$$(39) \quad \int_0^l \mu(s) \frac{\left[\eta(s_o) - \eta(s) \right] \xi'(s) - \left[\xi(s_o) - \xi(s) \right] \eta'(s)}{\left[\xi(s_o) - \xi(s) \right]^2 + \left[\eta(s_o) - \eta(s) \right]^2} ds + \pi \mu(s_o) = F(s_o).$$

Divide through by π . Then (39) becomes an integral equation for the determination of $\mu(s_o)$ with the kernel

$$(40) \quad K(s_o, s) = \frac{1}{\pi} \frac{\left[\eta(s_o) - \eta(s) \right] \xi'(s) - \left[\xi(s_o) - \xi(s) \right] \eta'(s)}{\left[\xi(s_o) - \xi(s) \right]^2 + \left[\eta(s_o) - \eta(s) \right]^2}.$$

Put $\frac{F(s_o)}{\pi} = f(s_o)$. Then the integral equation takes the standard form

$$(41) \quad \mu(s_o) = f(s_o) - \int_0^l K(s_o, s) \mu(s) ds.$$

This is a special case of the integral equation with a parameter λ , for which $\lambda = -1$.

Thus, in order that $u = w(x, y)$ may be a solution of Dirichlet's problem, it is *necessary* that the density $\mu(s)$ satisfy the integral equation (41). This condition is also *sufficient*. For, suppose $\mu(s)$ to be a continuous solution of (41). Then we have

$$\begin{aligned} K(0, s) &= K(l, s) \text{ on account of (35), and} \\ f(0) &= f(l) \text{ on account of (36), and hence} \\ \mu(0) &= \mu(l) \text{ from (41).} \end{aligned}$$

The function $w(x, y)$ formed with this function $\mu(s)$ will then satisfy (38) and, therefore, also (37) and, consequently, $u = w(x, y)$ will be a solution of Dirichlet's problem.

Second Method.—From equations (32) we find

$$(42) \quad w_i(x_o, y_o) - w_e(x_o, y_o) = 2\pi\mu(s_o), \text{ and}$$

$$w_i(x_o, y_o) + w_e(x_o, y_o) = 2w(x_o, y_o),$$

which by §41

$$= 2 \int_0^l \mu(s) \frac{\cos(r_o\nu)}{r_o} ds.$$

We now seek as a solution of Dirichlet's problem a function u which is harmonic everywhere except on C , and which upon C satisfies the condition

$$(43) \quad u_i + hu_e = F(s_o) + hG(s_o),$$

where h is an arbitrary parameter. For $h = 0$, we have the interior problem, and for $h = \infty$ the exterior problem.

The function $w(x, y)$ is harmonic everywhere except on C . This function will then be a solution of Dirichlet's problem, provided $\mu(s)$ can be so determined that $u = w(x, y)$ satisfies (43). Solve now the equations (42) for $u_i = w_i$ and $u_e = w_e$ and substitute in (43). We find

$$(44) \quad \mu(s_o) = f(s_o) + \lambda \int_0^l \mu(s) \frac{\cos(r_o\nu)}{\pi r_o} ds$$

where $f(s_o) = \frac{hG(s_o) + F(s_o)}{\pi(1-h)}$, $\lambda = \frac{h+1}{h-1}$.

This is an integral equation of the second kind with a parameter λ . The kernel is $\frac{\cos(r_o\nu)}{\pi r_o} \equiv K(s_o, s)$ [see (40)].

For $h = 0$, we have the interior problem, but for $h = 0$ we have $\lambda = -1$ and (44) reduces to (41). For $h = \infty$, we have the exterior problem, but for $h = \infty$ we have $\lambda = +1$, and (44) reduces to

$$(45) \quad \mu(s_o) = \frac{G(s_o)}{\pi} + \int_0^l K(s_o, s) \mu(s) ds.$$

We remark that, if $F(s_o) \equiv 0$ on C and $h = 0$, then (44) becomes a homogeneous linear integral equation with $\lambda = -1$.

46. Solution of the Integral Equation.—The kernel $K(s_o, s)$ is real, continuous, and $\not\equiv 0$ in the region G ,

$$G: \quad 0 \leq s_o \leq l, \quad 0 \leq s \leq l,$$

and thus the preceding theory of integral equations is applicable. From Fredholm's first fundamental theorem we know that if $D(-1) \neq 0$ then (41) has one and only one continuous solution.

We show first that $D(-1) \neq 0$. For this purpose we use Fredholm's second fundamental theorem, from which it follows that the corresponding homogeneous integral equation

$$(46) \quad \mu(s_o) + \int_0^l K(s_o, s)\mu(s)ds = 0$$

has no other solution than the trivial one $\mu \equiv 0$, if $D(-1) \neq 0$; while if $\lambda = -1$ is a root of $D(\lambda) = 0$ of index q , then (46) has ∞^q solutions. Hence, if (46) has no other solution than $\mu \equiv 0$, then $D(-1) \neq 0$.

Multiply the members of (46) by π . Then

$$\int_0^l \pi K(s_o, s)\mu(s)ds + \pi\mu(s_o) = 0,$$

uniformly as to s_o . This last equation, by (32), after taking account of our notations in §41 and equations (40), can be written

$$w_i(x_o, y_o) = 0, \text{ uniformly as to } s_o.$$

But, by Theorem XVI, if $w_i(x_o, y_o) = 0$ along C , then

$$\mu(s) \equiv 0.$$

Therefore, (46) has no other continuous solution than $\mu \equiv 0$ and, consequently,

$$D(-1) \neq 0.$$

Thus we see that $\lambda = -1$ is not a characteristic constant for the kernel $K(s_o, s)$. Hence (41) has one and only one

continuous solution $\mu(s_o)$ representable by Fredholm's formula

$$\mu(s_o) = f(s_o) + \int_0^l \frac{D(s_o, s; -1)}{D(-1)} f(s) ds.$$

Therefore, $w(x, y)$ with $\mu = \mu(s_o)$ is a solution of Dirichlet's problem. We state this result in the following theorem:

Theorem XVIII.—Given

1) A closed curve C

$$C: \quad x = \xi(s), \quad y = \eta(s), \quad 0 \leq s \leq l \\ \xi(0) = \xi(l), \quad \eta(0) = \eta(l)$$

with the properties 1) . . . 4) of §36.

2) A function $F(s)$, continuous on C for $0 \leq s \leq l$,

$$F(0) = F(l).$$

Then there exists a function $u(x, y)$ such that

a) u is harmonic on I' , and

b) $u_i(x_o, y_o) = F(s_o)$, uniformly as to s_o , where $u_i(x_o, y_o)$ is the limit approached by $u(x, y)$ as the point (x, y) approaches from the interior a point (x_o, y_o) of parameter s_o on the boundary C . This function is given by

$$u = \int_0^l \mu(s) \frac{\partial}{\partial \nu} \log \frac{1}{r} ds,$$

where $\mu(s)$ is the unique solution of (41) and is given by

$$\mu(s_o) = f(s_o) + \int_0^l \frac{D(s_o, s; -1)}{D(-1)} f(s) ds,$$

where $\pi f(s_o) = F(s_o)$.

47. Index of $\lambda = 1$ for $K(s_o, s)$.—We have seen that $\lambda = -1$ is not a characteristic constant of $K(s_o, s)$. We shall now prove the following theorem:

Theorem XIX.— $\lambda = +1$ is a characteristic constant of $K(s_o, s)$ of index 1.

Proof.—According to Fredholm's second fundamental theorem, the statement to be proved is equivalent to the statement that the homogeneous equation

$$(47) \quad \mu(s_o) = \int_0^l K(s_o, s)\mu(s)ds$$

has ∞^1 solutions. Now, if we make use of equations (40) and (39) and the explicit expression for $w(x_o, y_o)$ given in §41, we see that (47) is equivalent to

$$(48) \quad w(x_o, y_o) = \pi\mu(s_o).$$

But from the proof given for the equations (32) for the special case $\mu(s) \equiv 1$, we see that $\mu(s) \equiv 1$ is a solution of (48), for when $\mu(s) \equiv 1$, we have

$$w(x_o, y_o) = \pi = \pi\mu(s_o).$$

This is a solution of (47) which is continuous and not identically zero. Hence $\lambda = 1$ is a characteristic constant of $K(s_o, s)$.

We will now determine the index of $\lambda = +1$ for $K(s_o, s)$. To this end we determine all solutions of (47) or of its equivalent (48). Now, on account of the second of equations (32), our equation (48) above reduces to

$$w_e(x_o, y_o) = 0.$$

But, according to Theorem XVII, the most general solution of this equation, and, therefore, of (47), is

$$\mu(s_o) = \text{constant}.$$

Therefore, (47) has ∞^1 solutions, which shows that the index of the characteristic constant $\lambda = +1$ of $K(s_o, s)$ is 1.

VI. LOGARITHMIC POTENTIAL OF A SIMPLE LAYER

48. Definition.—We have given a closed curve C

$$C: \quad x = \xi(s), \quad y = \eta(s), \quad s = 0, \dots, l$$

having the properties 1) . . . 4) of §36. Let r represent the distance between a point (ξ, η) on C and a point (x, y) not on C , $k(s)$ a continuous function on C . Then the definite integral

$$\begin{aligned} v(x, y) &= \int_0^l k(s) \log \frac{1}{r} ds \\ &= -\frac{1}{2} \int_0^l k(s) \log \left\{ [x - \xi(s)]^2 + [y - \eta(s)]^2 \right\} ds \end{aligned}$$




FIG. 16.

is called the *logarithmic potential of a simple layer* of density $k(s)$ distributed over the curve C .

49. Properties of $v(x, y)$.—

The $\log \left\{ [x - \xi(s)]^2 + [y - \eta(s)]^2 \right\}$ is a regular analytic function of x and y at all points not on C and hence the definite integral as a function of the parameters x and y defines a function $v(x, y)$ which is continuous with all of its derivatives in I' and E' . Further, $v(x, y)$ is harmonic in the same region. For, from the theory of definite integrals

$$\begin{aligned} \Delta v &= \Delta \int_0^l k(s) \log \frac{1}{r} ds \\ &= \int_0^l k(s) \Delta \log \frac{1}{r} ds. \end{aligned}$$

But it has been previously shown that

$$\Delta \log \frac{1}{r} = 0.$$

Therefore

$$\Delta v(x, y) = 0.$$

Hence we have proved the following theorem:

Theorem XX.—*The logarithmic potential $v(x, y)$:*

$$(49) \quad v(x, y) = \int_0^l k(s) \log \frac{1}{r} ds$$

of a simple layer is continuous with its derivatives of all orders, and is harmonic, on I' and E' :

$$\Delta v(x, y) = 0.$$

Let us now investigate the behavior of $v(x, y)$ on the boundary C at any point (x_o, y_o) with parameter s_o .

$$x_o = \xi(s_o), \quad y_o = \eta(s_o).$$

From the explicit expression for $v(x, y)$:

$$v(x_o, y_o) = -\frac{1}{2} \int_0^l k(s) \times \log \left\{ \left[\xi(s_o) - \xi(s) \right]^2 + \left[\eta(s_o) - \eta(s) \right]^2 \right\} ds$$

we see that when $s = s_o$, the logarithm which appears in the integrand becomes infinite. Thus we see that we have to do with an improper definite integral. Apply Taylor's remainder theorem to the expression of which the logarithm is taken. We obtain

$$(s - s_o)^2 A,$$

where A does not become zero as s approaches s_o . Then the integrand becomes infinite like $\log(s - s_o)$ as $s \rightarrow s_o$. From the theory of improper definite integrals we know that then this definite integral remains definite and finite.

We state without proof¹ the two following theorems, in which $P_i(x, y)$ denotes an interior, $P_e(x, y)$ an exterior point in the vicinity of P_o .

¹ For a proof, see HORN, §53.

Theorem XXI.—*The limit of $v(x, y)$ as $P_i \rightarrow P_o$ exists:*

$$\lim_{P_i \rightarrow P_o} v(x, y) = v_i(x_o, y_o).$$

The limit of $v(x, y)$ as $P_e \rightarrow P_o$ exists:

$$\lim_{P_e \rightarrow P_o} v(x, y) = v_e(x_o, y_o);$$

and $v_i(x_o, y_o) = v_e(x_o, y_o) = v(x_o, y_o)$.

That is, $v(x, y)$ remains continuous as the point $P(x, y)$ crosses the boundary C . In this respect the behavior of the function $v(x, y)$ is simpler than that of $w(x, y)$ for a double layer.

Construct for the point $P_o(x_o, y_o)$ the interior normal n_i to C and take on it the interior point P_i and on its exterior prolongation the exterior point P_e ; then it follows from Theorem XX that $\frac{\partial v}{\partial n_i}$ exists at P_i and at P_e . For the approach of P_i and P_e toward P_o along the normal the following theorem holds:

Theorem XXII.—*The limit of $\frac{\partial v}{\partial n_i}$ as $P_i \rightarrow P_o$ exists:*

$$\lim_{P_i \rightarrow P_o} \frac{\partial v}{\partial n_i} = \frac{\partial v_i}{\partial n_i} = -\pi k(s_o) + \int_0^l k(s) \frac{\cos(r_o n_i)}{r_o} ds.$$

The limit of $\frac{\partial v}{\partial n_i}$ as $P_e \rightarrow P_o$ exists:

$$\lim_{P_e \rightarrow P_o} \frac{\partial v}{\partial n_i} = \frac{\partial v_e}{\partial n_i} = +\pi k(s_o) + \int_0^l k(s) \frac{\cos(r_o n_i)}{r_o} ds,$$




FIG. 17.

where r_o is the distance from $P_o(x_o, y_o)$ to a variable point $[\xi(s), \eta(s)]$ of C .

By referring to the figure, we obtain for $\cos(r_o n_i)$ the expression

$$\begin{aligned}\cos(r_o n_i) &= \cos(r_o x) \cos(n_i x) + \cos(r_o y) \cos(n_i y) \\ &= -\frac{\xi(s) - x_o}{r_o} \eta'(s_o) + \frac{\eta(s) - y_o}{r_o} \xi'(s_o),\end{aligned}$$

and hence

$$\frac{\cos(r_o n_i)}{r_o} = \frac{\left[\eta(s) - \eta(s_o) \right] \xi'(s_o) - \left[\xi(s) - \xi(s_o) \right] \eta'(s_o)}{\left[\xi(s) - \xi(s_o) \right]^2 + \left[\eta(s) - \eta(s_o) \right]^2}.$$

VII. FREDHOLM'S SOLUTION OF NEUMANN'S PROBLEM

50. Neumann's Problem.—We formulate the second boundary problem of the potential theory, or Neumann's problem, as follows:

Given.—

- 1) A closed curve C ,

$$C: \quad \begin{aligned}x &= \xi(s), \quad y = \eta(s), \quad 0 \leq s \leq l \\ \xi(0) &= \xi(l), \quad \eta(0) = \eta(l)\end{aligned}$$

with the properties 1) . . . 4) of §36.

- 2) A function $F(s)$, continuous on C for $0 \leq s \leq l$:

$$F(0) = F(l).$$

Required.—A function $u(x, y)$ such that

- α) $\Delta u = 0$ on I' .
- β) $\frac{\partial u_i}{\partial n_i} = F(s_o)$, uniformly as to s_o .

51. Reduction to an Integral Equation. First Method.—The function $v(x, y)$ given by (49) is a function satisfying condition (α) for any choice of $k(s)$ which is continuous on

C. If it is possible to determine the function $k(s)$ so as to satisfy the condition (β):

$$(50) \quad \frac{\partial v_i}{\partial n_i} = F(s_o), \text{ uniformly as to } s_o,$$

then we will have a solution of our problem.

Equation (50), written explicitly, according to Theorem XXII becomes

$$(51) \quad \int_0^l k(s) \frac{\left[\eta(s) - \eta(s_o) \right] \xi'(s_o) - \left[\xi(s) - \xi(s_o) \right] \eta'(s_o)}{\left[\xi(s) - \xi(s_o) \right]^2 + \left[\eta(s) - \eta(s_o) \right]^2} ds \\ = F(s_o) + \pi k(s_o).$$

This is an integral equation for the determination of $k(s)$. Divide through by π and put

$$(52) \quad \tilde{K}(s_o, s) = \frac{1}{\pi} \frac{\left[\eta(s) - \eta(s_o) \right] \xi'(s_o) - \left[\xi(s) - \xi(s_o) \right] \eta'(s_o)}{\left[\xi(s) - \xi(s_o) \right]^2 + \left[\eta(s) - \eta(s_o) \right]^2}$$

and $f(s_o) = -\frac{F(s_o)}{\pi}$.

Then (51) becomes

$$(53) \quad k(s_o) = f(s_o) + \int_0^l \tilde{K}(s_o, s) k(s) ds.$$

Introduce a parameter λ and write

$$(54) \quad k(s_o) = f(s_o) + \lambda \int_0^l \tilde{K}(s_o, s) k(s) ds,$$

which becomes identical with (53) for $\lambda = +1$.

From (51) we see that $k(0) = k(l)$, since, from our hypothesis, $F(0) = F(l)$ and $\tilde{K}(0, s) = \tilde{K}(l, s)$.

Second Method.—From Theorem XXII we have

$$(55) \quad \begin{aligned} \frac{\partial v_i}{\partial n_i} &= -\pi k(s_o) + \int_0^l k(s) \frac{\cos(r_o n_i)}{r_o} ds \\ \frac{\partial v_e}{\partial n_i} &= +\pi k(s_o) + \int_0^l k(s) \frac{\cos(r_o n_i)}{r_o} ds. \end{aligned}$$

We now seek, as a solution of Neumann's problem, a function u which is harmonic everywhere except on C , and which on C satisfies the condition

$$(56) \quad h \frac{\partial u_i}{\partial n_i} + \frac{\partial u_e}{\partial n_i} = hF(s_o) + G(s_o),$$

where h is an arbitrary parameter. For $h = \infty$, we have the so-called interior case, while for $h = 0$ we have the exterior case. This condition is more general than (50) and for $h = \infty$ reduces to (50).

The function $v(x, y)$ is harmonic everywhere except on C . This function will then be a solution of Neumann's problem provided $k(s)$ can be so determined that $u = v(x, y)$ satisfies (56). Substitute now from (55) in (56) for

$$\frac{\partial v_i}{\partial n_i} = \frac{\partial u_i}{\partial n_i} \text{ and } \frac{\partial v_e}{\partial n_i} = \frac{\partial u_e}{\partial n_i}.$$

We find

$$(57) \quad k(s_o) = f(s_o) + \lambda \int_0^l k(s) \frac{\cos(r_o n_i)}{\pi r_o} ds,$$

$$\text{where } f(s_o) = \frac{G(s_o) + hF(s_o)}{\pi(1-h)}, \lambda = \frac{h+1}{h-1}.$$

This is an integral equation of the second kind with a parameter λ . The kernel is $\frac{\cos(r_o n_i)}{\pi r_o} \equiv \tilde{K}(s_o, s)$. For $h = 0$, we have the exterior problem, but for $h = 0$ we have $\lambda = -1$. For $h = \infty$, we have the interior problem, but for $h = \infty$ we have $\lambda = +1$ and (57) reduces to (53).

52. Solution of the Integral Equation.—To the equation (54) the Fredholm theory which we have developed can be

applied. We look, then, for a solution of (54) when $\lambda = +1$. Now $\lambda = +1$ is a characteristic constant of $\tilde{K}(s_o, s)$, for $\tilde{K}(s_o, s)$ can be obtained from the kernel $K(s_o, s)$ for Dirichlet's problem by interchanging s and s_o . Thus \tilde{K} and K are adjoint kernels. Then from Theorem IX, Chap. III, $\tilde{K}(s_o, s)$ and $K(s_o, s)$ have the same characteristic constants with the same indices. But $\lambda = +1$ has been shown by Theorem XIX to be a characteristic constant of index 1 for $K(s_o, s)$. Therefore $\lambda = +1$ is a characteristic constant of index 1 for $\tilde{K}(s_o, s)$. According to Fredholm's third fundamental theorem, (54) has, in general, no solution for $\lambda = +1$. But if $f(s_o)$ satisfies certain $q = 1$ conditions, then there will be a solution. This condition is

$$(58) \quad \int_0^l f(s) \bar{\varphi}(s) ds = 0,$$

where $\bar{\varphi}(s_o)$ is a solution of the associated homogeneous integral equation

$$(59) \quad \bar{\varphi}(s_o) = \int_0^l K(s_o, s) \bar{\varphi}(s) ds,$$

the kernel of which is the adjoint of the kernel of the integral equation (54) under consideration. But $\lambda = +1$ is a characteristic constant of $K(s_o, s)$ of index 1 and $\bar{\varphi}(s) \equiv 1$ is a solution of (59), as shown in §47. Therefore, condition (58) becomes

$$(60) \quad \int_0^l f(s) ds = 0.$$

Unless this condition is satisfied, there is no solution. If this condition is satisfied, then there are ∞^1 solutions. By referring to the definition of $f(s)$, we see that (60) gives the condition

$$\int_0^l F(s) ds = 0.$$

Thus we see that $v(x, y)$ is a solution of Neumann's problem. Furthermore, by referring to Theorems XX, XXI, and XXII, we see that $v(x, y)$ satisfies the conditions (A) of §38. Let $\bar{v}(x, y)$ be any other solution of Neumann's problem which satisfies the conditions (A). Form the difference

$$\mathfrak{a}(x, y) = \bar{v}(x, y) - v(x, y).$$

Then it follows that

- 1) $\Delta \mathfrak{a} = 0$ on I' .
- 2) $\frac{\partial \mathfrak{a}_i}{\partial n_i} = 0$ along C .

From Corollary II to Theorem XI it follows that

$$\mathfrak{a} = C \text{ (a constant) on } I'$$

Therefore,

$$\bar{v} = v + c.$$

Hence, there are ∞^1 solutions of Neumann's problem which differ only by an additive constant. Furthermore, these are the only solutions of the problem which satisfy the conditions (A). We have now the following theorem:

Theorem XXIII.—*Given a closed curve C satisfying the conditions 1) . . . 4) of §36*

$$C: \quad x = \xi(s), \quad y = \eta(s), \quad 0 \leq s \leq l \\ \xi(0) = \xi(l), \quad \eta(0) = \eta(l),$$

and the function $F(s)$ continuous on C , $F(0) = F(l)$, and

$$(61) \quad \int_0^l F(s) ds = 0,$$

then Neumann's problem has ∞^1 solutions

$$\bar{u} = u + c$$

where

$$u = \int_0^l k(s) \log \frac{1}{r} \, ds$$

and $k(s)$ is a solution of

$$k(s_o) = f(s_o) + \lambda \int_0^l \tilde{K}(s_o, s) k(s) \, ds$$

for $\lambda = 1$, where $\tilde{K}(s_o, s)$ is defined by (52). If the condition (61) is not satisfied, the problem has no solution.

CHAPTER V

HILBERT-SCHMIDT THEORY OF INTEGRAL EQUATIONS WITH SYMMETRIC KERNELS

Solution Expressed in Terms of a Set of Fundamental Functions

I. EXISTENCE OF AT LEAST ONE CHARACTERISTIC CONSTANT

53. Introductory Remarks.—The Fredholm theory of a linear integral equation with a parameter λ :

$$(1) \quad u(x) = f(x) + \lambda \int_a^b K(x, t)u(t)dt$$

has been developed under the assumptions

- A) $K(x, t)$ real.
- B) $K(x, t)$ continuous.
- C) $K(x, t) \neq 0$ on $R : a \leq x \leq b, a \leq t \leq b$.

For the Hilbert-Schmidt theory a fourth assumption is made that the kernel is symmetric.

$$(2) \quad D) \quad K(x, t) = K(t, x).$$

It is clear that the results of the Fredholm theory still hold. But, besides, a number of new results will follow from the additional condition D). These results were first obtained by Hilbert.

Fredholm obtained the solution of (1) by considering it as the limit of a set of linear equations. He did not carry through the limiting processes, but guessed at the solution and then showed independently of the linear equations that the guess was correct. Hilbert started from the finite system of linear equations, of which equations (1) may be con-

sidered as a limit, and actually carried through the limiting process in detail. E. Schmidt, in 1906, obtained Hilbert's results directly without using the limiting process. Hilbert and Schmidt do not make use of Fredholm's results. We shall, however, avail ourselves of these results whenever they lead to simplifications of the proofs.

The first fundamental theorem of Hilbert's theory is the following:

Theorem I.—*Every symmetric kernel has at least one characteristic constant (real or imaginary).*

Our proof of this theorem involves several lemmas which we will give first.

We give an example due to Kowalewski, which illustrates that a kernel which is not symmetric does not necessarily have a characteristic constant. The kernel

$$K(x, t) = \sin(\pi x) \cos(\pi t), \quad [01]$$

has no characteristic constants, if

$$(3) \quad u(x) = \lambda \int_0^1 \sin(\pi x) \cos(\pi t) u(t) dt$$

has no solution other than $u \equiv 0$.

Equation (3) may be written

$$u(x) = \lambda \sin \pi x \int_0^1 \cos(\pi t) u(t) dt \equiv c \sin \pi x,$$

where c is a constant yet to be determined. Substitute this expression for $u(x)$ in (3). We obtain

$$c \sin \pi x = c \lambda \sin \pi x \int_0^1 \cos(\pi t) \sin(\pi t) dt.$$

But $\int_0^1 \cos(\pi t) \sin(\pi t) dt = 0.$

Therefore
whence
and

$$\begin{aligned} c \sin \pi x &\equiv 0, \\ c &= 0, \\ u &\equiv 0. \end{aligned}$$

Thus this particular kernel has no characteristic constants. This shows that Theorem I indeed states a property peculiar to *symmetric* kernels.

54. Power Series for $\frac{D'(\lambda)}{D(\lambda)}$.—The following proof of the theorem is due to Kneser. It is based upon a lemma concerning the expansion of $\frac{D'(\lambda)}{D(\lambda)}$, which holds under the assumptions A), B), C).

From Fredholm's first fundamental theorem, if $D(\lambda) \neq 0$, then (1) has one and only one solution given by

$$(4) \quad u(x) = f(x) + \int_a^b \frac{D(x, t; \lambda)}{D(\lambda)} f(t) dt.$$

Consider the expansion of $\frac{D(x, t; \lambda)}{D(\lambda)}$ as a power series in λ .

Now $D(0) = 1 \neq 0$ and, therefore, there exists a ρ such that if $|\lambda| < \rho$, then

$$(5) \quad D(\lambda) \neq 0, \quad |\lambda| < \rho.$$

Hence, since $D(\lambda)$ is permanently convergent, it can be expanded¹ into a power series convergent for $|\lambda| < \rho$. Then

$$(6) \quad \frac{1}{D(\lambda)} = d_0 + d_1\lambda + d_2\lambda^2 + \dots, \quad |\lambda| < \rho.$$

Now, $D(x, t; \lambda)$ is a permanently converging power series in λ . Hence, also, the product $D(x, t; \lambda) \cdot \frac{1}{D(\lambda)}$ can be expanded into a power series in λ :

$$(7) \quad \frac{D(x, t; \lambda)}{D(\lambda)} = \sum_{n=1}^{\infty} g_n(x, t) \lambda^n$$

¹ HARKNESS and MORLEY, "Theory of Functions," §83.

and the series will be uniformly convergent as to x and t in R for $|\lambda| < \rho$. The right-hand member of (7) has no term free from λ for

$$D(x, t; \lambda) = \lambda K(x, t) + \dots$$

Substitute from (7) in (4) and we obtain

$$(8) \quad u(x) = f(x) + \sum_{n=1}^{\infty} \lambda^n \int_a^b g_n(x, t) f(t) dt, \quad |\lambda| < \rho.$$

We have been able to interchange the order of the integration and summation in (8) on account of the uniform convergence of $\sum_{n=1}^{\infty} g_n(x, t) \lambda^n$.

The coefficients $g_n(x, t)$ can be determined by comparing (8) with the expression for the solution obtained by successive approximations (see §8):

$$(9) \quad u(x) = f(x) + \sum_{n=1}^{\infty} \lambda^n \int_a^b K_n(x, t) f(t) dt$$

where $K_n(x, t)$ are the iterated kernels given by

$$(10) \quad K_n(x, t) = \int_a^b K(x, s) K_{n-1}(s, t) ds.$$

This solution by successive approximations is valid if $|\lambda| < \frac{1}{M(b-a)}$, where M is the maximum of $|K(x, t)|$ on R . Denote by r the smaller of the two quantities ρ , $\frac{1}{M(b-a)}$. Then (8) and (9) hold simultaneously for $|\lambda| < r$. But (1) has one and only one solution; therefore (8) and (9) represent the same function, and coefficients of corresponding powers of λ must be equal. Therefore,

$$\int_a^b g_n(x, t) f(t) dt = \int_a^b K_n(x, t) f(t) dt, \quad (x)$$

where the notation (x) is used to mean uniformly as to x . Hence

$$(11) \quad \int_a^b \left\{ K_n(x, t) - g_n(x, t) \right\} f(t) dt = 0, \quad (x).$$

But $K_n(x, t)$ and $g_n(x, t)$ are independent of f and, for a given value of x ,

$$K_n(x, t) - g_n(x, t) = M(t)$$

is a real and continuous function of t . Now (11) holds for any choice of f which is continuous. Choose then $f(t) = M(t)$. Then (11) becomes

$$\int_a^b [M(t)]^2 dt = 0.$$

Hence

$$M(t) \equiv 0 \text{ on } [ab].$$

Therefore

$$g_n(x, t) \equiv K_n(x, t), \quad (x, t)$$

and (7) becomes

$$(12) \quad \frac{D(x, t; \lambda)}{D(\lambda)} = \sum_{n=1}^{\infty} K_n(x, t) \lambda^n, \quad |\lambda| < r.$$

Thus, under the assumptions A), B), C), we have proved the following theorem:

Theorem II.—For all sufficiently small values of λ

$$\frac{D(x, t; \lambda)}{D(\lambda)} = \sum_{n=1}^{\infty} K_n(x, t) \lambda^n,$$

the series converging uniformly as to x and t in R .

Theorem II holds when $t = x$. Then

$$\frac{D(x, x; \lambda)}{D(\lambda)} = \sum_{n=1}^{\infty} K_n(x, x) \lambda^n, \quad |\lambda| < r$$

and $\sum K_n(x, x) \lambda^n$ is uniformly convergent on $[ab]$

Integrate this equality with respect to x from a to b . We obtain

$$\frac{1}{D(\lambda)} \int_a^b D(x, x; \lambda) dx = \sum_{n=1}^{\infty} \lambda^n \int_a^b K_n(x, x) dx.$$

Let us introduce the permanent notation

$$(13) \quad \int_a^b K_n(x, x) dx = U_n, \text{ a constant,}$$

and recall from Chap. III, §20, equation (27) that

$$\int_a^b D(x, x; \lambda) = -\lambda D'(\lambda).$$

Then our equality above becomes

$$-\frac{\lambda D'(\lambda)}{D(\lambda)} = \sum_{n=1}^{\infty} U_n \lambda^n,$$

which may be written

$$\frac{D'(\lambda)}{D(\lambda)} = -\sum_{n=0}^{\infty} U_{n+1} \lambda^n, |\lambda| < r.$$

We have thus the following corollary to Theorem II:

Corollary.—

$$(14) \quad \frac{D'(\lambda)}{D(\lambda)} = -\sum_{n=0}^{\infty} U_{n+1} \lambda^n, |\lambda| < r.$$

55. Plan of Kneser's Proof.—Suppose that $K(x, t)$ has no characteristic constant; that is, that $D(\lambda) = 0$ has no roots, real or imaginary. Then the quotient $\frac{D'(\lambda)}{D(\lambda)}$ can be directly expanded into a power series

$$\frac{D'(\lambda)}{D(\lambda)} = \sum_{n=0}^{\infty} C_n \lambda^n,$$

and this series will be permanently converging.¹ But (14) holds for $|\lambda| < r$. Then

$$\sum_{n=0}^{\infty} C_n \lambda^n = \sum_{n=0}^{\infty} U_{n+1} \lambda^n, |\lambda| < r,$$

hence

$$C_n = U_{n+1}$$

and, therefore,

$$\sum_{n=0}^{\infty} U_{n+1} \lambda^n \text{ is permanently convergent.}$$

Then also² is

$$\sum_{n=0}^{\infty} |U_{n+1}| |\lambda|^n \text{ permanently convergent,}$$

and also

$$(15) \quad \sum_{n=0}^{\infty} |U_{n+1}| |\lambda|^{n+1} \text{ is permanently convergent.}$$

Then the series formed by omitting any number of terms from (15) is also permanently convergent. Hence

$$(16) \quad \sum_{n=0}^{\infty} |U_{2n}| |\lambda|^{2n} \text{ is permanently convergent.}$$

We have proved that (16) was permanently convergent by assuming that $K(x, t)$ had no characteristic constant. If, then, for a given kernel $K(x, t)$ we can show that the corresponding series (16) is *not* permanently convergent, we will have shown that $K(x, t)$ has at least one characteristic constant. We are then going to prove that if, in addition to the properties A) B) C), the kernel $K(x, t)$ is symmetric, then the corresponding series (16) is not permanently convergent. Theorem I will then be proved.

¹ HARKNESS and MORLEY, *Loc. cit.*, §83.

² HARKNESS and MORLEY, *Loc. cit.*, §76.

For this purpose we need some auxiliary lemmas on iterated kernels.

56. Lemmas on Iterations of a Symmetric Kernel.—The iterated kernel $K_n(x, t)$ was defined by

$$(17) \quad K_1(x, t) = K(x, t)$$

$$K_n(x, t) = \int_a^b K(x, s) K_{n-1}(s, t) ds.$$

By successive applications of this recursion formula we find

$$(18) \quad K_n(x, t) = \int_a^b \dots \int_a^b K(x, s_1) K(s_1, s_2) \dots$$

$$K(s_{n-1}, t) ds_1 \dots ds_{n-1}.$$

In Chap. II, §11, we showed that

$$(19) \quad K_{n+p}(x, t) = \int_a^b K_n(x, s) K_p(s, t) ds.$$

We now establish the following lemmas:

Lemma I.—*If $K(x, t)$ is symmetric, then $K_n(x, t)$ is symmetric.*

Proof.—By (18) we have

$$K_n(t, x) = \int_a^b \dots \int_a^b K(t, s_1) K(s_1, s_2) \dots K(s_{n-1}, x)$$

$$ds_1 \dots ds_{n-1}$$

$$= \int_a^b \dots \int_a^b K(s_1, t) K(s_2, s_1) \dots K(x, s_{n-1})$$

$$ds_1 \dots ds_{n-1}$$

on account of the symmetry of $K(x, t)$

$$= \int_a^b \dots \int_a^b K(x, s_{n-1}) \dots K(s_2, s_1) K(s_1, t)$$

$$ds_1 \dots ds_{n-1}.$$

Now make a change of notation.

For	s_1	s_2	\dots	s_{n-2}	s_{n-1}	
put	s_{n-1}	s_{n-2}	\dots	s_2	s_1	

Then

$$\begin{aligned} K_n(t, x) &= \int_a^b \dots \int_a^b K(x, s_1) K(s_1, s_2) \dots K(s_{n-1}, t) \\ &\quad ds_1 \dots ds_{n-1} \\ &= K_n(x, t) \text{ by (18).} \end{aligned}$$

Lemma II.—If $K(x, t)$ satisfies the conditions A) B) C) D), then $K_n(x, t) \not\equiv 0$ on R .

Proof.—Suppose $K_n(x, t) \equiv 0$ on R , and suppose that n is the lowest index for which an iterated kernel vanishes identically:

$$\begin{aligned} K(x, t) &\not\equiv 0, K_1(x, t) \not\equiv 0, \dots, K_{n-1}(x, t) \not\equiv 0, \\ K_n(x, t) &\equiv 0, R. \end{aligned}$$

Then certainly, by C), we have $n > 1$. Then from (19) it follows that all following iterated kernels are identically zero. Then

$$K_{n+1}(x, t) \equiv 0 \text{ on } R.$$

Now either n or $n + 1$ must be an even number $2m$:

$$2m = n, \text{ or } n + 1.$$

Then

$$K_{2m}(x, t) \equiv 0 \text{ on } R.$$

But by (19)

$$K_{2m}(x, t) = \int_a^b K_m(x, s) K_m(s, t) ds.$$

This equality holds when $t = x$. Then

$$\begin{aligned} K_{2m}(x, x) &= \int_a^b K_m(x, s) K_m(s, x) ds \\ (20) \qquad &= \int_a^b [K_m(x, s)]^2 ds \end{aligned}$$

on account of the symmetry of $K(x, t)$.

But $K_{2m}(x, t) \equiv 0$ on R . Therefore, since $K_m(x, t)$ is a real function of x and t , we have

$$K_m(x, t) \equiv 0 \text{ on } R.$$

But $m = \frac{n}{2}$, or $\frac{n+1}{2}$.

Hence $n - m = \frac{n}{2}$, or $\frac{n-1}{2} > 0$ for $n > 1$.

Therefore $m > n$.

Thus we have contradicted the assumption that $K_n(x, t)$ was the first iterated function to vanish and the lemma is proved.

It is now clear that if $K(x, t)$ satisfies the conditions A) B) C) D), then $K_n(x, t)$ satisfies the same conditions.

Corollary.— $U_{2m} > 0$.

Proof.— $U_{2m} = \int_a^b K_{2m}(x, x) dx$.

Now $K_{2m}(x, x) > 0$ by (20), since $K_m(x, t) \neq 0$.

57. Schwarz's Inequality.—Let $\varphi(x)$ and $\psi(x)$ be real and continuous on the interval $[ab]$, and u and v any two reals constant with respect to x . Then

$$\int_a^b [u\varphi(x) + v\psi(x)]^2 dx \geq 0,$$

since we have the square of a real function.

Expand and we obtain

$$u^2 \int_a^b \varphi^2(x) dx + 2uv \int_a^b \varphi(x)\psi(x) dx + v^2 \int_a^b \psi^2(x) dx \geq 0.$$

This is a definite quadratic form and hence

$$(21) \quad \left[\int_a^b \varphi(x)\psi(x) dx \right]^2 \leq \int_a^b \varphi^2(x) dx \times \int_a^b \psi^2(x) dx.$$

This is called the inequality of Schwarz. In like manner, we can prove that if $\varphi(x, y)$ and $\psi(x, y)$ are real and continuous on R in x and y , then

$$\begin{aligned} & \left[\int_a^b \int_a^b \varphi(x, y)\psi(x, y) dx dy \right]^2 \\ & \leq \left[\int_a^b \int_a^b \varphi^2(x, y) dx dy \right] \left[\int_a^b \int_a^b \psi^2(x, y) dx dy \right]. \end{aligned}$$

58. Application of Schwarz's Inequality.—We have obtained one expression for U_{2n} :

$$(13) \quad U_{2n} = \int_a^b K_{2n}(x, x) dx$$

$$(22) \quad = \int_a^b \int_a^b [K_n(x, t)]^2 dt dx > 0 \text{ by Lemma II.}$$

We now proceed to determine a second expression for U_{2n} . Now by (19) we have

$$K_{2n}(x, t) = \int_a^b K_{n-1}(x, s) K_{n+1}(s, t) dt.$$

If we put $t = x$, this equality becomes

$$\begin{aligned} K_{2n}(x, x) &= \int_a^b K_{n-1}(x, t) K_{n+1}(t, x) dt \\ &= \int_a^b K_{n-1}(x, t) K_{n+1}(x, t) dt, \end{aligned}$$

on account of the symmetry of $K_{n+1}(x, t)$. Then

$$(23) \quad U_{2n} = \int_a^b \int_a^b K_{n-1}(x, t) K_{n+1}(x, t) dt dx.$$

Apply Schwarz's inequality to the right member of (23). We obtain

$$\begin{aligned} &\left[\int_a^b \int_a^b K_{n-1}(x, t) K_{n+1}(x, t) dt dx \right]^2 \\ &\leq \left[\int_a^b \int_a^b K_{n-1}^2(x, t) dt dx \right] \left[\int_a^b \int_a^b K_{n+1}^2(x, t) dt dx \right], \end{aligned}$$

which, if we make use of (22) and (23), may be written

$$U_{2n}^2 \leq U_{2n-2} \cdot U_{2n+2}.$$

Divide both members of this inequality by $U_{2n-2} \cdot U_{2n}$. This is possible, since $U_{2n-2} \neq 0$, $U_{2n} \neq 0$. We obtain

$$(24) \quad \frac{U_{2n+2}}{U_{2n}} \geq \frac{U_{2n}}{U_{2n-2}}.$$

Putting successively $n = 2, 3, \dots, n$, we obtain in this way the sequence of inequalities

$$\frac{U_{2n+2}}{U_{2n}} \geq \frac{U_{2n}}{U_{2n-2}} \geq \dots \geq \frac{U_6}{U_4} \geq \frac{U_4}{U_2}.$$

Therefore

$$(25) \quad \frac{U_{2n+2}}{U_{2n}} \geq \frac{U_4}{U_2}, \quad (n).$$

Apply now the ratio test to the series

$$(16) \quad \sum_{n=1}^{\infty} U_{2n} |\lambda|^{2n} = \sum_{n=1}^{\infty} V_n$$

to find the radius of convergence in λ . We find

$$\frac{V_{n+1}}{V_n} = \frac{U_{2n+2}}{U_{2n}} |\lambda|^2 \geq \frac{U_4}{U_2} |\lambda|^2, \quad (n).$$

Therefore, the series diverges if

$$\frac{U_4}{U_2} |\lambda|^2 > 1,$$

that is, if

$$|\lambda| > \sqrt{\frac{U_2}{U_4}}.$$

Therefore, the series is not a permanently convergent power series in λ . This completes the proof of Theorem I.

Theorem I.—For a real, symmetric, continuous, non-identically vanishing kernel $K(x, t)$ there exists at least one characteristic constant λ_o .

We remark that for every value of λ in the λ -plane, without a circle C of radius

$\sqrt{\frac{U_2}{U_4}}$, $\sum_{n=1}^{\infty} U_{2n} |\lambda|^{2n}$ diverges. Hence, from

(14) and (16), $\frac{D'(\lambda)}{D(\lambda)}$, when expressed as

a series in λ , diverges for all values of λ without the circle C . Then, certainly, $D(\lambda) = 0$ has at least one root within the interior or on the boundary of C




FIG. 18.

II. ORTHOGONALITY

59. Orthogonality Theorem.—From the Fredholm theory

1) Two associated kernels $\tilde{K}(x, t)$, $K(x, t)$:

$$\tilde{K}(x, t) = K(t, x)$$

have the same characteristic constants with the same indices; if, then

2) λ_o and λ_1 ($\lambda_o \neq \lambda_1$) are two characteristic constants of $K(x, t)$, and hence of $K(t, x)$, and $\varphi_o(x)$ is a fundamental function of $K(x, t)$ for λ_o , and $\tilde{\varphi}_1(x)$ is a fundamental function of $\tilde{K}(x, t)$ for λ_1 , it follows that $\int_a^b \varphi_o(x) \tilde{\varphi}_1(x) dx = 0$.

But if the kernel $K(x, t)$ is symmetric, then $\tilde{\varphi}_1(x)$ is also a fundamental function of $K(x, t)$ for λ_1 . If, then, we write $\tilde{\varphi}_1 \equiv \varphi_1(x)$, we have

$$\int_a^b \varphi_o(x) \varphi_1(x) dx = 0.$$

Definition.—Two functions $\varphi(x)$, $\psi(x)$, continuous on the interval $[ab]$ are said to be orthogonal on $[ab]$ if

$$(\varphi\psi) = - \int_a^b \varphi(x) \psi(x) dx = 0.$$

We have thus proved the following theorem which is called the *orthogonality theorem*:

Theorem III.—If $K(x, t)$ is symmetric and $\varphi_o(x)$, $\varphi_1(x)$ are fundamental functions of $K(x, t)$ for λ_o and λ_1 respectively ($\lambda_o \neq \lambda_1$), then $\varphi_o(x)$ and $\varphi_1(x)$ are orthogonal on the interval $[ab]$.

$$(26) \quad \int_a^b \varphi_o(x) \varphi_1(x) dx = 0.$$

As an illustration of Theorem III, take the symmetric kernel $K(x, t)$:

$$K(x, t) = \begin{cases} (1-t)x, [0t], \\ (1-x)t, [t1] \end{cases}$$

which has the characteristic constants

$$\lambda_n = n^2\pi^2, \quad n = 1, 2, \dots$$

and the corresponding fundamental functions

$$\varphi_n(x) = \sin n\pi x.$$

It is then well known from the theory of definite integrals that

$$(27) \quad \int_0^1 \sin n\pi x \sin p\pi x dx = 0, \quad (n \neq p).$$

60. Reality of the Characteristic Constants.—By means of the preceding theorem we can now prove the following theorem due to Hilbert:

Theorem IV.—*If $K(x, t)$ is real and symmetric, continuous, and $\neq 0$, then all of the characteristic constants are real.*

Proof.—Suppose that there is a characteristic constant λ_o not real:

$$\lambda_o = \mu_o + i\nu_o, \quad \nu_o \neq 0.$$

Then the homogeneous equation

$$u(x) = \lambda_o \int_a^b K(x, t)u(t)dt$$

has at least one continuous solution $\varphi(x) \equiv 0$:

$$(28) \quad \varphi(x) = (\mu_o + i\nu_o) \int_a^b K(x, t)\varphi(t)dt.$$

1) Suppose $\varphi(x)$ real, then separating the real and imaginary parts of (28), we obtain

$$(29) \quad \varphi(x) = \mu_o \int_a^b K(x, t)\varphi(t)dt$$

$$(30) \quad 0 = \nu_o \int_a^b K(x, t)\varphi(t)dt.$$

From (30)

$$\int_a^b K(x, t)\varphi(t)dt = 0,$$

since $\nu_o \neq 0$. Then from (29), $\varphi(x) \equiv 0$, which contradicts the assumption that $\varphi(x) \not\equiv 0$. Therefore, $\varphi(x)$ cannot be real.

2) Suppose $\varphi(x) = v(x) + iw(x)$, where $v(x)$ and $w(x)$ are real. Then (28) becomes

$$(31) \quad v(x) + iw(x) = (\mu_o + i\nu_o) \int_a^b K(x, t) [v(t) + iw(t)] dt.$$

Separate (31) into its real and imaginary parts. We obtain

$$(32) \quad v(x) = \mu_o \int_a^b K(x, t)v(t)dt - \nu_o \int_a^b K(x, t)w(t)dt$$

$$(33) \quad w(x) = \mu_o \int_a^b K(x, t)w(t)dt + \nu_o \int_a^b K(x, t)v(t)dt.$$

Multiply both sides of (33) by $-i$ and add to (32). We obtain

$$v(x) - iw(x) = (\mu_o - i\nu_o) \int_a^b K(x, t) [v(t) - iw(t)] dt.$$

Therefore $v(x) - iw(x) \equiv \bar{\varphi}(x)$ is also a fundamental function of $K(x, t)$, belonging to $\bar{\lambda}_o \equiv \mu_o - i\nu_o$. If we now apply the orthogonality Theorem III, we obtain, since $\bar{\lambda}_o \neq \lambda_o$,

$$\int_a^b \varphi(x) \bar{\varphi}(x) dx = 0,$$

which may be written

$$\int_a^b [v^2(x) + w^2(x)] dx = 0.$$

But $v(x)$ and $w(x)$ are real functions. Therefore

$$v^2(x) + w^2(x) = 0, \quad (x)$$

and hence $v(x) \equiv 0, w(x) \equiv 0$

and therefore $\varphi(x) \equiv 0$,

which constitutes a contradiction. Therefore λ_o cannot be of the form $\lambda_o = \mu_o + i\nu_o (\nu_o \neq 0)$ and hence λ_o must be real.

This result might have been foreseen from the analogy with the finite system of linear equations:

$$u(t_i) - \lambda h \sum_{j=1}^n K(t_i, t_j) u(t_j) = f(t_i) \quad (i = 1, \dots, n)$$

with determinant

$$(34) \quad \Delta = \begin{vmatrix} 1 - \lambda h K_{11} & -\lambda h K_{12} & \dots & -\lambda h K_{1n} \\ -\lambda h K_{21} & 1 - \lambda h K_{22} & \dots & -\lambda h K_{2n} \\ \dots & \dots & \ddots & \dots \\ -\lambda h K_{n1} & -\lambda h K_{n2} & \dots & 1 - \lambda h K_{nn} \end{vmatrix}.$$

But it is well known¹ that, if K_{ij} is real and $K_{ij} = K_{ji}$, then all of the roots of (34) are real.

In the Fredholm theory we proved that $q \leq r$ where q is the index of λ_0 and r is the multiplicity of the root λ_0 of $D(\lambda_0) = 0$. For the case of a symmetric kernel, Hilbert has proved² the following more definite theorem which we state here without proof:

Theorem V.—*For a real symmetric kernel the index q of a characteristic constant λ_0 is always equal to its multiplicity r : $q = r$.*

61. Complete Normalized Orthogonal System of Fundamental Functions.—Let us suppose that λ_0 is a characteristic constant of the symmetric kernel $K(x, t)$ of index q . Then the equation

$$u(x) = \lambda_0 \int_a^b K(x, t) u(t) dt$$

has been shown to have q linearly independent solutions:

$$\varphi_\alpha(x) = \frac{D\left(x_1', \dots, x'_{\alpha-1}, x_\alpha, x'_{\alpha+1}, \dots, x'_q, \lambda_0\right)}{D\left(y_1', \dots, y'_{\alpha-1}, y_\alpha', y'_{\alpha+1}, \dots, y'_q, \lambda_0\right)}$$

$$\alpha = 1, \dots, q.$$

¹ See BOCHER, "Introduction to Higher Algebra," §59.

² For proof, see HORN, §39.

For a real symmetric kernel, λ_o and, therefore, also $\varphi_\alpha(x)$ are real, since the arguments in the numerator and denominator of the expression for $\varphi_\alpha(x)$ are all real. Any other solution $\psi_\alpha(x)$ is expressible linearly in terms of these q solutions, or in terms of q linearly independent linear combinations of the $\varphi_\alpha(x)$'s:

$$(35) \quad \psi_\alpha(x) = c_{\alpha 1}\varphi_1(x) + \dots + c_{\alpha q}\varphi_q(x),$$

where the determinant of the coefficients $|c_{\alpha\beta}| \neq 0$, $\left[\alpha, \beta = 1, \dots, q \right]$ does not vanish.

We are going to choose such a system of fundamental functions in such a way that they form what is called a *normalized orthogonal system*.

a) *Normalized Fundamental Functions*.—A function ψ is said to be *normalized* if

$$(\psi\psi) \equiv \int_a^b \psi^2(x)dx = 1.$$

If $\varphi(x)$ is a fundamental function belonging to λ_o :

$$\varphi(x) = \lambda_o \int_a^b K(x, t)\varphi(t)dt, \quad \left[\varphi(x) \neq 0 \right]$$

then $\psi(x) = c\varphi(x)$ ($c \neq 0$) is clearly again a fundamental function belonging to λ_o , and we can choose c so that

$$(36) \quad (\psi\psi) = 1,$$

that is

$$c^2 \int_a^b \varphi^2(x)dx = 1,$$

whence

$$(37) \quad c = \pm \frac{1}{\sqrt{\int_a^b \varphi^2(x)dx}}.$$

This value for c is finite, real, and $\neq 0$. Hence we obtain

$$(38) \quad \psi(x) = \frac{\varphi(x)}{\sqrt{\int_a^b \varphi^2(x) dx}}.$$

b) *Normalized Orthogonal Systems.*—We now propose to determine the constants $c_{\alpha\beta}$ in (35) in such a way that the ψ_α 's satisfy the following two conditions:

1) All ψ_α 's are normalized:

$$(39) \quad \int_a^b \psi_\alpha^2(x) dx = 1, \alpha = 1, \dots, q$$

2) Two ψ_α 's with different subscripts are orthogonal:

$$(40) \quad \int_a^b \psi_\alpha(x) \psi_\beta(x) dx = 0, \alpha \neq \beta, (\alpha, \beta = 1, \dots, q).$$

For this purpose we put $\psi_1(x) = c\varphi_1(x)$, where

$$c = \frac{1}{\sqrt{\int_a^b \varphi_1^2(x) dx}}.$$

Then the condition $\int_a^b \psi_1^2(x) dx = 1$ is satisfied.

Next choose $\psi_2(x)$ as a linear function of φ_1 and φ_2 . On account of (38) it can then be expressed linearly in terms of ψ_1 and ψ_2 :

$$\psi_2(x) = \alpha_1 \psi_1(x) + \alpha_2 \varphi_2(x).$$

We now determine α_1 and α_2 so that the conditions 1) and 2) are satisfied for ψ_1 and ψ_2 .

From (40)

$$(42) \quad (\psi_1 \psi_2) = \alpha_1 \int_a^b \psi_1^2 dx + \alpha_2 \int_a^b \psi_1(x) \varphi_2(x) dx = 0.$$

Therefore, on account of (39),

$$\alpha_1 = -\alpha_2 (\psi_1 \varphi_2)$$

and (41) becomes

$$\psi_2(x) = \alpha_2 \left[\varphi_2(x) - (\psi_1 \varphi_2) \psi_1(x) \right].$$

We can now determine α_2 in such a way that $\psi_2(x)$ is normalized provided

$$(43) \quad \varphi_2(x) - (\psi_1 \varphi_2) \psi_1(x) \not\equiv 0.$$

But $(\psi_1 \varphi_2)$ is a constant and if (43) is not true, then

$$\varphi_2(x) + c_1 \varphi_1(x) \equiv 0$$

which is a linear relation between two fundamental functions φ_1 and φ_2 and, therefore, φ_1 and φ_2 would be linearly dependent, which is contrary to hypothesis. Therefore, $\psi_2(x)$ is completely determined as a linear function of φ_1 and φ_2 or of ψ_1 and φ_2 . Now choose $\psi_3(x)$ as a linear function of φ_1 , φ_2 , φ_3 , or, what amounts to the same thing, as a linear function of ψ_1 , ψ_2 , φ_3 :

$$\psi_3(x) = \beta_1 \psi_1 + \beta_2 \psi_2 + \beta_3 \varphi_3.$$

Apply condition (2). We obtain

$$(\psi_1 \psi_3) = \beta_1 (\psi_1 \psi_1) + \beta_2 (\psi_1 \psi_2) + \beta_3 (\psi_1 \varphi_3) = 0.$$

But $(\psi_1 \psi_1) = 1$ and $(\psi_1 \psi_2) = 0$, therefore

$$(\psi_1 \psi_3) = \beta_1 + \beta_2 (\psi_1 \varphi_3) = 0.$$

Hence

$$\beta_1 = -\beta_3 (\psi_1 \varphi_3).$$

Also by condition (2) we have

$$(\psi_2 \psi_3) = \beta_1 (\psi_2 \psi_1) + \beta_2 (\psi_2 \psi_2) + \beta_3 (\psi_2 \varphi_3) = 0,$$

from which we derive

$$\beta_2 = -\beta_3 (\psi_2 \varphi_3).$$

The expression for $\psi_3(x)$ now becomes

$$\psi_3(x) = \beta_3 \left[\varphi_3 - (\psi_1 \varphi_3) \psi_1 - (\psi_2 \varphi_3) \psi_2 \right].$$

We can now determine β_3 by means of condition 1), provided it is not true that

$$(44) \quad \varphi_3 - (\psi_1\varphi_3)\psi_1 - (\psi_2\varphi_3)\psi_2 \equiv 0.$$

But (44) cannot be satisfied, otherwise $\varphi_1, \varphi_2, \varphi_3$ would be linearly dependent, which is contrary to hypothesis. Hence β_3 can be so determined that $\psi_3(x)$ satisfies the conditions 1) and 2). This process can be continued until q functions $\psi_\alpha(x)$ are obtained, satisfying conditions 1) and 2), or, as we say, form a *normalized orthogonal system*.

The ψ_α 's so obtained are real, they are furthermore linearly independent. For, suppose we had

$$c_1\psi_1(x) + c_2\psi_2(x) + \dots + c_q\psi_q(x) \equiv 0.$$

Multiply by $\psi_\alpha(x)$ and integrate from a to b . On account of (39) and (40) we would obtain $c_\alpha = 0$.

A linear transformation

$$y_i = c_{i1}x_1 + \dots + c_{in}x_n \quad (i = 1, \dots, n)$$

is said to be orthogonal if

$$\sum_{i=1}^n x_i^2 = \sum_{i=1}^n y_i^2.$$

The transformation of rectangular coordinates with fixed origin is a transformation of this kind.

The condition on the coefficients which insures this property of the transformation is

$$(45) \quad \sum_{j=1}^n c_{ij}c_{kj} = \begin{cases} 1, & i = K \\ 0, & i \neq K. \end{cases}$$

Now consider the definite integral

$$\int_a^b \psi_i(x)\psi_k(x)dx$$

as the limit of a sum

$$(46) \quad \lim_{h \rightarrow 0} \sum_j h \psi_i(x_j) \psi_k(x_j) = \int_a^b \psi_i(x) \psi_k(x) dx.$$

Then the analogy between equations (39), (40), and (45) becomes apparent.

c) *Complete Normalized Orthogonal System of Fundamental Functions.*—To each root of $D(\lambda) = 0$ there belongs such a normalized orthogonal system of fundamental functions. Now we have shown that $D(\lambda) = 0$ has at least one root. There may be a finite or an infinite number of such roots. If they are infinite in number it follows from the theory of permanently convergent power series that they constitute a *denumerable* set and they may be arranged in the order of the magnitude of their absolute values:

$$|\lambda_1| \leq |\lambda_2| \leq \dots \leq |\lambda_n| \leq |\lambda_{n+1}| \leq \dots$$

and each λ_i has a definite index q_i . We have then the following table:

Characteristic constant	Index	Normalized orthogonal fundamental functions
λ_1	q_1	$\psi_1^1, \dots, \psi_{q_1}^1$
λ_2	q_2	$\psi_1^2, \dots, \psi_{q_2}^2$
.	.	.
.	.	.
.	.	.
λ_n	q_n	$\psi_1^n, \dots, \psi_{q_n}^n$

We now change the notation and denote the functions ψ in the order in which they stand in the following line:

$\psi_1^1, \dots, \psi_{q_1}^1, \psi_1^2, \dots, \psi_{q_2}^2, \dots, \psi_1^n, \dots, \psi_{q_n}^n$ by the symbols $\psi_1, \psi_2, \dots, \psi_r, \dots$ and the characteristic constants to which they belong by $\lambda_1, \lambda_2, \dots,$

λ_r, \dots , where some of the λ 's may be equal, for instance $\lambda_1 = \lambda_2 = \dots = \lambda_{q_1}$; $\lambda_{q_1+1} = \lambda_{q_1+2} = \dots = \lambda_{q_1+q_2}$. Such a system of fundamental functions ψ_r is called a complete normalized orthogonal system of fundamental functions. We sum up the preceding discussion in the following theorem:

Theorem VI.—*To every real symmetric kernel there belongs a complete normalized orthogonal system of fundamental functions $\psi_r(x)$, with the following properties:*

1) $\psi_r(x)$ is a fundamental function belonging to λ_r .

$$\psi_r(x) = \lambda_r \int_a^b K(x, t) \psi_r(t) dt.$$

$$2) \int_a^b \psi_r^2(x) dx = 1.$$

$$3) \int_a^b \psi_r(x) \psi_s(x) dx = 0, \quad (r \neq s).$$

4) $\psi_r(x)$ is real.

5) Every fundamental function $\varphi(x)$ is expressible in the form

$$\varphi(x) = c_{r_1} \psi_{r_1}(x) + \dots + c_{r_m} \psi_{r_m}(x).$$

Proof of 3).—If ψ_r, ψ_s belong to the same λ , they are orthogonal by construction, if to different λ 's by Theorem III.

Proof of 5).— $\varphi(x)$ belongs to a certain characteristic constant λ_r and is, therefore, linearly expressible in terms of the fundamental functions $\psi_1(x), \dots, \psi_{q_r}$, if we use the notation of the table.

Example.—For the problem of the vibrating string we had the kernel:

$$(47) \quad K(x, t) = \begin{cases} (1-t)x, & 0 \leq x \leq t \\ (1-x)t, & t \leq x \leq 1 \end{cases}$$

with the characteristic constants $\lambda_n = n^2\pi^2$, $n = 1, 2, \dots$ of index 1, and the corresponding fundamental functions $\varphi_n(x) = \beta \sin n\pi x$. These functions $\varphi_n(x)$ will form a

complete normalized orthogonal system of fundamental functions, if

$$\beta = \frac{1}{\sqrt{\int_0^1 \sin^2 n\pi x dx}} = \pm \sqrt{2}.$$

Our fundamental functions are then

$$\psi_n(x) = \sqrt{2} \sin n\pi x.$$

III. EXPANSION OF AN ARBITRARY FUNCTION ACCORDING TO THE FUNDAMENTAL FUNCTIONS OF A COMPLETE NORMALIZED ORTHOGONAL SYSTEM

62. a) Problem of the Vibrating String Resumed.—The problem was to determine $y(x, t)$ so as to satisfy the following conditions (see §26):

$$(48) \quad 1) \frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}.$$

$$(49) \quad 2) y(0, t) = 0, y(1, t) = 0.$$

$$(50) \quad 3) y(x, 0) = f(x), f(x) \text{ an arbitrary given function.}$$

$$y_t(x, 0) = F(x), F(x) \text{ an arbitrary given function.}$$

We attempted to find a solution in the form

$$y = \varphi(t)u(x)$$

and found that $u(x)$ must satisfy

$$1) \frac{d^2 u}{dx^2} + \lambda u = 0$$

$$2) u(0) = 0, u(1) = 0,$$

and that $\varphi(t)$ must satisfy the equation

$$\frac{d^2 \varphi}{dt^2} + \lambda c^2 \varphi = 0.$$

The boundary problem in u had non-trivial solutions only when

$$\lambda = \lambda_n = n^2\pi^2,$$

n a positive integer, the solutions being

$$u = \beta \sin n\pi x.$$

To complete the solution of the partial differential equation (48) with the conditions (49) and (50), we must next integrate the differential equation for $\varphi(t)$ for the values $\lambda = n^2\pi^2$:

$$\frac{d^2\varphi}{dt^2} + n^2\pi^2 c^2 \varphi = 0.$$

The most general solution of this equation is

$$\varphi(t) = A_n \cos n\pi ct + B_n \sin n\pi ct$$

(A_n, B_n , arbitrary constants), which gives for a solution of the problem of the vibrating string

$$y = (A_n \cos n\pi ct + B_n \sin n\pi ct) \sin n\pi x.$$

This expression for y will satisfy (48) and (49), but, in general, it will not satisfy (50). In order to obtain a solution of (48) which will satisfy both (49) and (50) we notice that, owing to the linear character of (48), the series

$$(51) \quad y = \sum_{n=1}^{\infty} (A_n \cos n\pi ct + B_n \sin n\pi ct) \sin n\pi x$$

will also satisfy (48) and (49), provided it is convergent and admits of two successive term-by-term differentiations with respect to t and x .

Assuming this condition satisfied, it remains, then, so to determine the constants A_n and B_n that (50) is satisfied:

$$y(x, 0) = \sum_{n=1}^{\infty} A_n \sin n\pi x = f(x)$$

$$y_t(x, 0) = \sum_{n=1}^{\infty} B_n n\pi c \sin n\pi x = F(x).$$

But this is equivalent to asking us to develop the arbitrarily given functions $f(x)$ and $F(x)$ into sine series, or, since

$\psi_n(x) = \sqrt{2} \sin n\pi x$, into a series proceeding according to the complete normalized orthogonal system of fundamental functions of the special kernel $K(x, t)$ defined by (47).

b) *Determination of the Coefficients in the General Problem.* The problem just considered is a particular case of the following more general problem: Given a symmetric kernel $K(x, t)$ with its complete normalized orthogonal system of fundamental functions ψ , with corresponding characteristic constants λ_ν , it is possible to expand an arbitrary continuous function $f(x)$ in the form

$$f(x) = \sum_\nu c_\nu \psi_\nu(x).$$

Theorem VII.—*If $f(x)$ is a continuous function and is expressible in the form*

$$(52) \quad f(x) = \sum_\nu c_\nu \psi_\nu(x)$$

and if this series, if infinite, is uniformly convergent on $[ab]$, then the coefficients c_ν are given by

$$c_n = \int_a^b f(x) \psi_n(x) dx \equiv (f \psi_n)$$

Proof.—Multiply both sides of (52) by $\psi_n(x)$ and integrate with respect to x from a to b :

$$\begin{aligned} \int_a^b f(x) \psi_n(x) dx &= \int_a^b \sum_\nu c_\nu \psi_\nu(x) \psi_n(x) dx \\ &= \sum_\nu \int_a^b c_\nu \psi_\nu(x) \psi_n(x) dx \\ &= c_n, \end{aligned}$$

since

$$\int_a^b \psi_\nu(x) \psi_n(x) dx = \begin{cases} 1, & \nu = n \\ 0, & \nu \neq n. \end{cases}$$

IV. EXPANSION OF THE KERNEL ACCORDING TO THE FUNDAMENTAL FUNCTIONS OF A COMPLETE NORMALIZED ORTHOGONAL SYSTEM

63. a) Determination of the Coefficients.— $K(x, t)$ with t fixed is a function of x continuous on $[ab]$. Hence, if we assume that $K(x, t)$ can be expanded into a uniformly convergent series on $[ab]$:

$$K(x, t) = \sum_{\nu} c_{\nu} \psi_{\nu}(x),$$

then, by Theorem VII above, we have

$$(53) \quad c_{\nu} = \int_a^b K(x, t) \psi_{\nu}(x) dx.$$

But

$$\begin{aligned} \psi_{\nu}(t) &= \lambda_{\nu} \int_a^b K(t, x) \psi_{\nu}(x) dx \\ &= \lambda_{\nu} \int_a^b K(x, t) \psi_{\nu}(x) dx, \end{aligned}$$

on account of the symmetry of $K(x, t)$. Therefore

$$c_{\nu} = \frac{\psi_{\nu}(t)}{\lambda_{\nu}}$$

and

$$K(x, t) = \sum_{\nu} \frac{\psi_{\nu}(x) \psi_{\nu}(t)}{\lambda_{\nu}}.$$

Thus we obtain the following corollary to Theorem VII.

Corollary.—If $K(x, t) = \sum_{\nu} c_{\nu} \psi_{\nu}(x)$ and this series, if infinite, is uniformly convergent, then

$$(54) \quad K(x, t) = \sum_{\nu} \frac{\psi_{\nu}(x) \psi_{\nu}(t)}{\lambda_{\nu}}.$$

Equation (54) is what Kneser calls the *bilinear formula*.

If we apply this bilinear formula to

$$K(x, t) = \begin{cases} (1-t)x, & 0 \leq x \leq t \\ (1-x)t, & t \leq x \leq 1, \end{cases}$$

we obtain the hypothetical expansion:¹

$$K(x, t) = \sum_{n=1}^{\infty} \frac{2 \sin n\pi x \sin n\pi t}{n^2 \pi^2}.$$

b) *The Bilinear Formula for the Case of a Finite Number of Fundamental Functions.*—The bilinear formula holds always if the complete normalized orthogonal system contains only a finite number, m , of fundamental functions ψ_ν . The formula to be proved then reads

$$(55) \quad K(x, t) = \sum_{\nu=1}^m \frac{\Psi_\nu(x)\Psi_\nu(t)}{\lambda_\nu}.$$

To show that (55) holds, we show that the difference

$$(56) \quad H(x, t) = K(x, t) - \sum_{\nu=1}^m \frac{\Psi_\nu(x)\Psi_\nu(t)}{\lambda_\nu} \equiv 0, \text{ on } R.$$

Now $H(x, t)$ is continuous, real, and symmetric. If, in addition $H \not\equiv 0$, then there would exist at least one characteristic constant of $H(x, t)$ considered as a kernel of an integral equation. Hence, if we can show that $H(x, t)$ has no characteristic constant, then it follows that

$$H(x, t) \equiv 0.$$

Suppose that $\varphi(x)$ is a fundamental function of H belonging to a characteristic constant ρ of H , then φ is continuous and $\not\equiv 0$, and

$$\varphi(x) = \rho \int_a^b H(x, t)\varphi(t)dt.$$

¹ This formula is actually true. For a direct proof, see KNESER, "Die Integralgleichungen und ihre Anwendung," §4.

Whence, substituting for $H(x, t)$ its value from (56),

$$(57) \quad \varphi(x) = \rho \int_a^b K(x, t) \varphi(t) dt - \rho \sum_{\nu=1}^m \frac{\psi_\nu(x)}{\lambda_\nu} \int_a^b \psi_\nu(t) \varphi(t) dt.$$

Multiply both sides of (57) by $\psi_n(x)$ and integrate with respect to x from a to b , then

$$\begin{aligned} \int_a^b \varphi(x) \psi_n(x) dx &= \rho \int_a^b \int_a^b K(x, t) \psi_n(x) \varphi(t) dt dx \\ &\quad - \rho \sum_{\nu=1}^m \frac{1}{\lambda_\nu} \int_a^b \psi_\nu(x) \psi_n(x) dx \int_a^b \psi_\nu(t) \varphi(t) dt \\ &= \frac{\rho}{\lambda_n} \int_a^b \psi_n(t) \varphi(t) dt - \frac{\rho}{\lambda_n} \int_a^b \psi_n(t) \varphi(t) dt \\ &= 0, \quad (n = 1, \dots, m). \end{aligned}$$

Therefore, from (57),

$$\varphi(x) = \rho \int_a^b K(x, t) \varphi(t) dt.$$

By hypothesis $\varphi(x)$ is continuous and $\not\equiv 0$, and so this equation shows that $\varphi(x)$ is a fundamental function of $K(x, t)$ belonging to ρ . Therefore

$$\varphi(x) = c_1 \psi_1(x) + \dots + c_m \psi_m(x),$$

whence

$$\begin{aligned} \int_a^b \varphi(x) \psi_n(x) dx &= \int_a^b \psi_n(x) \sum_{\nu=1}^m c_\nu \psi_\nu(x) dx \\ &= c_n = 0. \end{aligned}$$

Therefore $\varphi(x) \equiv 0$, which constitutes a contradiction. Therefore $H(x, t)$ has no characteristic constant and $H(x, t) \equiv 0$, and hence the bilinear formula holds for m finite. This gives us the following theorem:

Theorem IX.—*If there are a finite number m of characteristic constants λ_ν , then*

$$K(x, t) = \sum_{\nu=1}^m \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu}.$$

c) *The Bilinear Formula for Kernels Having an Infinite Number of Characteristic Constants.*—We proceed to prove the following theorem:

Theorem X.—*If there are an infinite number of characteristic constants and $\sum_{\nu=1}^{\infty} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu}$ is uniformly convergent in x and t on R , then*

$$K(x, t) = \sum_{\nu=1}^{\infty} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu}.$$

Proof.—The proof is entirely analogous to the proof of Theorem IX. Form

$$(58) \quad H(x, t) = K(x, t) - \sum_{\nu=1}^{\infty} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu}.$$

H is continuous, for K is continuous, and the sum of the infinite series in (58) is a continuous function, being a uniformly convergent series the terms of which are continuous.¹ Furthermore, H is real, for K and ψ_ν are real. Finally, H is symmetric, for K is symmetric, and each term of the infinite series in (58) is symmetric. Hence it follows, as under b), that if $H(x, t)$ has no characteristic constant, then $H \equiv 0$. Suppose $H(x, t)$ had a characteristic constant ρ . Let $\varphi(x)$ be a fundamental function for H , belonging to the characteristic constant ρ , which implies that $\varphi(x)$ is continuous and $\not\equiv 0$ and satisfies

$$\varphi(x) = \rho \int_a^b H(x, t)\varphi(t)dt,$$

which, on account of (58), may be written

$$(59) \quad \varphi(x) = \rho \int_a^b K(x, t)\varphi(t)dt - \rho \int_a^b \sum_{\nu=1}^{\infty} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu} \varphi(t)dt.$$

¹ GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §173.

The series in (59) is again uniformly convergent as to x in $[ab]$. Multiply both sides of (59) by $\psi_n(x)$ and integrate with respect to x from a to b . We obtain

$$\int_a^b \psi_n(x)\varphi(x)dx = \rho \int_a^b \int_a^b K(x, t)\varphi(t)\psi_n(x)dtdx \\ - \rho \sum_{\nu=1}^{\infty} \frac{1}{\lambda_{\nu}} \int_a^b \psi_{\nu}(x)\psi_n(x)dx \int_a^b \psi_{\nu}(t)\varphi(t)dt,$$

which reduces to

$$\int_a^b \psi_n(x)\varphi(x)dx = \frac{\rho}{\lambda_n} \int_a^b \psi_n(t)\varphi(t)dt - \frac{\rho}{\lambda_n} \int_a^b \psi_n(t)\varphi(t)dt \\ = 0, \quad (n).$$

Therefore, from (59),

$$\varphi(x) = \rho \int_a^b K(x, t)\varphi(t)dt.$$

Hence we infer, as under b), that $\varphi(x) \equiv 0$, which contradicts our assumption $\varphi(x) \not\equiv 0$. Therefore, $H(x, t)$ has no characteristic constant and, hence, $H(x, t) \equiv 0$.

Since a series with a finite number of terms, functions of x and t , is always uniformly convergent, we may combine Theorems IX and X into the following

Theorem XI.—If $\left\{ \psi_{\nu}(x) \right\}$ are a complete normalized orthogonal system of fundamental functions belonging to the characteristic constant λ_{ν} , for the real symmetric kernel $K(x, t)$, and

$$\frac{1}{\lambda_{\nu}} \sum_{\nu} \psi_{\nu}(x)\psi_{\nu}(t)$$

is uniformly convergent on R , then

$$K(x, t) = \sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}}.$$

64. The Complete Normalized Orthogonal System for the Iterated Kernel $K_n(x, t)$.—The series $\sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}}$ is not always uniformly convergent and therefore the bilinear formula does not hold for every kernel $K(x, t)$. But we shall be able to show that it does always hold for the iterated kernels. For this purpose we prove first the following:

Theorem XII.—*If $\{\psi_{\nu}(x)\}$ are a complete normalized orthogonal system of fundamental functions for $K(x, t)$ belonging to the characteristic constants $\{\lambda_{\nu}\}$, then $\{\psi_{\nu}(x)\}$ are a complete normalized orthogonal system of fundamental functions for $K_n(x, t)$ belonging to the characteristic constants $\{\lambda_{\nu^n}\}$.*

Proof.—a) By assumption, $\psi_{\nu}(x)$ is continuous and $\not\equiv 0$ and

$$(61) \quad \psi_{\nu}(x) = \lambda_{\nu} \int_a^b K(x, t)\psi_{\nu}(t)dt,$$

whence we derive

$$\int_a^b K(s, x)\psi_{\nu}(x)dx = \lambda_{\nu} \int_a^b \int_a^b K(s, x)K(x, t)\psi_{\nu}(t)dtdx,$$

which, from (61) and our definitions of iterated kernels, may be written

$$\frac{\psi_{\nu}(s)}{\lambda_{\nu}} = \lambda_{\nu} \int_a^b K_2(s, t)\psi_{\nu}(t)dt,$$

or, if we multiply by λ_{ν} and put x in place of s ,

$$(62) \quad \psi_{\nu}(x) = \lambda_{\nu}^2 \int_a^b K_2(x, t)\psi_{\nu}(t)dt.$$

From (62) we derive

$$\psi_{\nu}(x) = \lambda_{\nu}^3 \int_a^b K_3(x, t)\psi_{\nu}(t)dt$$

in the same way that (62) was derived from (61). Continuing this process, it is easy to show by mathematical induction that

$$(63) \quad \psi_\nu(x) = \lambda_\nu^n \int_a^b K_n(x, t) \psi_\nu(t) dt.$$

Therefore, λ_ν^n is a characteristic constant of $K_n(x, t)$, and $\psi_\nu(x)$ is a fundamental function of $K_n(x, t)$ belonging to λ_ν^n .

b) It is still necessary to show that $K_n(x, t)$ has no other characteristic constants than the λ_ν^n , and that every fundamental function $\varphi(x)$ of $K_n(x, t)$ can be expressed in the form

$$\varphi(x) = C_{\nu_1} \psi_{\nu_1}(x) + \dots + C_{\nu_r} \psi_{\nu_r}(x).$$

Let ρ be any characteristic constant of $K_n(x, t)$ and $\varphi(x)$ be a fundamental function of $K_n(x, t)$ belonging to ρ :

$$(64) \quad \varphi(x) = \rho \int_a^b K_n(x, t) \varphi(t) dt, \quad \varphi(x)^c \not\equiv 0.$$

We must then prove that

$$1) \quad \rho = \lambda_\nu^n.$$

$$2) \quad \varphi(x) = \sum_{i=1}^r C_{\nu_i} \psi_{\nu_i}(x).$$

Let h_1, h_2, \dots, h_n be the n th roots of ρ :

$$(65) \quad h_\nu^n = \rho, \quad (\nu = 1, \dots, n).$$

Build up the functions $nG_i(x)$ as follows:

$$(66) \quad \begin{aligned} nG_i(x) &= \varphi(x) + h_i \int_a^b K_1(x, t) \varphi(t) dt \\ &\quad + h_i^2 \int_a^b K_2(x, t) \varphi(t) dt + \dots \\ &\quad + h_i^{n-1} \int_a^b K_{n-1}(x, t) \varphi(t) dt, \end{aligned}$$

for $i = 1, \dots, n$. Add these equations and cancel a factor n . We obtain

$$(67) \quad G_1(x) + \dots + G_n(x) = \varphi(x),$$

since $h_1^s + h_2^s + \dots + h_n^s = 0$,
 $(s = 1, \dots, n - 1)$.

Multiply both sides of (66) by $K(s, x)$ and integrate. We obtain

$$n \int_a^b K(s, x) G_i(x) dx = \int_a^b K(s, x) \varphi(x) dx + \\ \sum_{s=1}^{n-1} h_i^s \int_a^b \int_a^b K(s, x) K_s(x, t) \varphi(t) dt dx.$$

Multiply the members of this equation by h_i , put $h_i^n = \rho$ by (65), make use of (64) for $x = s, t = x$, and of the definitions of the iterated kernels. We obtain

$$nh_i \int_a^b K(s, x) G_i(x) dx = h_i \int_a^b K(s, t) \varphi(t) dt \\ + h_i^2 \int_a^b K_2(s, t) \varphi(t) dt + \dots \\ + h_i^{n-1} \int_a^b K_{n-1}(s, t) \varphi(t) dt + \varphi(s),$$

which, on account of (66), may be written, after canceling a factor n ,

$$h_i \int_a^b K(s, x) G_i(x) dx = G_i(s).$$

Suppose for some value of i that $G_i(x) \not\equiv 0$, then h_i is a characteristic constant of $K(x, t)$ and $G_i(x)$ is a fundamental function of $K(x, t)$ belonging to h_i . Therefore, there exists a value m of ν , such that $h_i = \lambda_m$ and hence $\rho = \lambda_m^n$. Since $G_i(x)$ is a fundamental function of $K(x, t)$, it can be written in the form

$$(68) \quad G_i(x) = C_{i1}\psi_{i1} + \dots + C_{ik}\psi_{ik}.$$

But the $G_i(x)$ do not all vanish identically, otherwise from (67) we would have $\varphi(x) \equiv 0$, which cannot be. Hence

$$\rho = \lambda_m^n,$$

and from (67) and (68) we have

$$\varphi(x) = C_{\nu_1} \psi_{\nu_1}(x) + \dots + C_{\nu_r} \psi_{\nu_r}(x),$$

which completes the proof of the theorem.

V. AUXILIARY THEOREMS

65. Bessel's Inequality.—For the further discussion of the bilinear formula we need the following theorem:

Theorem XIII.—*Given*

1) $f(x)$, real and continuous.

2) ψ_s , ($s = 1, \dots, m$), real and continuous, constituting a normalized orthogonal set:

$$\int_a^b \psi_r \psi_s dx = \begin{cases} 1, & r = s \\ 0, & r \neq s, \end{cases}$$

then

$$(69) \quad \sum_{r=1}^m \left[\int_a^b f(x) \psi_r(x) dx \right]^2 \leq \int_a^b [f(x)]^2 dx.$$

Proof.—Let C_s be any real constants, then

$$\int_a^b \left[f(x) - \sum_{s=1}^m C_s \psi_s(x) \right]^2 dx \geq 0,$$

whence, after squaring the expression in the bracket,

$$(70) \quad \int_a^b [f(x)]^2 dx - 2 \sum_{s=1}^m C_s \int_a^b f(x) \psi_s(x) dx + \int_a^b \left[\sum_{s=1}^m C_s \psi_s(x) \right]^2 dx \geq 0.$$

But

$$\begin{aligned} \int_a^b \left[\sum_{s=1}^m C_s \psi_s(x) \right]^2 dx &= \sum_{s=1}^m C_s^2 \int_a^b \psi_s^2 dx \\ &\quad + 2 \sum_{(r,s)} C_r C_s \int_a^b \psi_r \psi_s dx. \end{aligned}$$

Now

$$\int_a^b \psi_s^2 dx = 1 \text{ and } \int_a^b \psi_r \psi_s dx = 0$$

by hypothesis. Therefore,

$$\int_a^b \left[\sum_{s=1}^m C_s \psi_s(x) \right]^2 dx = \sum_{s=1}^m C_s^2.$$

But C_s are any real constants and so we may choose

$$C_s = \int_a^b f(x) \psi_s(x) dx.$$

For this choice of C_s , equation (70) becomes

$$\begin{aligned} \int_a^b [f(x)]^2 dx - 2 \sum_{s=1}^m \left(\int_a^b f(x) \psi_s(x) dx \right)^2 \\ + \sum_{s=1}^m \left(\int_a^b f(x) \psi_s(x) dx \right)^2 \geq 0, \end{aligned}$$

whence we obtain (69).

We now apply (69) to the particular case where

$$f(x) = K(x, t)$$

for a fixed t , and the functions ψ_s are a normalized orthogonal system of fundamental functions of $K(x, t)$ belonging to λ_s .

Equation (69) now becomes

$$(71) \quad \sum_{s=1}^m \left(\int_a^b K(x, t) \psi_s(x) dx \right)^2 \leq \int_a^b [K(x, t)]^2 dx.$$

But

$$\psi_s(x) = \lambda_s \int_a^b K(x, t) \psi_s(t) dt,$$

whence

$$\begin{aligned} \psi_s(t) &= \lambda_s \int_a^b K(t, x) \psi_s(x) dx \\ (72) \quad &= \lambda_s \int_a^b K(x, t) \psi_s(x) dx, \end{aligned}$$

on account of the symmetry of $K(x, t)$. Now $K(x, t)$ is continuous on R and, therefore,

$$(73) \quad |K(x, t)| \leq G \quad (\text{a constant}).$$

Applying (72) and (73) to (71), we obtain

$$\sum_{s=1}^m \left(\frac{\psi_s(t)}{\lambda_s} \right)^2 \leq G^2(b - a).$$

Thus we have the following corollary to Theorem XIII:

Corollary.—If $\psi_s(x)$, ($s = 1, \dots, m$) are normalized orthogonal fundamental functions of the kernel $K(x, t)$, belonging to the characteristic constant λ_s , then

$$(74) \quad \sum_{s=1}^m \left(\frac{\psi_s(x)}{\lambda_s} \right)^2 \leq G^2(b - a)$$

From (74) we obtain by integrating from a to b

$$\sum_{s=1}^m \frac{1}{\lambda_s^2} \int_a^b [\psi_s(x)]^2 dx \leq G^2(b - a)^2.$$

But $\int_a^b [\psi_s(x)]^2 dx = 1$ by hypothesis.

Therefore

$$(75) \quad \sum_{s=1}^m \frac{1}{\lambda_s^2} \leq G^2(b - a)^2,$$

the inequality holding for any finite number of λ_s and hence for the first m of the λ_s 's.

Theorem XIV.— $\sum_{s=1}^{\infty} \frac{1}{\lambda_s^2}$ is convergent.

Proof.—The proof follows from (75) by applying the principle of monotony: if u_i are real and positive and

$$u_1 + u_2 + \dots + u_n \leq A, \quad (n)$$

that is bounded, then

$$\sum_{n=1}^{\infty} u_n$$

is convergent.

66. Proof of the Bilinear Formula for the Iterated Kernel $K_n(x, t)$ for $n \leq 4$.—In Theorem XI it was stated that if $\sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}}$ was uniformly convergent, then

$$\sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}} = K(x, t).$$

We have stated previously that this series is not always uniformly convergent and, therefore, the bilinear formula does not hold for every kernel. But we shall be able to prove that it does always hold for the iterated kernels.

Since, according to Theorem XII, $\psi_{\nu}(x)$ are a complete normalized orthogonal system of fundamental functions for $K_n(x, t)$ belonging to λ_{ν}^n , the bilinear formula for the iterated kernel, if true, would read

$$(76) \quad \sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}^n} = K_n(x, t), \quad n > 1.$$

By referring to Theorem XI just quoted, we see that (76) will be proved if we can show that $\sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}^n}$ is uniformly convergent. We prove this first for the case $n = 4$. We desire then to show the uniform convergence of

$$(77) \quad \sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}^4} \text{ on } R.$$

We apply the general test for the uniform convergence of the series

$$\sum_{\nu=1}^{\infty} u_{\nu}(x, t), \text{ on } R.$$

For any $\epsilon > 0$ we can assign an N depending on ϵ , but not on x and t (N_{ϵ}) such that

$|u_{n+1}(x, t) + u_{n+2}(x, t) + \dots + u_{n+p}(x, t)| < \epsilon, \quad (R)$
for every $n > N_{\epsilon}$ for any integer $p > 0$.

Use the notation

$$\Delta_{np} = \sum_{\nu=n+1}^{n+p} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu^4}.$$

We have the inequalities

$$(78) \quad |\Delta_{np}| \leq \sum_{\nu=n+1}^{n+p} \frac{|\psi_\nu(x)\psi_\nu(t)|}{\lambda_\nu^4} \leq \sum_{\nu} \frac{|\psi_\nu(x)||\psi_\nu(t)|}{|\lambda_\nu||\lambda_\nu|} \sum_{\nu} \frac{1}{\lambda_\nu^2}.$$

We now make use of the inequality

$$(79) \quad \sum_{\nu} a_\nu b_\nu \leq \sqrt{\sum a_\nu^2} \sqrt{\sum b_\nu^2}$$

(a_ν, b_ν real and positive), which arises from the following consideration: given the matrix

$\begin{vmatrix} a_1 & a_2 & \dots & a_p \\ b_1 & b_2 & \dots & b_p \end{vmatrix}$, form the product

$$(80) \quad \begin{aligned} \left\| \begin{vmatrix} a_1 & a_2 & \dots & a_p \\ b_1 & b_2 & \dots & b_p \end{vmatrix} \right\|^2 &= \begin{vmatrix} \sum_{\nu=1}^p a_\nu^2 & \sum_{\nu=1}^p a_\nu b_\nu \\ \sum_{\nu=1}^p a_\nu b_\nu & \sum_{\nu=1}^p b_\nu^2 \end{vmatrix} \\ &= \sum_{(\mu, \nu)=1}^p \begin{vmatrix} a_\mu & a_\nu \\ b_\mu & b_\nu \end{vmatrix}^2 \geq 0. \end{aligned}$$

From the inequality in (80) follows (79), which is the algebraic analogue of Schwarz's inequality (21) of §57.

Now apply (79) to (78) with $a_\nu = \left| \frac{\psi_\nu(x)}{\lambda_\nu} \right|$, $b_\nu = \left| \frac{\psi_\nu(t)}{\lambda_\nu} \right|$. We obtain

$$\sum_{\nu=n+1}^{n+p} \left| \frac{\psi_\nu(x)}{\lambda_\nu} \right| \left| \frac{\psi_\nu(t)}{\lambda_\nu} \right| \leq \sqrt{\sum_{\nu=n+1}^{n+p} \frac{\psi_\nu^2(x)}{\lambda_\nu^2}} \sqrt{\sum_{\nu=n+1}^{n+p} \frac{\psi_\nu^2(t)}{\lambda_\nu^2}}.$$

But by (74)

$$(81) \quad \sum_{\nu=n+1}^{n+p} \frac{\psi_\nu^2(x)}{\lambda_\nu^2} \leq G^2(b-a), \quad \sum_{\nu=n+1}^{n+p} \frac{\psi_\nu^2(t)}{\lambda_\nu^2} \leq G^2(b-a).$$

By Theorem XIV, $\sum \frac{1}{\lambda_\nu^2}$ is convergent, and thus, by the general theorem on convergent series,

$$\lim_{n \rightarrow \infty} d_{np} = 0,$$

where

$$(82) \quad d_{np} = \sum_{\nu=n+1}^{n+p} \frac{1}{\lambda_\nu^2}.$$

Substitution from (81) and (82) in (78) gives

$$|\Delta_{np}| \leq G^2(b-a)d_{np}.$$

The right member of this inequality is independent of x and t , therefore (77) is absolutely and uniformly convergent in x and t on R . Thus we have proved the following theorem:

Theorem XV. — $\sum_{\nu} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu^4}$ is absolutely and uniformly convergent in x and t on R , and

$$\sum_{\nu} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu^4} = K_4(x, t).$$

We can now show that (76) holds for $n > 4$. Suppose that (76) holds in the n th instance:

$$(83) \quad K_n(x, t) = \sum_{\nu} \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu^n},$$

the series being uniformly convergent in x and t on R . Multiply both sides of (83) by $K(s, x)$ and integrate with respect to x from a to b . We obtain

$$(84) \quad \int_a^b K(s, x)K_n(x, t)dx = \sum_{\nu} \frac{\psi_\nu(t)}{\lambda_\nu^n} \int_a^b K(s, x)\psi_\nu(x)dx$$

and the series on the right is again uniformly convergent as to x and t . But

$$\int_a^b K(s, x)\psi_\nu(x)dx = \frac{\psi_\nu(s)}{\lambda_\nu}$$

and

$$\int_a^b K(s, x) K_n(x, t) dx = K_{n+1}(s, t).$$

We obtain, then, from (84), if we put $s = x$,

$$K_{n+1}(x, t) = \sum_{\nu} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu}^{n+1}}.$$

The induction is now complete and (76) holds for $n \geq 4$. Kowalewski¹ gives a proof that (76) holds for $n = 2$, whence from our induction proof, (76) holds for $n = 3$, and therefore for $n > 1$.

67. An Auxiliary Theorem of Schmidt.—Before we can take up the problem of the expansion of an arbitrary function according to the fundamental functions of a symmetric kernel, we need another auxiliary theorem due to E. Schmidt.

Theorem XVI.—*Let $h(x)$ be continuous on $[ab]$, $K(x, t)$ real and symmetric with the complete normalized orthogonal system of fundamental functions $\{\psi_{\nu}(x)\}$ belonging to the characteristic constants $\{\lambda_{\nu}\}$, then*

A) If $\int_a^b K(x, t)h(t)dt = 0$, uniformly as to x , then

$$\int_a^b \psi_n(x)h(x)dx = 0, \text{ uniformly as to } n.$$

B) If $\int_a^b \psi_n(x)h(x)dx = 0$, uniformly as to n , then

$$\int_a^b K(x, t)h(t)dt = 0, \text{ uniformly as to } x.$$

Proof.—A) Multiply both members of

$$\int_a^b K(x, t)h(t)dt = 0, (x)$$

¹ KOWALEWSKI, "Einführung in die Determinanten Theorie," Teubner, p. 533, §200, 1900.

by $\psi_n(x)$ and integrate. We obtain

$$(85) \quad \int_a^b \int_a^b K(x, t) \psi_n(x) h(t) dt dx = 0,$$

which may be written

$$\int_a^b h(t) \left(\int_a^b K(x, t) \psi_n(x) dx \right) dt = 0.$$

$$\text{But} \quad \int_a^b K(x, t) \psi_n(x) dx = \frac{\psi_n(t)}{\lambda_n}.$$

Therefore (85) may be written

$$\frac{1}{\lambda_n} \int_a^b \psi_n(t) h(t) dt = 0, \quad (n).$$

$$\text{Therefore} \quad \int_a^b \psi_n(t) h(t) dt = 0, \quad (n).$$

B) Our hypothesis is that

$$\int_a^b \psi_n(x) h(x) dx = 0, \quad (n).$$

We have previously shown that

$$K_4(x, t) = \sum_{\nu} \frac{\psi_{\nu}(x) \psi_{\nu}(t)}{\lambda_{\nu}^4}$$

is uniformly convergent on R . Multiply both members of this equality by $h(x)h(t)$ and integrate. We obtain

$$\int_a^b \int_a^b K_4(x, t) h(x) h(t) dx dt = \sum_{\nu} \frac{1}{\lambda_{\nu}^4} \int_a^b \int_a^b \psi_{\nu}(x) \times \\ h(x) \cdot \psi_{\nu}(t) h(t) dt dx.$$

The right member of this equality may be written

$$\sum_{\nu} \frac{1}{\lambda_{\nu}^4} \int_a^b \psi_{\nu}(x) h(x) dx \cdot \int_a^b \psi_{\nu}(t) h(t) dt,$$

which, on account of our hypothesis, vanishes. Therefore

$$(86) \quad \int_a^b \int_a^b K_4(x, t) h(x) h(t) dx dt = 0.$$

$$\text{But} \quad K_4(x, t) = \int_a^b K_2(x, s) K_2(t, s) ds,$$

since $K_2(t, s) = K_2(s, t)$ from the symmetry of K . Substitute this value of $K_4(x, t)$ in (86). We obtain

$$\int_a^b \int_a^b \int_a^b K_2(x, s) K_2(t, s) h(x) h(t) ds dx dt = 0,$$

which may be written

$$\int_a^b \left[\int_a^b K_2(x, s) h(x) dx \times \int_a^b K_2(t, s) h(t) dt \right] ds = 0,$$

or

$$\int_a^b \left[\int_a^b K_2(x, s) h(x) dx \right]^2 ds = 0.$$

But $\int_a^b K_2(x, s) h(x) ds$ is a real continuous function of s .

Therefore

$$\int_a^b K_2(x, s) h(x) dx \equiv 0, (s),$$

which, if we change our notation and, in place of s and x , put t and x , and remember that $K_2(x, t) = K_2(t, x)$, becomes

$$\int_a^b K_2(x, t) h(t) dt \equiv 0, (x).$$

Multiply both members of this last equation by $h(x)$ and integrate. We obtain

$$(87) \quad \int_a^b \int_a^b K_2(x, t) h(x) h(t) dx dt = 0.$$

But

$$K_2(x, t) = \int_a^b K(x, s) K(t, s) ds.$$

Treat (87) in the same manner as we have just done with (86) and we will finally obtain

$$\int_a^b K(x, t) h(t) dt \equiv 0, (x).$$

This completes the proof of Schmidt's auxiliary theorem.

VI. EXPANSION OF AN ARBITRARY FUNCTION ACCORDING TO THE COMPLETE NORMALIZED ORTHOGONAL SYSTEM OF FUNDAMENTAL FUNCTIONS OF A SYMMETRIC KERNEL

68. We are now in a position to solve the fundamental problem stated in §62: Given an arbitrary function $f(x)$, to represent it, if possible, in the form

$$f(x) = \sum_{\nu} C_{\nu} \psi_{\nu}(x),$$

when the $\psi_{\nu}(x)$ are the fundamental functions of a complete normalized orthogonal system of the symmetric kernel $K(x, t)$.

We have seen in §62, that if such an expansion is possible and, moreover, uniformly convergent, then the coefficients C_{ν} must have the values

$$C_{\nu} = \int_a^b f(x) \psi_{\nu}(x) dx \equiv (f\psi_{\nu}),$$

that is

$$f(x) = \sum_{\nu} (f\psi_{\nu}) \psi_{\nu}(x).$$

We now prove the following theorem:

Theorem XVII.—*If $f(x)$ can be represented in the form*

$$(88) \quad f(x) = \int_a^b K(x, t) g(t) dt,$$

where $g(x)$ is a function continuous on $[ab]$, then

$$f(x) = \sum_{\nu} (f\psi_{\nu}) \psi_{\nu}(x) \text{ on } [ab]$$

and the series is uniformly and absolutely convergent on $[ab]$.

Proof.—a) Convergence Proof. Use the notation

$$\Delta_{n,p} = \sum_{\nu=n+1}^{n+p} (f\psi_{\nu}) \psi_{\nu}(x), \quad p > 0.$$

Then, from the general convergence principle stated in §66, the series

$$(89) \quad \sum_{\nu=1}^{\infty} (f\psi_{\nu})\psi_{\nu}(x)$$

converges uniformly on $[ab]$ if for every $\epsilon > 0$ there exists an N dependent upon ϵ , but not on $x(N_{\epsilon})$, such that for every $n > N_{\epsilon}$ it is true that $|\Delta_{np}| < \epsilon$ on $[ab]$. Now

$$(90) \quad \begin{aligned} (f\psi_{\nu}) &= \int_a^b f(x)\psi_{\nu}(x)dx \\ &= \int_a^b \int_a^b K(x, t)g(t)\psi_{\nu}(x)dt dx \end{aligned}$$

on account of (88).

The right member of (90) may be written

$$\int_a^b g(t) \left(\int_a^b K(x, t)\psi_{\nu}(x)dx \right) dt.$$

But $\int_a^b K(x, t)\psi_{\nu}(x)dx = \frac{\psi_{\nu}(t)}{\lambda_{\nu}}$.

Therefore (90) becomes

$$(91) \quad (f\psi_{\nu}) = \frac{1}{\lambda_{\nu}} (g\psi_{\nu}).$$

Now

$$|\Delta_{np}| \leq \sum_{\nu=n+1}^{n+p} |(f\psi_{\nu})| |\psi_{\nu}(x)|.$$

Therefore, by (91),

$$(92) \quad |\Delta_{np}| \leq \sum_{\nu=n+1}^{n+p} |(g\psi_{\nu})| \cdot \left| \frac{\psi_{\nu}(x)}{\lambda_{\nu}} \right|.$$

But from the algebraic analogue (79) of Schwarz's inequality (21), proved in §57, we have

$$\sum_{\nu} |(g\psi_{\nu})| \left| \frac{\psi_{\nu}(x)}{\lambda_{\nu}} \right| \leq \sqrt{\sum_{\nu} (g\psi_{\nu})^2} \sqrt{\sum_{\nu} \frac{\psi_{\nu}^2(x)}{\lambda_{\nu}^2}},$$

and from (74) in the corollary to Bessel's inequality

$$\sum_{\nu} \frac{\psi_{\nu}^2(x)}{\lambda_{\nu}^2} \leq G^2(b-a),$$

G being the maximum of $|K(x, t)|$ on R .

If, now, we use the notation

$$d_{np} = \sum_{\nu=n+1}^{n+p} (g\psi_{\nu})^2$$

we may write

$$(93) \quad |\Delta_{np}| \leq G\sqrt{(b-a)d_{np}}.$$

But

$$(94) \quad \sum_{\nu=1}^{\infty} (g\psi_{\nu})^2$$

is convergent, for

$$\sum_{\nu=1}^n (g\psi_{\nu})^2 = \sum_{\nu=1}^n \left(\int_a^b g(x)\psi_{\nu}(x)dx \right)^2 \leq \int_a^b [g(x)]^2 dx,$$

from Bessel's inequality (69), §65, and $\int_a^b [g(x)]^2 dx$ is a finite fixed constant, independent of n , and therefore

$$\sum_{\nu=1}^n (g\psi_{\nu})^2$$

is bounded for every n , and hence, by the monotony principle (94), is convergent. Therefore, $\lim_{n \rightarrow \infty} d_{np} = 0$.

Hence, from (93), $\lim_{n \rightarrow \infty} |\Delta_{np}| = 0$, and therefore (89) is uniformly convergent on $[ab]$, and from (92) we see that it is also absolutely convergent. This completes the first part of the proof. It remains to determine the sum.

b) *Summation.*—Put

$$h(x) \equiv f(x) - \sum_{\nu} (f\psi_{\nu})\psi_{\nu}(x).$$

From (88), $f(x)$ is continuous, since $g(x)$ is supposed to be continuous and the ψ_ν 's are continuous and the series is uniformly convergent, hence h is continuous. We will have proved that the series represents $f(x)$, if we can show

that $\int_a^b [h(x)]^2 dx = 0,$

for then

$$h(x) \equiv 0.$$

Now

$$(95) \quad \int_a^b [h(x)]^2 dx = (fh) - \sum_\nu (f\psi_\nu)(h\psi_\nu).$$

We compute

$$(h\psi_n) = (f\psi_n) - \sum_\nu (f\psi_\nu)(\psi_\nu\psi_n).$$

But $(\psi_\nu\psi_n) = \begin{cases} 1, & \nu = n \\ 0, & \nu \neq n. \end{cases}$

Therefore

$$(96) \quad (h\psi_n) = (f\psi_n) - (f\psi_n) = 0.$$

Whence, from (95),

$$(97) \quad \int_a^b [h(x)]^2 dx = (fh).$$

But $(fh) = \int_a^b f(x)h(x)dx$

and, by hypothesis, as given by (88),

$$f(x) = \int_a^b K(x, t)g(t)dt.$$

Therefore,

$$(98) \quad \begin{aligned} (fh) &= \int_a^b \int_a^b K(x, t)g(t)h(x)dt dx \\ &= \int_a^b g(t) \left(\int_a^b K(x, t)h(x)dx \right) dt. \end{aligned}$$

Now $\int_a^b \psi_n(x)h(x)dx = 0, (n) \text{ by (96),}$

hence, by Theorem XVI, §67,

$$\int_a^b K(x, t)h(t)dt = 0, \quad (x)$$

and, therefore,

$$\int_a^b K(x, t)h(x)dx = 0, \quad (t)$$

whence, from (98),

$$(fh) \equiv 0, \quad (x)$$

and, therefore, from (97),

$$h(x) \equiv 0, \quad (x),$$

which completes the proof of the theorem. We have the following corollaries to Theorem XVII.

Corollary I.— $(f\psi_\nu) = \frac{(g\psi_\nu)}{\lambda_\nu}$ by (91).

Corollary II.—If $g(x)$ is continuous, then

$$(99) \quad \int_a^b K(x, t)g(t)dt = \sum_{\nu=1}^{\infty} \frac{(g\psi_\nu)}{\lambda_\nu} \psi_\nu(x)$$

and the series is uniformly and absolutely convergent.

This follows from Theorem XVII and Corollary I.

Corollary III.—If $g(x)$ and $h(x)$ are continuous, then

$$\int_a^b \int_a^b K(x, t)g(t)h(x)dtdx = \sum_{\nu=1}^{\infty} \frac{(g\psi_\nu)(h\psi_\nu)}{\lambda_\nu}.$$

This is sometimes called *Hilbert's formula*. It follows from (99), if we multiply by $h(x)$ and integrate.

VII. SOLUTION OF THE INTEGRAL EQUATION

69. Schmidt's Solution of the Non-homogeneous Integral Equation When λ Is Not a Characteristic Constant.—As an application of the last theorem, we will now give Schmidt's solution of the equation

$$(100) \quad f(x) = u(x) - \lambda \int_a^b K(x, t)u(t)dt.$$

A) Necessary Form of the Solution.—Let $u(x)$ be a continuous solution of (100), then

$$(101) \quad v(x) = \frac{u(x) - f(x)}{\lambda} = \int_a^b K(x, t)u(t)dt.$$

The function $v(x)$ satisfies the condition (88) of Theorem XVII and, therefore,

$$v(x) = \sum_{\nu} (v\psi_{\nu})\psi_{\nu}(x),$$

the series being absolutely and uniformly convergent on $[ab]$. On account of (101) and Corollary I §68, we have

$$(102) \quad v(x) = \sum_{\nu} \frac{(u\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x).$$

Multiply (100) by $\psi_{\nu}(x)$ and integrate. We obtain

$$\begin{aligned} (f\psi_{\nu}) &= (u\psi_{\nu}) - \lambda \int_a^b \int_a^b K(x, t)u(t)\psi_{\nu}(x)dtdx \\ &= (u\psi_{\nu}) - \lambda \int_a^b \frac{\psi_{\nu}(t)u(t)}{\lambda_{\nu}} dt \\ &= (u\psi_{\nu}) - \frac{\lambda}{\lambda_{\nu}}(u\psi_{\nu}), \end{aligned}$$

which may be written

$$(\lambda_{\nu} - \lambda)(u\psi_{\nu}) = \lambda_{\nu}(f\psi_{\nu}).$$

When $\lambda \neq \lambda_{\nu}$, we have

$$(103) \quad (u\psi_{\nu}) = \frac{\lambda_{\nu}(f\psi_{\nu})}{\lambda_{\nu} - \lambda}.$$

If λ is not a characteristic constant, then $\lambda \neq \lambda_1, \lambda_2, \dots$, and for such values of λ (103) holds for all values of ν and hence, from (102),

$$v(x) = \sum_{\nu=1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x).$$

Whence from (100) and (101)

$$(104) \quad u(x) = f(x) + \lambda \sum_{\nu=1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x).$$

Hence, if (100) has a continuous solution $u(x)$, this solution is unique and is given by (104). Replacing the symbol $(f\psi_{\nu})$ by its explicit expression, we have for the solution of (100)

$$(105) \quad u(x) = f(x) + \lambda \sum_{\nu=1}^{\infty} \left[\frac{1}{\lambda_{\nu} - \lambda} \int_a^b f(t) \psi_{\nu}(t) dt \right] \psi_{\nu}(x).$$

If for a given value of λ , the series

$$\sum_{\nu=1}^{\infty} \frac{\psi_{\nu}(t)\psi_{\nu}(x)}{\lambda_{\nu} - \lambda}$$

is uniformly convergent in t , we can write the solution in a form previously obtained:

$$u(x) = f(x) + \lambda \int_a^b K(x, t; \lambda) f(t) dt$$

where

$$K(x, t; \lambda) = \sum_{\nu=1}^{\infty} \frac{\psi_{\nu}(x)\psi_{\nu}(t)}{\lambda_{\nu} - \lambda}.$$

The solution given by (105) has an advantage over that given by Fredholm, in that it shows the meromorphic character of the solution with respect to λ and indicates the principal part of $u(x)$ with respect to each pole λ .

B) Sufficiency Proof.—It remains to show that (104) is absolutely and uniformly convergent and satisfies (100).

1) *Uniform and Absolute Convergence.*—The series in (104) may be written as follows

$$(106) \quad \sum_{\nu=1}^{\infty} \frac{1}{1 - \frac{\lambda}{\lambda_{\nu}}} \frac{(f\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x).$$

But by Corollary II to Theorem XVII

$$\sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x)$$

is absolutely and uniformly convergent; and if there is an infinitude of characteristic constants λ_{ν} , then as $\nu \rightarrow \infty$ also $\lambda_{\nu} \rightarrow \infty$, and therefore $\frac{1}{1 - \frac{\lambda}{\lambda_{\nu}}} \rightarrow 1$, whence also (106) is

absolutely and uniformly convergent.

2) (104) *Satisfies (100).*—From (104) we obtain

$$u(x) - \lambda \int_a^b K(x, t) u(t) dt = f(x) + \lambda \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x) \\ - \lambda \int_a^b K(x, t) \left\{ f(t) + \lambda \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(t) \right\} dt.$$

The right-hand member may be written

$$(107) \quad f(x) + \lambda \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x) - \lambda \int_a^b K(x, t) f(t) dt \\ - \lambda^2 \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \int_a^b K(x, t) \psi_{\nu}(t) dt.$$

$$\text{But } \int_a^b K(x, t) \psi_{\nu}(t) dt = \frac{\psi_{\nu}(x)}{\lambda_{\nu}},$$

therefore (107) may be written

$$f(x) + \lambda \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x) \left\{ 1 - \frac{\lambda}{\lambda_{\nu}} \right\} - \lambda \int_a^b K(x, t) f(t) dt$$

or

$$f(x) + \lambda \sum_{\nu} \frac{(f\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x) - \lambda \int_a^b K(x, t) f(t) dt.$$

By (99) this last expression reduces to $f(x)$ and hence

$$u(x) - \lambda \int_a^b K(x, t) u(t) dt = f(x).$$

We have thus proved the following theorem:

Theorem XVIII.—*If $f(x)$ is continuous and λ is not a characteristic constant, then*

$$u(x) = f(x) + \lambda \int_a^b K(x, t)u(t)dt$$

has one and only one continuous solution $u(x)$ given by

$$u(x) = f(x) + \lambda \sum_{\nu} \left[\frac{1}{\lambda_{\nu} - \lambda} \int_a^b f(t)\psi_{\nu}(t)dt \right] \psi_{\nu}(x),$$

the series being absolutely and uniformly convergent.

70. Schmidt's Solution of the Non-homogeneous Integral Equation When λ Is a Characteristic Constant.—Let us suppose that λ is a characteristic constant, for example, $\lambda = \lambda_1$ of index q , so that

$$\lambda = \lambda_1 = \lambda_2 = \dots = \lambda_q.$$

A) As before, we find that

$$(\lambda_{\nu} - \lambda)(u\psi_{\nu}) = \lambda_{\nu}(f\psi_{\nu}).$$

Now, since $\lambda_{\nu} = \lambda$, $\nu = 1, 2, \dots, q$,

we have

$$(108) \quad (f\psi_{\nu}) = 0, \nu = 1, 2, \dots, q.$$

We have thus q necessary conditions on f . If $\nu > q$, then $\lambda_{\nu} - \lambda \neq 0$ and

$$(u\psi_{\nu}) = \frac{\lambda_{\nu}(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \text{ as before.}$$

Now $v(x)$ is expressible as before by the formula (102), but for $\nu = 1, \dots, q$ the constant coefficients $\frac{(u\psi_{\nu})}{\lambda_{\nu}}$ are as yet undetermined. So we write

$$v(x) = C_1\psi_1(x) + \dots + C_q\psi_q(x) + \sum_{\nu=q+1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x)$$

whence

$$(109) \quad u(x) = f(x) + C_1\psi_1(x) + \dots + C_q\psi_q(x) \\ + \lambda \sum_{\nu=q+1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x).$$

B) That (109) converges is seen as before. To show that (109) satisfies (100) we first show that

$$u_o(x) = f(x) + \lambda \sum_{\nu=q+1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu} - \lambda} \psi_{\nu}(x)$$

satisfies (100). For this proceed as before and we find, after some reductions,

$$u_o(x) - \lambda \int_a^b K(x, t)u_o(t)dt = f(x) \\ + \lambda \sum_{\nu=q+1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x) - \lambda \sum_{\nu=1}^{\infty} \frac{(f\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x).$$

But $(f\psi_{\nu}) = 0$ for $\nu = 1, 2, \dots, q$, by (108) and thus we get

$$(110) \quad u_o(x) - \lambda \int_a^b K(x, t)u_o(t)dt = f(x).$$

It remains to show that

$$u(x) = u_o(x) + C_1\psi_1(x) + \dots + C_q\psi_q(x) \\ \equiv u_o(x) + \varphi(x)$$

satisfies (100). Now

$$u(x) - \lambda \int_a^b K(x, t)u(t)dt = u_o(x) - \lambda \int_a^b K(x, t)u_o(t)dt \\ + \varphi(x) - \lambda \int_a^b K(x, t)\varphi(t)dt = f(x) \text{ by (110),}$$

for

$$\varphi(x) - \lambda \int_a^b K(x, t)\varphi(t)dt = 0,$$

since

$$\psi_{\alpha}(x) - \lambda \int_a^b K(x, t)\psi_{\alpha}(t)dt = 0 (\alpha = 1, \dots, q).$$

We have thus proved the following theorem:

Theorem XIX.—*If $\lambda = \lambda_1$ is a characteristic constant of index q , then*

$$u(x) = f(x) + \lambda_1 \int_a^b K(x, t)u(t)dt$$

has in general no solution. Solutions exist only if

$$(f\psi_\alpha) = 0, (\alpha = 1, \dots, q).$$

Then there are q^∞ solutions given by

$$(111) \quad u(x) = f(x) + \lambda \sum_{\nu=q+1}^{\infty} \left[\frac{1}{\lambda_\nu - \lambda} \int_a^b f(t)\psi_\nu(t)dt \right] \psi_\nu(x) \\ + C_1\psi_1(x) + \dots + C_q\psi_q(x),$$

where C_1, \dots, C_q are arbitrary constants.

71. Remarks on Obtaining a Solution.—To obtain the solution of any numerical problem, one must compute $D(\lambda)$ in order to obtain λ_ν . To obtain the ψ_ν , one must solve the corresponding homogeneous equation. Having obtained the ψ_ν and the λ_ν , one writes down the solution at once by direct substitution in the proper formula, (104) or (111).

In general, $D(\lambda)$ is an infinite series given by (5) in Chap. III. However, for many simple kernels all but a few of the first terms vanish identically and we obtain $D(\lambda)$ as a polynomial in λ . By the method of §25, Chap. III, $D(\lambda)$ can be obtained in finite form as a polynomial in λ for every kernel $K(x, t)$ which is a polynomial in x and t .

The ψ_ν can be computed from the formula for $\varphi_\alpha(x)$ given in Theorem VIII, §21, Chap. III. By the method of §25, Chap. III, ψ_ν can be obtained in finite form for every kernel $K(x, t)$ which is a polynomial in x and t .

EXERCISES

Compute $D(\lambda)$ and obtain the characteristic constants λ , for the following symmetric kernels for the specified interval $[ab]$.

	<i>Ans.</i>
1. $K(x, t) = 1, [01].$	$D(\lambda) = 1 - \lambda.$
2. $K(x, t) = -1, [01].$	$D(\lambda) = 1 + \lambda.$
3. $K(x, t) = xt, [01].$	$D(\lambda) = 1 - \frac{\lambda}{3}.$
4. $K(x, t) = \sin x \sin t [0, 2\pi].$	$D(\lambda) = 1 - \pi\lambda.$
5. $K(x, t) = e^x \cdot e^t [0, \log_e 2].$	$D(\lambda) = 1 - \frac{3}{2}\lambda.$
6. $K(x, t) = x + t, [01].$	$D(\lambda) = 1 - \lambda - \frac{\lambda^2}{12}.$
7. $K(x, t) = x^2 + t^2, [01].$	$D(\lambda) = 1 - \frac{2}{3}\lambda - \frac{4}{45}\lambda^2.$
8. $K(x, t) = x^2t + xt^2, [01].$	$D(\lambda) = 1 - \frac{\lambda}{2} - \frac{1}{240}\lambda^2.$
9. $K(x, t) = x^2 + xt + t^2, [01].$	$D(\lambda) = 1 - \lambda - \frac{7}{60}\lambda^2 + \frac{1}{2,160}\lambda^3.$

Solve the following equations:

10. $u(x) = \int_0^1 u(t)dt.$ $u(x) = C.$
11. $u(x) = x + \lambda \int_0^1 u(t)dt, (\lambda \neq 1).$ $u(x) = x + \frac{\lambda}{2(1 - \lambda)}.$
12. $u(x) = \frac{1}{2} - x + \int_0^1 u(t)dt.$ $u(x) = \frac{1}{2} - x + C.$
13. $u(x) = \lambda \int_0^1 (x + t)u(t)dt$ for $\lambda = -6 \pm 4\sqrt{3}.$
Ans. $u_1(x) = C_1(1 + \sqrt{3}x)$ for $\lambda = \lambda_1 = -6 + 4\sqrt{3}.$
 $u_2(x) = C_2(1 - \sqrt{3}x)$ for $\lambda = \lambda_2 = -6 - 4\sqrt{3}.$
14. $u(x) = x + \lambda \int_0^1 (x + t)u(t)dt, \lambda \neq \lambda_1, \lambda_2.$
Ans. $u(x) = x - \frac{\frac{\lambda}{2}x - 2 - \sqrt{3}}{\lambda + 6 + 4\sqrt{3}} - \frac{\frac{\lambda}{2}x + 2 - \sqrt{3}}{\lambda + 6 - 4\sqrt{3}}$
 or $u(x) = \frac{(6\lambda - 12)x - 4\lambda}{\lambda^2 + 12\lambda - 12}.$
15. $u(x) = (1 - \sqrt{3}x) + (-6 + 4\sqrt{3}) \int_0^1 (x + t)u(t)dt.$
Ans. $u(x) = (1 - \sqrt{3}x) + B(1 + \sqrt{3}x) - \left(1 + \frac{3}{2}x\right).$
16. $u(x) = (1 + \sqrt{3}x) + (-6 - 4\sqrt{3}) \int_0^1 (x + t)u(t)dt.$
Ans. $u(x) = (1 + \sqrt{3}x) + B(1 - \sqrt{3}x) - \left(1 + \frac{3}{2}x\right).$

CHAPTER VI

APPLICATIONS OF THE HILBERT-SCHMIDT THEORY

I. BOUNDARY PROBLEMS FOR ORDINARY LINEAR DIFFERENTIAL EQUATIONS

72. Introductory Remarks.—*a) Formulation of the Problem.*—We have previously considered the boundary problem

$$\frac{d^2u}{dx^2} + \lambda u = 0, \quad u(0) = 0, \quad u(1) = 0.$$

We shall now discuss the general type of boundary problem of which this is a special case, namely, we take the general homogeneous linear differential equation of the second order with a parameter λ contained linearly in the coefficients of u :

$$P(x) \frac{d^2u}{dx^2} + Q(x) \frac{du}{dx} + [R(x) + \lambda S(x)] u = 0.$$

We first reduce this equation to a self-adjoint linear differential equation by multiplying it by $\frac{1}{P(x)} e^{\int \frac{Q(x)}{P(x)} dx}$

Putting $p = e^{\int \frac{Q}{P} dx}$, we obtain

$$p \left(u'' + \frac{Q}{P} u' + \frac{R + \lambda S}{P} u \right) = 0,$$

which may be written

$$(1) \quad pu'' + p'u' + (q + \lambda g)u = 0$$

or

$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + (q + \lambda g)u = 0.$$

If we make use of the permanent notation

$$(2) \quad L(u) = \frac{d}{dx} \left(p \frac{du}{dx} \right) + qu,$$

equation (1) may be written in the form

$$(3) \quad L(u) + \lambda g(u) = 0.$$

We use also more general boundary conditions, and for the general interval $[ab]$:

$$(4) \quad \begin{aligned} R_0(u) &\equiv Au(a) + Bu'(a) = 0 \\ R_1(u) &\equiv Cu(b) + Du'(b) = 0, \end{aligned}$$

where A, B, C, D are given constants, of which A and B are not both zero, and C and D are not both zero.

Our problem then is to determine all solutions of class C'' of the differential equation (3) which satisfy the boundary conditions (4).

For this generalized boundary problem we make the hypotheses:

$$(H_1) \quad p \text{ is of class } C', p \neq 0 \text{ on } [ab]$$

$$q \text{ and } g \text{ are of class } C \text{ on } [ab].$$

Then (1) has two linearly independent solutions of class C'' on $[ab]$: u_1 and u_2 . Any other solution of (1) of class C'' is linearly expressible in terms of u_1 and u_2 :

$$u = C_1 u_1 + C_2 u_2.$$

The boundary problem has the trivial solution $u = 0$. Every value of λ for which our boundary problem has a non-trivial solution is called a characteristic constant and those solutions of class C'' , non-trivial, which exist for these values of λ are called fundamental functions of the boundary problem.

To the hypothesis (H_1) we add for the present a second hypothesis (H_2) which will be dropped later:

(H_2) $\lambda = 0$ is not a characteristic constant.

That is, $L(u) = 0$, $R_0(u) = 0$, $R_1(u) = 0$
has no solution of class C'' other than $u \equiv 0$.

b) *Green's Formula*.—From our hypothesis (H_1) we obtain the following theorem:

Theorem I.—If $u(x)$ and $v(x)$ are any two functions of x of class C'' , then

$$(5) \quad uL(v) - vL(u) = \frac{d}{dx} p(uv' - vu').$$

Proof.—Given two functions $u(x)$, $v(x)$ of class C'' , form

$$\begin{aligned} uL(v) - vL(u) &= \begin{vmatrix} u & pu'' + p'u' + qu \\ v & pv'' + p'v' + qv \end{vmatrix} \\ &= \begin{vmatrix} u & pu'' + p'u' \\ v & pv'' + p'v' \end{vmatrix} \\ &= p(uv'' - vu'') + p'(uv' - vu') \\ &= \frac{d}{dx} p(uv' - vu'). \end{aligned}$$

Corollary.—If u and v are of class C'' and $L(u) = 0$, $L(v) = 0$, then

$$(6) \quad p(uv' - vu') = \text{constant} \neq 0,$$

if u and v are independent.

For if $C = 0$, then $uv' - vu' \equiv 0$ and

$$\frac{uv' - vu'}{u^2} \equiv 0.$$

Whence $\frac{d}{dx}\left(\frac{v}{u}\right) \equiv 0$,

and, therefore, $v = Cu$ and v and u would be linearly dependent, contrary to hypothesis.

c) *Consequences of Hypothesis (H_2)*.—We now establish the following lemmas:

Lemma I.—If u_1, u_2 are of class C'' and $L(u_1) = 0$, $L(u_2) = 0$, and u_1, u_2 are linearly independent, then

$R_0(u_1), R_0(u_2)$ are not both zero and also

$R_1(u_1), R_1(u_2)$ are not both zero.

Proof.—The general solution of class C'' of $L(u) = 0$ is

$$u = c_1 u_1 + c_2 u_2.$$

Now $R_0(u) = c_1 R_0(u_1) + c_2 R_0(u_2)$.

Therefore, if the lemma is not true, then

$$R_0(u) = 0 \text{ for every } c_1, c_2.$$

But

$$R_1(u) = c_1 R_1(u_1) + c_2 R_1(u_2),$$

and, if c_1, c_2 are properly chosen, we can make $R_1(u) = 0$.

But then we would have

$$L(u) = 0, R_0(u) = 0, R_1(u) = 0,$$

which is contrary to hypothesis (H_2) . Therefore not both $R_0(u_1), R_0(u_2)$ can vanish at once. Likewise, we show that $R_1(u_1), R_1(u_2)$ cannot both vanish simultaneously.

Lemma II.—There exist functions u and v of class C'' , determined except for a constant factor, such that $L(u) = 0$, $R_0(u) = 0$, $L(v) = 0$, $R_1(v) = 0$. u and v are linearly independent and the constant factors can be determined so that $p(uv' - vu') = -1$.

Proof.—Let u_1, u_2 be two linearly independent solutions of $L(u) = 0$. Let $u = c_1 u_1 + c_2 u_2$ and determine c_1 and c_2 so that $R_0(u) = c_1 R_0(u_1) + c_2 R_0(u_2) = 0$. Now $R_0(u_1), R_0(u_2)$ cannot both be zero. Therefore

$$c_1 = \rho_0 R_0(u_2) \text{ and } c_2 = -\rho_0 R_0(u_1),$$

and hence

$$u = \rho_0 \left[u_1 R_0(u_2) - u_2 R_0(u_1) \right].$$

This is the most general expression for a function u for which $L(u) = 0, R_0(u) = 0$.

In like manner, we show that the most general expression for a function v for which

$$L(v) = 0, R_1(v) = 0$$

is

$$v = \rho_1 \left[v_1 R_1(v_2) - v_2 R_1(v_1) \right].$$

Suppose u and v were not independent, then

$$v = Cu, C \neq 0,$$

whence

$$R_1(u) = \frac{1}{C} R_1(v) = 0.$$

But by hypothesis $R_0(u) = 0$. This is in contradiction to our hypothesis (H_2).

According to Green's formula

$$p(uv' - vu') = C \neq 0;$$

hence the constant factors ρ_0, ρ_1 can be determined so that

$$p(uv' - vu') = -1.$$

73. Construction of Green's Function.—Let us take between a and b a fixed value t : $a < t < b$. Then the following theorem holds:

Theorem II.—*For our boundary problem there exists under the assumptions (H_1) and (H_2) one and but one function $K(x, t)$, which, as a function of x , has the following properties:*

A) K is continuous on $[ab]$.

B) K is of class C'' on each of its subintervals $[at][tb]$,

and $L(K) = 0$ on each subinterval.

C) $R_0(K) = 0, R_1(K) = 0$.

D) $K'(x, t - 0) - K'(x, t + 0) = \frac{i}{p(t)}$.

where

$$K'(x, t) = \frac{\partial}{\partial x} K(x, t).$$

This function $K(x, t)$ is called *Green's function*, belonging to the boundary problem.

Proof.—If $K(x, t)$ is to satisfy $L(K) = 0$, it must be linearly expressible in terms of the two functions u and v the existence of which have been shown in Lemma II of §72:

$$K(x, t) = \begin{cases} A_0 u + B_0 v, & \begin{bmatrix} at \\ tb \end{bmatrix} \\ A_1 u + B_1 v, & \end{cases}$$

Now demand that $K(x, t)$ satisfy C):

$$R_0(K) = R_0(A_0 u + B_0 v) = A_0 R_0(u) + B_0 R_0(v) = 0.$$

$$R_1(K) = R_1(A_1 u + B_1 v) = A_1 R_1(u) + B_1 R_1(v) = 0.$$

But $R_0(u) = 0$, and not both $R_0(u)$, $R_0(v)$ can vanish simultaneously, whence $B_0 = 0$. Also, $R_1(v) = 0$, and not both $R_1(u)$, $R_1(v)$ can vanish simultaneously, whence $A_1 = 0$. Therefore

$$K(x, t) = \begin{cases} A_0 u, & \begin{bmatrix} at \\ tb \end{bmatrix} \\ B_1 v, & \end{cases}$$

We now impose the condition A):

$$A_0 u(t) = B_1 v(t),$$

whence

$$A_0 = \rho v(t), \quad B_1 = \rho u(t),$$

and

$$K(x, t) = \begin{cases} \rho v(t) u(x), & \begin{bmatrix} at \\ tb \end{bmatrix} \\ \rho u(t) v(x), & \end{cases}$$

Now impose the condition D):

$$K'(t - 0) - K'(t + 0) = \frac{1}{p(t)},$$

that is

$$\rho v(t)u'(t) - \rho u(t)v'(t) = \frac{1}{p(t)}.$$

But

$$p(uv' - vu') = -1.$$

Therefore

$$\rho \left[v(t)u'(t) - u(t)v'(t) \right] = \frac{\rho}{p(t)},$$

whence

$$\rho = +1.$$

Therefore

$$(7) \quad K(x, t) = \begin{cases} v(t)u(x) \equiv K_0(x, t), & a \leq x \leq t \\ u(t)v(x) \equiv K_1(x, t), & t \leq x \leq b. \end{cases}$$

This is *Green's function* and it is uniquely determined.

Corollary.—*Green's function is symmetric. That is*

$$K(x, t) = K(t, x).$$

To show this, compute $K(z_1, z_2)$ and $K(z_2, z_1)$ where

$$0 \leq z_1 < z_2 \leq b.$$

$$K(z_1, z_2) = v(z_2)u(z_1),$$

for here we identify z_1 and x , z_2 and t ; and, since $z_1 < z_2$, we use the first expression for K , namely, $K_0(x, t)$. Similarly

$$K(z_2, z_1) = u(z_1)v(z_2).$$

A comparison shows that

$$K(z_1, z_2) = K(z_2, z_1),$$

that is,

$$K(x, t) = K(t, x).$$

74. Equivalence Between the Boundary Problem and a Homogeneous Linear Integral Equation. *a) Hilbert's Fundamental Theorem.*—Under the assumptions (H_1) , (H_2) let us consider the non-homogeneous linear differential equation

$$L(u) + f = 0,$$

where f is supposed to be continuous on $[ab]$. We can then prove the following theorem:

Theorem IIIA.—*If F is of class C'' and $L(F) + f = 0$, $R_0(F) = 0$, $R_1(F) = 0$, then*

$$F(x) = \int_a^b K(x, t)f(t)dt.$$

Proof.—We have

$$L(F) = -f \text{ on } [ab], \text{ and}$$

$$L(K) = 0 \quad \text{on } [at], [tb] \text{ separately.}$$

Multiply the first by $-K$ and the last by F and add. We obtain

$$(8) \quad FL(K) - KL(F) = KF \text{ on } [at], [tb] \text{ separately.}$$

Apply Green's formula to the left-hand side of (8). We obtain

$$(9) \quad \frac{d}{dx} p(FK' - KF') = Kf.$$

Integrate (9) from a to t and from t to b . We have

$$\begin{aligned} \left[p(FK' - KF') \right]_a^{t=0} &= \int_a^{t=0} K(x, t)f(x)dx \\ \left[p(FK' - KF') \right]_{t+0}^b &= \int_{t+0}^b K(x, t)f(x)dx. \end{aligned}$$

Add the last two equations. We obtain

$$(10) \quad \begin{aligned} \left[p(FK' - KF') \right]_{t+0}^{t=0} - \left[p(FK' - KF') \right]_{x=a}^{x=b} \\ + \left[p(FK' - KF') \right]_{x=a}^{x=b} = \int_a^b K(x, t)f(x)dx, \end{aligned}$$

since $K(x, t)$ is continuous at $x = t$. Now p, F, F' , and K are continuous at $x = t$.

Hence $\left. -KF' \right|_{t+0}^{t=0} = 0.$

While

$$\begin{aligned} p(FK') \Big|_{t=0}^{t=0} &= p(t-0)F(t-0)K'(t-0) \\ &\quad - p(t+0)F(t+0)K'(t+0) \\ &= p(t)F(t) \left[K'(t-0) - K'(t+0) \right] \\ \text{which by } D) \\ &= p(t)F(t) \frac{1}{p(t)}. \end{aligned}$$

Now write the second term in (10) as a determinant

$$p(a) \begin{vmatrix} F(a) & F'(a) \\ K(a) & K'(a) \end{vmatrix}.$$

If $A \neq 0$, then

$$\begin{aligned} \left[p(FK' - F'K) \right]_{x=a}^{x=a} &= \frac{p(a)}{A} \begin{vmatrix} AF(a) + BF'(a) & F'(a) \\ AK(a) + BK'(a) & K'(a) \end{vmatrix} \\ &\equiv \frac{p(a)}{A} \begin{vmatrix} R_0(F) & F'(a) \\ R_0(K) & K'(a) \end{vmatrix} = 0, \end{aligned}$$

since $R_0(F) = 0, R_0(K) = 0$.

If $A = 0$, then $B \neq 0$ and, as before,

$$\left[p(FK' - KF') \right]_{x=a}^{x=a} = \begin{vmatrix} F(a) & R_0(F) \\ K(a) & R_0(K) \end{vmatrix} \frac{p(a)}{B} = 0.$$

Likewise $\left[p(FK' - KF') \right]_{x=b}^{x=b} = 0$.

Therefore, $F(t) = \int_a^b K(x, t)f(x)dx$.

Whence, interchanging x and t , on account of the symmetry of $K(x, t)$, we have

$$(11) \quad F(x) = \int_a^b K(x, t)f(t)dt.$$

Corollary.—If $F(x)$ is of class C'' and $R_0(F) = 0, R_1(F) = 0$, and $f = -L(F)$, then

$$(12) \quad F(x) = \int_a^b K(x, t) \left[-L(F) \right] dt.$$

Equation (11) is an integral equation of the *first kind*, with f as the function to be determined. Equation (12) shows us that (11) has a solution:

$$f = -L(F).$$

Theorem III B.—If $F(x) = \int_a^b K(x, t)f(t)dt$, then F is of class C'' , $L(F) + f = 0$, $R_0(F) = 0$, $R_1(F) = 0$.

Proof.—In order to form the derivatives of $F(x)$, we break the integral into two parts, inasmuch as K' is not continuous for $x = t$.

$$(13) \quad F(x) = \int_a^x K_1(x, t)f(t)dt + \int_x^b K_0(x, t)f(t)dt$$

$$F'(x) = \int_a^x K_1'(x, t)f(t)dt + \int_x^b K_0'(x, t)f(t)dt$$

$$+ K_1(x, x)f(x) - K_0(x, x)f(x).$$

But $K_1(x, x) - K_0(x, x) = 0$, since K is continuous. Therefore

$$(14) \quad F'(x) = \int_a^x K_1'(x, t)f(t)dt + \int_x^b K_0'(x, t)f(t)dt.$$

$$F''(x) = \int_a^x K_1''(x, t)f(t)dt + \int_x^b K_0''(x, t)f(t)dt$$

$$+ K_1'(x, x)f(x) - K_0'(x, x)f(x).$$

$$\text{But } K_1'(x, x) - K_0'(x, x) = \frac{-1}{p(x)}.$$

Therefore

$$(15) \quad F''(x) = \int_a^x K_1''(x, t)f(t)dt + \int_x^b K_0''(x, t)f(t)dt$$

$$- \frac{f(x)}{p(x)}.$$

Multiply (13), (14), (15) by $q(x)$, $p'(x)$, $p(x)$ respectively and add. We get

$$L(F) = \int_a^x L(K_1)f(t)dt + \int_x^b L(K_0)f(t)dt - f(x).$$

But $L(K_1) = 0$ and $L(K_0) = 0$, therefore

$$(16) \quad L(F) = -f(x).$$

Now form $R_0(F)$.

$$\begin{aligned} R_0(F) &= AF(a) + BF'(a) \\ &= \int_a^b R_0(K_0)f(t)dt = 0, \end{aligned}$$

since $R_0(K_0) = 0$. Similarly, we show that $R_1(F) = 0$.

That $F(x)$ is of class C'' follows from (13), (14), and (15), if we substitute for K_0 , K_1 their explicit expressions from (7).

We combine Theorems IIIA and IIIB into the following, which is Hilbert's third fundamental theorem:

Theorem III.—If $f(x)$ is continuous, then

$$F(x) = \int_a^b K(x, t)f(t)dt$$

implies and is implied by $F(x)$ is of class C'' ,

$$L(F) + f = 0, R_0(F) = 0, \text{ and } R_1(F) = 0.$$

b) *Equivalence of Boundary Problem and Integral Equation.*—If in Theorem III we put

$$f(x) = \lambda g(x)u(x), F(x) = u(x)$$

we obtain the theorem:

Theorem IV.—If $u(x)$ is continuous, then

$$(17) \quad u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt$$

implies and is implied by $u(x)$ is of class C'' ,

$$L(u) + \lambda gu = 0, R_0(u) = 0, \text{ and } R_1(u) = 0.$$

For λ is constant and by (H_1) g is continuous, therefore the hypothesis that $f(x)$ is continuous becomes $u(x)$ is continuous.

We notice that (17) is a homogeneous linear integral equation of the second kind with, in general, an unsymmetric kernel:

$$(18) \quad H(x, t) \equiv K(x, t)g(t).$$

We remark further that the condition that $u(x)$ be of class C'' carries with it the condition that $u(x)$ is of class C . Thus we drop the explicit statement that $u(x)$ be continuous and obtain from Theorem IV the two following theorems:

Theorem IVA.—*The conditions*

- a) $u(x)$ is of class C'' .
- b) $L(u) + \lambda gu = 0, R_0(u) = 0, R_1(u) = 0$

imply that

$$u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt.$$

Theorem IVB.—*The conditions*

- a) $u(x)$ is continuous.
- b) $u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt$

imply that

- a) $u(x)$ is of class C'' .
- b) $L(u) + \lambda gu = 0, R_0(u) = 0, R_1(u) = 0$.

Theorems IVA and IVB establish the equivalence of the boundary problem and of the integral equation. Hence,

1) If λ_0 is a characteristic constant of the boundary problem, then λ_0 is a characteristic constant of the integral equation and *vice versa*.

2) If $u(x)$ is a fundamental function for the boundary problem, belonging to λ_0 , then $u(x)$ is a fundamental function, belonging to λ_0 , for the integral equation and *vice versa*.

We remark that for this particular kernel (18), the fundamental functions are of class C'' .

75. Special Case $g(x) \equiv 1$.—For the special case $g(x) \equiv 1$, the equivalence between the boundary problem and the integral equation becomes:

$$(19) \quad \left\{ \begin{array}{l} L(u) + \lambda u = 0, R_0(u) = 0, R_1(u) = 0 \\ \text{implies and is implied by} \\ u(x) = \lambda \int_a^b K(x, t)u(t)dt. \end{array} \right.$$

The kernel of the integral equation in (19) is now symmetric and thus the results of both the Fredholm and the Hilbert-Schmidt theory apply.

Let us state some of the results of the Hilbert-Schmidt theory:

- 1) There exists at least one characteristic constant.
- 2) All of the characteristic constants are real.
- 3) The index $q =$ the multiplicity r .
- 4) There exists a complete normalized orthogonal system of fundamental functions $\{\psi_\nu(x)\}$ with corresponding characteristic constants $\{\lambda_\nu\}$, $\nu = 1, 2, \dots$.

- 5) If $\sum_\nu \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu}$ is uniformly convergent on R , then

$$(20) \quad \sum_\nu \frac{\psi_\nu(x)\psi_\nu(t)}{\lambda_\nu} = K(x, t).$$

- 6) If $u(x)$ is expressible in the form

$$u(x) = \lambda \int_a^b K(x, t)u(t)dt$$

and if $u(x)$ is continuous, then

$$(21) \quad u(x) = \sum_\nu C_\nu \psi_\nu(x),$$

where $C_\nu = (u\psi_\nu)$, and the series is absolutely and uniformly convergent on $[ab]$.

For the special case under consideration, each of these six statements can be made for the boundary problem on account of the equivalence established.

Further results follow, for the kernel under consideration, besides satisfying the conditions $K(x, t)$, is continuous, $\neq 0$, real, and symmetric of the Hilbert-Schmidt theory, satisfies now, in addition, the four conditions A) B) C) D) of §73.

Theorem V.—*There exists always an infinite number of characteristic constants $\{\lambda_\nu\}$ and a corresponding complete normalized orthogonal system of fundamental functions $\{\psi_\nu(x)\}$.*

Proof.—Suppose that there is a finite number of characteristic constants. Then there are a finite number of fundamental functions:

$$\begin{aligned}\psi_1(x), & \dots, \psi_m(x) \\ \lambda_1, & \dots, \lambda_m.\end{aligned}$$

For these the bilinear formula (20) holds. But $\psi_\nu(x)$ is of class C'' on R and, therefore, $K(x, t)$ would be of class C'' on R , which contradicts D), namely

$$K'(x, t - 0) - K'(x, t + 0) = \frac{1}{p(t)}.$$

Theorem VI.—*If $u(x)$ is of class C'' and $R_0(u) = 0$,*

$$R_1(u) = 0,$$

then

$$u(x) = \sum_\nu C_\nu \psi_\nu(x)$$

where $C_\nu = (u\psi_\nu)$, and the series is absolutely and uniformly convergent.

Proof.—From the corollary to Theorem IIIA

$$u(x) = \int_a^b K(x, t) \left[-L(u) \right] dx.$$

But $L(u)$ is continuous and, therefore, from (21)

$$u(x) = \sum_\nu C_\nu \psi_\nu(x).$$

Theorem VII.—*For every characteristic constant λ the index q is unity: $q = 1$.*

Proof.—Suppose $q > 2$, then at least three of the characteristic constants are equal, say, $\lambda_1 = \lambda_2 = \lambda_3$. Let the

corresponding fundamental functions be $\psi_1(x)$, $\psi_2(x)$, $\psi_3(x)$. We would then have three different independent solutions for $\lambda = \lambda_1$, of the same integral equation, that is, by the equivalence theorem three different solutions of the same differential equation of the second order. Hence one of them, say, ψ_3 , is linearly expressible in terms of the other two:

$$\psi_3(x) = C_1\psi_1 + C_2\psi_2.$$

This is a contradiction. Therefore $q > 2$.

Suppose $q = 2$. Then there would be two fundamental functions $\psi_1(x)$, $\psi_2(x)$ belonging to $\lambda = \lambda_1$. Now, by Green's formula,

$$(22) \quad p \left[\psi_1(x)\psi_2'(x) - \psi_2(x)\psi_1'(x) \right] = C \neq 0.$$

But

$$\begin{aligned} R_0(\psi_1) &= A\psi_1(a) + B\psi_1'(a) = 0 \\ R_0(\psi_2) &= A\psi_2(a) + B\psi_2'(a) = 0. \end{aligned}$$

These are two linear homogeneous equations for the determination of A , B , not both zero. Hence

$$\psi_1(a)\psi_2'(a) - \psi_2(a)\psi_1'(a) = 0,$$

which contradicts (22), therefore $q \neq 2$. Therefore we have $q = 1$.

We shall now discuss the more special boundary problem in which it follows from the boundary conditions that

$$(23) \quad \left[puu' \right]_a^b = 0.$$

For instance, if we have

$u(a) = 0$, $u(b) = 0$, or $u(a) = 0$, $u'(b) = 0$,
or $u'(a) = 0$, $u(b) = 0$, or $u'(a) = 0$, $u'(b) = 0$,
equation (23) is satisfied.

For this more special boundary problem we have the theorem:

Theorem VIII.—*If, as a consequence of the boundary conditions, we have*

$$\left[p u u' \right]_a^b = 0,$$

then there exists a $\left\{ \begin{array}{l} \text{smallest} \\ \text{largest} \end{array} \right\}$ characteristic constant if $\begin{cases} p(x) > 0 \\ p(x) < 0 \end{cases}$.

Proof.—The assumptions $p(x) > 0$, $p(x) < 0$ are in harmony with our previous assumption that $p(x) \neq 0$ on $[ab]$, for then $p(x)$ is always of the same sign. We have previously shown that there are an infinite number of characteristic constants, but, so far as we know, at present they may be infinite in number in both the positive and negative directions. Let $\psi_\nu(x)$ be a fundamental function belonging to λ_ν , then

$$(24) \quad L(\psi_\nu) + \lambda_\nu \psi_\nu = 0, \\ R_0(\psi_\nu) = 0, R_1(\psi_\nu) = 0,$$

ψ_ν is of class C'' , and $\psi_\nu \neq 0$. Multiply both members of (24) by ψ_ν and integrate. We obtain

$$(25) \quad \lambda_\nu = - \int_a^b \psi_\nu(x) L(\psi_\nu) dx$$

since

$$\int_a^b \psi_\nu^2(x) dx = 1.$$

In (25) put the explicit expression for $L(\psi_\nu)$. We have

$$\lambda_\nu = - \int_a^b \left[\psi_\nu \frac{d}{dx} (p \psi_\nu') + q \psi_\nu^2 \right] dx,$$

or $\lambda_\nu = - \int_a^b \psi_\nu \frac{d}{dx} (p \psi_\nu') dx + \int_a^b (-q) \psi_\nu^2 dx.$

Integrate the first integral by parts. We find

$$\lambda_r = - \left[p\psi_r \psi_r' \right]_a^b + \int_a^b p\psi_r'^2 dx + \int_a^b (-q)\psi_r^2 dx.$$

Now $-q$ is a continuous function on $[ab]$ and hence has a finite maximum M and a finite minimum m :

$$m \leq -q \leq M \text{ on } [ab].$$

Therefore

$$(26) \quad \begin{cases} \lambda_r \leq \int_a^b p\psi_r'^2 dx + M, \text{ and} \\ \lambda_r \geq \int_a^b p\psi_r'^2 dx + m. \end{cases}$$

If $p > 0$ on $[ab]$, then the integral in (26) is a positive number and therefore $\lambda_r > m$.

If $p < 0$ on $[ab]$, then the integrand in (26) is a negative number and therefore $\lambda_r < M$.

Combining these results with the fact that a finite interval can contain only a finite number of the λ_r 's, we obtain the result:

When $p(x) > 0$, the characteristic constants can be arranged in a sequence increasing towards plus infinity:

$$p(x) > 0; \lambda_1 < \lambda_2 < \lambda_3 < \dots < \lambda_r < \dots$$

and when $p(x) < 0$, they may be arranged in a sequence decreasing towards minus infinity:

$$p(x) < 0; \lambda_1 > \lambda_2 > \lambda_3 > \dots > \lambda_r > \dots$$

Corollary.—If $p(x) > 0$ and $q \leq 0$ on $[ab]$, then all of the characteristic constants are positive; if $p(x) < 0$ and $q \geq 0$ on $[ab]$, then all of the characteristic constants are negative.

For if $-q \geq 0$, then $m \geq 0$, so that, from (26), since $p > 0$, we get

$$m + (\text{a positive quantity}) \leq \lambda_p.$$

Therefore $\lambda_p > 0$.

If $-q \leq 0$, then $M \leq 0$, so that, from (26), since $p < 0$, we obtain

$$\lambda_p \leq M + (\text{a negative quantity not zero}).$$

Therefore $\lambda_p < 0$.

Definition.—A real symmetric kernel $K(x, t)$ is said to be closed if there exists no continuous function $h(x)$ other than $h(x) \equiv 0$, for which

$$\int_a^b K(x, t)h(t)dt = 0, \quad (x).$$

Theorem IX.—Green's function $K(x, t)$ for our boundary problem is always closed.

Proof.—We make use of Hilbert's third fundamental theorem, which states the equivalence of the boundary problem and the homogeneous linear integral equation:

If $f(x)$ is continuous, then

$$F(x) = \int_a^b K(x, t)f(t)dt$$

implies and is implied by $F(x)$ is of class C'' ,

$$L(F) + f = 0, \quad R_0(F) = 0, \quad \text{and} \quad R_1(F) = 0.$$

We apply this theorem for $F(x) \equiv 0$, whence if $f(x)$ is continuous and

$$\int_a^b K(x, t)f(t)dt = 0, \quad (x),$$

then $f = -L(0) \equiv 0$.

76. Miscellaneous Remarks. *a) The General Case $g(x) \not\equiv 1$.*—In the previous article we considered the special case $g(x) \equiv 1$. We now consider the problem for $g(x) \not\equiv 1$. Hilbert's third fundamental theorem now reads:

If $u(x)$ is continuous, then

$$u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt$$

implies and is implied by $u(x)$ is of class C'' ,

$$L(u) + \lambda gu = 0, R_0(u) = 0, R_1(u) = 0.$$

There are two cases to be considered:

Case I.— $g(x) \neq 0$ on $[ab]$. This case can be reduced to the case $g(x) \equiv 1$. Since $g(x) \neq 0$ on $[ab]$, we have either $g(x) > 0$ or $g(x) < 0$ on $[ab]$. Consider the first case.

Multiply both sides of

$$(27) \quad u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt$$

by $\sqrt{g(x)}$, and put

$$\begin{aligned} \bar{u}(x) &= \sqrt{g(x)} \cdot u(x). \\ \bar{K}(x, t) &= K(x, t)\sqrt{g(x)}\sqrt{g(t)}. \end{aligned}$$

We then obtain

$$(28) \quad \bar{u}(x) = \lambda \int_a^b \bar{K}(x, t)\bar{u}(t)dt.$$

This is the reduction desired, for $\bar{K}(x, t)$ is symmetric.

If $g(x) < 0$ on $[ab]$, multiply both sides of (27) by $\sqrt{-g(x)}$, and put

$$\begin{aligned} \bar{u}(x) &= \sqrt{-g(x)} \cdot u(x). \\ \bar{K}(x, t) &= K(x, t)\sqrt{-g(x)}\sqrt{-g(t)}. \end{aligned}$$

We obtain

$$\bar{u}(x) = -\lambda \int_a^b \bar{K}(x, t)\bar{u}(t)dt,$$

which is again the reduction desired.

Case II.— $g(x)$ vanishes at some point of $[ab]$. This case has been treated by Hilbert [“Gott. Nach.” 5, p. 462, 1906] for functions $K(x, t)$ which are *definite*, by means of the theory of quadratic forms with infinitely many variables. J. Marty [*Comptes Rendus*, vol. 150, p. 515; *Ibid.*, p. 605, 1910] has reached the same results without the use of quadratic forms of infinitely many variables.

Definition.—A real symmetric¹ kernel $K(x, t)$ is said to be definite if no continuous function exists, other than $h(x) \equiv 0$, for which

$$\int_a^b \int_a^b K(x, t)h(x)h(t)dxdt = 0.$$

The name “definite” has been used on account of the analogy with a *definite* quadratic form $\sum K_{ij}y_iy_j$ which vanishes only when all of the y ’s vanish.

Integral equations of the form

$$(29) \quad u(x) = f(x) + \lambda \int_a^b K(x, t)g(t)u(t)dt$$

with $K(x, t)$ *definite* have been called by Hilbert *polar* integral equations or integral equations of the third kind. If we multiply the equation (29) by $g(x)$ and put

$$\begin{aligned}\bar{f}(x) &= f(x)g(x) \\ \bar{K}(x, t) &= K(x, t)g(x)g(t),\end{aligned}$$

we obtain the equation in the form in which Hilbert considered it:

$$g(x)u(x) = \bar{f}(x) + \lambda \int_a^b \bar{K}(x, t)u(t)dt.$$

¹ For a discussion of this and other special kernels consult LALESCO, T., “Théorie Des Équations Intégrales,” pp. 64ff.

Definition.—A kernel $H(x, t)$ is said to be symmetrizable if there exists a definite symmetric kernel $G(x, t)$ such that either

$$K_1(x, t) = \int_a^b G(x, s)H(s, t)ds$$

or

$$K_2(x, t) = \int_a^b H(x, s)G(s, t)ds$$

is symmetrical.

In this instance the kernel

$$H(x, t) = K(x, t)g(t)$$

is symmetrizable if $K(x, t)$ is definite, for

$$K_2(x, t) = \int_a^b K(x, s)g(s)K(s, t)ds$$

is symmetric, since

$$\begin{aligned} K_2(t, x) &= \int_a^b K(t, s)g(s)K(s, x)ds \\ &= \int_a^b K(s, t)g(s)K(x, s)ds \\ &= K_2(x, t). \end{aligned}$$

For symmetrizable kernels Marty has shown that

- 1) There exists at least one characteristic constant.
- 2) All of the characteristic constants are real.

b) *Non-homogeneous Boundary Problems.*—We next consider the non-homogeneous boundary problem:

$$L(u) + \lambda gu + r = 0, \quad R_0(u) = 0, \quad R_1(u) = 0,$$

where r is a given function of x , continuous on $[ab]$, and

$R_0(u), R_1(u)$ are defined by (4).

We are going to show that the boundary problem is equivalent to a non-homogeneous integral equation. To do

this we again make use of Hilbert's third fundamental theorem for

$$f = \lambda gu + r, F = u.$$

Since λ, g, r are continuous by hypothesis, this theorem now reads: if $u(x)$ is continuous, then the combined statement u is of class C'' , $L(u) + \lambda gu + r = 0$, $R_0(u) = 0$, $R_1(u) = 0$ is equivalent to the statement that

$$\begin{aligned} u(x) &= \int_a^b K(x, t) \left[\lambda g(t)u(t) + r(t) \right] dt \\ &= \lambda \int_a^b K(x, t)g(t)u(t)dt + \int_a^b K(x, t)r(t)dt. \end{aligned}$$

Hence we have the theorem:

Theorem X.—The non-homogeneous boundary problem:

$$L(u) + \lambda gu + r = 0, R_0(u) = 0, R_1(u) = 0$$

with the further condition that $u(x)$ is of class C'' , is equivalent to the non-homogeneous integral equation

$$u(x) = f(x) + \lambda \int_a^b K(x, t)g(t)u(t)dt$$

where $u(x)$ is continuous and $f(x) = \int_a^b K(x, t)r(t)dt$.

c) *The Exceptional Case $\lambda = 0$ Is a Characteristic Constant.*—All of the preceding developments have been made under the hypothesis (H_2) that $\lambda = 0$ is not a characteristic constant of the boundary problem. This hypothesis, however, is not satisfied in certain problems of mathematical physics. For example, the following boundary problem in the theory of heat:

$$\frac{d^2u}{dx^2} + \lambda u = 0, u'(a) = 0, u'(b) = 0$$

has the non-trivial solution $u = \text{constant}$ for $\lambda = 0$.

The exceptional case in which the hypothesis (H_2) is not satisfied can be treated, according to Hilbert,¹ by introducing a modified Green's function.

¹ See HILBERT, "Gott. Nach.", p. 213, 1904.

In most cases, however, the following simple artifice, due to Kneser will be sufficient.

Let us replace the assumption (H_2) by the much milder assumption (H_2') :

(H_2') *There exists at least one value c of λ which is not a characteristic constant for the boundary problem.*

Let us write the differential equation

$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + (q + \lambda g)u = 0$$

in the form

$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + \left[q + cg + (\lambda - c)g \right] u = 0$$

or
$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + (\bar{q} + \bar{\lambda}g)u = 0,$$

where $\bar{q} = q + cg, \bar{\lambda} = \lambda - c.$

Then $\bar{\lambda} = 0$ is certainly not a characteristic constant of the boundary problem

$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + (\bar{q} + \bar{\lambda}g)u = 0, R_0(u) = 0, R_1(u) = 0,$$

since, by hypothesis (H_2') , the boundary problem

$$\frac{d}{dx} \left(p \frac{du}{dx} \right) + qu + cgu = 0, R_0(u) = 0, R_1(u) = 0$$

has no solution other than $u \equiv 0$.

II. APPLICATIONS TO SOME PROBLEMS OF THE CALCULUS OF VARIATIONS

77. Some Auxiliary Theorems of the Calculus of Variations. *a) Formulation of the Simplest Type of Problem.*—For the simplest type of problems of the calculus of variations we have given

- 1) Two points $P_0(x_0, y_0), P_1(x_1, y_1).$
- 2) A function $F(x, y, y')$ of three independent variables.

Required: to find among all curves

$$(30) \quad y = y(x)$$

joining P_0 and P_1 that one which furnishes for the definite integral

$$J(y) = \int_{x_0}^{x_1} F \left[x, y(x), y'(x) \right] dx, \left[y'(x) = \frac{dy}{dx} \right]$$

the smallest value.

Concerning the admissible curves (30) we make the assumption that they satisfy the following conditions:

A) $y(x)$ of class C'' .

(31)

B) $y(x_0) = y_0; y(x_1) = y_1$.

We assume that the function F is of class C''' for all systems of values $x, y(x), y'(x)$ furnished by all of the admissible curves.

b) *Euler's Differential Equation*.—Suppose we have found the minimizing curve C_0

$$C_0: \quad y = f(x).$$

We replace it by a neighboring admissible curve of the special form

$$C: \quad y = f(x) + \epsilon\eta(x),$$

where ϵ is a small constant and $\eta(x)$ a function of x satisfying the conditions:

A') $\eta(x)$ is of class C'' .

(32)

B') $\eta(x_0) = 0, \eta(x_1) = 0$.

Since C_0 minimizes the integral, we have

$$\int_{x_0}^{x_1} F \left[x, f + \epsilon\eta, f' + \epsilon\eta' \right] dx \geq \int_{x_0}^{x_1} F \left[x, f, f' \right] dx,$$

which we shall write in the form

$$I(\epsilon) \geq I(0).$$

Considered as a function of ϵ , $I(\epsilon)$ has, therefore, a minimum for $\epsilon = 0$ and hence

$$I'(0) = 0, I''(0) \geq 0.$$

These are *necessary* conditions for a minimum. It is customary to write

$$\delta I \equiv \epsilon I'(0), \quad \delta^2 I \equiv \epsilon^2 I''(0), \quad (\eta)$$

and to call δI , $\delta^2 I$, the first and second variation respectively.

We first consider the condition $I'(0) = 0$. By definition

$$I(\epsilon) = \int_{x_0}^{x_1} F \left[x, y + \epsilon\eta, y' + \epsilon\eta' \right] dx,$$

whence, by the rules for differentiating a definite integral with respect to a parameter,

$$(33) \quad I'(\epsilon) = \int_{x_0}^{x_1} \left[\bar{F}_{y\eta} + \bar{F}_{y'\eta'} \right] dx, \quad (\eta)$$

where the dash indicates that we use the arguments, $x, f(x) + \epsilon\eta(x), f'(x) + \epsilon\eta'(x)$. Therefore

$$I'(0) = \int_{x_0}^{x_1} \left(F_{y\eta} + F_{y'\eta'} \right) dx, \quad (\eta)$$

the arguments now being $x, f(x), f'(x)$.¹

An integration by parts gives

$$\begin{aligned} I'(0) &= \left[\eta F_{y'} \right]_{x_0}^{x_1} + \int_{x_0}^{x_1} \eta \left(F_y - \frac{d}{dx} F_{y'} \right) dx, \quad (\eta) \\ &= \int_{x_0}^{x_1} \eta \left(F_y - \frac{d}{dx} F_{y'} \right) dx, \quad (\eta) \end{aligned}$$

on account of (32).

But $I'(0) = 0$, and, hence, by the fundamental theorem of the calculus of variations,²

$$(34) \quad F_y - \frac{d}{dx} F_{y'} = 0.$$

¹ Consult BOLZA, "Lectures on the Calculus of Variations," pp. 16ff., University of Chicago Press, 1904.

² See BOLZA, "Lectures on the Calculus of Variations," §5.

This is a differential equation of the second order for the determination of $y = f(x)$. It is the first necessary condition and is known as Euler's equation. Its solution involves two arbitrary constants which have to be determined by the two conditions (31).

From (33) we get

$$I''(\epsilon) = \int_{x_0}^{x_1} \left(\bar{F}_{yy}\eta^2 + 2\bar{F}_{yy'}\eta\eta' + \bar{F}_{y'y'}\eta'^2 \right) dx,$$

whence

$$I''(0) = \int_{x_0}^{x_1} \left(F_{yy}\eta^2 + 2F_{yy'}\eta\eta' + F_{y'y'}\eta'^2 \right) dx, (\eta).$$

c) *Euler's Rule for Isoperimetric Problems.*—For isoperimetric problems the admissible curves are subject to a third condition C in addition to A) and B):

$$C: \quad \int_{x_0}^{x_1} G[x, y(x), y'(x)] = l, \text{ a constant.}$$

The problem is, then, to determine, among all curves

$$y = y(x)$$

satisfying the conditions A) B) C), that one which will furnish the smallest value for the integral

$$J(y) = \int_{x_0}^{x_1} F(x, y, y')dx.$$

The conditions on G are the same as those on F , viz., F and G are continuous and possess continuous partial derivatives of the first, second, and third order in the region under consideration. Put

$$H = F + \lambda G,$$

λ an arbitrary constant. Then, by Euler's rule, the first necessary condition for a minimum is the same as if it were required to minimize the integral

$$\int_{x_0}^{x_1} H(x, y, y')dx$$

with respect to the totality of curves satisfying the conditions *A*) and *B*), that is,

$$H_y - \frac{d}{dx}H_{y'} = 0,$$

or, more explicitly,

$$F_y - \lambda G_y - \frac{d}{dx}(F_{y'} - \lambda G_{y'}) = 0.$$

The solution of this equation involves three arbitrary constants (α , β , λ) which are to be determined by the three conditions *B*) and *C*).

78. Dirichlet's Problem.—We now propose to minimize the integral

$$(35) \quad D(u) = \int_a^b \left[p \left(\frac{du}{dx} \right)^2 - qu^2 \right] dx$$

with respect to the totality M of curves satisfying the conditions

$$(36) \quad \begin{cases} u \text{ is of class } C'', u(a) = 0, u(b) = 0, \\ \int_a^b u^2(x) dx = 1, \end{cases}$$

with the further hypotheses

$$\left. \begin{array}{l} p > 0, p \text{ of class } C' \\ q \text{ continuous} \end{array} \right\} \text{on } [ab].$$

Hilbert, to whom the following developments are due, calls this problem *Dirichlet's problem*. The problem is an isoperimetrical problem of the type considered in §77, with

$$F = pu'^2 - qu^2, G = u^2.$$

Hence

$$H(x, u, u') = pu'^2 - (q + \lambda)u^2$$

and, therefore, Euler's differential equation for H is

$$H_u - \frac{d}{dx}H_{u'} = -2(q + \lambda)u - \frac{d}{dx}(2pu') = 0.$$

Whence

$$\frac{d}{dx}(pu') + (q + \lambda)u = 0,$$

which may be written

$$(37) \quad L(u) + \lambda u = 0.$$

Every solution of this problem must satisfy this equation and the conditions (36). Now (36) and (37) constitute a boundary problem of the type previously discussed in which

$$g(x) \equiv 1, \left[puu' \right]_a^b = 0,$$

hence the Theorems I to IX hold.

We can, therefore, state at once that this problem (36), (37) has no solutions except when λ is a characteristic constant. We know that

- 1) The characteristic constants are real.
- 2) The characteristic constants are infinite in number.
- 3) For each characteristic constant the index $q = 1$.

Therefore the λ 's are distinct, and, since $p > 0$, we have, by Theorem VIII,

$$(38) \quad \lambda_1 < \lambda_2 < \lambda_3 < \dots,$$

with a corresponding complete set,

$$\psi_1(x), \psi_2(x), \psi_3(x), \dots$$

of normalized orthogonal fundamental functions of class C'' . We have, then, for $n = 1, 2, \dots$

$$(39) \quad \begin{aligned} L(\psi_n) + \lambda_n \psi_n &= 0, \\ \psi_n(a) &= 0, \psi_n(b) = 0, \\ \int_a^b \psi_n^2(x) dx &= 1. \end{aligned}$$

If, then, Dirichlet's problem has a solution, it must be of the form

$$u = C\psi_n(x), \lambda = \lambda_n.$$

But from (36) we obtain

$$C^2 \int_a^b \psi_n^2(x) dx = 1.$$

Comparing with (39) we find that $C = \pm 1$. The only possible solutions, therefore, are

$$u = \pm \psi_n(x), \lambda = \lambda_n.$$

But in the proof of Theorem VIII we showed that

$$\lambda_n = \int_a^b (p\psi_n'^2 - q\psi_n^2) dx.$$

Therefore

$$\lambda_n = D(\pm \psi_n), \text{ by (35).}$$

Hence, it follows from (38) that

$$D(\pm \psi_1) < D(\pm \psi_n), (n = 2, 3, \dots).$$

Hence we infer that: *If there exists at all a function which minimizes $D(u)$ with respect to the totality M of all admissible curves, it must be the function*

$$u = \pm \psi_1(x)$$

and λ must have the value λ_1 .

b) *Sufficiency Proof.* a) *Transformation of $D(u)$.*—By an integration by parts we obtain

$$\int_a^b (pu')u' dx = upu' \Big|_a^b - \int_a^b u \frac{d}{dx} \left(p \frac{du}{dx} \right) dx.$$

But $upu' \Big|_a^b = 0$, since $u(a) = 0, u(b) = 0$.

Therefore

$$\begin{aligned} D(u) &= \int_a^b \left[p \left(\frac{du}{dx} \right)^2 - qu^2 \right] dx \\ &= - \int_a^b u \left[\frac{d}{dx} \left(p \frac{du}{dx} \right) + qu \right] dx. \end{aligned}$$

Therefore

$$D(u) = - \int_a^b u L(u) dx.$$

Let us put

$$-L(u) = \omega(x).$$

Then ω is continuous and we have

$$D(u) = + \int_a^b u(x)\omega(x)dx.$$

We now apply Hilbert's third fundamental theorem, whence from

$$u \text{ is of class } C'', L(u) + \omega = 0, u(a) = 0, u(b) = 0$$

it follows that

$$u(x) = \int_a^b K(x, t)\omega(t)dt.$$

Multiply both members of this equation by $\omega(x)$ and integrate. We obtain

$$(40) \quad D(u) = \int_a^b \int_a^b K(x, t)\omega(t)\omega(x)dtdx.$$

β) Applications of the Expansion Theorem.—From Corollary II to Theorem XVII of §68, we have

$$u(x) = \int_a^b K(x, t)\omega(t)dt = \sum_{\nu=1}^{\infty} \frac{(\omega\psi_{\nu})}{\lambda_{\nu}} \psi_{\nu}(x),$$

the series being absolutely and uniformly convergent. Let us put

$$C_{\nu} = (\omega\psi_{\nu}), \text{ constant,}$$

then

$$(41) \quad u(x) = \sum_{\nu=1}^{\infty} \frac{C_{\nu}}{\lambda_{\nu}} \psi_{\nu}(x).$$

Form

$$u^2(x) = \sum_{\mu, \nu} \frac{C_{\mu} C_{\nu}}{\lambda_{\mu} \lambda_{\nu}} \psi_{\mu}(x) \psi_{\nu}(x).$$

This series is also absolutely and uniformly convergent. We obtain, therefore,

$$\int_a^b u^2(x)dx = \sum_{\mu, \nu} \frac{C_\mu C_\nu}{\lambda_\mu \lambda_\nu} \int_a^b \psi_\mu(x) \psi_\nu(x) dx = 1.$$

But $\int_a^b \psi_\mu(x) \psi_\nu(x) dx = \begin{cases} 1, & \mu = \nu \\ 0, & \mu \neq \nu. \end{cases}$

Therefore

$$(42) \quad \sum_\nu \frac{C_\nu^2}{\lambda_\nu^2} = 1.$$

We now apply Corollary III to Theorem XVII of §68, to the double integral in (40). We obtain

$$D(u) = \sum_{\nu=1}^{\infty} \frac{(\omega \psi_\nu)^2}{\lambda_\nu} = \sum_{\nu=1}^{\infty} \frac{C_\nu^2}{\lambda_\nu}.$$

γ) Computation of $D(u) - D(\pm \psi_1)$.—Let us now compute the difference

$$D(u) - D(\pm \psi_1) = D(u) - \lambda_1.$$

On account of (42)

$$\lambda_1 = \sum_{\nu=1}^{\infty} \frac{C_\nu^2 \lambda_1}{\lambda_\nu^2}.$$

Therefore

$$\begin{aligned} D(u) - \lambda_1 &= \sum_\nu \frac{C_\nu^2}{\lambda_\nu} - \sum_\nu \frac{C_\nu^2 \lambda_1}{\lambda_\nu^2} \\ &= \sum_{\nu=2}^{\infty} \frac{C_\nu^2}{\lambda_\nu^2} (\lambda_\nu - \lambda_1) \geq 0, \end{aligned}$$

since $\frac{C_\nu^2}{\lambda_\nu^2} \geq 0$, and $\lambda_\nu - \lambda_1 > 0$ for $\nu \geq 2$.

Therefore we have

$$D(u) - D(\pm \psi_1) \geq 0.$$

The equality holds only if

$$C_\nu = 0 \text{ for } \nu = 2, 3, \dots, \infty.$$

But in this event from (42) we obtain $C_1 = \pm \lambda_1$, whence from (41), $u(x) = \pm \psi_1(x)$. Therefore, whenever $u(x) \neq \psi_1(x)$, it results that

$$D(u) - D(\pm \psi_1(x)) \geq 0.$$

Thus we have proved the following theorem:

Theorem XI.— $u = \pm \psi_1(x)$ furnishes for

$$D(u) = \int_a^b \left[p \left(\frac{du}{dx} \right)^2 - qu^2 \right] dx$$

a smaller value (viz., λ_1) than any other function satisfying the conditions

$$u \text{ is of class } C'', u(a) = 0, u(b) = 0, \int_a^b u^2(x) dx = 1.$$

c) *Dropping the Assumption (H_2).*—The results for the Dirichlet problem seem to presuppose (H_2), that is, that $\lambda = 0$ is not a characteristic constant of the boundary problem

$$L(u) + \lambda u = 0, u(a) = 0, u(b) = 0,$$

but they are independent of this assumption.

Proof.—α) We notice first that the assumption, $\lambda = 0$ is not a characteristic constant, was necessary for the construction of a Green's function. But the Green's function did not occur in the proof of Theorem VIII. In this theorem, we made the special assumption that

$$R_0(u) = 0, R_1(u) = 0, \text{ imply } \left[puu' \right]_a^b = 0.$$

Let then λ_0 be any characteristic constant of the boundary problem:

$$(43) \quad L(u) + \lambda u = 0, R_0(u) = 0, R_1(u) = 0$$

and $\psi_0(x)$ a corresponding normalized fundamental function of the boundary problem:

ψ_0 of class C'' , $L(\psi_0) + \lambda_0\psi_0 = 0$, $R_0(\psi_0) = 0$, $R_1(\psi_0) = 0$,

$$\int_a^b \psi_0^2(x)dx = 1.$$

Hence we derived, since $\left[p\psi_0\psi_0' \right]_a^b = 0$, without using the Green's function, that is, independently of (H_2) the equality

$$\lambda_0 = \int_a^b (p\psi_0'^2 - q\psi_0^2)dx \equiv D(\psi_0)$$

Hence followed from the corollary to Theorem VIII that $p > 0$, $q \leq 0$, imply $\lambda_0 > 0$;

that is, all of the characteristic constants of the boundary problem are greater than zero. Thus the hypothesis (H_2) is satisfied and the Theorems V to VIII certainly hold.

$\beta)$ Suppose now that the condition $q \leq 0$ is not satisfied.

Denote, as before, by m the minimum of $-q$ on $[ab]$, that is,

$$q + m \leq 0.$$

Then the differential equation

$$L(u) + \lambda u \equiv \frac{d}{dx} \left(p \frac{du}{dx} \right) + (q + \lambda)u = 0$$

becomes

$$\bar{L}(u) + \bar{\lambda}u \equiv \frac{d}{dx} \left(p \frac{du}{dx} \right) + \bar{q}u + \lambda u = 0$$

where

$$q + m = \bar{q} \text{ and } \lambda - m = \bar{\lambda}.$$

Since $p > 0$, $\bar{q} \leq 0$, the Theorems V to VIII hold for the boundary problem

$$(44) \quad \bar{L}(u) + \bar{\lambda}u = 0, R_0(u) = 0, R_1(u) = 0.$$

In particular, it has a single infinitude of real, distinct characteristic constants:

$$0 < \bar{\lambda}_1 < \bar{\lambda}_2 < \bar{\lambda}_3 < \dots$$

with corresponding normalized fundamental functions

$$\psi_1(x), \psi_2(x), \psi_3(x), \dots$$

But since

$$\bar{L}(u) = L(u) + mu$$

it follows that, if λ_0 is a characteristic constant and $\psi_0(x)$ a fundamental function for (43), then $\bar{\lambda}_0 = \lambda_0 - m$ is a characteristic constant and $\psi_0(x)$ the corresponding fundamental function for (44), and *vice versa*.

Hence also the boundary problem (43) has an infinitude of real, distinct characteristic constants forming an increasing sequence:

$$\begin{aligned}\lambda_1 &= \bar{\lambda}_1 + m, \lambda_2 = \bar{\lambda}_2 + m, \dots, \\ \lambda_1 &< \lambda_2 < \lambda_3 < \dots,\end{aligned}$$

with the corresponding normalized orthogonal fundamental functions $\psi_1(x), \psi_2(x), \dots$

γ) Consider now the problem

$$D(u) = \text{minimum on } (M).$$

This problem is evidently equivalent to

$$\bar{D}(u) = D(u) - m = \text{minimum on } (M), \text{ i.e.,}$$

$$\bar{D}(u) = \int_a^b \left[p \left(\frac{du}{dx} \right)^2 - (q + m)u^2 \right] dx = \text{minimum on } (M).$$

Since $q + m \leq 0$, our former results hold and

$$u = \pm \psi_1(x)$$

furnishes the minimum for $\bar{D}(u)$, that is,

$$\bar{D} \left[\pm \psi_1(x) \right] = \bar{\lambda}_1.$$

On account of the equivalence of the two problems, the same function $u = \pm \psi_1(x)$ furnishes the minimum for $D(u)$ and

$$D(\pm \psi_1) = \bar{D}(\pm \psi_1) + m = \bar{\lambda}_1 + m = \lambda_1.$$

But by $\beta)$ $\lambda_1 = \bar{\lambda}_1 + m$ is the smallest characteristic constant for the boundary problem

$$L(u) + \lambda u = 0, \quad u(a) = 0, \quad u(b) = 0.$$

This shows that Theorem XI is true also when the assumption (H_2) is not satisfied.

79. Applications to the Second Variation.—Hilbert has made an application of the preceding result to the discussion of the second variation for the simplest type of variation problem considered in §77.

a) *Reduction of the Problem.*—The problem is to find the condition under which

$$(45) \quad \int_{x_0}^{x_1} \left(F_{yy} \eta^2 + 2F_{yy'} \eta \eta' + F_{y'y'} \eta'^2 \right) dx \geq 0$$

for all functions $\eta(x)$ satisfying the conditions

A') $\eta(x)$ is of class C'' .

B') $\eta(x_0) = 0, \eta(x_1) = 0$.

The arguments of $F_{yy}, F_{yy'}, F_{y'y'}$ are $x, y = f(x), y' = f'(x)$ where $f(x)$ is a solution of Euler's differential equation (34) satisfying the initial conditions. We adopt the notation

$$F_{yy} \left[x, f(x), f'(x) \right] = P(x), \quad F_{yy'} \left[x, f(x), f'(x) \right] = Q(x), \\ F_{y'y'} \left[x, f(x), f'(x) \right] = R(x).$$

Then (45) becomes

$$(46) \quad \int_{x_0}^{x_1} \left[P\eta^2 + 2Q\eta\eta' + R\eta'^2 \right] dx \geq 0, (\eta).$$

It is easily shown¹ that a first necessary condition for the inequality (46) is that $R \geq 0$ on $[ab]$, which is Legendre's condition.

¹ Compare BOLZA, "Lectures on the Calculus of Variations," §11.

We suppose this condition to be satisfied in the stronger form $R > 0$ on $[ab]$.

We now transform the integral by integrating the second term by parts

$$\begin{aligned}\int_{x_0}^{x_1} 2Q\eta\eta' dx &= \int_{x_0}^{x_1} Q \frac{d}{dx} \eta^2 dx \\ &= \eta^2 \Big|_{x_0}^{x_1} - \int_{x_0}^{x_1} \eta^2 Q' dx \\ &= - \int_{x_0}^{x_1} \eta^2 Q' dx.\end{aligned}$$

Therefore (45) becomes

$$\int_x^{x_1} \left[(P - Q')\eta^2 + R\eta'^2 \right] dx \geq 0, (\eta).$$

If for $\eta, x_0, x_1, R, P - Q'$
we put $u, a, b, p, -q,$

then the above inequality becomes

$$D(u) \geq 0$$

for the totality N of all functions u satisfying the conditions

$$u \text{ is of class } C'', u(a) = 0, u(b) = 0.$$

b) *Connection with Dirichlet's Problem.*—We now show the equivalence of the two statements

$$(47) \quad D(u) \geq 0, \text{ for all curves of class } N, \text{ and}$$

$$(48) \quad D(u) \geq 0, \text{ for all curves of class } M.$$

A) The class of curves M is contained within the class N and hence

$$D(u) \geq 0, (N) \text{ implies } D(u) \geq 0, (M).$$

B) Suppose $u \not\equiv 0$ belongs to the class (N) . Construct $u_1 = \rho u$ such that

$$\int_a^b u_1^2 dx = 1.$$

Then u_1 belongs to M , for u_1 is of class C'' and $u_1(a) = 0$, $u_1(b) = 0$, since $u(a) = 0$, $u(b) = 0$ and by construction

$$\int_a^b u_1^2 dx = 1.$$

Therefore $D(u_1) \geq 0$.

But $D(u_1) = D(\rho u) = \rho^2 D(u)$.

Therefore $D(u) \geq 0$.

Hence, if u belongs to the class (N) and $u \not\equiv 0$, then $D(u) \geq 0$, while if $u \equiv 0$ also $D(u) = 0$. Therefore, if u belongs to (N) , then $D(u) \geq 0$.

The equivalence between the two inequalities (47) and (48) being established, we can now apply the results of §78:

The smallest value which $D(u)$ can take in M is λ_1 and this value is furnished by $u = \pm \psi_1(x)$ and by no other function of M .

Hence if

- 1) $\lambda_1 > 0$, then $D(u) > 0$ (M).
- 2) $\lambda_1 = 0$, then $D(u) > 0$ (M).

except when $u = \pm \psi_1(x)$, in which case $D(u) = 0$.

- 3) $\lambda_1 < 0$, then $D(u)$ can be made negative in M .

Hence, $\lambda_1 \geq 0$ is the necessary and sufficient condition that $D(u) \geq 0$ for all curves of the class M and therefore also the necessary and sufficient condition that $D(u) \geq 0$ for all curves of the class N .

Returning now to the notation of the calculus of variations, we have the theorem:

Theorem XII.—Suppose $R > 0$ and let λ_1 denote the smallest characteristic constant of the boundary problem

$$\frac{d}{dx} \left(R \frac{du}{dx} \right) - (P - Q')u = 0, \quad u(x_0) = 0, \quad u(x_1) = 0,$$

then $\lambda_1 \geq 0$ is the necessary and sufficient condition that

$$\delta^2 I \geq 0, \quad (\eta).$$

80. Connection with Jacobi's Condition. *a) Sturm's Oscillation Theorem.*—It is *a priori* clear that the condition $\lambda_1 \geq 0$ must be equivalent to Jacobi's condition.¹ The connection between the two can be established by means of Sturm's oscillation theorem.

Since the differential equation

$$L(u) + \lambda u = 0$$

has no singularities on $[ab]$, there exists one and only one solution for which

$$u(a) = 0, u'(a) = 1.$$

Let us call this $u = V(x, \lambda)$, so that

$$V(a, \lambda) = 0, V'(a, \lambda) = 1.$$

Any other solution $u(x)$, for which $u(a) = 0$, is then of the form

$$u = C \cdot V(x, \lambda).$$

Now the boundary problem

$$L(u) + \lambda u = 0, u(a) = 0, u(b) = 0$$

has for $\lambda = \lambda_1$ a solution $u = \psi_1(x)$, for which

$$\psi_1(a) = 0, \psi_1(b) = 0.$$

Hence

$$\psi_1(x) = C \cdot V(x, \lambda_1)$$

and, therefore,

$$V(a, \lambda_1) = 0, V(b, \lambda_1) = 0.$$

Let us designate by $\xi(\lambda)$ the root of $V(x, \lambda)$ next greater than a . Then Sturm's oscillation theorem states² that

¹ Cf. BOLZA, "Lectures," §16.

² See BÖCHER, *Bull. Am. Math. Soc.*, 4, 1898.

- 1) as λ increases, $\xi(\lambda)$ decreases
 (49) as λ decreases, $\xi(\lambda)$ increases
 2) $V(x, \lambda_1) \neq 0$ between a and b and, therefore,
 $\xi(\lambda_1) = b$.

b) The Conjugate Point.—Consider the solution

$$u = V(x, 0)$$

and put

$$a' = \xi(0).$$

Case I.— $\lambda_1 > 0$. Then, by (49), as λ decreases from λ_1 to 0, $\xi(\lambda)$ increases from $\xi(\lambda_1) = b$ to $\xi(0) = a'$ and, therefore, $a' > b$.




FIG. 19.

Case II.— $\lambda_1 = 0$. Then $a' = b$.

Case III.— $\lambda_1 < 0$. Then, by (49), as λ increases from λ_1 to 0, $\xi(\lambda)$ decreases from $\xi(\lambda_1) = b$ to $\xi(0) = a'$ and, therefore, $a' < b$.

Let us now return to the notation of the calculus of variations, that is

from $a, b, \frac{d}{dx} \left(p \frac{du}{dx} \right) + qu$

to $x_0, x_1, \frac{d}{dx} \left(R \frac{du}{dx} \right) - (P - Q')u$.

Then $u = V(x, 0)$ is defined as that integral of Jacobi's differential equation

$$\frac{d}{dx} \left(R \frac{du}{dx} \right) - (P - Q')u = 0$$

which satisfies the initial conditions.

$$u(x_0) = 0, u'(x_0) = 1.$$

Hence $V(x, 0)$ is, up to a constant factor, identical with the function denoted in the calculus of variations by $\Delta(x, x_0)$ and, therefore, a' is identical with x_0' , the abscissa of the point conjugate to x_0 .

This establishes the equivalence between Hilbert's condition

$$\lambda_1 \geq 0$$

and Jacobi's condition


$$x_1 \leq x_0'$$

for a non-negative sign of $\delta^2 I$.

III. VIBRATION PROBLEMS

81. Vibrating String. *a) Reduction to Boundary Problem.* For the homogeneous string we had

$$(50) \quad \frac{\partial^2 \eta}{\partial t^2} = C^2 \frac{\partial^2 \eta}{\partial x^2}.$$



$$(51) \quad \eta(0, t) = 0, \eta(1, t) = 0, (t)$$

$$(52) \quad \eta(x, 0) = f(x), \eta_t(x, 0) = F(x), (x)$$

where

$$\begin{aligned} f(0) &= 0, f(1) = 0 \\ F(0) &= 0, F(1) = 0. \end{aligned}$$

C was a real positive constant and

$$C^2 = \frac{P}{\kappa\sigma}, \text{ where}$$

P = normal tension, κ = density, σ = area of cross-section.

The differential equation (50) also holds¹ when κ, σ are given functions of x , i.e., the string is non-homogeneous. C^2 is then an always positive, given function of x .

¹ See WEBER, "Differentialgleichungen," vol. 2, p. 201.

We try for a solution of the form

$$\eta = u(x) \cdot \phi(t).$$

Substitution of this expression for η in (50) gives

$$u(x)\phi''(t) = C^2 u''(x)\phi(t)$$

whence

$$\frac{\phi''(t)}{\phi(t)} = \frac{C^2 u''(x)}{u(x)}.$$

But the left-hand member is a function of t alone, the right-hand member is a function of x alone; they are equal, hence equal to the same constant, say $-\lambda$. We are thus led to two ordinary differential equations:

$$(53) \quad \frac{d^2u}{dx^2} + \frac{\lambda}{C^2}u = 0$$

with the boundary conditions

$$(54) \quad u(0) = 0, u(1) = 0,$$

and

$$(55) \quad \frac{d^2\phi}{dt^2} + \lambda\phi = 0.$$

b) *The Boundary Problem.*—The problem (53) (54) is of the type

$$(56) \quad \begin{cases} L(u) + \lambda g u = 0 \\ R_0(u) = 0, R_1(u) = 0, \text{ with } \left[p u u' \right]_a^b = 0 \end{cases}$$

with $p = 1, q = 0, g = \frac{1}{C^2} > 0$ on $[ab]$.

We shall show first that every characteristic constant of (56) is positive if

$$(57) \quad p > 0, q \leq 0, g > 0.$$

Let λ_0 be a characteristic constant and $\varphi_0(x)$ a corresponding fundamental function of (56):

$$\varphi_0(x) \text{ of class } C'', \varphi_0(x) \not\equiv 0,$$

$$L(\varphi_0) + \lambda_0 g \varphi_0 = 0, R_0(\varphi_0) = 0, R_1(\varphi_0) = 0.$$

Then

$$\begin{aligned}\lambda_0 \int_a^b g\varphi_0^2 dx &= - \int_a^b \varphi_0 L(\varphi_0) dx \\ &= \int_a^b \left[p \left(\frac{d\varphi_0}{dx} \right)^2 - q\varphi_0^2 \right] dx.\end{aligned}$$

Whence, on account of (57),

$$\lambda_0 = \frac{\int_a^b \left[p \left(\frac{d\varphi_0}{dx} \right)^2 - q\varphi_0^2 \right] dx}{\int_a^b g\varphi_0^2 dx} > 0.$$

Hence, under the assumption (57) the condition (H_2) is always satisfied.

Applying this result to the special case (53) (54), we obtain the following lemma:

Lemma I.—For the boundary problem (53) (54), all of the characteristic constants are positive.

Green's function $K(x, t)$ is the same as for the homogeneous string, since it depends only on $L(u)$ and not on g . Therefore

$$K(x, t) = \begin{cases} (1-t)x, & x \leq t \\ (1-x)t, & x \geq t. \end{cases}$$

The boundary problem (53) (54) is equivalent to

$$u(x) = \lambda \int_0^1 \frac{K(x, t)}{C^2(t)} u(t) dt.$$

In this integral equation the kernel $\frac{K(x, t)}{C^2(t)}$ is not symmetric. But, since $C(x) \neq 0$, this integral equation is reducible to one in which the kernel is symmetric by putting
(58) $u(x) = C(x)\bar{u}(x)$ and $K(x, t) = C(x)C(t)\bar{K}(x, t)$.

Then

$$\bar{u}(x) = \lambda \int_0^1 \bar{K}(x, t) \bar{u}(t) dt.$$

By the same transformation (58) the differential equation (53) is transformed into

$$(59) \quad \frac{d}{dx} \left(C^2 \frac{d\bar{u}}{dx} \right) + CC''\bar{u} + \lambda \bar{u} = 0$$

with the boundary conditions

$$(60) \quad \bar{u}(0) = 0, \bar{u}(1) = 0$$

and it is easily shown that $\bar{K}(x, t)$ is Green's function belonging to the boundary problem (59) (60).

We can now apply the Theorems *V* to *VIII*, with the following result:

The boundary problem (59) (60) has an infinitude of real characteristic constants all of index 1, forming an increasing sequence

$$\lambda_1 < \lambda_2 < \lambda_3 < \dots$$

with corresponding normalized fundamental functions

$$\bar{\psi}_1(x), \bar{\psi}_2(x), \bar{\psi}_3(x), \dots$$

But, if for a given λ , the boundary problem (59) (60) has a non-trivial solution, then for the same λ , (53) (54) has a non-trivial solution $u = C\bar{u}$. Therefore we have the following lemma:

Lemma II.—The boundary problem (53) (54) has an infinitude of real positive characteristic constants, all of index 1, forming an increasing sequence:

$$0 < \lambda_1 < \lambda_2 < \dots,$$

with corresponding normalized fundamental functions

$$\psi_1(x), \psi_2(x), \dots$$

where

$$\psi_n(x) = C\bar{\psi}_n(x).$$

c) *The Generalized Fourier Series.*—We return now to (55) with $\lambda = \lambda_n$:

$$(61) \quad \frac{d^2\phi}{dt^2} + \lambda_n \phi = 0.$$

Since $\lambda_n > 0$, the general solution of (61) is

$$\phi = A_n \cos \sqrt{\lambda_n} t + B_n \sin \sqrt{\lambda_n} t.$$

Therefore, presumably a solution of (50) which satisfies (51) is

$$(62) \quad \eta = \sum_{n=1}^{\infty} (A_n \cos \sqrt{\lambda_n} t + B_n \sin \sqrt{\lambda_n} t) \psi_n(x).$$

In order to be a solution, the condition (52) must be satisfied. Imposing condition (52), we obtain

$$(63) \quad \begin{cases} \eta(x, 0) = \sum_{n=1}^{\infty} A_n \psi_n(x) = f(x) \\ \eta_t(x, 0) = \sum_{n=1}^{\infty} \sqrt{\lambda_n} B_n \psi_n(x) = F(x). \end{cases}$$

But $f(x)$ and $F(x)$ are given functions. Hence (63) can be satisfied if A_n and B_n can be determined so that the series in (63) represents $f(x)$ and $F(x)$. Since $f(0) = 0$, $f(1) = 0$, $F(0) = 0$, $F(1) = 0$, it follows, from Theorem VI, that $f(x)$ and $F(x)$ can be so represented if f and F are of class C'' , and then

$$A_n = (f\psi_n), \sqrt{\lambda_n} B_n = (F\psi_n).$$

We can be sure that (62) satisfies the differential equation if the series is twice differentiable, term by term, with respect to x and t . Since

$$\psi_n''(x) = -\frac{\lambda_n}{C^2} \psi_n(x)$$

this means if

$$\sum \lambda_n (A_n \cos \sqrt{\lambda_n} t + B_n \sin \sqrt{\lambda_n} t) \psi_n(x)$$

is uniformly convergent in x and t .

d) Special Case of Homogeneous String.—If c is constant, then

$$\lambda_n = n^2 \pi^2 c^2$$

and $\psi_n(x) = \sqrt{2} \sin n\pi x$,

while

$$\begin{aligned}\eta &= \sum_{n=1}^{\infty} (A_n \cos n\pi ct + B_n \sin n\pi ct) \sin n\pi x \\ &\equiv \sum_{n=1}^{\infty} \eta_n, \text{ say.}\end{aligned}$$

For $f(x)$ and $F(x)$ we obtain:

$$\begin{aligned}f(x) &= \sum_{n=1}^{\infty} \sqrt{2} A_n \sin n\pi x \\ F(x) &= \sum_{n=1}^{\infty} \sqrt{2} n\pi c B_n \sin n\pi x.\end{aligned}$$

These are sine series for $f(x)$ and $F(x)$. For the development of an arbitrary function in trigonometric series we need know only that the function is continuous and has a finite number of maxima and minima. These conditions are not so strong as those obtained by means of the theory of integral equations which were demanded for the development in series of fundamental functions.

η_1 is periodic in t with period $T_1 = \frac{2}{c}$, this being the period of the fundamental tone. η_n is periodic with period $T_n = \frac{2}{nc} = \frac{T_1}{n}$, and with intensity $\sqrt{A_n^2 + B_n^2}$. Upon the intensity of the different harmonics depends the quality of the tone. The tone of period $\frac{T}{n}$ is called the n th harmonic overtone, or simply the n th harmonic.

For the non-homogeneous string η_n is also periodic with period $\frac{2\pi}{\sqrt{\lambda_n}}$.

T_n decreases with n ; the different periods are not fractions of T_1 , hence the total motion is not periodic.

82. Vibrations of a Rope. *a) Differential Equation of the Problem.*—Let us consider a heavy rope of length 1

$$AB = 1$$

suspended at one end A . It is given a small initial displacement in a vertical plane through AB and then each particle is given an initial velocity. The rope is supposed to vibrate in a given vertical plane and the displacement is so small that each particle is supposed to move horizontally; the cross-section is constant; the density constant; the cross-section infinitesimal compared to the length.

Let AB' be the position of the rope at time t and P any point on AB' . Draw PM horizontal and put

$$MP = \eta, BM = x.$$

Then the differential equation¹ of the motion is given by

$$(64) \quad \frac{\partial^2 \eta}{\partial t^2} = C^2 \frac{\partial}{\partial x} \left(x \frac{\partial \eta}{\partial x} \right)$$

where $C^2 = \text{constant}$,

with the boundary conditions

$$(65) \quad \eta(1, t) = 0, \quad \eta(0, t) \text{ finite}$$

$$(66) \quad \eta(x, 0) = f(x), \quad \eta_t(x, 0) = F(x).$$

We try for a solution of the form

$$\eta = u(x) \cdot \phi(t).$$

Substitute this expression for η in (64). We obtain

$$u(x)\phi''(t) = C^2\phi(t) \frac{d}{dx} \left(x \frac{du}{dx} \right).$$

¹ See KNESER, "Die Integralgleichungen und ihre Anwendung in der Mathematischen Physik," §11.




FIG. 21.

Whence

$$\frac{\phi''(t)}{\phi(t)} = C^2 \frac{d}{dx} \left(x \frac{du}{dx} \right) = -\lambda C^2, \text{ constant.}$$

That is

$$\frac{d^2\phi}{dt^2} + \lambda C^2 \phi = 0,$$

and

$$(67) \quad \frac{d}{dx} \left(x \frac{du}{dx} \right) + \lambda u = 0$$

with the boundary condition derived from (65):

$$(68) \quad u(0) \text{ finite}, \quad u(1) = 0.$$

Equation (67) is of the form

$$L(u) + \lambda u = 0$$

with

$$p = x, \quad q = 0.$$

In the general case $p \neq 0$ on $[ab]$. This condition is not

fulfilled in (67), for $p(0) = 0$. We have also the condition $u(0)$ finite, which did not appear in the general case. The differential equation (67) has a singular point for $x = 0$.

b) *Solution of the Boundary Problem.*—We solve the boundary problem (67), (68) directly. Put

$$x = \frac{t^2}{4\lambda},$$

then (67) becomes

$$(69) \quad t \frac{d^2u}{dt^2} + \frac{du}{dt} + tu = 0.$$

We try to find a solution of the form

$$u = \sum_{n=0}^{\infty} C_n t^n.$$

If we substitute this series for u in (69), we obtain

$$C_1 + \sum_{n=1}^{\infty} \left[C_{n-1} + (n+1)^2 C_{n+1} \right] t^n \equiv 0.$$

Therefore

$$C_1 = 0, C_{n-1} + (n+1)^2 C_{n+1} = 0, n = 1, 2, \dots.$$

That is,

$$C_{n+1} = -\frac{C_{n-1}}{(n+1)^2}.$$

Whence

$$C_1 = C_3 = C_5 = \dots = C_{2n+1} = 0 \dots$$

$$C_2 = -\frac{C_0}{2^2}, C_4 = \frac{C_0}{2^2 \cdot 4^2}, C_6 = -\frac{C_0}{2^2 \cdot 4^2 \cdot 6^2}, \dots$$

Therefore

$$u = C_0 \left[1 - \frac{t^2}{2^2} + \frac{t^4}{2^2 \cdot 4^2} - \frac{t^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots \right].$$

By comparison with the series for e^x , which is permanently convergent, we show that this series for u is permanently convergent. This series also satisfies (69). Put

$$(70) \quad J(t) = 1 - \frac{t^2}{2^2} + \frac{t^4}{2^2 \cdot 4^2} - \frac{t^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$

$J(t)$ is called Bessel's function of order zero.

$J(t)$ is then a solution of (69). Knowing a particular solution, we can, by means of Green's formula, find the general solution:

$$t \left[J(t)V' - VJ'(t) \right] = C_1 \neq 0 (C_1 \text{ constant}).$$

Divide both members by $J^2(t)$ and integrate. We find

$$\frac{V}{J(t)} = C_1 \int \frac{dt}{tJ^2(t)} + C_0.$$

But from (70)

$$\frac{1}{J^2(t)} = 1 + t^2 P(t^2),$$

$P(t^2)$ being a power series in t^2 , whence

$$\int \frac{dt}{tJ^2(t)} = \log t + P_1(t^2)$$

and

$$V = C_0 J(t) + C_1 J(t) \left\{ \log t + P_1(t^2) \right\}$$

where $P_1(t^2)$ is a power series in t^2 .

This is the general solution of (69). The solution of (67) is given by putting

$$t = 2\sqrt{\lambda} \sqrt{x}.$$

Whence

$$u = C_0 J(2\sqrt{\lambda}x) + C_1 J(2\sqrt{\lambda}x) \left\{ \log 2\sqrt{\lambda}x + P_1(4\lambda x) \right\}.$$

But, $u(0)$ finite, is a condition upon the solution and this is impossible unless $C_1 = 0$. Therefore

$$(71) \quad u = C_0 J(2\sqrt{\lambda}x)$$

is the most general solution of (67), which satisfies the first initial condition. We have the further condition $u(1) = 0$, whence

$$J(2\sqrt{\lambda}) = 0.$$

The solution of this equation gives us the characteristic constants.

c) *Construction of Green's Function.*—We construct for the boundary problem (67) (68) the Green's function $K(x, t)$ satisfying the following conditions:

A) K continuous on $[0, 1]$.

B) K of class C'' on $[0, t] [t, 1]$ separately.

$$\frac{d}{dx} \left(x \frac{dK}{dx} \right) = 0 \text{ on } [0, t] [t, 1] \text{ separately.}$$

C) $K(0, t)$ finite, $K(1) = 0$.

$$D) \quad K'(t - 0) - K'(t + 0) = \frac{1}{t}.$$

Integrating the differential equation in B), we obtain

$$K(x, t) = \begin{cases} \alpha_0 \log x + \beta_0, & [0, t] \\ \alpha_1 \log x + \beta_1, & [t, 1]. \end{cases}$$

But $K(0, t)$ is finite, therefore $\alpha_0 = 0$,
 and $K(1, t) = 0$, therefore $\beta_1 = 0$.

Hence

$$K(x, t) = \begin{cases} K_0(x, t) = \beta_0 & , \begin{bmatrix} 0t \\ \end{bmatrix} \\ K_1(x, t) = \alpha_1 \log x, \begin{bmatrix} t1 \\ \end{bmatrix} \end{cases}$$

From condition A)

$$\beta_0 = \alpha_1 \log t.$$

From condition D), since $K'(t - 0) = 0$ and $K'(t + 0) = \frac{\alpha_1}{t}$, we obtain $\alpha_1 = -1$.

Therefore

$$(72) \quad K(x, t) = \begin{cases} K_0(x, t) = -\log t, \begin{bmatrix} 0t \\ \end{bmatrix} \\ K_1(x, t) = -\log x, \begin{bmatrix} t1 \\ \end{bmatrix} \end{cases}$$

We observe that $K(x, t)$ is symmetric. The graph of $K(x, t)$ for t fixed is shown by the full line in the accompanying figure. As a function of the two variables x and t , $K(x, t)$ is continuous on $[01]$ except at $x = t = 0$. For in the region marked I we have $t > x$ and

$$K(x, t) = \log \frac{1}{t},$$




FIG. 22.




FIG. 23.

while in the region marked II we have $t < x$ and

$$K(x, t) = \log \frac{1}{x}.$$

For $x = 0, t > 0$, we have $K = -\log t$, finite.

For $t = 0, x > 0$, we have $K = -\log x$, finite.

d) *Equivalence with a Homogeneous Integral Equation.*—
We have

$$\begin{aligned} L(u) &= -\lambda u \\ L(K) &= 0. \end{aligned}$$

Multiply the first of these by $-K$ and the second by u and add. We obtain

$$uL(K) - KL(u) = \lambda u K.$$

Integrate both members of this expression from 0 to $t = 0$ and from $t + 0$ to 1 with respect to t and add. We obtain

$$u(x) = \lambda \int_0^1 K(x, t)u(t)dt.$$

The details of the integration are the same as those given several times previously, except for the term

$$x \left[uK' - Ku' \right] \text{ for } x = 0.$$

But here u and K are finite by hypothesis. From (71) we see that if u is finite so also is u' . From the explicit expression (72) for K we find $K'(0, t) \equiv 0$. Therefore

$$x \left[uK' - Ku' \right] = 0 \text{ for } x = 0.$$

The kernel

$$K(x, t) = \begin{cases} -\log t, & \begin{bmatrix} 0t \\ t \\ \hline \end{bmatrix} \\ -\log x, & \begin{bmatrix} t \\ 1 \\ \hline \end{bmatrix} \end{cases}$$

is symmetric, but it is discontinuous at one point, *viz.*, $x = t = 0$. Schmidt (page 21 of his dissertation) has shown that the results of the Hilbert-Schmidt theory of

continuous symmetric kernels still hold for a discontinuous kernel if

- 1) $\int_a^b K(x, t)f(t)dt$ for f continuous, is continuous in x on $[ab]$.

- 2) The iterated kernel $K_2(x, t)$ is continuous and does not vanish identically.

These conditions are satisfied in the present instance and thus all of the Hilbert-Schmidt theory, as well as the Theorems V to VIII on boundary problems, remain true. Therefore we know that there exists an infinite sequence of positive characteristic constants

$$0 < \lambda_1 < \lambda_2 < \lambda_3 < \dots,$$

with a corresponding complete normalized orthogonal system of fundamental functions. Therefore

$$J(2\sqrt{\lambda}) = 0$$

has an infinitude of positive roots λ_n . Put

$$2\sqrt{\lambda} = k.$$

Then $J(k) = 0$, and the roots are

$$k_n = 2\sqrt{\lambda_n}.$$

The first four values of k_n are¹

$$k_1 = 2.405, k_2 = 5.520, k_3 = 8.654, k_4 = 11.792$$

and generally $(n - \frac{1}{2})\pi < k_n < n\pi$. Therefore

$$\varphi_n(x) = C_0 J(2\sqrt{\lambda_n}x) = C_0 J(k_n\sqrt{x}).$$

These fundamental functions $\varphi_n(x)$ will become orthogonalized if we choose

$$C_0 = \frac{1}{\sqrt{\int_0^1 J^2(k_n\sqrt{x})dx}}$$

¹ See FRICKE, "Analytische-Funktionentheoretische Vorlesungen," p. 74.

But¹ $\int_0^1 J^2(k_n \sqrt{x}) dx = \frac{1}{[J'(k_n)]^2}.$

Therefore

$$\psi_n(x) = \frac{J(k_n \sqrt{x})}{J'(k_n)}, \text{ and}$$

$$\eta = \sum_{n=1}^{\infty} \left(A_n \cos \frac{Ck_n}{2} t + B_n \sin \frac{Ck_n}{2} t \right) \psi_n(x).$$

This expression for η satisfies (64) and (65). We now determine A_n , B_n , if possible, in order that (66) may be satisfied. This gives us the two equations

$$\sum A_n \psi_n(x) = f(x)$$

$$\sum \frac{Ck_n}{2} B_n \psi_n(x) = F(x).$$

Since $f(0)$ is finite, $f(1) = 0$,
 $F(0)$ is finite, $F(1) = 0$,

$f(x)$ and $F(x)$ can be expanded as series in $\psi_n(x)$, provided f and F are of class C'' , and then

$$A_n = \int_0^1 f(x) \psi_n(x) dx = \int_0^1 f(x) \frac{J(k_n \sqrt{x})}{J'(k_n)} dx,$$

while $\frac{Ck_n}{2} B_n = \int_0^1 F(x) \psi_n(x) dx.$

83. The Rotating Rope.—(See KNESER, page 46.)

a) *The Problem and Its Differential Equation.*—Let FG be an axis around which a plane is rotating with constant velocity; a rope AB is attached at a point A of the axis and constrained to remain in the rotating plane. The velocity of the rotation is so large that the weight of the rope can be neglected. Then the straight line AB perpendicular to FG




FIG. 24.

¹ See FRICKE, *Loc. cit.*, p. 65.

is a relative position of equilibrium for the rope. Displace the rope slightly from this position AB , then let it go after imparting to its particles initial velocities perpendicular to AB . The rope will then describe small vibrations around the position of equilibrium.

Let APB' be the position of the rope at the time t , P one of its points, $PM \perp AB$. Put $AM = x$, $MP = \eta$, and suppose $AB = 1$. Then the function $\eta(x, t)$ must satisfy the partial differential equation

$$\frac{\partial^2 \eta}{\partial t^2} = C^2 \frac{\partial}{\partial x} \left[(1 - x^2) \frac{\partial \eta}{\partial x} \right], \quad C \text{ constant},$$

the boundary conditions

$$\eta(0, t) = 0, \quad \eta(1, t) = \text{finite}, \quad (t),$$

and the initial conditions

$$\eta(x, 0) = f(x), \quad \eta_t(x, 0) = F(x), \quad (x).$$

Putting $\eta = u(x) \phi(t)$

we obtain

$$(73) \quad \begin{aligned} & \frac{d}{dx} \left[(1 - x^2) \frac{du}{dx} \right] + \lambda u = 0 \\ & u(0) = 0, \quad u(1) = \text{finite} \\ & \frac{d^2 \phi}{dt^2} + \lambda C^2 u = 0. \end{aligned}$$

b) *Solution of the Boundary Problem.*— $x = 1$ is a singular point of (73). By Fuch's theory¹ (73) has for every λ one and only one solution, which in the vicinity of $x = 1$ is given by

$$u_1 = 1 + a_1(x - 1) + a_2(x - 1)^2 + \dots$$

Every other solution u_2 is obtained from Green's formula

$$(1 - x^2)(u_1 u_2' - u_2 u_1') = C_1 \neq 0.$$

¹ See GOURSAT, "Cours D'Analyse," tome 2, §412.

Therefore

$$\frac{d}{dx} \frac{u_2}{u_1} = \frac{C_1}{(x^2 - 1)u_1^2} = C_1 \left\{ \frac{1}{x-1} + S_1(x-1) \right\}.$$

Whence

$$u_2 = C_0 u_1 + C_1 u_1 \left\{ \log(x-1) + S(x-1) \right\}.$$

Therefore the condition, $u(1)$ finite, leads to $C_1 = 0$ and hence

$$(74) \quad u = C_0 u_1$$

is the most general solution of (73) which satisfies the second boundary condition. Put

$$u_1 = U(x, \lambda).$$

Then the first boundary condition gives

$$U(0, \lambda) = 0$$

for the determination of the characteristic constants. From (74) we see that each characteristic constant is of index 1.

Let λ_0 be a characteristic constant and $\varphi_0(x) = U(x, \lambda_0)$ a corresponding fundamental function. Then $\varphi_0(x)$ is an odd function:

$$\varphi_0(-x) = -\varphi_0(x).$$

For u_1 can be expanded according to powers of x , say

$$u_1 = \sum_{r=1}^{\infty} C_r x^r.$$

The substitution of this series in the differential equation leads to the following recurrent formula

$$C_{r+2} = \frac{r(r+1) - \lambda_0}{(r+1)(r+2)} C_r, \quad r = 0, 1, 2, \dots,$$

which shows that if $C_0 = 0$, then

$$C_{2\mu} = 0, \quad \mu = 0, 1, 2, \dots.$$

But from $U(0, \lambda) = 0$, we obtain $\varphi_0(0) = 0$, hence $C_0 = 0$, and, therefore, $\varphi_0(x)$ is an odd function.

We can at once indicate some of the characteristic constants and fundamental functions from the theory of Legendre's polynomials:

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} \left[(x^2 - 1)^n \right]$$

$$P_0(x) = 1.$$

$P_n(x)$ is a rational integral function of degree n , satisfying the differential equation

$$\frac{d}{dx} \left[(1 - x^2) \frac{dy}{dx} \right] + n(n + 1)y = 0.$$

Furthermore, $P_{2n}(x)$ is an even function and $P_{2n}(0) \neq 0$, $P_{2n-1}(x)$ is an odd function and $P_{2n-1}(0) = 0$. Therefore

$$\lambda_n = 2n(2n - 1), \quad n = 1, 2, 3, \dots$$

is a characteristic constant and $P_{2n-1}(x)$ a corresponding fundamental function.

The characteristic constants are of index 1 since the condition $u(1) = 0$ determines u up to a constant factor.

We will now show that there are no other characteristic constants than

$$\lambda_n = 2n(2n - 1), \quad n = 1, 2, 3, \dots$$

Suppose λ_0 to be a characteristic constant and $\lambda_0 \neq \lambda_n$ and $\psi_0(x)$ a corresponding normalized fundamental function, then, according to the orthogonality theorem (§59),

$$\int_0^1 \psi_0(x) P_{2n-1}(x) dx = 0.$$

Now $\psi_0(x)$ is continuous on the interval $[-1, +1]$ and, therefore, by a theorem due to Weierstrass,¹ can be expanded in an infinite series of polynomials uniformly convergent on $[-1, +1]$:

¹ GOURSAT-HEDRICK, "Mathematical Analysis," vol. 1, §199.

$$(75) \quad \psi_0(x) = \sum_{r=1}^{\infty} G_r(x).$$

But, as shown under b),

$$\psi_0(-x) = -\psi_0(x).$$

Therefore, from

$$\psi_0(-x) = \sum_{r=1}^{\infty} G_r(-x)$$

and (75) above, we obtain

$$\psi_0(x) = \frac{1}{2} \sum_{r=1}^{\infty} \left[G_r(x) - G_r(-x) \right] \equiv \sum_{r=1}^{\infty} H_r(x)$$

and $\sum H_r(x)$ is uniformly convergent on $[-1, +1]$. From the definition of $H_r(x)$ we obtain

$$H_r(-x) = -H_r(x),$$

that is, $H_r(x)$ is an odd function.

From the uniform convergence of $\sum H_r(x)$ we have that for every positive ϵ it is possible to find an m such that

$$(76) \quad \left| \psi_0(x) - \sum_{r=1}^m H_r(x) \right| < \epsilon, \quad [-1, +1].$$

Let us choose

$$(77) \quad 0 < \epsilon < \frac{1}{\int_0^1 |\psi_0(x)| dx}.$$

Since $H_r(x)$ is an odd function we have

$$(78) \quad \sum_{r=1}^m H_r(x) = \sum_{r=0}^n C_r x^{2r-1}.$$

But $P_{2n-1}(x)$ are odd functions:

$$P_1(x) = a_{11}x$$

$$P_3(x) = a_{22}x^3 + a_{21}x$$

$$P_{2n-1}(x) = a_{nn}x^{2n-1} + \dots + a_{n1}x$$

with $a_{11}, a_{22}, \dots, a_{nn} \neq 0$.

These equations can be solved sequentially for x , x^3 , \dots , x^{2n-1} in terms of $P_1, P_3, \dots, P_{2n-1}$. Put these values of x^{2r-1} in (78) above and we will have

$$\sum_{r=1}^m H_r(x) = \sum_{r=0}^n C_r P_{2r-1}(x).$$

This expression for $\sum H_r(x)$, substituted in (76), gives

$$\left| \psi_0(x) - \sum_{r=0}^n C_r P_{2r-1}(x) \right| < \epsilon.$$

Put

$$(79) \quad \psi_0(x) = \sum_{r=0}^n C_r P_{2r-1}(x) + r(x),$$

whence

$$|r(x)| < \epsilon \text{ on } [-1, +1].$$

From (79) we obtain

$$\int_0^1 \psi_0^2(x) dx = \sum_{r=0}^n C_r \int_0^1 \psi_0(x) P_{2r-1}(x) dx + \int_0^1 r(x) \psi_0(x) dx.$$

Part

$$\int_0^1 \psi_0(x) P_{2r-1}(x) dx = 0$$

and

$$\int_0^1 \psi_0^2(x) dx = 1.$$

Therefore

$$1 = \int_0^1 r(x) \psi_0(x) dx.$$

$$\text{Now } \left| \int_0^1 r(x) \psi_0(x) dx \right| \leq \int_0^1 \left| r(x) \psi_0(x) \right| dx \\ < \epsilon \int_0^1 \left| \psi_0(x) \right| dx.$$

$$\text{Therefore } 1 < \epsilon \int_0^1 \left| \psi_0(x) \right| dx,$$

which, on account of our choice (77) of ϵ , gives

$$1 < \epsilon \int_0^1 \left| \psi_0(x) \right| dx < 1,$$

which constitutes a contradiction. Therefore

$$\lambda_n = 2n(2n - 1)$$

are the only characteristic constants, and the only fundamental functions are

$$\varphi_n(x) = \frac{P_{2n-1}(x)}{\sqrt{\int_0^1 P_{2n-1}^2(x) dx}}.$$

$$\text{But } \int_0^1 P_{2n-1}^2(x) dx = \frac{1}{4n-1},$$

as shown in the theory of Legendre's polynomials. Therefore

$$\varphi_n(x) = \sqrt{4n-1} P_{2n-1}(x).$$

C) Equivalence with Integral Equation.—We construct the Green's function as before and obtain

$$K(x, t) = \begin{cases} \frac{1}{2} \log \frac{1+x}{1-x}, & 0 \leq x \leq t \\ \frac{1}{2} \log \frac{1+t}{1-t}, & t \leq x \leq 1. \end{cases}$$

$K(x, t)$ is symmetric and has but one point of discontinuity and that is for $x = t = 1$.

Schmidt's conditions for a discontinuous kernel hold, however, and so the theorems of the Hilbert-Schmidt theory apply. Proceeding as in the previous problems, we find

that the boundary problem is equivalent to the following integral equation:

$$u(x) = \lambda \int_0^1 K(x, t)u(t)dt.$$

The Theorems V to IX hold for this boundary problem as well as the orthogonality theorem, §59. Hence

$$\int_0^1 P_{2n-1}(x)P_{2m-1}(x)dx = 0, m \neq n$$

This last result agrees with a result of the theory of Legendre polynomials. Applying Theorem VI, we see that if f is of class C'' and $f(0) = 0$, then

$$f(x) = \sum_{r=0}^{\infty} C_r P_{2n-1}(x)$$

where

$$C_r = (fP_{2n-1}) \frac{4n - 1}{1}.$$

IV. APPLICATIONS OF THE HILBERT-SCHMIDT THEORY TO THE FLOW OF HEAT IN A BAR

84. The Partial Differential Equations of the Problem.¹

a) *General Hypotheses.*—The theory of the flow of heat in a bar is based upon the following hypotheses:

A) Let dm denote the mass of an element of a conductor of temperature θ , C its specific heat, then the amount of heat dQ necessary to increase the temperature of the element from θ to $\theta + d\theta$ is given by the formula

$$dQ = Cdmd\theta.$$

B) Let $d\omega$ be the area of an element of surface through an interior point P of a conductor, n one of the two normals to the element, k the inner conductivity at point P , then the amount of heat which flows through the inner surface $d\omega$ in time dt is

$$dQ = -k \frac{\partial \theta}{\partial n} d\omega dt.$$

¹ See WEBER, "Partielle Differentialgleichungen," §§32–34.

C) Let $d\omega$ be an element of the outer surface of the conductor, h the outer conductivity at P , θ the temperature of the conductor at P , Θ the temperature of the surrounding medium at P , then the amount of heat which flows through the outer surface $d\omega$ in time dt is

$$dQ = h(\theta - \Theta)d\omega dt.$$

b) We now apply these principles to the determination of the flow of heat in a straight bar placed along the x -axis of a rectangular system of coordinates, with a cross-section σ , infinitesimal compared to the length, so that the temperature θ may be considered as constant; accordingly, θ will be a function of x and t : $\theta(x, t)$. The bar is imbedded in a medium of given temperature which is also supposed to be a function of x and t : $\Theta(x, t)$. The bar has a given initial temperature $\theta(x, 0) = f(x)$. Consider an element of the bar of length dx ; we may regard it as a cylinder.




FIG. 25.

1) By B) the quantity of heat which enters at the left-hand end is $-k\sigma \frac{\partial \theta}{\partial x} \Big|_x dt$.

2) Similarly, the quantity of heat which flows out through the right-hand end is $-k\sigma \frac{\partial \theta}{\partial x} \Big|_{x+dx} dt$,

or, if we expand according to powers of dx and neglect higher powers of dx ,

$$-k\sigma \frac{\partial \theta}{\partial x} \Big|_x dt - \frac{\partial}{\partial x} \left(k\sigma \frac{\partial \theta}{\partial x} \right) \Big|_x dt dx.$$

3) The quantity of heat which flows out through the cylindrical surface is, by C),

$$h(\theta - \Theta)ldxdt,$$

where l is the periphery of the cross-section. The total amount of heat which enters the element of the bar in the time dt is, therefore,

$$\left\{ \frac{\partial}{\partial x} \left(k\sigma \frac{\partial \theta}{\partial x} \right) - h(\theta - \Theta)l \right\} dxdt.$$

This amount of heat increases the temperature of the element in the time dt by $\frac{\partial \theta}{\partial t} dt$ and, therefore, by A), is equal to $C\rho\sigma dx \frac{\partial \theta}{\partial t} dt$, where ρ is the density. Hence we obtain the partial differential equation

$$(80) \quad C\rho\sigma \frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left(k\sigma \frac{\partial \theta}{\partial x} \right) - h(\theta - \Theta)l.$$

If, moreover, heat is produced in the interior of the bar by electric currents or other sources of energy, let

$$A(x)dxdt$$

be the quantity of heat produced in the time dt in the interior of the element of the bar between the cross-sections, at x and $x + dx$. Then the partial differential equation (80) becomes

$$c\rho\sigma \frac{\partial \theta}{\partial x} = \frac{\partial}{\partial x} \left(k\sigma \frac{\partial \theta}{\partial x} \right) - h(\theta - \Theta)l + A(x).$$

In addition, the temperature θ has to satisfy certain boundary conditions, which are obtained as follows:

The amount of heat which leaves at a is, by C),

$$(81) \quad h(\theta - \Theta)\sigma \Big|_a dt.$$

On the other hand, the quantity of heat which flows through the cross-section $a + h$ in the direction of the negative x -axis in the time dt is, by $B)$,

$$k\sigma \left. \frac{\partial \theta}{\partial x} \right]_{a+h} dt.$$

The limit of this expression as h approaches zero must be equal to the quantity (81). Hence we obtain the first boundary condition

$$h(\theta - \Theta) - k \left. \frac{\partial \theta}{\partial x} \right]_a = 0.$$

The same reasoning applied to the cross-section b gives us the second boundary condition

$$h(\theta - \Theta) + k \left. \frac{\partial \theta}{\partial x} \right]_b = 0.$$

c) *The Special Case $\Theta \equiv 0, A \equiv 0$.*—We now consider the differential equation under the assumption that

$$\Theta \equiv 0, A \equiv 0.$$

The differential equation is, then,

$$c\sigma\rho \frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left(k\sigma \frac{\partial \theta}{\partial x} \right) - hl\theta.$$

Put $c\sigma\rho = g(x), k\sigma = p(x), hl = -q(x)$,
then $g(x) > 0, p(x) > 0, q(x) \leq 0$,
and $q(x) = 0$ only when $h = 0$.

The differential equation of the problem now becomes

$$(82) \quad g \frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left(p \frac{\partial \theta}{\partial x} \right) + q\theta.$$

Put $\frac{h(a)}{k(a)} = H_0 \geq 0, \frac{h(b)}{k(b)} = H_1 \geq 0$,

then the boundary conditions become

$$(83) \quad \begin{cases} \theta_x(a, t) - H_0\theta(a, t) = 0 \\ \theta_x(b, t) + H_1\theta(b, t) = 0 \end{cases} \quad (t).$$

The initial condition is

$$(84) \quad \theta(x, 0) = f(x).$$

$f(x)$ is not entirely arbitrary, for from (83) we obtain for $t = 0$

$$\begin{aligned} \theta_x(a, 0) - H_0\theta(a, 0) &= 0 \\ \theta_x(b, 0) + H_1\theta(b, 0) &= 0 \end{aligned}$$

whence

$$(85) \quad \begin{aligned} f'(a) - H_0f(a) &= 0 \\ f'(b) + H_1f(b) &= 0. \end{aligned}$$

To solve (82), put

$$\theta = u(x)\phi(t).$$

Then, in the usual manner, (82) breaks up into the two ordinary differential equations:

$$(86) \quad \frac{d}{dx} \left(p \frac{du}{dx} \right) + qu + \lambda gu = 0$$

$$(87) \quad \frac{d\phi}{dt} + \lambda\phi = 0,$$

while from (83) we get

$$(88) \quad \begin{aligned} u'(a) - H_0u(a) &= 0 \\ u'(b) + H_1u(b) &= 0 \end{aligned}$$

as boundary conditions on (86).

We have, as in our hypothesis (H_1), §72, that p is of class C' , $p > 0$. In addition, we have $g > 0$ and $q \leq 0$.

We shall now show that all of the characteristic constants of (86) are ≥ 0 . This proof is analogous to that given in the proof of Theorem VIII.

We suppose that λ_0 is a characteristic constant and φ_0 a corresponding fundamental function. Therefore

$$(89) \quad L(\varphi_0) + \lambda_0\varphi_0g = 0$$

$$(90) \quad \varphi_0'(a) - H_0\varphi_0(a) = 0, \quad \varphi_0'(b) + H_1\varphi_0(b) = 0$$

and $\varphi_0 \not\equiv 0$ is of class C'' .

From (89) we obtain

$$\lambda_0 \int_a^b g\varphi_0^2 dx = - \int_a^b \varphi_0 L(\varphi_0) dx.$$

Integration by parts gives

$$\begin{aligned} \lambda_0 \int_a^b g\varphi_0^2 dx &= - p(b)\varphi_0(b)\varphi_0'(b) + p(a)\varphi_0(a)\varphi_0'(a) \\ &\quad + \int_a^b (p\varphi_0'^2 - q\varphi_0^2) dx, \end{aligned}$$

which, by (90), reduces to

$$(91) \quad \lambda_0 \int_a^{b1} g\varphi_0^2 dx = p(b)\varphi_0^2(b)H_1 + p(a)\varphi_0^2(a)H_0 + \int_a^b (p\varphi_0'^2 - q\varphi_0^2) dx.$$

Now $p(b)$, $\varphi_0^2(b)$, $p(a)$, $\varphi_0^2(a)$, and p are positive and not zero, and H_1 , H_0 , $\varphi_0'^2$, φ_0^2 , $-q$ are ≥ 0 . Whence, from (91), we conclude

$$\lambda_0 \geq 0.$$

The equality will hold only when $H_1 = 0$, $H_0 = 0$, $q \equiv 0$ simultaneously. But $-q = hl$ and $l > 0$. Therefore $h \equiv 0$, which means that no heat escapes through the cylindrical surface. It also follows that no heat escapes through the ends for $h(a) = 0$, $h(b) = 0$, and hence the equality holds only when no heat escapes to the surrounding medium.

We can now construct the Green's function $K(x, t)$ and establish the equivalence of the boundary problem with the integral equation

$$u(x) = \lambda \int_a^b K(x, t)g(t)u(t)dt.$$

The kernel is not symmetric but the substitution

$$u(x)\sqrt{g(x)} = \bar{u}(x), \quad K(x, t)\sqrt{g(x)g(t)} = \bar{K}(x, t)$$

transforms the problem into one with a symmetric kernel. This transformation leads the given differential equation

into one in which $g \equiv 1$. We can then apply the results of the general Hilbert-Schmidt theory. Further, we can apply the Theorems V to VIII. We know then the existence of an infinitude of real characteristic constants each of index 1:

$$0 \leq \lambda_1 < \lambda_2 < \lambda_3 < \dots$$

with corresponding normalized fundamental functions

$$\psi_1(x), \psi_2(x), \psi_3(x), \dots$$

We now turn to the solution of (87) for $\lambda = \lambda_n$. The solution is

$$\phi(t) = C_n e^{-\lambda_n t}$$

whence

$$\theta = C_n e^{-\lambda_n t} \psi_n(x)$$

is a solution of (82) which satisfies the boundary conditions but in general will not satisfy the initial conditions (84). Construct

$$(92) \quad \theta = \sum_{n=1}^{\infty} C_n e^{-\lambda_n t} \psi_n(x).$$

If we assume that (92) is convergent and admits one term-by-term partial differentiation with respect to t and two term-by-term partial differentiations with respect to x , then (92) satisfies (82) and (83), and (84) will be satisfied if

$$\sum_{n=1}^{\infty} C_n \psi_n(x) = f(x),$$

that is, if $f(x)$ can be expanded into a series of fundamental functions. That this is possible follows from Theorem VI when we take account of (85), provided f is of class C'' , and C_n will be given by

$$C_n = (f \psi_n).$$

85. Application to an Example.—We consider now a special case where C, σ, ρ, k, h, l are constants and $a = 0$,

$b = 1$. Hence, p , q , g in (86) are constant. Put $\frac{q}{p} = -b^2$, and $\frac{\lambda g}{p} = \text{constant}$, which we use as a new parameter and again designate by λ . Then (86) becomes

$$\frac{d^2u}{dx^2} + (\lambda - b^2)u = 0.$$

If we now assume that no heat escapes through the two end surfaces, then $H_0 = H_1 = 0$ and the boundary conditions become

$$u'(0) = 0, u'(1) = 0.$$

For this boundary problem the characteristic constants are easily found to be

$$\lambda_0 = b^2, \lambda_n = n^2\pi^2 + b^2, n = 1, 2, 3, \dots.$$

and the normalized fundamental functions are $\psi_0(x) = 1$, $\psi_n(x) = \sqrt{2} \cos n\pi x$.

For the further discussion we have to distinguish two cases:

Case I.— $b \neq 0$. Then all of the characteristic constants are positive. Therefore $\lambda = 0$ is not a characteristic constant and condition (H_2) is satisfied. Hence for this boundary problem we can construct a Green's function satisfying the following conditions:

- A) K is continuous on $[01]$.
- B) K is of class C'' on $[0t] [t1]$ separately.
- $\frac{d^2K}{dx^2} - b^2K = 0$ on $[0t] [t1]$ separately.
- C) $K'(0) = 0, K'(1) = 0$.
- D) $K'(t - 0) - K'(t + 0) = 1$.

The Green's function satisfying these conditions is easily found to be

$$K(x, t) = \begin{cases} \frac{\cosh bx \cosh b(1-t)}{b \sinh b}, & 0 \leq x \leq t \\ \frac{\cosh bt \cosh b(1-x)}{b \sinh b}, & t \leq x \leq 1. \end{cases}$$

It is symmetric and the boundary problem is equivalent to the integral equation

$$u(x) = \lambda \int_0^1 K(x, t) u(t) dt.$$

Theorem VI of this chapter assures us that every fundamental function of class C'' of this boundary problem, for which $u'(0) = 0, u'(1) = 0$ can be expanded into a cosine series

$$u(x) = \sum_{n=0}^{\infty} C_n \cos n\pi x$$

convergent on the interval $0 \leq x \leq 1$.

Case II.— $b = 0$. The boundary problem now becomes

$$(93) \quad \frac{d^2u}{dx^2} + \lambda u = 0, \quad u'(0) = 0, \quad u'(1) = 0,$$

for which the characteristic constants are

$$\lambda_0 = 0, \quad \lambda_n = n^2\pi^2, \quad n = 1, 2, 3, \dots$$

with the corresponding normalized fundamental functions

$$\psi_0(x) = 1, \quad \psi_n(x) = \sqrt{2} \cos n\pi x.$$

Let us consider

$$\bar{K}(x, t) = K(x, t) - \frac{1}{b^2}.$$

From the conditions on K we find that \bar{K} satisfies the same conditions. Now let

$$\lim_{b \rightarrow 0} \bar{K}(x, t) = H(x, t).$$

We find

$$H(x, t) = \begin{cases} \frac{x^2 + t^2}{2} + \frac{1}{3} - t \equiv H_0, & 0 \leq x \leq t \\ \frac{x^2 + t^2}{2} + \frac{1}{3} - x \equiv H_1, & t \leq x \leq 1. \end{cases}$$

We see that $H(x, t)$ is symmetric and satisfies the following conditions:

- A) H is continuous on $[0, 1]$.
- B) H is of class C'' on $[0, t] \cup [t, 1]$ separately.
- $\frac{d^2H}{dx^2} = 1$ on $[0, t] \cup [t, 1]$ separately.
- C) $H'(0) = 0, H'(1) = 0$.
- D) $H'(t - 0) - H'(t + 0) = 1$.
- E) $\int_0^1 H(x, t)dt = 0, (t)$.

We will now show that the boundary problem (93) is equivalent to

$$u(x) = \lambda \int_0^1 H(x, t)u(t)dt.$$

A) Multiply both members of $\frac{d^2u}{dx^2} = -\lambda u$ by $-H$, and both members of $\frac{d^2H}{dx^2} = 1$ by u and add. We obtain

$$\frac{d}{dx}(uH' - Hu') = \lambda uH + u,$$

whence

$$(uH' - Hu') \Big|_0^{t=0} + (uH' - Hu') \Big|_{t=0}^1 = \int_0^1 u(x)dx + \lambda \int_0^1 H(x, t)u(x)dx,$$

which, on account of (93) and condition C) on H , reduces to

$$(uH' - Hu') \Big|_{t=0}^{t=0} = \lambda \int_0^1 u(x)dx + \lambda \int_0^1 H(x, t)u(x)dx,$$

which, by conditions A) and D) on H , reduces to

$$u(t) = \lambda \int_0^1 u(x)dx + \lambda \int_0^1 H(x, t)u(x)dx.$$

Now from

$$\frac{d^2u}{dx^2} = -\lambda u(x)$$

we obtain, by integration,

$$-\int_0^1 u(x)dx = \int_0^1 \frac{d^2u}{dx^2}dx = \frac{du}{dx} \Big|_0^1 = 0.$$

Therefore

$$(94) \quad u(x) = \lambda \int_0^1 H(x, t)u(t)dt.$$

Thus we have shown that every solution of the boundary problem (93) with $\lambda \neq 0$ is a solution of the integral equation (94).

B) Conversely, suppose that u is continuous and

$$(95) \quad u(x) = \lambda \int_0^1 H(x, t)u(t)dt, \text{ and } \lambda \neq 0.$$

Then

$$u(x) = \lambda \int_0^x H_1(x, t)u(t)dt + \lambda \int_x^1 H_0(x, t)u(t)dt,$$

whence

$$(96) \quad u'(x) = \lambda \int_0^x H_1'(x, t)u(t)dt + \lambda \int_x^1 H_0'(x, t)u(t)dt$$

and

$$u''(x) = \lambda \int_0^x H_1''(x, t)u(t)dt + \lambda \int_x^1 H_0''(x, t)u(t)dt \\ + \lambda H_1'(x, x)u(x) - \lambda H_0'(x, x)u(x).$$

But $H'' = 1$ and $H'(t = 0) - H'(t = 1) = 1$.

Therefore

$$u''(x) + \lambda u(x) = \lambda \int_0^1 u(t)dt.$$

But from (95)

$$\begin{aligned}\int_0^1 u(x)dx &= \lambda \int_0^1 \int_0^1 H(x, t)u(t)dtdx \\ &= \lambda \int_0^1 u(t)dt \int_0^1 H(x, t)dx = 0\end{aligned}$$

on account of E). Therefore

$$u''(x) + \lambda u(x) = 0.$$

Further, from (96) we obtain

$$\begin{aligned}u'(0) &= \lambda \int_0^1 H_0'(0, t)u(t)dt = 0 \\ u'(1) &= \lambda \int_0^1 H_1'(1, t)u(t)dt = 0,\end{aligned}$$

on account of C). Therefore we have proved the equivalence of the boundary problem and integral equation.

86. General Theory of the Exceptional Case.—The method just applied to the special boundary problem (93) is a special case of the general method which Hilbert uses in the exceptional case where $\lambda = 0$ is a characteristic constant of the boundary problem:

$$(97) \quad L(u) + \lambda u = 0$$

$$(98) \quad R_0(u) = 0, R_1(u) = 0.$$

Let $\psi_0(x)$ be a normalized fundamental function belonging to $\lambda = 0$, so that ψ_0 is of class C'' ,

$$\begin{aligned}L(\psi_0) &= 0, R_0(\psi_0) = 0, R_1(\psi_0) = 0, \\ \int_a^b \psi_0^2(x)dx &= 1.\end{aligned}$$

Then, according to Hilbert,¹ the equivalence of (97) (98) with an integral equation can be established as follows.

a) *The Modified Green's Function.*—

¹ Gött. Nach., 2, p. 219, 1904.

Theorem I.—Under the above assumptions there exists one and only one function $H(x, t)$ satisfying as a function of x the following conditions:

A) H is continuous on $[ab]$.

B) H is of class C'' on $[at] [tb]$ separately.

$$L(H) = \psi_0(x)\psi_0(t) \text{ on } [at] [tb] \text{ separately.}$$

C) $R_0(H) = 0, R_1(H) = 0.$

D) $H'(t - 0) - H'(t + 0) = \frac{1}{p(t)}.$

E) $\int_a^b H(x, t)\psi_0(x)dx = 0, (t).$

$H(x, t)$ is called the modified Green's function. An outline of the proof of this theorem is as follows:

Let $u_0(x)$ be a particular solution of

$$L(u) = \psi_0(x)\psi_0(t),$$

then the general solution is

$$u = u_0(x) + \alpha\psi_0(x) + \beta V(x),$$

where V is a particular solution of $L(u) = 0$, independent of $\psi_0(x)$.

Now, by Green's theorem,

$$p(\psi_0 V' - \psi_0' V) = C$$

and we can select V in such a way that $C = 1$.

We now put

$$H(x, t) = \begin{cases} u_0(x) + \alpha_0\psi_0(x) + \beta_0 V(x), & x \leq t \\ u_0(x) + \alpha_1\psi_0(x) + \beta_1 V(x), & t \leq x. \end{cases}$$

Then

$$(99) \quad \begin{aligned} R_0(H) &= R_0(u_0) + \beta_0 R_0(V) = 0 \\ R_1(H) &= R_1(u_0) + \beta_1 R_1(V) = 0 \end{aligned}$$

determine β_0, β_1 , since $R_0(V) \neq 0, R_1(V) \neq 0$.

The conditions $A)$ to $D)$ determine $\alpha_0 - \alpha_1$ and $\beta_0 - \beta_1$:

$$\alpha_0 - \alpha_1 = -V(t), \quad \beta_0 - \beta_1 = \psi_0(t).$$

Since β_0, β_1 have already been determined, this furnishes an apparent contradiction, but it will be found that the determination of $\beta_0 - \beta_1$ is a consequence of the determination of β_0 and β_1 separately. Thus the conditions $A)$ to $D)$ determine $\beta_0, \beta_1, \alpha_0, \alpha_1$, except for an additive constant which is determined by E .

Theorem II. *The function $H(x, t)$ is symmetric.*—

For let $a \leq t \leq s \leq b$.

Multiply each member of

$$L \left\{ H(x, t) \right\} = \psi_0(x)\psi_0(t)$$

by $-H(x, s)$ and each member of

$$L \left\{ H(x, s) \right\} = \psi_0(x)\psi_0(s)$$

by $H(x, t)$. Add and integrate from a to t , t to s , s to b and add the results. We obtain

$$H(t, s) = H(s, t).$$

b) Hilbert's fundamental Theorem III now takes the form

Theorem III.—*If f is continuous, then the statements F is of class C'' , $L(F) + f = 0$, $R_0(F) = 0$, $R_1(F) = 0$ imply and are implied by*

$$F(x) = \int_a^b H(x, t)f(t)dt.$$

Hence the expansion Theorem VI has to be modified as follows:

If F is of class C'' , $R_0(F) = 0$, $R_1(F) = 0$, then

$$F(x) = \sum C_\nu \psi_\nu(x)$$

where

$$C_\nu = (F\psi_\nu).$$

Applying this theorem when $F = u$, $f = \lambda u$, $\lambda \neq 0$ we obtain: if u is continuous, then the statements, u is of class C'' , $L(u) + \lambda u = 0$, $R_0(u) = 0$, $R_1(u) = 0$, $\lambda \neq 0$, imply and are implied by

$$u(x) = \lambda \int_a^b H(x, t)u(t)dt.$$

This establishes the equivalence of the boundary problem and the integral equation.

87. Flow of Heat in a Ring.—The deductions which we made in the previous problem for a straight bar hold for any linear conductor if we take for the independent variable x the length of arc from a fixed point O . The results also hold if the ring is closed and the two end points A and B coincide. When A and B coincide, the boundary conditions are modified as shown later.



FIG. 26.

Let us suppose $\Theta \equiv 0$, $A \equiv 0$, total length of ring = 1. Then the boundary problem becomes

$$(100) \quad g \frac{\partial \theta}{\partial t} = \frac{\partial}{\partial x} \left(p \frac{\partial \theta}{\partial x} \right) + q\theta$$

$$\theta(0, t) = \theta(1, t); \theta_x(0, t) = \theta_x(1, t)$$

from the continuity of the temperature and of the flow of heat at A . Also

$$\theta(x, 0) = f(x), \text{ initial temperature.}$$

Put $\theta = u(x)\phi(t)$ in (100).

We obtain

$$(101) \quad \frac{d^2u}{dx^2} + (\lambda - b^2)u = 0$$

$$(102) \quad u(0) = u(1), \quad u'(0) = u'(1).$$

Case I.— $\lambda - b^2 > 0$, say $\lambda - b^2 = \mu^2$.

The general solution of (101) is

$$u = \alpha \cos \mu x + \beta \sin \mu x.$$

The boundary conditions (102) give the following two equations for the determination of α, β .

$$(103) \quad \begin{aligned} \alpha(\cos \mu - 1) + \beta \sin \mu &= 0 \\ -\alpha \sin \mu + \beta(\cos \mu - 1) &= 0. \end{aligned}$$

These two equations are compatible for values of α, β not both zero, if and only if

$$\begin{vmatrix} \cos \mu - 1 & \sin \mu \\ -\sin \mu & \cos \mu - 1 \end{vmatrix} = 0.$$

That is, if $2(1 - \cos \mu) = 0$,

whence

$$\mu = 2n\pi.$$

For these values of μ , equations (103) are satisfied by all values of α and β . For other values of μ , the only solution is the trivial one $u \equiv 0$. The characteristic constants are then

$$\lambda_n = b^2 + 4n^2\pi^2, n = 1, 2, \dots,$$

each of index 2.

The corresponding fundamental functions are

$$u = \alpha \cos 2n\pi x + \beta \sin 2n\pi x.$$

Case II.— $\lambda - b^2 = 0$. Then $\lambda = b^2$ is the only characteristic constant. $u = \text{constant}$ is the only solution.

Case III.— $\lambda - b^2 < 0$. Then the only solution is the trivial one $u \equiv 0$. The characteristic constants of the problem are then $\lambda = b^2, \lambda_n = b^2 + 4n^2\pi^2, n = 1, 2, \dots$ with normalized fundamental functions

$$1, \sqrt{2} \cos 2n\pi x, \sqrt{2} \sin 2n\pi x$$

and, since $\int_0^1 \cos 2n\pi x \sin 2n\pi x \, dx = 0$, the complete normalized orthogonal system of fundamental functions is

$$1, \sqrt{2} \cos 2n\pi x, \sqrt{2} \sin 2n\pi x$$

with the characteristic constants

$$b^2, b^2 + 4n^2\pi^2, b^2 + 4n^2\pi^2.$$

This seems to contradict Theorem VII of §75, which stated that the index is always 1. The assumptions of this theorem were, however, that the boundary conditions were of the form

$$\begin{aligned} Au(a) + Bu'(a) &= 0 \\ Cu(b) + Du'(b) &= 0. \end{aligned}$$

The boundary conditions of the above problem are of a more general type:

$$\begin{aligned} A_0 u(a) + B_0 u'(a) + C_0 u(b) + D_0 u'(b) &= 0 \\ A_1 u(a) + B_1 u'(a) + C_1 u(b) + D_1 u'(b) &= 0, \end{aligned}$$

which explains the apparent contradiction.

If $b \neq 0$, (H_2) is satisfied and we can proceed to construct Green's function:

$$K(x, t) = \begin{cases} \frac{\cosh b(t - x - \frac{1}{2})}{2b \sinh \frac{b}{2}}, & x \leq t \\ \frac{\cosh b(x - t - \frac{1}{2})}{2b \sinh \frac{b}{2}}, & x \geq t \end{cases}$$

and the boundary problem is equivalent to the integral equation

$$u(x) = \lambda \int_a^b K(x, t)u(t)dt.$$

The expansion theorem can now be stated as follows: If F is of class C'' , $F(0) = F(1)$ and $F'(0) = F'(1)$ then

$$F(x) = A_0 + \sum (A_n \cos 2n\pi x + B_n \sin 2n\pi x).$$

This is an ordinary Fourier series for the expansion of $F(x)$ for the interval $[0, 1]$.

If $b = 0$, then (H_2) is not satisfied. Then, as in §86, we construct¹ a modified kernel:

$$H(x, t) = \begin{cases} \frac{1}{2}(t - x - \frac{1}{2})^2 + \frac{1}{24}, & x \leq t \\ \frac{1}{2}(x - t - \frac{1}{2})^2 + \frac{1}{24}, & t \leq x. \end{cases}$$

¹See KNESER, "Integralgleichungen," §7.

88. Stationary Flow of Heat Produced by an Interior Source.—We have so far been considering the problem of the flow of heat under the assumption that no heat was produced in the interior of the conductor: $A \equiv 0$.

a) *Solution of the Problem of Stationary Flow of Heat.*—Let us now consider the case $A \not\equiv 0$. The equations of the problem now are

$$\begin{aligned} q \frac{\partial \theta}{\partial t} &= \frac{\partial}{\partial x} \left(p \frac{\partial \theta}{\partial x} \right) + q\theta + A(x) \\ H_0 \theta - \frac{\partial \theta}{\partial x} &= 0 \text{ for } x = a, H_0 \geq 0 \\ H_1 \theta + \frac{\partial \theta}{\partial x} &= 0 \text{ for } x = b, H_1 \geq 0. \end{aligned}$$

Instead of the initial condition $\theta(x, 0) = f(x)$, we require that the flow of heat shall be stationary, that is, independent of the time, that is, $\theta = w(x)$ and, therefore, $\frac{\partial \theta}{\partial t} \equiv 0$.

Then the equations of the problem are

$$(104) \quad \frac{d}{dx} \left(p \frac{dw}{dx} \right) + qw + A(x) = 0 \quad \left[L(w) + A = 0 \right]$$

$$(105) \quad \begin{aligned} R_0(w) &\equiv H_0 w(a) - w'(a) = 0 \\ R_1(w) &\equiv H_1 w(b) + w'(b) = 0. \end{aligned}$$

This is a non-homogeneous boundary problem. From Hilbert's third fundamental theorem we can write down at once the following:

If A is continuous, then the statements, w is of class C'' , $L(w) + A = 0$, $R_0(w) = 0$, $R_1(w) = 0$, imply and are implied by

$$(106) \quad w(x) = \int_a^b K(x, t) A(t) dt.$$

That is, the boundary problem (104) (105) has one and only one solution given by (106). Thus we have the theorem: *Every continuous source of heat $A(x)$ produces one and only one stationary flow of heat expressed by (106).*

b) *Physical Interpretation of Green's Function.*—From (106) Kneser obtains a physical interpretation for Green's function.




FIG. 27.

Take t_0 between a and b and let us suppose that

$$A(x) \begin{cases} = 0 & \text{for } a \leq x \leq t_0 - h, t_0 + h \leq x \leq b \\ \geq 0 & \text{for } t_0 - h \leq x \leq t_0 + h. \end{cases}$$

Then, from the first mean value theorem for definite integrals, there exists a t_1 , $t_0 - h \leq t_1 \leq t_0 + h$ such that

$$w(x, h) = \int_{t_0-h}^{t_0+h} K(x, t)A(t)dt = K(x, t_1) \int_{t_0-h}^{t_0+h} A(t)dt.$$

Now $A(x)dx$ is the quantity of heat produced in an element dx in unit of time, and hence $\int_a^b A(t)dt$ is the total amount of heat produced in the bar in unit time. This is called the *strength of source of heat*.

Let us suppose that the strength of the source of heat is 1:

$$(107) \quad \int_{t_0-h}^{t_0+h} A(t)dt = 1.$$

Let now $h \rightarrow 0$ and let at the same time $A(x)$, which depends upon h , so vary that (107) remains satisfied. Then

$$K(x, t_0) = \lim_{h \rightarrow 0} w(x, h).$$

Thus we have the theorem: *In the general case, Green's function $K(x, t)$ represents the stationary temperature produced by a point source of strength unity placed at $x = t$.*

89. Direct Computation of the Characteristic Constants and Fundamental Functions.—In all of the examples in which we have been able actually to determine the charac-

teristic constants and the fundamental functions, we have determined the latter by means of a boundary problem.

The question arises: How could we directly determine the characteristic constants, and the fundamental functions for an integral equation which does not correspond to a boundary problem? We might obtain the characteristic constants by the solution of the equation $D(\lambda) = 0$, and determine the index and the fundamental functions by means of the Fredholm minors. For some simple kernels this is easily done, but in the general case this method is hardly practicable.

Another direct method has been developed by Schmidt ("Diss.," pages 18 to 21) and Kneser ("Integralgleichungen," pages 190 to 197), at least for a symmetric kernel.

For simplicity, let us suppose that we knew *a priori* that all of the characteristic constants are positive and of index 1.

a) *Determination of λ_1 .*—It was established in the general theory that if $0 < \lambda_1 < \lambda_2 < \dots$ were the characteristic constants for $K(x, t)$ and $\psi_1(x), \psi_2(x), \dots$ the corresponding complete normalized orthogonal system of fundamental functions, then the kernel $K_2(x, t)$ had the same fundamental functions and they belong to the characteristic constants $0 < \lambda_1^2 < \lambda_2^2 < \dots$, for which we write $\mu_1 < \mu_2 < \dots$. Now we have the following expansion for the logarithmic derivative of Fredholm's determinant $D(\lambda)$ derived from the kernel $K(x, t)$:

$$\frac{D'(\lambda)}{D(\lambda)} = - \sum_{n=0}^{\infty} U_{n+1} \lambda^n \quad [(14), §54].$$

where $U_n = \int_a^b K_n(x, x) dx$, convergent for sufficiently small values of λ .

Hence if we call $D_2(\mu)$ the Fredholm determinant for $K_2(x, t)$, we obtain the corresponding expression

$$(108) \quad \frac{D'_2(\mu)}{D_2(\mu)} = - \sum_{n=0}^{\infty} U_{2n+2} \mu^n.$$

Now $\frac{D'_2(\mu)}{D_2(\mu)}$ is a meromorphic function with simple poles, the poles being the roots of $D_2(\mu) = 0$. Arrange the roots $0 < \mu_1 < \mu_2 < \mu_3 < \dots$ in order of magnitude. Hence the circle of convergence of the expression (106) for $\frac{D'_2}{D_2}$ passes through μ_1 and hence the radius of convergence R is equal to μ_1 .

Now there is a theorem¹ on power series to the effect that the radius of convergence R for the series $\sum A_n z^n$ is given by

$$\lim_{n \rightarrow \infty} \frac{A_n}{A_{n+1}} = R,$$

provided this ratio has a determinate limit as $n \rightarrow \infty$.

This theorem applied to the present problem gives

$$\lim_{n \rightarrow \infty} \frac{U_{2n}}{U_{2n+2}} = \mu_1,$$

provided the limit exists. That this ratio has a determinate limit follows from the monotony principle applied to the inequality

$$\frac{U_{2n-2}}{U_{2n}} \geq \frac{U_{2n}}{U_{2n+2}} > 0$$

which we proved in §58.

b) *Computation of the Fundamental Function $\psi_1(x)$.*—From

$$K_{2n+2}(x, y) = \int_a^b K_2(x, t) K_{2n}(t, y) dt$$

we obtain

$$(109) \quad \mu_1^{n+1} K_{2n+2}(x, y) = \mu_1 \int_a^b K_2(x, t) \left(\mu_1^n K_{2n}(t, y) \right) dt.$$

Now Schmidt proves in his existence proof for the characteristic constants that

$$\lim_{n \rightarrow \infty} \mu_1^n K_{2n}(x, y) = f(x, y)$$

¹ HARKNESS and MORLEY, "Theory of Functions," §76.

a definite limit function which is approached uniformly with respect to x and y in the region R . Therefore, by passing to the limit in (109), we find

$$f(x, y) = \mu_1 \int_a^b K_2(x, t)f(t, y)dt.$$

Further Schmidt proves that $f(x, x) \not\equiv 0$. Hence, if for some quantity c we have $f(c, c) \not\equiv 0$, then the function $f(x, c) = \varphi(x) \not\equiv 0$ and, therefore, $\varphi(x)$ is a fundamental function belonging to μ_1 since it satisfies the equation.

This function $\varphi(x)$, normalized in the usual way, gives us the fundamental function $\psi_1(x)$ belonging to λ_1 .

c) Computation of the Other Characteristic Constants and Fundamental Functions.—The other characteristic constants and fundamental functions can be obtained successively as follows. Let

$$(110) \quad K_2'(x, t) = K_2(x, t) - \frac{\psi_1(x)\psi_1(t)}{\mu_1}$$

be a new kernel, then we can show that its characteristic constants and fundamental functions are

$$\lambda_2^2 < \lambda_3^2 < \dots ; \psi_2(x), \psi_3(x), \dots .$$

For from (110) we obtain

$$(111) \quad \int_a^b K_2'(x, t)\psi_\nu(t)dt = \int_a^b K_2(x, t)\psi_\nu(t)dt + \frac{\psi_1(x)}{\mu_1} \delta_{1\nu},$$

where $\delta_{1\nu}$ is the Kronecker symbol. When $\nu = 1$, (111) gives

$$\int_a^b K_2'(x, t)\psi_1(t)dt = 0.$$

When $\nu \neq 1$, (111) gives

$$\psi_\nu(x) = \mu_\nu \int_a^b K_2'(x, t)\psi_\nu(t)dt,$$

which proves the above statement.

Applying now to the kernel $K_2'(x, t)$ the method described under *a*) and *b*), we obtain λ_2^2 and $\psi_2(x)$, and so on.

INDEX

A

- Abel's problem, 2
Associated equation, 70
fundamental functions of, 60
homogeneous, 56
Associated kernel, determinant
for, 57
minors for, 58
the function $H(x, y)$ for, 61

B

- Baltzer, 30, 31
Bar, flow of heat in, 229
Bessel's inequality, 149, 160
function, 217
Bilinear formula, 141, 144, 152
Bôcher, 2, 20, 69, 207

C

- Calculus of variations, 192
Cauchy, 5
Characteristic constants, 7, 56,
60, 183, 185, 186, 202, 211,
212, 235, 244, 247
reality of, 129
Conjugate point, 208
Cramer, 25, 69
Curve, smooth, 85

D

- Derivative, directional, 87
Dirichlet's problem, 100, 196, 205
solution of, 105

E

- Elastic string, free vibrations of,
73
constrained vibrations of, 81
differential equation of, 73
reduction to one-dimensional
boundary problem, 74
solution of boundary problem,
74
Equations, differential, relation
with integral equation, 4
homogeneous, solution of when
 $D(\lambda) = 0, D'(\lambda) \neq 0$, 42,
43, 45
completeness proof of solu-
tions of, 54
 q independent solutions of,
52, 55
solution when $D(\lambda) = 0$, 46
integral (see Integral equation).
system of linear, replacing
integral equation, 23
Euler's differential equation, 193
Euler's rule for isoperimetric
problems, 195

F

- Fourier series, 212. 245
Fréchet, 22, 46
Fredholm, 7, 100
determinant, 25, 38
convergence proof, 32
equation, 9, 69
Volterra's solution of, 19

Fredholm, first fundamental relation, 27, 34
 generalization of, 47, 49
 first fundamental theorem, 41
 first minor, 26, 38
 convergence proof, 33
 minors for the associated kernel, 58
 second fundamental relation, 28, 34
 generalization of, 47, 49
 second fundamental theorem, 55
 theory, application of, 73
 third fundamental theorem, 66
Functions, iterated, 17
 reciprocal, 17, 19
Fundamental functions, 7, 56, 247
 normalized, 132, 136, 137, 138,
 141, 183, 212, 235, 237, 244
 of the associated equation, 60
Fundamental numbers, 7

G

Green's formula, 172, 217
 function, 76, 174, 176, 187,
 191, 201, 211, 218, 228,
 234, 237, 241, 245
 physical interpretation of,
 247
 theorem, 86, 241

H

Hadamard's theorem, 28, 31, 32
Harmonic functions, 85
Heywood-Frèchet, 22, 46
Hilbert, 7, 116, 189, 191, 196,
 204, 240
Hilbert's fundamental theorem,
 176
 third fundamental theorem,
 180, 187, 191, 199

Homogeneous integral equation,
 3
 associated, 56
 completeness proof of solutions
 of, 54
 q independent solutions of, 52,
 55
 solution when $D(\lambda) = 0, D'(\lambda)$
 $\neq 0$, 42, 43, 45
 solution when $D(\lambda) = 0$, 46
Horn, 96, 98

I

Index, 212, 244
Index of λ_0 , 52, 131
Integral equation, linear, first
 kind, 1
 homogeneous, 3
 more than one variable, 6
 non-homogeneous, when $D(\lambda)$
 = 0, 62
 non-linear, 5
 Schmidt's solution, 162, 166
 second kind, 3
 Volterra, 3, 5
 singular, 6
 third kind, 4
 with a parameter, 3
Iterated functions, 17

J

Jacobi's condition, 207

K

Kernel, 1
 closed, 187
 expansion of, 141
 symmetric, 117, 127, 131, 137,
 145, 234
 characteristic constants of,
 129

Kernel, symmetric, iterations of,
123
symmetrizable, 190
Kneser, 118, 121, 141, 142, 192,
215, 222, 245, 247, 248
Kowalewski, 117, 155

L

Lalesco, 5, 189
Liouville, 7

M

Marty, 189, 190
Minors, Fredholm first, 26, 38
for the associated kernels, 58
 p -th minors, 46
Multiplicity of λ_0 , 51, 131

N

Neumann, 7
Neumann's problem, 110
solution of, 114

O

Orthogonality theorem, 61, 128

P

Pierpont, 44
Potential, logarithmic, of a
double layer, 92
of a simple layer, 107

R

Reciprocal functions, 17
Ring, flow of heat in, 243
Rope, rotating, 222

S

Schmidt, 7, 116, 162
auxiliary theorem of, 155
Schwarz's inequality, 125, 159
application of, 126
Solutions, discontinuous, 20
Fredholm's solution when $D(\lambda)$
 $\neq 0$, 39
of homogeneous equation, 42
types of, 7
uniqueness of, 16
String, elastic, constrained vibra-
tions of, 81
free vibrations of, 73
vibrating, 138, 209
Sturm's oscillation theorem, 207
Successive approximations, 15
substitutions, 9

V

Vibrations, of a rope, 215
of a string, 209
Volterra, 3, 7
equation, 13
solution of Fredholm's equa-
tion, 19

W

Weber, 209, 229

Catalogue of Dover SCIENCE BOOKS

DIFFERENTIAL EQUATIONS (ORDINARY AND PARTIAL DIFFERENTIAL)

INTRODUCTION TO THE DIFFERENTIAL EQUATIONS OF PHYSICS, L. Hopf. Especially valuable to engineer with no math beyond elementary calculus. Emphasizes intuitive rather than formal aspects of concepts. Partial contents: Law of causality, energy theorem, damped oscillations, coupling by friction, cylindrical and spherical coordinates, heat source, etc. 48 figures. 160pp. 5 $\frac{1}{2}$ x 8. \$120 Paperbound \$1.25

INTRODUCTION TO BESSEL FUNCTIONS, F. Bowman. Rigorous, provides all necessary material during development, includes practical applications. Bessel functions of zero order, of any real order, definite integrals, asymptotic expansion, circular membranes, Bessel's solution to Kepler's problem, much more. "Clear . . . useful not only to students of physics and engineering, but to mathematical students in general," Nature. 226 problems. Short tables of Bessel functions. 27 figures. x + 135pp. 5 $\frac{1}{2}$ x 8. \$462 Paperbound \$1.35

DIFFERENTIAL EQUATIONS, F. R. Moulton. Detailed, rigorous exposition of all non-elementary processes of solving ordinary differential equations. Chapters on practical problems; more advanced than problems usually given as illustrations. Includes analytic differential equations; variations of a parameter; integrals of differential equations; analytic implicit functions; problems of elliptic motion; sine-amplitude functions; deviation of formal bodies; Cauchy-Lipshitz process; linear differential equations with periodic coefficients; much more. Historical notes. 10 figures. 222 problems. xv + 395pp. 5 $\frac{1}{2}$ x 8. \$451 Paperbound \$2.00

PARTIAL DIFFERENTIAL EQUATIONS OF MATHEMATICAL PHYSICS, A. G. Webster. Valuable sections on elasticity, compression theory, potential theory, theory of sound, heat conduction, wave propagation, vibration theory. Contents include: deduction of differential equations, vibrations, normal functions, Fourier's series. Cauchy's method, boundary problems, method of Riemann-Volterra, spherical, cylindrical, ellipsoidal harmonics, applications, etc. 97 figures. vii + 440pp. 5 $\frac{1}{2}$ x 8. \$263 Paperbound \$2.00

ORDINARY DIFFERENTIAL EQUATIONS, E. L. Ince. A most compendious analysis in real and complex domains. Existence and nature of solutions, continuous transformation groups, solutions in an infinite form, definite integrals, algebraic theory. Sturmian theory, boundary problems, existence theorems, 1st order, higher order, etc. "Deserves highest praise, a notable addition to mathematical literature," Bulletin, Amer. Math. Soc. Historical appendix. 18 figures. viii + 558pp. 5 $\frac{1}{2}$ x 8. \$349 Paperbound \$2.55

ASYMPTOTIC EXPANSIONS, A. Erdélyi. Only modern work available in English; unabridged reproduction of monograph prepared for Office of Naval Research. Discusses various procedures for asymptotic evaluation of integrals containing a large parameter; solutions of ordinary linear differential equations. vi + 108pp. 5 $\frac{1}{2}$ x 8. \$318 Paperbound \$1.35

LECTURES ON CAUCHY'S PROBLEM, J. Hadamard. Based on lectures given at Columbia, Rome, discusses work of Riemann, Kirchhoff, Volterra, and author's own research on hyperbolic case in linear partial differential equations. Extends spherical cylindrical waves to apply to all (normal) hyperbolic equations. Partial contents: Cauchy's problem, fundamental formula, equations with odd number, with even number of independent variables; method of descent. 32 figures. iii + 316pp. 5 $\frac{1}{2}$ x 8. \$105 Paperbound \$1.75

CATALOGUE OF

PRINCIPLES OF MECHANICS, Heinrich Hertz. A classic of great interest in logic of science. Last work by great 19th century physicist, created new system of mechanics based upon space, time, mass; returns to axiomatic analysis, understanding of formal, structural aspects of science, taking into account logic, observation, a priori elements. Of great historical importance to Poincaré, Carnap, Einstein, Milne. 20 page introduction by R. S. Cohen, Wesleyan U., analyzes implications of Hertz's thought and logic of science. 13 page introduction by Helmholtz. xlii + 274pp. 5 1/2 x 8.

S316 Clothbound \$3.50
S317 Paperbound \$1.75

MATHEMATICAL FOUNDATIONS OF STATISTICAL MECHANICS, A. I. Khinchin. A thoroughly up-to-date introduction, offering a precise and mathematically rigorous formulation of the problems of statistical mechanics. Provides analytical tools to replace many commonly used cumbersome concepts and devices. Partial contents: Geometry, kinematics of phase space; ergodic problem; theory of probability; central limit theorem; ideal monatomic gas; foundation of thermodynamics; dispersion, distribution of sum functions; etc. "Excellent introduction . . . clear, concise, rigorous," Quarterly of Applied Mathematics. viii + 179pp. 5 1/2 x 8.

S146 Clothbound \$2.95
S147 Paperbound \$1.35

MECHANICS OF THE GYROSCOPE, THE DYNAMICS OF ROTATION, R. F. Deimel, Prof. of Mechanical Engineering, Stevens Inst. of Tech. Elementary, general treatment of dynamics of rotation, with special application of gyroscopic phenomena. No knowledge of vectors needed. Velocity of a moving curve, acceleration to a point, general equations of motion, gyroscopic horizon, free gyro, motion of discs, the damped gyro, 103 similar topics. Exercises. 75 figures. 208pp. 5 1/2 x 8.

S66 Paperbound \$1.65

MECHANICS VIA THE CALCULUS, P. W. Norris, W. S. Legge. Wide coverage, from linear motion to vector analysis; equations determining motion, linear methods, compounding of simple harmonic motions, Newton's laws of motion, Hooke's law, the simple pendulum, motion of a particle in 1 plane, centers of gravity, virtual work, friction, kinetic energy of rotating bodies, equilibrium of strings, hydrostatics, shearing stresses, elasticity, etc. Many worked-out examples. 550 problems. 3rd revised edition. xii + 367pp.

S207 Clothbound \$3.95

A TREATISE ON THE MATHEMATICAL THEORY OF ELASTICITY, A. E. H. Love. An indispensable reference work for engineers, mathematicians, physicists, the most complete, authoritative treatment of classical elasticity in one volume. Proceeds from elementary notions of extension to types of strain, cubical dilatation, general theory of strains. Covers relation between mathematical theory of elasticity and technical mechanics; equilibrium of isotropic elastic solids and aelotropic solid bodies; nature of force transmission, Volterra's theory of dislocations; theory of elastic spheres in relation to tidal, rotational, gravitational effects on earth; general theory of bending; deformation of curved plates; buckling effects; much more. "The standard treatise on elasticity," American Math. Monthly. 4th revised edition. 76 figures. xviii + 643pp. 6 1/8 x 9 1/4.

S174 Paperbound \$2.95

NUCLEAR PHYSICS, QUANTUM THEORY, RELATIVITY

MESON PHYSICS, R. E. Marshak. Presents basic theory, and results of experiments with emphasis on theoretical significance. Phenomena involving mesons as virtual transitions avoided, eliminating some of least satisfactory predictions of meson theory. Includes production study of π mesons at nonrelativistic nucleon energies contracts between π and μ mesons, phenomena associated with nuclear interaction of π mesons, etc. Presents early evidence for new classes of particles, indicates theoretical difficulties created by discovery of heavy mesons and hyperons. viii + 378pp. 5 1/2 x 8.

S500 Paperbound \$1.95

THE FUNDAMENTAL PRINCIPLES OF QUANTUM MECHANICS, WITH ELEMENTARY APPLICATIONS, E. C. Kemble. Inductive presentation, for graduate student, specialists in other branches of physics. Apparatus necessary beyond differential equations and advanced calculus developed as needed. Though general exposition of principles, hundreds of individual problems fully treated. "Excellent book . . . of great value to every student . . . rigorous and detailed mathematical discussion . . . has succeeded in keeping his presentation clear and understandable," Dr. Linus Pauling, J. of American Chemical Society. Appendices: calculus of variations, math. notes, etc. 611pp. 5 1/2 x 8 1/2.

T472 Paperbound \$2.95

WAVE PROPAGATION IN PERIODIC STRUCTURES, L. Brillouin. General method, application to different problems: pure physics—scattering of X-rays in crystals, thermal vibration in crystal lattices, electronic motion in metals; problems in electrical engineering. Partial contents: elastic waves along 1-dimensional lattices of point masses. Propagation of waves along 1-dimensional lattices. Energy flow. 2, 3 dimensional lattices. Mathieu's equation. Matrices and propagation of waves along an electric line. Continuous electric lines. 131 illustrations. xii + 253pp. 5 1/2 x 8.

S34 Paperbound \$1.85

DOVER SCIENCE BOOKS

THEORY OF ELECTRONS AND ITS APPLICATION TO THE PHENOMENA OF LIGHT AND RADIANT HEAT, H. Lorentz. Lectures delivered at Columbia Univ., by Nobel laureate. Unabridged, form historical coverage of theory of free electrons, motion, absorption of heat, Zeeman effect, optical phenomena in moving bodies, etc. 109 pages notes explain more advanced sections. 9 figures. 352pp. 5 $\frac{3}{8}$ x 8. S173 Paperbound \$1.85

SELECTED PAPERS ON QUANTUM ELECTRODYNAMICS, edited by J. Schwinger. Facsimiles of papers which established quantum electrodynamics; beginning to present position as part of larger theory. First book publication in any language of collected papers of Bethe, Bloch, Dirac, Dyson, Fermi, Feynman, Heisenberg, Kusch, Lamb, Oppenheimer, Pauli, Schwinger, Tomonaga, Weisskopf, Wigner, etc. 34 papers: 29 in English, 1 in French, 3 in German, 1 in Italian. Historical commentary by editor. xvii + 423pp. 6 $\frac{1}{8}$ x 9 $\frac{1}{4}$. S444 Paperbound \$2.45

FOUNDATIONS OF NUCLEAR PHYSICS, edited by R. T. Beyer. 13 of the most important papers on nuclear physics reproduced in facsimile in the original languages; the papers most often cited in footnotes, bibliographies. Anderson, Curie, Joliot, Chadwick, Fermi, Lawrence, Cockcroft, Hahn, Yukawa. Unparalleled bibliography: 122 double columned pages, over 4,000 articles, books, classified. 57 figures. 288pp. 6 $\frac{1}{8}$ x 9 $\frac{1}{4}$. S19 Paperbound \$1.75

THE THEORY OF GROUPS AND QUANTUM MECHANICS, H. Weyl. Schroedinger's wave equation, de Broglie's waves of a particle, Jordan-Hoelder theorem, Lie's continuous groups of transformations, Pauli exclusion principle, quantization of Maxwell-Dirac field equations, etc. Unitary geometry, quantum theory, groups, application of groups to quantum mechanics, symmetry permutation group, algebra of symmetric transformations, etc. 2nd revised edition. xxii + 422pp. 5 $\frac{3}{8}$ x 8. S268 Clothbound \$4.50
S269 Paperbound \$1.95

PHYSICAL PRINCIPLES OF THE QUANTUM THEORY, Werner Heisenberg. Nobel laureate discusses quantum theory; his own work, Compton, Schroedinger, Wilson, Einstein, many others. For physicists, chemists, not specialists in quantum theory. Only elementary formulae considered in text; mathematical appendix for specialists. Profound without sacrificing clarity. Translated by C. Eckart, F. Hoyt. 18 figures. 192pp. 5 $\frac{3}{8}$ x 8. S113 Paperbound \$1.25

INVESTIGATIONS ON THE THEORY OF THE BROWNIAN MOVEMENT, Albert Einstein. Reprints from rare European journals, translated into English. 5 basic papers, including Elementary Theory of the Brownian Movement, written at request of Lorentz to provide a simple explanation. Translated by A. D. Cowper. Annotated, edited by R. Fürth. 33pp. of notes elucidate, give history of previous investigations. 62 footnotes. 124pp. 5 $\frac{3}{8}$ x 8. S304 Paperbound \$1.25

THE PRINCIPLE OF RELATIVITY, E. Einstein, H. Lorentz, M. Minkowski, H. Weyl. The 11 basic papers that founded the general and special theories of relativity, translated into English. 2 papers by Lorentz on the Michelson experiment, electromagnetic phenomena. Minkowski's "Space and Time," and Weyl's "Gravitation and Electricity." 7 epoch-making papers by Einstein: "Electromagnetics of Moving Bodies," "Influence of Gravitation in Propagation of Light," "Cosmological Considerations," "General Theory," 3 others. 7 diagrams. Special notes by A. Sommerfeld. 224pp. 5 $\frac{3}{8}$ x 8. S93 Paperbound \$1.75

STATISTICS

ELEMENTARY STATISTICS, WITH APPLICATIONS IN MEDICINE AND THE BIOLOGICAL SCIENCES, F. E. Croxton. Based primarily on biological sciences, but can be used by anyone desiring introduction to statistics. Assumes no prior acquaintance, requires only modest knowledge of math. All basic formulas carefully explained, illustrated; all necessary reference tables included. From basic terms and concepts, proceeds to frequency distribution, linear, non-linear, multiple correlation, etc. Contains concrete examples from medicine, biology. 101 charts. 57 tables. 14 appendices. iv + 376pp. 5 $\frac{3}{8}$ x 8. S506 Paperbound \$1.95

ANALYSIS AND DESIGN OF EXPERIMENTS, H. B. Mann. Offers method for grasping analysis of variance, variance design quickly. Partial contents: Chi-square distribution, analysis of variance distribution, matrices, quadratic forms, likelihood ratio tests, test of linear hypotheses, power of analysis, Galois fields, non-orthogonal data, interblock estimates, etc. 15pp. of useful tables. x + 195pp. 5 x 7 $\frac{1}{2}$. S180 Paperbound \$1.45

FREQUENCY CURVES AND CORRELATION, W. P. Elderton. 4th revised edition of standard work on classical statistics. Practical, one of few books constantly referred to for clear presentation of basic material. Partial contents: Frequency Distributions; Pearson's Frequency Curves; Theoretical Distributions; Standard Errors; Correlation Ratio—Contingency; Corrections for Moments, Beta, Gamma Functions; etc. Key to terms, symbols. 25 examples. 40 tables. 16 figures. xi + 272pp. 5 $\frac{1}{2}$ x 8 $\frac{1}{2}$. Clothbound \$1.49

HYDRODYNAMICS, ETC.

HYDRODYNAMICS, Horace Lamb. Standard reference work on dynamics of liquids and gases. Fundamental theorems, equations, methods, solutions, background for classical hydrodynamics. Chapters: Equations of Motion, Integration of Equations in Special Gases, Vortex Motion, Tidal Waves, Rotating Masses of Liquids, etc. Excellently planned, arranged, Clear, lucid presentation. 6th enlarged, revised edition. Over 900 footnotes, mostly bibliographical. 119 figures. xv + 738pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$. S256 Paperbound \$2.95

HYDRODYNAMICS, A STUDY OF LOGIC, FACT, AND SIMILITUDE, Garrett Birkhoff. A stimulating application of pure mathematics to an applied problem. Emphasis is on correlation of theory and deduction with experiment. Examines recently discovered paradoxes, theory of modelling and dimensional analysis, paradox and error in flows and free boundary theory. Classical theory of virtual mass derived from homogenous spaces; group theory applied to fluid mechanics. 20 figures, 3 plates. xiii + 186pp. 5 $\frac{1}{2}$ x 8. S22 Paperbound \$1.85

HYDRODYNAMICS, H. Dryden, F. Murhaghian, H. Bateman. Published by National Research Council, 1932. Complete coverage of classical hydrodynamics, encyclopedic in quality. Partial contents: physics of fluids, motion, turbulent flow, compressible fluids, motion in 1, 2, 3 dimensions; laminar motion, resistance of motion through viscous fluid, eddy viscosity, discharge of gases, flow past obstacles, etc. Over 2900-item bibliography. 23 figures. 634pp. 5 $\frac{1}{2}$ x 8. S303 Paperbound \$2.75

ACOUSTICS AND OPTICS

PRINCIPLES OF PHYSICAL OPTICS, Ernst Mach. Classical examination of propagation of light, color, polarization, etc. Historical, philosophical treatment unequalled for breadth and readability. Contents: Rectilinear propagation, reflection, refraction, dioptrics, composition of light, periodicity, theory of interference, polarization, mathematical representation of properties, etc. 279 illustrations. 10 portraits. 324pp. 5 $\frac{1}{2}$ x 8. S170 Paperbound \$1.75

THE THEORY OF SOUND, Lord Rayleigh. Written by Nobel laureate, classical methods here will cover most vibrating systems likely to be encountered in practice. Complete coverage of experimental, mathematical aspects. Partial contents: Harmonic motions, lateral vibrations of bars, curved plates or shells, applications of Laplace's functions to acoustical problems, fluid friction, etc. First low-priced edition of this great reference-study work. Historical introduction by R. B. Lindsay. 1040pp. 97 figures. 5 $\frac{1}{2}$ x 8. S292, S293, Two volume set, paperbound \$4.00

THEORY OF VIBRATIONS, N. W. McLachlan. Based on exceptionally successful graduate course, Brown University. Discusses linear systems having 1 degree of freedom, forced vibrations of simple linear systems, vibration of flexible strings, transverse vibrations of bars and tubes, of circular plate, sound waves of finite amplitude, etc. 99 diagrams. 160pp. 5 $\frac{1}{2}$ x 8. S190 Paperbound \$1.35

APPLIED OPTICS AND OPTICAL DESIGN, A. E. Conrady. Thorough systematic presentation of physical and mathematical aspects, limited mostly to "real optics." Stresses practical problem of maximum aberration permissible without affecting performance. Ordinary ray tracing methods; complete theory ray tracing methods, primary aberrations; enough higher aberration to design telescopes, low powered microscopes, photographic equipment. Covers fundamental equations, extra-axial image points, transverse chromatic aberration, angular magnification, similar topics. Tables of functions of N. Over 150 diagrams. x + 518pp. 5 $\frac{1}{2}$ x 8 $\frac{1}{2}$. S366 Paperbound \$2.98

RAYLEIGH'S PRINCIPLE AND ITS APPLICATIONS TO ENGINEERING, G. Temple, W. Bickley. Rayleigh's principle developed to provide upper, lower estimates of true value of fundamental period of vibrating system, or condition of stability of elastic system. Examples, rigorous proofs. Partial contents: Energy method of discussing vibrations, stability. Perturbation theory, whirling of uniform shafts. Proof, accuracy, successive approximations, applications of Rayleigh's theory. Numerical, graphical methods. Ritz's method. 22 figures. ix + 156pp. 5 $\frac{1}{2}$ x 8. S307 Paperbound \$1.50

OPTICKS, Sir Isaac Newton. In its discussion of light, reflection, color, refraction, theories of wave and corpuscular theories of light, this work is packed with scores of insights and discoveries. In its precise and practical discussions of construction of optical apparatus, contemporary understanding of phenomena, it is truly fascinating to modern scientists. Foreword by Albert Einstein. Preface by I. B. Cohen, Harvard. 7 pages of portraits, facsimile pages, letters, etc. cxvi + 414pp. 5 $\frac{1}{2}$ x 8. S205 Paperbound \$2.00

DOVER SCIENCE BOOKS

ON THE SENSATIONS OF TONE, Hermann Helmholtz. Using acoustical physics, physiology, experiment, history of music, covers entire gamut of musical tone: relation of music science to acoustics, physical vs. physiological acoustics, vibration, resonance, tonality, progression of parts, etc. 33 appendices on various aspects of sound, physics, acoustics, music, etc. Translated by A. J. Ellis. New introduction by H. Margenau, Yale. 68 figures. 43 musical passages analyzed. Over 100 tables. xix + 576pp. 6 1/4 x 9 1/4.

S114 Clothbound \$4.95

ELECTROMAGNETICS, ENGINEERING, TECHNOLOGY

INTRODUCTION TO RELAXATION METHODS, F. S. Shaw. Describes almost all manipulative resources of value in solution of differential equations. Treatment is mathematical rather than physical. Extends general computational process to include almost all branches of applied math and physics. Approximate numerical methods are demonstrated, although high accuracy is obtainable without undue expenditure of time. 48pp. of tables for computing irregular star first and second derivatives, irregular star coefficients for second order equations, for fourth order equations. "Useful. . . exposition is clear, simple . . . no previous acquaintance with numerical methods is assumed," Science Progress. 253 diagrams. 72 tables. 400pp. 5 3/8 x 8.

S244 Paperbound \$2.45

THE ELECTROMAGNETIC FIELD, M. Mason, W. Weaver. Used constantly by graduate engineers. Vector methods exclusively; detailed treatment of electrostatics, expansion methods, with tables converting any quantity into absolute electromagnetic, absolute electrostatic, practical units. Discrete charges, ponderable bodies. Maxwell field equations, etc. 416pp. 5 3/8 x 8.

S185 Paperbound \$2.00

ELASTICITY, PLASTICITY AND STRUCTURE OF MATTER, R. Houwink. Standard treatise on rheological aspects of different technically important solids: crystals, resins, textiles, rubber, clay, etc. Investigates general laws for deformations; determines divergences. Covers general physical and mathematical aspects of plasticity, elasticity, viscosity. Detailed examination of deformations, internal structure of matter in relation to elastic, plastic behaviour, formation of solid matter from a fluid, etc. Treats glass, asphalt, balata, proteins, baker's dough, others. 2nd revised, enlarged edition. Extensive revised bibliography in over 500 footnotes. 214 figures. xvii + 368pp. 6 x 9 1/4.

S385 Paperbound \$2.45

DESIGN AND USE OF INSTRUMENTS AND ACCURATE MECHANISM, T. N. Whitehead. For the instrument designer, engineer; how to combine necessary mathematical abstractions with independent observations of actual facts. Partial contents: instruments and their parts, theory of errors, systematic errors, probability, short period errors, erratic errors, design precision, kinematic, semikinematic design, stiffness, planning of an instrument, human factor, etc. 85 photos, diagrams. xii + 288pp. 5 3/8 x 8.

S270 Paperbound \$1.95

APPLIED HYDRO- AND AEROMECHANICS, L. Prandtl, O. G. Tietjens. Presents, for most part, methods valuable to engineers. Flow in pipes, boundary layers, airfoil theory, entry conditions, turbulent flow, boundary layer, determining drag from pressure and velocity, etc. "Will be welcomed by all students of aerodynamics," Nature. Unabridged, unaltered. An Engineering Society Monograph, 1934. Index. 226 figures. 28 photographic plates illustrating flow patterns. xvi + 311pp. 5 3/8 x 8.

S375 Paperbound \$1.85

FUNDAMENTALS OF HYDRO- AND AEROMECHANICS, L. Prandtl, O. G. Tietjens. Standard work, based on Prandtl's lectures at Goettingen. Wherever possible hydrodynamics theory is referred to practical considerations in hydraulics, unifying theory and experience. Presentation extremely clear. Though primarily physical, proofs are rigorous and use vector analysis to a great extent. An Engineering Society Monograph, 1934. "Still recommended as an excellent introduction to this area," Physikalische Blätter. 186 figures. xvi + 270pp. 5 3/8 x 8.

S374 Paperbound \$1.85

GASEOUS CONDUCTORS: THEORY AND ENGINEERING APPLICATIONS, J. D. Cobine. Indispensable text, reference, to gaseous conduction phenomena, with engineering viewpoint prevailing throughout. Studies kinetic theory of gases, ionization, emission phenomena; gas breakdown, spark characteristics, glow, discharges; engineering applications in circuit interrupters, rectifiers, etc. Detailed treatment of high pressure arcs (Suits); low pressure arcs (Langmuir, Tonks). Much more. "Well organized, clear, straightforward," Tonks, Review of Scientific Instruments. 83 practice problems. Over 600 figures. 58 tables. xx + 606pp. 5 3/8 x 8.

S442 Paperbound \$2.75

PHOTOELASTICITY: PRINCIPLES AND METHODS, H. T. Jessop, F. C. Harris. For engineer, specific problems of stress analysis. Latest time-saving methods of checking calculations in 2-dimensional design problems, new techniques for stresses in 3 dimensions, lucid description of optical systems used in practical photoelectricity. Useful suggestions, hints based on on-the-job experience included. Partial contents: strain, stress-strain relations, circular disc under thrust along diameter, rectangular block with square hold under vertical thrust, simply supported rectangular beam under central concentrated load, etc. Theory held to minimum, no advanced mathematical training needed. 164 illustrations. viii + 184pp. 6 1/4 x 9 1/4.

S137 Clothbound \$3.75

CATALOGUE OF

MICROWAVE TRANSMISSION DESIGN DATA, T. Moreno. Originally classified, now rewritten, enlarged (14 new chapters) under auspices of Sperry Corp. Of immediate value or reference use to radio engineers, systems designers, applied physicists, etc. Ordinary transmission line theory; attenuation; parameters of coaxial lines; flexible cables; tuneable wave guide impedance transformers; effects of temperature, humidity; much more. "Packed with information . . . theoretical discussions are directly related to practical questions," U. of Royal Naval Scientific Service. Tables of dielectrics, flexible cable, etc. ix + 248pp. 5 $\frac{1}{2}$ x 8.
S549 Paperbound \$1.50

THE THEORY OF THE PROPERTIES OF METALS AND ALLOYS, H. F. Mott, H. Jones. Quantum methods develop mathematical models showing interrelationship of fundamental chemical phenomena with crystal structure, electrical, optical properties, etc. Examines electron motion in applied field, cohesion, heat capacity, refraction, noble metals, transition and di-valent metals, etc. "Exposition is as clear . . . mathematical treatment as simple and reliable as we have become used to expect of . . . Prof. Mott," Nature. 138 figures. xiii + 320pp. 5 $\frac{1}{2}$ x 8.
S456 Paperbound \$1.85

THE MEASUREMENT OF POWER SPECTRA FROM THE POINT OF VIEW OF COMMUNICATIONS ENGINEERING, R. B. Blackman, J. W. Tukey. Pathfinding work reprinted from "Bell System Technical Journal." Various ways of getting practically useful answers in power spectra measurement, using results from both transmission and statistical estimation theory. Treats: Autocovariance, Functions and Power Spectra, Distortion, Heterodyne Filtering, Smoothing, Decimation Procedures, Transversal Filtering, much more. Appendix reviews fundamental Fourier techniques. Index of notation. Glossary of terms. 24 figures. 12 tables. 192pp. 5 $\frac{1}{2}$ x 8 $\frac{1}{2}$.
S507 Paperbound \$1.85

TREATISE ON ELECTRICITY AND MAGNETISM, James Clerk Maxwell. For more than 80 years a seemingly inexhaustible source of leads for physicists, mathematicians, engineers. Total of 1082pp. on such topics as Measurement of Quantities, Electrostatics, Elementary Mathematical Theory of Electricity, Electrical Work and Energy in a System of Conductors, General Theorems, Theory of Electrical Images, Electrolysis, Conduction, Polarization, Dielectrics, Resistance, much more. "The greatest mathematical physicist since Newton," Sir James Jeans. 3rd edition. 107 figures, 21 plates. 1082pp. 5 $\frac{1}{2}$ x 8.
S186 Clothbound \$4.95

CHEMISTRY AND PHYSICAL CHEMISTRY

THE PHASE RULE AND ITS APPLICATIONS, Alexander Findlay. Covers chemical phenomena of 1 to 4 multiple component systems, the "standard work on the subject" (Nature). Completely revised, brought up to date by A. N. Campbell, N. O. Smith. New material on binary, tertiary liquid equilibria, solid solutions in ternary systems, quinary systems of salts, water, etc. Completely revised to triangular coordinates in ternary systems, clarified graphic representation, solid models, etc. 9th revised edition. 236 figures. 505 footnotes, mostly bibliographic. xii + 449pp. 5 $\frac{1}{2}$ x 8.
S92 Paperbound \$2.45

DYNAMICAL THEORY OF GASES, James Jeans. Divided into mathematical, physical chapters for convenience of those not expert in mathematics. Discusses mathematical theory of gas in steady state, thermodynamics, Boltzmann, Maxwell, kinetic theory, quantum theory, exponentials, etc. "One of the classics of scientific writing . . . as lucid and comprehensive an exposition of the kinetic theory as has ever been written," J. of Institute of Engineers. 4th enlarged edition, with new material on quantum theory, quantum dynamics, etc. 28 figures. 444pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$.
S136 Paperbound \$2.45

POLAR MOLECULES, Pieter Debye. Nobel laureate offers complete guide to fundamental electrostatic field relations, polarizability, molecular structure. Partial contents: electric intensity, displacement, force, polarization by orientation, molar polarization, molar refraction, halogen-hydrides, polar liquids, ionic saturation, dielectric constant, etc. Special chapter considers quantum theory. "Clear and concise . . . coordination of experimental results with theory will be readily appreciated," Electronics Industries. 172pp. 5 $\frac{1}{2}$ x 8.
S63 Clothbound \$3.50
S64 Paperbound \$1.50

ATOMIC SPECTRA AND ATOMIC STRUCTURE, G. Herzberg. Excellent general survey for chemists, physicists specializing in other fields. Partial contents: simplest line spectra, elements of atomic theory; multiple structure of line spectra, electron spin; building-up principle, periodic system of elements; finer details of atomic spectra; hyperfine structure of spectral lines; some experimental results and applications. 80 figures. 20 tables. xiii + 257pp. 5 $\frac{1}{2}$ x 8.
S115 Paperbound \$1.95

TREATISE ON THERMODYNAMICS, Max Planck. Classic based on his original papers. Brilliant concepts of Nobel laureate make no assumptions regarding nature of heat, rejects earlier approaches of Helmholtz, Maxwell, to offer uniform point of view for entire field. Seminal work by founder of quantum theory, deducing new physical, chemical laws. A standard text, an excellent introduction to field for students with knowledge of elementary chemistry, physics, calculus. 3rd English edition. xvi + 297pp. 5 $\frac{1}{2}$ x 8.
S219 Paperbound \$1.75

DOVER SCIENCE BOOKS

KINETIC THEORY OF LIQUIDS, J. Frenkel. Regards kinetic theory of liquids as generalization, extension of theory of solid bodies, covers all types of arrangements of solids; thermal displacements of atoms; interstitial atoms, ions; orientational, rotational motion of molecules; transition between states of matter. Mathematical theory developed close to physical subject matter. "Discussed in a simple yet deeply penetrating fashion . . . will serve as seeds for a great many basic and applied developments in chemistry." J. of the Amer. Chemical Soc. 216 bibliographical footnotes. 55 figures. xi + 485pp. 5 $\frac{1}{2}$ x 8.

S94 Clothbound \$3.95
S95 Paperbound \$2.45

ASTRONOMY

OUT OF THE SKY, H. H. Nininger. Non-technical, comprehensive introduction to "meteoritics" —science concerned with arrival of matter from outer space. By one of world's experts on meteorites, this book defines meteors and meteorites; studies fireball clusters and processions, meteorite composition, size, distribution, showers, explosions, origins, much more. viii + 336pp. 5 $\frac{1}{2}$ x 8.

T519 Paperbound \$1.85

AN INTRODUCTION TO THE STUDY OF STELLAR STRUCTURE, S. Chandrasekhar. Outstanding treatise on stellar dynamics by one of greatest astro-physicists. Examines relationship between loss of energy, mass, and radius of stars in steady state. Discusses thermodynamic laws from Caratheodory's axiomatic standpoint; adiabatic, polytropic laws; work of Ritter, Emden, Kelvin, etc.; Stroemgren envelopes as starter for theory of gaseous stars; Gibbs statistical mechanics (quantum); degenerate stellar configuration, theory of white dwarfs; etc. "Highest level of scientific merit," Bulletin. Amer. Math. Soc. 33 figures. 509pp. 5 $\frac{1}{2}$ x 8.

S413 Paperbound \$2.75

LES MÉTHODES NOUVELLES DE LA MÉCANIQUE CÉLESTE, H. Poincaré. Complete French text of one of Poincaré's most important works. Revolutionized celestial mechanics; first use of integral invariants, first major application of linear differential equations, study of periodic orbits, lunar motion and Jupiter's satellites, three body problem, and many other important topics. "Started a new era . . . so extremely modern that even today few have mastered his weapons," E. T. Bell. 3 volumes. Total 1282pp. 6 $\frac{1}{8}$ x 9 $\frac{1}{4}$.

Vol. 1 S401 Paperbound \$2.75
Vol. 2 S402 Paperbound \$2.75
Vol. 3 S403 Paperbound \$2.75
The set \$7.50

THE REALM OF THE NEBULAE, E. Hubble. One of the great astronomers of our time presents his concept of "island universes," and describes its effect on astronomy. Covers velocity-distance relation; classification, nature, distances, general field of nebulae; cosmological theories; nebulae in the neighborhood of the Milky way; etc. 39 photos, including velocity-distance relations shown by spectrum comparison. "One of the most progressive lines of astronomical research," The Times, London. New Introduction by A. Sandage. 55 illustrations. xxiv + 201pp. 5 $\frac{1}{2}$ x 8.

S455 Paperbound \$1.50

HOW TO MAKE A TELESCOPE, Jean Texereau. Design, build an f/6 or f/8 Newtonian type reflecting telescope, with altazimuth Couder mounting, suitable for planetary, lunar, and stellar observation. Covers every operation step-by-step, every piece of equipment. Discusses basic principles of geometric and physical optics (unnecessary to construction), comparative merits of reflectors, refractors. A thorough discussion of eyepieces, finders, grinding, installation, testing, etc. 241 figures, 38 photos, show almost every operation and tool. Potential errors are anticipated. Foreword by A. Couder. Sources of supply. xiii + 191pp. 6 $\frac{1}{4}$ x 10.

T464 Clothbound \$3.50

BIOLOGICAL SCIENCES

THE BIOLOGY OF THE AMPHIBIA, G. K. Noble, Late Curator of Herpetology at Am. Mus. of Nat. Hist. Probably most used text on amphibia, most comprehensive, clear, detailed. 19 chapters, 85 page supplement: development; heredity; life history; speciation; adaptation; sex, integument, respiratory, circulatory, digestive, muscular, nervous systems; instinct, intelligence, habits, economic value classification, environment relationships, etc. "Nothing comparable to it," C. H. Pope, curator of Amphibia, Chicago Mus. of Nat. Hist. 1047 item bibliography. 174 illustrations. 600pp. 5 $\frac{1}{2}$ x 8.

S206 Paperbound \$2.98

THE ORIGIN OF LIFE, A. I. Oparin. A classic of biology. This is the first modern statement of theory of gradual evolution of life from nitrocarbon compounds. A brand-new evaluation of Oparin's theory in light of later research, by Dr. S. Margulis, University of Nebraska. xxv + 270pp. 5 $\frac{1}{2}$ x 8.

S213 Paperbound \$1.75

CATALOGUE OF

THE BIOLOGY OF THE LABORATORY MOUSE, edited by G. D. Snell. Prepared in 1941 by staff of Roscoe B. Jackson Memorial Laboratory, still the standard treatise on the mouse, assembling enormous amount of material for which otherwise you spend hours of research. Embryology, reproduction, histology, spontaneous neoplasms, gene and chromosomes mutations, genetics of spontaneous tumor formations, of tumor transplantation, endocrine secretion and tumor formation, milk influence and tumor formation, inbred, hybrid animals, parasites, infectious diseases, care and recording. "A wealth of information of vital concern. . . recommended to all who could use a book on such a subject," Nature. Classified bibliography of 1122 items. 172 figures, including 128 photos. ix + 497pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$.
S248 Clothbound \$6.00

THE TRAVELS OF WILLIAM BARTRAM, edited by Mark Van Doran. Famous source-book of American anthropology, natural history, geography, is record kept by Bartram in 1770's on travels through wilderness of Florida, Georgia, Carolinas. Containing accurate, beautiful descriptions of Indians, settlers, fauna, flora, it is one of finest pieces of Americana ever written. 13 original illustrations. 448pp. 5 $\frac{1}{2}$ x 8.
T13 Paperbound \$2.00

BEHAVIOR AND SOCIAL LIFE OF THE HONEYBEE, Ronald Ribbands. Outstanding scientific study; a compendium of practically everything known of social life of honeybee. Stresses behaviour of individual bees in field, hive. Extends von Frisch's experiments on communication among bees. Covers perception of temperature, gravity, distance, vibration; sound production; glands; structural differences; wax production; temperature regulation; recognition, communication; drifting, mating behaviour, other highly interesting topics. "This valuable work is sure of a cordial reception by laymen, beekeepers and scientists," Prof. Karl von Frisch, Brit. J. of Animal Behaviour. Bibliography of 690 references. 127 diagrams, graphs, sections of bee anatomy, fine photographs. 352pp.
S410 Clothbound \$4.50

ELEMENTS OF MATHEMATICAL BIOLOGY, A. J. Lotka. Pioneer classic, 1st major attempt to apply modern mathematical techniques on large scale to phenomena of biology, biochemistry, psychology, ecology, similar life sciences. Partial contents: Statistical meaning of irreversibility; Evolution as redistribution; Equations of kinetics of evolving systems; Chemical, inter-species equilibrium; parameters of state; Energy transformers of nature, etc. Can be read with profit by even those having no advanced math; unsurpassed as study-reference. Formerly titled "Elements of Physical Biology." 72 figures. xxx + 460pp. 5 $\frac{1}{2}$ x 8.
S346 Paperbound \$2.45

TREES OF THE EASTERN AND CENTRAL UNITED STATES AND CANADA, W. M. Harlow. Serious middle-level text covering more than 140 native trees, important escapes, with information on general appearance, growth habit, leaf forms, flowers, fruit, bark, commercial use, distribution, habitat, woodlore, etc. Keys within text enable you to locate various species easily, to know which have edible fruit, much more useful, interesting information. "Well illustrated to make identification very easy," Standard Cat. for Public Libraries. Over 600 photographs, figures. xiii + 288pp. 5 $\frac{1}{2}$ x 6 $\frac{1}{2}$.
T395 Paperbound \$1.35

FRUIT KEY AND TWIG KEY TO TREES AND SHRUBS (Fruit key to Northeastern Trees, Twig key to Deciduous Woody Plants of Eastern North America), W. M. Harlow. Only guides with photographs of every twig, fruit described. Especially valuable to novice. Fruit key (both deciduous trees, evergreens) has introduction on seeding, organs involved, types, habits. Twig key introduction treats growth, morphology. In keys proper, identification is almost automatic. Exceptional work, widely used in university courses, especially useful for identification in winter, or from fruit or seed only. Over 350 photos, up to 3 times natural size. Index of common, scientific names, in each key. xvii + 125pp. 5 $\frac{1}{2}$ x 8.
T511 Paperbound \$1.25

INSECT LIFE AND INSECT NATURAL HISTORY, S. W. Frost. Unusual for emphasizing habits, social life, ecological relations of insects rather than more academic aspects of classification, morphology. Prof. Frost's enthusiasm and knowledge are everywhere evident as he discusses insect associations, specialized habits like leaf-rolling, leaf mining, case-making, the gall insects, boring insects, etc. Examines matters not usually covered in general works: insects as human food; insect music, musicians; insect response to radio waves; use of insects in art, literature. "Distinctly different, possesses an individuality all its own," Journal of Forestry. Over 700 illustrations. Extensive bibliography. x + 524pp. 5 $\frac{1}{2}$ x 8.
T519 Paperbound \$2.49

A WAY OF LIFE, AND OTHER SELECTED WRITINGS, Sir William Osler. Physician, humanist, Osler discusses brilliantly Thomas Browne, Gui Patin, Robert Burton, Michael Servetus, William Beaumont, Laennec. Includes such favorite writing as title essay, "The Old Humanities and the New Science," "Books and Men," "The Student Life," 6 more of his best discussions of philosophy, literature, religion. "The sweep of his mind and interests embraced every phase of human activity," G. L. Keynes. 5 photographs. Introduction by G. L. Keynes, M.D., F.R.C.S. xx + 278pp. 5 $\frac{1}{2}$ x 8.
T488 Paperbound \$1.50

THE GENETICAL THEORY OF NATURAL SELECTION, R. A. Fisher. 2nd revised edition of vital reviewing of Darwin's Selection Theory in terms of particulate inheritance, by one of greatest authorities on experimental, theoretical genetics. Theory stated in mathematical form. Special features of particulate inheritance are examined: evolution of dominance, maintenance of specific variability, mimicry, sexual selection, etc. 5 chapters on man's special circumstances as a social animal. 16 photographs. x + 310pp. 5 $\frac{1}{2}$ x 8.
S466 Paperbound \$1.85

DOVER SCIENCE BOOKS

THE AUTOBIOGRAPHY OF CHARLES DARWIN, AND SELECTED LETTERS, edited by Francis Darwin. Darwin's own record of early life; historic voyage aboard "Beagle;" furore surrounding evolution, his replies; reminiscences of his son. Letters to Henslow, Lyell, Hooker, Huxley, Wallace, Kingsley, etc., and thoughts on religion, vivisection. We see how he revolutionized geology with concepts of ocean subsidence; how his great books on variation of plants and animals, primitive man, expression of emotion among primates, plant fertilization, carnivorous plants, protective coloration, etc., came into being. 365pp. 5 $\frac{1}{2}$ x 8.

T479 Paperbound \$1.65

ANIMALS IN MOTION, Eadweard Muybridge. Largest, most comprehensive selection of Muybridge's famous action photos of animals, from his "Animal Locomotion." 3919 high-speed shots of 34 different animals, birds, in 123 types of action; horses, mules, oxen, pigs, goats, camels, elephants, dogs, cats, guanacos, sloths, lions, tigers, jaguars, raccoons, baboons, deer, elk, gnus, kangaroos, many others, walking, running, flying, leaping. Horse alone in over 40 ways. Photos taken against ruled backgrounds; most actions taken from 3 angles at once: 90°, 60°, rear. Most plates original size. Of considerable interest to scientists as biology classic, records of actual facts of natural history, physiology. "Really marvelous series of plates," Nature. "Monumental work," Waldemar Kaempfert. Edited by L. S. Brown, 74 page introduction on mechanics of motion. 340pp. of plates. 3919 photographs. 416pp. Deluxe binding, paper. (Weight: 4 $\frac{1}{2}$ lbs.) 7 $\frac{1}{2}$ x 10 $\frac{1}{2}$.

T203 Clothbound \$10.00

THE HUMAN FIGURE IN MOTION, Eadweard Muybridge. New edition of great classic in history of science and photography, largest selection ever made from original Muybridge photos of human action: 4789 photographs, illustrating 163 types of motion: walking, running, lifting, etc. in time-exposure sequence photos at speeds up to 1/6000th of a second. Men, women, children, mostly undraped, showing bone, muscle positions against ruled backgrounds, mostly taken at 3 angles at once. Not only was this a great work of photography, acclaimed by contemporary critics as work of genius, but it was also a great 19th century landmark in biological research. Historical introduction by Prof. Robert Taft, U. of Kansas. Plates original size, full of detail. Over 500 action strips. 407pp. 7 $\frac{3}{4}$ x 10 $\frac{1}{2}$. Deluxe edition.

T7204 Clothbound \$10.00

AN INTRODUCTION TO THE STUDY OF EXPERIMENTAL MEDICINE, Claude Bernard. 90-year old classic of medical science, only major work of Bernard available in English, records his efforts to transform physiology into exact science. Principles of scientific research illustrated by specified case histories from his work; roles of chance, error, preliminary false conclusion, in leading eventually to scientific truth; use of hypothesis. Much of modern application of mathematics to biology rests on foundation set down here. "The presentation is polished . . . reading is easy," Revue des questions scientifiques. New foreword by Prof. I. B. Cohen, Harvard U. xxv + 266pp. 5 $\frac{1}{2}$ x 8.

T400 Paperbound \$1.50

STUDIES ON THE STRUCTURE AND DEVELOPMENT OF VERTEBRATES, E. S. Goodrich. Definitive study by greatest modern comparative anatomist. Exhaustive morphological, phylogenetic expositions of skeleton, fins, limbs, skeletal visceral arches, labial cartilages, visceral clefts, gills, vascular, respiratory, excretory, peripheral nervous systems, etc., from fish to higher mammals. "For many a day this will certainly be the standard textbook on Vertebrate Morphology in the English language," Journal of Anatomy. 754 illustrations. 69 page biographical study by C. C. Hardy. Bibliography of 1186 references. Two volumes, total 906pp. 5 $\frac{1}{2}$ x 8.

Two vol. set \$449, 450 Paperbound \$5.00

EARTH SCIENCES

THE EVOLUTION OF IGNEOUS ROCKS, N. L. Bowen. Invaluable serious introduction applies techniques of physics, chemistry to explain igneous rock diversity in terms of chemical composition, fractional crystallization. Discusses liquid immiscibility in silicate magmas, crystal sorting, liquid lines of descent, fractional resorption of complex minerals, petrogen, etc. Of prime importance to geologists, mining engineers; physicists, chemists working with high temperature, pressures. "Most important," Times, London. 263 bibliographic notes. 82 figures. xviii + 334pp. 5 $\frac{1}{2}$ x 8.

S311 Paperbound \$1.85

GEOGRAPHICAL ESSAYS, M. Davis. Modern geography, geomorphology rest on fundamental work of this scientist. 26 famous essays present most important theories, field researches. Partial contents: Geographical Cycle; Plains of Marine, Subaerial Denudation; The Peneplain; Rivers, Valleys of Pennsylvania; Outline of Cape Cod; Sculpture of Mountains by Glaciers; etc. "Long the leader and guide," Economic Geography. "Part of the very texture of geography . . . models of clear thought," Geographic Review. 130 figures. vi + 777pp. 5 $\frac{1}{2}$ x 8.

S383 Paperbound \$2.95

URANIUM PROSPECTING, H. L. Barnes. For immediate practical use, professional geologist considers uranium ores, geological occurrences, field conditions, all aspects of highly profitable occupation. "Helpful information . . . easy-to-use, easy-to-find style," Geotimes. x + 117pp. 5 $\frac{1}{2}$ x 8.

T309 Paperbound \$1.00

CATALOGUE OF

DE RE METALLICA, Georgius Agricola. 400 year old classic translated, annotated by former President Herbert Hoover. 1st scientific study of mineralogy, mining, for over 200 years after its appearance in 1556 the standard treatise. 12 books, exhaustively annotated, discuss history of mining, selection of sites, types of deposits, making pits, shafts, ventilating, pumps, crushing machinery; assaying, smelting, refining metals; also salt alum, nitre, glass making. Definitive edition, with all 289 16th century woodcuts of original. Biographical, historical introductions. Bibliography, survey of ancient authors. Indexes. A fascinating book for anyone interested in art, history of science, geology, etc. Deluxe Edition. 289 illustrations. 672pp. 6½ x 10. Library cloth.

\$6 Clothbound \$10.00

INTERNAL CONSTITUTION OF THE EARTH, edited by Beno Gutenberg. Prepared for National Research Council, this is a complete, thorough coverage of earth origins, continent formation, nature and behaviour of earth's core, petrology of crust, cooling forces in core, seismic and earthquake material, gravity, elastic constants, strain characteristics, similar topics. "One is filled with admiration . . . a high standard . . . there is no reader who will not learn something from this book." London, Edinburgh, Dublin, Philosophic Magazine. Largest Bibliography in print: 1127 classified items. Table of constants. 43 diagrams. 439pp. 6½ x 9¾.

S414 Paperbound \$2.45

THE BIRTH AND DEVELOPMENT OF THE GEOLOGICAL SCIENCES, F. D. Adams. Most thorough history of earth sciences ever written. Geological thought from earliest times to end of 19th century, covering over 300 early thinkers and systems; fossils and their explanation, vulcanists vs. neptunists, figured stones and paleontology, generation of stones, dozens of similar topics. 91 illustrations, including Medieval, Renaissance woodcuts, etc. 632 footnotes, mostly bibliographical. 511pp. 5¾ x 8.

T5 Paperbound \$2.00

HYDROLOGY, edited by O. E. Meinzer, prepared for the National Research Council. Detailed, complete reference library on precipitation, evaporation, snow, snow surveying, glaciers, lakes, infiltration, soil moisture, ground water, runoff, drought, physical changes produced by water hydrology of limestone terranes, etc. Practical in application, especially valuable for engineers. 24 experts have created "the most up-to-date, most complete treatment of the subject," Am. Assoc. of Petroleum Geologists. 165 illustrations. xi + 712pp. 6½ x 9¼.

S191 Paperbound \$2.95

LANGUAGE AND TRAVEL AIDS FOR SCIENTISTS

SAY IT language phrase books

"SAY IT" in the foreign language of your choice! We have sold over ½ million copies of these popular, useful language books. They will not make you an expert linguist overnight, but they do cover most practical matters of everyday life abroad.

Over 1000 useful phrases, expressions, additional variants, substitutions.

Modern! Useful! Hundreds of phrases not available in other texts: "Nylon," "air-conditioned," etc.

The ONLY inexpensive phrase book completely indexed. Everything is available at a flip of your finger, ready to use.

Prepared by native linguists, travel experts.

Based on years of travel experience abroad.

May be used by itself, or to supplement any other text or course. Provides a living element. Used by many colleges, institutions: Hunter College; Barnard College; Army Ordnance School, Aberdeen; etc.

Available, 1 book per language:

Danish (T818) 75¢
Dutch (T817) 75¢
English (for German-speaking people) (T801) 60¢
English (for Italian-speaking people) (T816) 60¢
English (for Spanish-speaking people) (T802) 60¢
Esperanto (T820) 75¢
French (T803) 60¢
German (T804) 60¢
Modern Greek (T813) 75¢
Hebrew (T805) 60¢

Italian (T806) 60¢
Japanese (T807) 75¢
Norwegian (T814) 75¢
Russian (T810) 75¢
Spanish (T811) 60¢
Turkish (T821) 75¢
Yiddish (T815) 75¢
Swedish (T812) 75¢
Polish (T808) 75¢
Portuguese (T809) 75¢

DOVER SCIENCE BOOKS

MONEY CONVERTER AND TIPPING GUIDE FOR EUROPEAN TRAVEL, C. Vomacka. Purse-size handbook crammed with information on currency regulations, tipping for every European country, including Israel, Turkey, Czechoslovakia, Rumania, Egypt, Russia, Poland. Telephone, postal rates; duty-free imports, passports, visas, health certificates; foreign clothing sizes; weather tables. What, when to tip. 5th year of publication. 128pp. 3½ x 5¼. T260 Paperbound 60¢

NEW RUSSIAN-ENGLISH AND ENGLISH-RUSSIAN DICTIONARY, M. A. O'Brien. Unusually comprehensive guide to reading, speaking, writing Russian, for both advanced, beginning students. Over 70,000 entries in new orthography, full information on accentuation, grammatical classifications. Shades of meaning, idiomatic uses, colloquialisms, tables of irregular verbs for both languages. Individual entries indicate stems, transiteness, perfective, imperfective aspects, conjugation, sound changes, accent, etc. Includes pronunciation instruction. Used at Harvard, Yale, Cornell, etc. 738pp. 5¾ x 8. T208 Paperbound \$2.00

PHRASE AND SENTENCE DICTIONARY OF SPOKEN RUSSIAN, English-Russian, Russian-English. Based on phrases, complete sentences, not isolated words—recognized as one of best methods of learning idiomatic speech. Over 11,500 entries, indexed by single words, over 32,000 English, Russian sentences, phrases, in immediately useable form. Shows accent changes in conjugation, declension; irregular forms listed both alphabetically, under main form of word. 15,000 word introduction covers Russian sounds, writing, grammar, syntax. 15 page appendix of geographical names, money, important signs, given names, foods, special Soviet terms, etc. Originally published as U.S. Gov't Manual TM 30-944. iv + 573pp. 5¾ x 8. T496 Paperbound \$2.75

PHRASE AND SENTENCE DICTIONARY OF SPOKEN SPANISH, Spanish-English, English-Spanish. Compiled from spoken Spanish, based on phrases, complete sentences rather than isolated words—not an ordinary dictionary. Over 16,000 entries indexed under single words, both Castilian, Latin-American. Language in immediately useable form. 25 page introduction provides rapid survey of sounds, grammar, syntax, full consideration of irregular verbs. Especially apt in modern treatment of phrases, structure. 17 page glossary gives translations of geographical names, money values, numbers, national holidays, important street signs, useful expressions of high frequency, plus unique 7 page glossary of Spanish, Spanish-American foods. Originally published as U.S. Gov't Manual TM 30-900. iv + 513pp. 5¾ x 8¾. T495 Paperbound \$1.75

SAY IT CORRECTLY language record sets

The best inexpensive pronunciation aids on the market. Spoken by native linguists associated with major American universities, each record contains:

14 minutes of speech—12 minutes of normal, relatively slow speech, 2 minutes of normal conversational speed.

120 basic phrases, sentences, covering nearly every aspect of everyday life, travel—introducing yourself, travel in autos, buses, taxis, etc., walking, sightseeing, hotels, restaurants, money, shopping, etc.

32 page booklet containing everything on record plus English translations easy-to-follow phonetic guide.

Clear, high-fidelity recordings.

Unique bracketing systems, selection of basic sentences enabling you to expand use of SAY IT CORRECTLY records with a dictionary, to fit thousands of additional situations.

Use this record to supplement any course or text. All sounds in each language illustrated perfectly—imitate speaker in pause which follows each foreign phrase in slow section, and be amazed at increased ease, accuracy of pronunciation. Available, one language per record for

French	Spanish	German
Italian	Dutch	Modern Greek
Japanese	Russian	Portuguese
Polish	Swedish	Hebrew
English (for German-speaking people)		English (for Spanish-speaking people)

7" (33 1/3 rpm) record, album, booklet. \$1.00 each.

SPEAK MY LANGUAGE: SPANISH FOR YOUNG BEGINNERS, M. Ahlman, Z. Gilbert. Records provide one of the best, most entertaining methods of introducing a foreign language to children. Within framework of train trip from Portugal to Spain, an English-speaking child is introduced to Spanish by native companion. (Adapted from successful radio program of N.Y. State Educational Department.) A dozen different categories of expressions, including greeting, numbers, time, weather, food, clothes, family members, etc. Drill is combined with poetry and contextual use. Authentic background music. Accompanying book enables a reader to follow records, includes vocabulary of over 350 recorded expressions. Two 10" 33 1/3 records, total of 40 minutes. Book. 40 illustrations. 69pp. 5 1/4 x 10 1/2. T890 The set \$4.95

LISTEN & LEARN language record sets

LISTEN & LEARN is the only extensive language record course designed especially to meet your travel and everyday needs. Separate sets for each language, each containing three 33 1/3 rpm long-playing records—1 1/2 hours of recorded speech by eminent native speakers who are professors at Columbia, New York U., Queens College.

Check the following features found only in LISTEN & LEARN:

Dual language recording. 812 selected phrases, sentences, over 3200 words, spoken first in English, then foreign equivalent. Pause after each foreign phrase allows time to repeat expression.

128-page manual (196 page for Russian)—everything on records, plus simple transcription. Indexed for convenience. Only set on the market completely indexed.

Practical. No time wasted on material you can find in any grammar. No dead words. Covers central core material with phrase approach. Ideal for person with limited time. Living, modern expressions, not found in other courses. Hygienic products, modern equipment, shopping, "air-conditioned," etc. Everything is immediately useable.

High-fidelity recording, equal in clarity to any costing up to \$6 per record.

"Excellent . . . impress me as being among the very best on the market," Prof. Mario Pei, Dept. of Romance Languages, Columbia U. "Inexpensive and well done . . . ideal present," Chicago Sunday Tribune. "More genuinely helpful than anything of its kind," Sidney Clark, well-known author of "All the Best" travel books.

UNCONDITIONAL GUARANTEE. Try LISTEN & LEARN, then return it within 10 days for full refund, if you are not satisfied. It is guaranteed after you actually use it.

6 modern languages—FRENCH, SPANISH, GERMAN, ITALIAN, RUSSIAN, or JAPANESE *—one language to each set of 3 records (33 1/3 rpm). 128 page manual. Album.

Spanish	the set \$4.95	German	the set \$4.95	Japanese*	the set \$5.95
French	the set \$4.95	Italian	the set \$4.95	Russian	the set \$5.95

* Available Oct. 1959.

TRÜBNER COLLOQUIAL SERIES

These unusual books are members of the famous Trübner series of colloquial manuals. They have been written to provide adults with a sound colloquial knowledge of a foreign language, and are suited for either class use or self-study. Each book is a complete course in itself, with progressive, easy to follow lessons. Phonetics, grammar, and syntax are covered, while hundreds of phrases and idioms, reading texts, exercises, and vocabulary are included. These books are unusual in being neither skimpy nor overdetailed in grammatical matters, and in presenting up-to-date, colloquial, and practical phrase material. Bilingual presentation is stressed, to make thorough self-study easier for the reader.

COLLOQUIAL HINDUSTANI, A. H. Harley, formerly Nizam's Reader in Urdu, U. of London. 30 pages on phonetics and scripts (devanagari & Arabic-Persian) are followed by 29 lessons, including material on English and Arabic-Persian influences. Key to all exercises. Vocabulary. 5 x 7 1/2. 147pp. Clothbound \$1.75

COLLOQUIAL ARABIC, DeLacy O'Leary. Foremost Islamic scholar covers language of Egypt, Syria, Palestine, & Northern Arabia. Extremely clear coverage of complex Arabic verbs & noun plurals; also cultural aspects of language. Vocabulary. xviii + 192pp. 5 x 7 1/2. Clothbound \$1.75

COLLOQUIAL GERMAN, P. F. Doring. Intensive thorough coverage of grammar in easily-followed form. Excellent for brush-up, with hundreds of colloquial phrases. 34 pages of bilingual texts. 224pp. 5 x 7 1/2. Clothbound \$1.75

COLLOQUIAL SPANISH, W. R. Patterson. Castilian grammar and colloquial language, loaded with bilingual phrases and colloquialisms. Excellent for review or self-study. 164pp. 5 x 7 1/2. Clothbound \$1.75

COLLOQUIAL FRENCH, W. R. Patterson. 16th revised edition of this extremely popular manual. Grammar explained with model clarity, and hundreds of useful expressions and phrases; exercises, reading texts, etc. Appendixes of new and useful words and phrases. 223pp. 5 x 7 1/2. Clothbound \$1.75

DOVER SCIENCE BOOKS

COLLOQUIAL PERSIAN, L. P. Elwell-Sutton. Best introduction to modern Persian, with 90 page grammatical section followed by conversations, 35 page vocabulary. 139pp. Clothbound \$1.75

COLLOQUIAL CZECH, J. Schwarz, former headmaster of Lingua Institute, Prague. Full easily followed coverage of grammar, hundreds of immediately useable phrases, texts. Perhaps the best Czech grammar in print. "An absolutely successful textbook." JOURNAL OF CZECHOSLOVAK FORCES IN GREAT BRITAIN. 252pp. 5 x 7½. Clothbound \$2.50

COLLOQUIAL RUMANIAN, G. Nandris, Professor of University of London. Extremely thorough coverage of phonetics, grammar, syntax; also included 70 page reader, and 70 page vocabulary. Probably the best grammar for this increasingly important language. 340pp. 5 x 7½. Clothbound \$2.50

COLLOQUIAL ITALIAN, A. L. Hayward. Excellent self-study course in grammar, vocabulary, idioms, and reading. Easy progressive lessons will give a good working knowledge of Italian in the shortest possible time. 5 x 7½. Clothbound \$1.75

MISCELLANEOUS

TREASURY OF THE WORLD'S COINS, F. Seinfeld. Finest general introduction to numismatics; non-technical, thorough, always fascinating. Coins of Greece, Rome, modern countries of every continent, primitive societies, such oddities as 200-lb stone money of Yap, nail coinage of New England; all mirror man's economy, customs, religion, politics, philosophy, art. Entertaining, absorbing study; novel view of history. Over 750 illustrations. Table of value of coins illustrated. List of U.S. coin clubs. 224pp. 6½ x 9¼. T433 Paperbound \$1.75

ILLUSIONS AND DELUSIONS OF THE SUPERNATURAL AND THE OCCULT, D. H. Rawcliffe. Rationally examines hundreds of persistent delusions including witchcraft, trances, mental healing, peyotl, poltergeists, stigmata, lycanthropy, live burial, auras, Indian rope trick, spiritualism, dowsing, telepathy, ghosts, ESP, etc. Explains, exposes mental, physical deceptions involved, making this not only an exposé of supernatural phenomena, but a valuable exposition of characteristic types of abnormal psychology. Originally "The Psychology of the Occult." Introduction by Julian Huxley. 14 illustrations. 551pp. 5¾ x 8. T503 Paperbound \$2.00

HOAXES, C. D. MacDougall. Shows how art, science, history, journalism can be perverted for private purposes. Hours of delightful entertainment, a work of scholarly value, often shocking. Examines nonsense news, Cardiff giant, Shakespeare forgeries, Loch Ness monster, biblical frauds, political schemes, literary hoaxers like Chatterton, Ossian, disumbrationist school of painting, lady in black at Valentino's tomb, over 250 others. Will probably reveal truth about few things you've believed, will help you spot more easily the editorial "gander" or planted publicity release. "A stupendous collection . . . and shrewd analysis," New Yorker. New revised edition. 54 photographs. 320pp. 5¾ x 8. T465 Paperbound \$1.75

YOGA: A SCIENTIFIC EVALUATION, Kovoov T. Behanan. Book that for first time gave Western readers a sane, scientific explanation, analysis of yoga. Author draws on laboratory experiments, personal records of year as disciple of yoga, to investigate yoga psychology, physiology, "supernatural" phenomena, ability to plumb deepest human powers. In this study under auspices of Yale University Institute of Human Relations, strictest principles of physiological, psychological inquiry are followed. Foreword by W. A. Miles, Yale University. 17 photographs. xx + 270pp. 5¾ x 8. T505 Paperbound \$1.65

Write for free catalogs!

Indicate your field of interest. Dover publishes books on physics, earth sciences, mathematics, engineering, chemistry, astronomy, anthropology, biology, psychology, philosophy, religion, history, literature, mathematical recreations, languages, crafts, art, graphic arts, etc.

Write to Dept. catr
Dover Publications, Inc.
180 Varick St., N. Y. 14, N. Y.

