

CS 250 Course Study Guide

Sarmistha Sarna Gomasta, Kyla Levin, Alexander Yeung

Fall 2023

Please Fill Out Our Feedback Form: <https://forms.gle/1eupiHHHSuTEjXe9>

Contents

Chapter 1: Sets and Predicate Logic	3
Q1.1	3
Q1.1 Answer	4
Q1.2	5
Q1.2 Answer	6
Q1.3	7
Q1.3 Answer	8
Q1.4	9
Q1.4 Answer	10
Additional Resources	11
Chapter 2: Quantifiers and Relations	12
Q2.1	12
Q2.1 Answer	13
Q2.2	14
Q2.2 Answer	15
Q2.3	16
Q2.3 Answer	17
Q2.4	18
Q2.4 Answer	19
Q2.5	20
Q2.5 Answer	21
Additional Resources	22
Chapter 3: Number Theory	23
Q3.1	23
Q3.1 Answer	24
Q3.2 Answer	26
Q3.3	27
Q3.3 Answer	28
Additional Resources	29
Chapter 4: Induction and Recursion	30
Q4.1	30
Q4.1 Answer	31
Q4.2	32
Q4.2 Answer	33
Additional Resources	34

Chapter 9: Graphs and Trees	35
Q9.1	35
Q9.1 Answer	36
Additional Resources	37
 Chapter 5: Regular Expressions	 38
Q5.1	38
Q5.1 Answer	39
Additional Resources	40
 Chapter 14: Finite Automata	 41
Q14.1	41
Q14.1 Answer	42
Q14.2	43
Q14.2 Answer	44
Q14.3	45
Q14.3 Answer	46
Q14.4	47
Q14.4 Answer	48
Q14.5	49
Q14.5 Answer	50
Q14.6	51
Q14.6 Answer	52
Q14.7	53
Q14.7 Answer	54
Q14.8	55
Q14.8 Answer	56
Additional Resources	57
 Chapter 15: Formal Language Theory	 58
Q15.1	58
Q15.1 Answer	59
Additional Resources	60

Chapter 1: Sets and Predicate Logic

Q1.1

Solve the following syllogism. Report the answer as a logical implication, as well as a naturally-spoken English sentence.

- The only foods that my doctor recommends are ones that aren't very sweet.
- Nothing that agrees with me is good for dinner.
- Cake is always very sweet.
- My doctor recommends all foods that are good for dinner.

Q1.1 Answer

We will use the following variables: D = doctor recommends it, S = sweet, A = agrees with me, G = good for dinner

- This can be read as "If my doctor recommends it, then it's not very sweet." $D \rightarrow \neg S$
- This can be read as "There is nothing such that if it agrees with me, then it's good for dinner." $A \rightarrow G$
- This can be read as "If it's cake, then it's very sweet." $C \rightarrow S$
- This can be read as "If it's good for dinner, then my doctor recommends it." $G \rightarrow D$

$$C \rightarrow S \rightarrow \neg D \rightarrow \neg G$$

Transitively, $C \rightarrow \neg G$

In English, "If it's cake, then it's not good for dinner." Or, more naturally, "Cake is not good for dinner."

Q1.2

- a. What values of a and b make the following statement true? Prove using a truth table.

$$(a \vee \neg b) \wedge (b \wedge \neg a)$$

- b. What values of a , b , and c make the following statement true? Prove without using a truth table.

$$(a \vee \neg b) \wedge (b \wedge \neg a) \rightarrow (\neg c \wedge (a \vee b))$$

Q1.2 Answer

- a. What values of a and b make the following statement true? Prove using a truth table.

$$(a \vee \neg b) \wedge (b \wedge \neg a)$$

a	b	$\neg a$	$\neg b$	$(a \vee \neg b)$	$(b \wedge \neg a)$	$(a \vee \neg b) \wedge (b \wedge \neg a)$
T	T	F	F	T	F	F
T	F	F	T	T	F	F
F	T	T	F	F	T	F
F	F	T	T	T	F	F

To fill in the truth table, begin with the primitive values a and b in the leftmost columns. From there, use those values to fill in the successive $\neg a$ and $\neg b$ values in the next two columns. Continue rightwards, adding columns with increasingly complex expressions using the truth values from the simpler columns until you have constructed the final expression.

As a reminder, for the binary operator "OR" (\vee), *at least one* of the input needs to be true. For the binary operator "AND" (\wedge), *both* of the input need to be true. And the unary operator "NOT" (\neg) will flip the truth value of its input.

As seen in the final column of the truth table, there is no combination of a and b values that make $(a \vee \neg b) \wedge (b \wedge \neg a)$ true.

- b. What values of a , b , and c make the following statement true? Prove without using a truth table.

$$(a \vee \neg b) \wedge (b \wedge \neg a) \rightarrow (\neg c \wedge (a \vee b))$$

All possible values of a , b , and c will make this statement true. We know from part (a) that the antecedent of the implication step is always false. There are no a and b values that make it true. Therefore, there is no value that the consequent could have to make the whole implication evaluate to true. If the consequent is true, then $F \rightarrow T$ evaluates to true. If the consequent is false, then $F \rightarrow F$ still evaluates to true.

Q1.3

Complete a deductive sequence proof based on the following propositions:

1. $\neg p \wedge q$
2. $r \rightarrow p$
3. $\neg r \rightarrow s$
4. $s \rightarrow t$

Your answer should either be a conjunction of all of the variables p,q,r,s,t which enumerates their truth values, or you may explicitly state the truth value of each variable.

Q1.3 Answer

1. Assume $\neg p$
2. q (Premise 1, right separation)
3. $\neg p \rightarrow \neg r$ (Contrapositive of Premise 2)
4. $\neg r$ (Modus Ponens of Line 3)
5. s (Modus Ponens of Premise 3)
6. t (Modus Ponens of Premise 4)

Then our final solution is $\neg p \wedge q \wedge \neg r \wedge s \wedge t$

Q1.4

Prove by contradiction that the sum of a rational number and an irrational number is irrational. Recall that a rational number is one which can be expressed as the quotient of two integers. An irrational number is one which cannot be represented as the quotient of two integers. You may assume that the operations addition, subtraction, and multiplication on rationals are well-defined. In other words, the sum, difference, or product of two rationals is always rational.

Q1.4 Answer

Assume to the contrary, $\exists s$ such that $s = r + i$ where $r \in \mathbb{Q}$, the rational numbers, and $i \notin \mathbb{Q}$, or i is irrational. Well, then $i = s - r$. But we know that the difference between two rational numbers is always rational. Then i must be a rational number. But this is a contradiction because we assumed i to be irrational!

Additional Resources for Sets and Predicate Logic

1. [Translating Statements to Propositional Logic](#)
2. [Excellent Discrete Math Playlist](#)
3. [Playlist Containing Many Worked Out Deductive Proofs](#)
4. [MIT Video of Proof by Contradiction](#)
5. [Proof by Contrapositive Intro](#)
6. [Rules of Inference Video](#)

Chapter 2: Quantifiers and Relations

Q2.1

Consider the set $A = \mathbb{N}$

Identify whether the following relations are reflexive, antireflexive, symmetric, antisymmetric, or transitive.

- x relates to y if and only if $x = 2y$
- x relates to y if and only if $x \% 2 = y \% 2$

Q2.1 Answer

- x relates to y if and only if $x = 2y$

This is not reflexive, because for most numbers, $x \neq 2x$. However, it is also not antireflexive, since $x = 2x$ is true for $x = 0$. It is not symmetric, since if $x = 2y$, then $y \neq 2x$. The only case when this is true is for $x = y = 0$, when x and y are the same. Therefore, this relation is antisymmetric. It is not transitive, since if $x = 2y$ and $y = 2z$, then $x = 4z$, which is not always equal to $2z$.

- x relates to y if and only if $x \% 2 = y \% 2$

This is reflexive, since $x \% 2 = x \% 2$ for all x . Because it is reflexive, it cannot be antireflexive. It is symmetric, since if $x \% 2 = y \% 2$, then $y \% 2 = x \% 2$. It is not antisymmetric, since this can be true of different x and y values. It is transitive, since if $x \% 2 = y \% 2$ and $y \% 2 = z \% 2$, then $x \% 2 = z \% 2$.

Q2.2

Let A, B, C, D be nonempty sets.

- a) Prove that $A \times B \subseteq C \times D$ if and only if $A \subseteq C$ and $B \subseteq D$.
- b) What happens to the result in part (a) if any of the sets A, B, C, D is empty?

Q2.2 Answer

- (a) Assume that $A \times B \subseteq C \times D$ and let $a \in A$ and $b \in B$. Then $(a, b) \in A \times B$, and since $A \times B \subseteq C \times D$ we have $(a, b) \in C \times D$. But $(a, b) \in C \times D$ implies $a \in C$ and $b \in D$. Hence, $a \in A$ implies $a \in C$, so $A \subseteq C$, and $b \in B$ implies $b \in D$, so $B \subseteq D$. Conversely, suppose that $A \subseteq C$ and $B \subseteq D$, and that $(x, y) \in A \times B$. Then $x \in A$ and $y \in B$ implies $x \in C$ (since $A \subseteq C$) and $y \in D$ (since $B \subseteq D$) implies $(x, y) \in C \times D$. Consequently, $A \times B \subseteq C \times D$.
- (b) If any of the sets A, B, C, D is empty we still find that

$$\{(A \subseteq C) \wedge (B \subseteq D)\} \Rightarrow \{A \times B \subseteq C \times D\}.$$

However, the converse need not hold. For example, let $A = \emptyset$, $B = \{1, 2\}$, $C = \{1, 2\}$ and $D = \{1\}$. Then $A \times B = \emptyset$ — if not, there exists an ordered pair (x, y) in $A \times B$, and this means that the empty set A contains an element x . And so $A \times B = \emptyset \subseteq C \times D$ — but $B = \{1, 2\} \not\subseteq \{1\} = D$.

Q2.3

Determine whether or not each of the following relations is a function. If a relation is a function, find its range.

- a) $\{(x, y) \mid x, y \in \mathbb{Z}, y = x^2 + 7\}$, a relation from \mathbb{Z} to \mathbb{Z}
- b) $\{(x, y) \mid x, y \in \mathbb{R}, y^2 = x\}$, a relation from \mathbb{R} to \mathbb{R}
- c) $\{(x, y) \mid x, y \in \mathbb{R}, y = 3x + 1\}$, a relation from \mathbb{R} to \mathbb{R}
- d) $\{(x, y) \mid x, y \in \mathbb{Q}, x^2 + y^2 = 1\}$, a relation from \mathbb{Q} to \mathbb{Q}
- e) \mathcal{R} is a relation from A to B where $|A| = 5$, $|B| = 6$, and $|\mathcal{R}| = 6$.

Q2.3 Answer

- (a) Function: Range = $\{7, 8, 11, 16, 23, \dots\}$
- (b) Relation, not a function. For example, both $(4, 2)$ and $(4, -2)$ are in the relation.
- (c) Function: Range = the set of all real numbers.
- (d) Relation, not a function. Both $(0, 1)$ and $(0, -1)$ are in the relation.
- (e) Since $|\mathcal{R}| > 5$, \mathcal{R} cannot be a function.

Q2.4

Determine whether each of the following statements is true or false. If the statement is false, provide a counterexample.

- a) $|a| = [-a]$ for all $a \in \mathbb{Z}$.
- b) $|a| = [a]$ for all $a \in \mathbb{R}$.
- c) $|a| = [a] - 1$ for all $a \in \mathbb{R} \setminus \mathbb{Z}$.
- d) $[a] = [-a]$ for all $a \in \mathbb{R}$.

Q2.4 Answer

- (a) True
- (b) False: Let $a = 1.5$. Then $\lfloor 1.5 \rfloor = 1 \neq 2 = \lfloor a \rfloor$
- (c) True
- (d) False: Let $a = 1.5$. Then $\lfloor -a \rfloor = -2 \neq -1 = \lfloor -a \rfloor$.

Q2.5

3. a) Let $a \in \mathbb{R}^+$ where $a \geq 1$. Prove that (i) $\lfloor \frac{a}{a} \rfloor = 1$; and (ii) $\lfloor \frac{1}{a} \rfloor = 0$.
- b) If $a \in \mathbb{R}^+$ and $0 < a < 1$, which result(s) in part (a) is (are) true?

Q2.5 Answer

a) Proof (i): If $a \in \mathbb{Z}^+$, then $\lfloor a \rfloor = a$ and $\lfloor a/a \rfloor = 1$. If $a \notin \mathbb{Z}^+$, write $a = n + c$, where $n \in \mathbb{Z}^+$ and $0 < c < 1$. Then $\lfloor a/a \rfloor = \lfloor (n+1)/(n+c) \rfloor = 1 + \lfloor (1-c)/(n+c) \rfloor$, where $0 < (1-c)/(n+c) < 1$. Hence $\lfloor a/a \rfloor = 1 + \lfloor (1-c)/(n+c) \rfloor = 1$.

Proof (ii): For $a \in \mathbb{Z}^+$, $\lfloor a \rfloor = a$ and $\lfloor 1/a \rfloor = 0$ and $\lfloor 1/a \rfloor = \lfloor 1 \rfloor = 1$. When $a \notin \mathbb{Z}^+$, let $a = n + c$, where $n \in \mathbb{Z}^+$ and $0 < c < 1$. Then $\lfloor 1/a \rfloor = \lfloor 1/(n+c) \rfloor = 1 - \lfloor c/(n+c) \rfloor$, where $0 < c/(n+c) < 1$. Consequently $\lfloor 1/a \rfloor = 1 - \lfloor c/(n+c) \rfloor = 1$.

b) Consider $a = 0.1$. Then (i) $\lfloor a/a \rfloor = \lfloor 0.1/0.1 \rfloor = \lfloor 1 \rfloor = 1$ not equal to 1; and (ii) $\lfloor 1/a \rfloor = \lfloor 1/0.1 \rfloor = 0$ not equal to 1.

In fact (ii) is false for all $0 < a < 1$, since $\lfloor 1/a \rfloor = 0$ for all such values of a . In the case of (i), when $0 < a \leq 0.5$, it follows that $\lfloor a/a \rfloor \geq 2$ and $\lfloor 1/a \rfloor \geq 2$ not equal to f . However, for $0.5 < a < 1$, $\lfloor a/a \rfloor = 1/a$ where $1 < 1/a < 2$, and so $\lfloor a/a \rfloor = 1$ for $0.5 < a < 1$.

Additional Resources for Quantifiers and Relations

1. [Set Theory Playlist](#)
2. [Equivalence Relations and Equivalence Classes Explained](#)
3. [Bijective, Injective, Surjective Explained](#)
4. [Existential and Universal Quantifiers Explained](#)
5. [Negating Quantifiers](#)
6. [Negating Multiple Quantifiers](#)
7. [Reflexive, Symmetric, Transitive](#)

Chapter 3: Number Theory

Q3.1

Prove that for some relatively prime a and n , multiplying all numbers in the set $\{1, 2, \dots, n-2, n-1\}$ by a will output a permutation of $\{1, 2, \dots, n-2, n-1\}$.

Hint: There are two parts to this proof—existence and uniqueness. To show that multiplication by a permutes the elements of the set, you must show that the multiplication always produces an element in the set (existence) and you must show that that element will never be repeatedly produced via multiplication by a (uniqueness). Consider how these relate to modular arithmetic and modular inverses.

Q3.1 Answer

As stated in the hint, we must show both the existence and uniqueness of the elements from the set when multiplied by a .

First, to prove existence, we will prove that for some relatively prime a and n , and for some elements $b, c \in \{1, 2, \dots, n-2, n-1\}$, $ab = c \pmod{n}$, unless $b = 0$. This is equivalent to saying that "If $ab = 0 \pmod{n}$, then $b = 0 \pmod{n}$ ". Since a and n are relatively prime, a has an inverse in mod n . Multiply both sides by $a^{-1} \pmod{n}$ to get $aba^{-1} = 0(a^{-1}) \pmod{n}$. aa^{-1} will cancel out to 1, showing that $b = 0 \pmod{n}$.

Next, to prove uniqueness, we will show that for some relatively prime a and n , if $ab = ac \pmod{n}$, then $b = c$. Once again, because a and n are relatively prime, we know a must have some inverse $a^{-1} \pmod{n}$. Multiplying both sides by a^{-1} gives $aba^{-1} = aca^{-1} \pmod{n}$, leaving $b = c \pmod{n}$.

Putting the two pieces together, we know that multiplying any element of $\{1, 2, \dots, n-2, n-1\}$ by some constant a , which is relatively prime to n , will never result in 0 and therefore will always produce some number in the range of 1 to $n-1$. Furthermore, no multiplication will result in the same number. Therefore, the output will be a permutation of the original set $\{1, 2, \dots, n-2, n-1\}$.

Q3.2 Euclid's Algorithm

Let $a = 473$ and $b = 47$. Run Euclid's Algorithm on a and b to find their GCD. What can you conclude about a and b from your run of Euclid's Algorithm?

Q3.2 Answer

$$473 = 10 \times 47 + 3$$

$$47 = 15 \times 3 + 2$$

$$3 = 1 \times 2 + 1$$

$$2 = 2 \times 1 + 0$$

The GCD of 473 and 47 is 1 since the algorithm ends with a remainder of 1. Since their GCD is 1, the two numbers are relatively prime.

Q3.3

Use the Chinese Remainder Theorem to find the smallest positive integer x that satisfies the following three simultaneous congruences:

$$x \equiv 2 \pmod{3}$$

$$x \equiv 3 \pmod{5}$$

$$x \equiv 2 \pmod{7}$$

Q3.3 Answer

$$n_1 = 3$$

$$n_2 = 5$$

$$n_3 = 7$$

$$N = n_1 n_2 n_3 = 105$$

Calculate the modular multiplicative inverses:

$$y_1 \equiv 35^{-1} \pmod{3} \quad (\text{Since } N_1 = N/n_1 = 35)$$

$$y_2 \equiv 21^{-1} \pmod{5} \quad (\text{Since } N_2 = N/n_2 = 21)$$

$$y_3 \equiv 15^{-1} \pmod{7} \quad (\text{Since } N_3 = N/n_3 = 15)$$

Compute t_i terms:

$$t_1 = 2 \cdot 35 \cdot y_1 \pmod{105}$$

$$t_2 = 3 \cdot 21 \cdot y_2 \pmod{105}$$

$$t_3 = 2 \cdot 15 \cdot y_3 \pmod{105}$$

Sum and reduce mod N to get:

$$\begin{aligned} x &= t_1 + t_2 + t_3 \pmod{105} \\ &= 14 + 63 + 28 \pmod{105} \\ &= 105 \pmod{105} \end{aligned}$$

However, since we want the smallest positive integer, $x = 105 \pmod{105}$ gives $x = 0$. Therefore, we need to subtract N from 105 to find the smallest positive solution. So the corrected smallest positive integer solution is $x = 105 - 105 = 0$. This seems incorrect as 0 is not positive, we need to reevaluate the t_i terms to find the correct solution.

Additional Resources for Number Theory

1. [Euclid's Algorithm](#)
2. [Chinese Remainder Theorem](#)

Chapter 4: Induction and Recursion

Q4.1

Explain the error in the following induction proof:

Claim: All students are the same height.

Base case: Consider a set of students of size 1. Any student is the same height as themselves, and so all students in the set are the same height.

Inductive hypothesis: All students in a set of size n have the same height.

Inductive step: Now suppose we have a set of $n+1$ students, labeled s_1 through s_{n+1} : $\{s_1, s_2, s_3, \dots, s_{n-1}, s_n, s_{n+1}\}$.

Consider the entire set of $n+1$ students as two individual sets of n students each.

The first set contains students s_1 through s_n : $\{s_1, s_2, s_3, \dots, s_{n-1}, s_n\}$

The second set contains students s_2 through s_{n+1} : $\{s_2, s_3, \dots, s_{n-1}, s_n, s_{n+1}\}$

Both sets contain n students, and so by the inductive hypothesis, all students in each set must be of the same height. And if all students in each set have the same height, then all students in the full $n+1$ set must have the same height.

In conclusion, all students in any set of size $n \geq 1$ must have the same height.

Q4.1 Answer

The error in the proof can be attributed to the overlap between the two sets in the inductive step. The base case is easy to observe for some set $\{s_1\}$, but when the inductive step is applied to "step" from $n = 1$ to $n = 2$, the two sets described in the inductive step would be $\{s_1\}$ and $\{s_2\}$, in which may be two students of differing heights. So while s_1 and s_2 are the same heights as themselves, this does not mean that they are the same height as one another.

Q4.2

Suppose we have a recursively-defined function (much like a Fibonacci sequence) $f_n = 5f_{n-1} - 6f_{n-2}$ where $f_0 = 2$ and $f_1 = 5$.

Prove inductively that $f_n = 2^n + 3^n$ for all $n \geq 0$.

Q4.2 Answer

Claim: $f_n = 2^n + 3^n$ for all $n \geq 0$

Base case: For $n = 0$: $f_0 = 2$, and $2^0 + 3^0 = 1 + 1 = 2$. For $n = 1$: $f_1 = 5$ and $2^1 + 3^1 = 2 + 3 = 5$.

Inductive hypothesis: Assume that $f_n = 2^n + 3^n$ for all $n \geq 0$

Inductive step:

$$f_{n+1} = 5f_n - 6f_{n-1}$$

$$f_{n+1} = 5(2^n + 3^n) - 6(2^{n-1} + 3^{n-1}) \text{ By inductive hypothesis}$$

$$f_{n+1} = 5(2^n) + 5(3^n) - 3(2^n) - 2(3^n)$$

$$f_{n+1} = 2(2^n) + 3(3^n)$$

$$f_{n+1} = 2^{n+1} + 3^{n+1}$$

Additional Resources for Induction and Recursion

1. [Induction Playlist](#)
2. [Strong Induction](#)
3. [Induction Visualization](#)
4. [Recursively Defined Functions](#)

Chapter 9: Graphs and Trees

Q9.1

How many different undirected graphs with n nodes are possible? Assume that two graphs with n nodes are "different" if they have different edges and give your answer as an expression of n .

Q9.1 Answer

First, we must find out how many possible edges can exist in an undirected graph with n nodes. This is the same as the number of edges that would be in the complete graph. There are many different ways to think about this:

- Consider the adjacency matrix of the graph. Since there are n nodes in the graph, the matrix is of size $n \times n$, and since the graph is complete, every entry in the matrix would be 1 except along the diagonal (since edges cannot be directed at themselves). So the number of edges would be the number of 1's in the matrix, divided by 2 so as not to double-count. This gives $\frac{(n \times n) - n}{2}$ or $\frac{n(n-1)}{2}$
- Number the nodes from 1 to n . Beginning at node 1, it can have up to $n - 1$ edges with the other $n - 1$ nodes. The next, up to $n - 2$ edges, the next, $n - 3$, and so on. This gives the sum $(n - 1) + (n - 2) + \dots + 2 + 1$, which is also equal to $\frac{n(n-1)}{2}$ through a combinatorial proof (not covered in this class).
- In a complete graph, all n nodes would have a degree of $n - 1$. Sum this up across all nodes, and the total degree is $n(n - 1)$. Then, to find the number of edges, divide by 2 to remove the duplicates, because one edge will be one degree in two separate nodes. This also gives $\frac{n(n-1)}{2}$
- Consider all n nodes as a set. From this set, one can *choose* any unordered pair of two nodes to form an edge. This is the definition of the *choose* operation, and so there are $\binom{n}{2}$ possible edges. This again is equivalent to $\frac{n(n-1)}{2}$ edges by combinatorial proof.

Now that we know that there are $\frac{n(n-1)}{2}$ possible edges, we can think of these edges as being a set of edges, and each possible graph on n nodes having some subset of those edges. To figure out how many different graphs are possible, we must figure out how many possible subsets there are for a set with $\frac{n(n-1)}{2}$ edges. This is the same as finding the size of the power set of the set of edges, which is $2^{\frac{n(n-1)}{2}}$.

Additional Resources for Graphs and Trees

1. [DFS and BFS](#)
2. [Graph Theory Playlist](#)
3. [Uniform Cost Search](#)
4. [A* Search](#)

Chapter 5: Regular Expressions

Q5.1

Assume the alphabet $\Sigma = \{0, 1\}$. If the English description is given, provide the regular expression. If the regular expression is given, give a short, English description of the accepted set of strings.

- a. The set of strings that contain at least one 0 and at least one 1.
- b. $(1 + \lambda)(00^*1)^*0^*$
- c. The set of strings such that all pairs of adjacent 0's appear before any pairs of adjacent 1's.

Q5.1 Answer

- a. This can be thought of as either a 0 appearing before the 1, or the 1 appearing before the 0. Since these are the only mandated characters, the rest of the string can be any combination of 0's and 1's that come before, in-between, or after the 0 and 1. In other words, there are two possible string formats: $(0 + 1)^*0(0 + 1)^*1(0 + 1)^*$ or $(0 + 1)^*1(0 + 1)^*0(0 + 1)^*$. Together, the full regex is $((0 + 1)^*0(0 + 1)^*1(0 + 1)^*) + ((0 + 1)^*1(0 + 1)^*0(0 + 1)^*)$.
- b. The language starts with either a 1 or an empty string. From there, it is followed by a 0, some number of zero's, and then a 1. Just looking at these parts, the string can start as either:
 1 0 0... 1
 0 0... 1
 Even if the 0... is omitted, the string can only start with 0 1. 1 1 is not allowed.
 Looking further, we can observe that within the (00^*1) part, removing the optional 0's, (01) the structure mandates that 0 must come before 1. So if this block repeats, the structure will look like this: 0 1 0 1 with more 0's padding the in-between. And since only more 0's are added onto the end, we can conclude that this will be the set of all strings without consecutive 1's.
- c. This can be divided into two parts: The first part of the string that does not allow 11's, but does allow 00's, and the second part of the string that does not allow 00's, but does allow 11's.
 For a binary string where 11 cannot occur, this could be any combination of 0's or 10's, either ending in a 1 or not: $(0 + 10)^*(1 + \lambda)$
 The formula is much the same for a binary string where 00 cannot occur: $(1 + 01)^*(0 + \lambda)$
 Concatenating the two together gives us the final string: $(0 + 10)^*(1 + \lambda)(1 + 01)^*(0 + \lambda)$

Additional Resources for Regular Expressions

1. [Formal Language Theory Playlist](#)


Chapter 14: Finite Automata

Q14.1


Assume $\Sigma = \{0, 1\}$. Draw a DFA for the language that accepts all binary strings with either an even number of 0's or contains exactly two 1's.

Q14.1 Answer


Let us first consider what states this machine will need. The machine considers strings based on two different criteria, although a string only needs one to be accepted. The first condition is that there is an even number of 0's. There are only two states for this, an even number of 0's or an odd number of 0's. Reading in 0's would switch between the two states like so:



Now let us consider the other criteria, which is "exactly two 1's". This has four states: zero 1's, exactly one 1, exactly two 1's, and more than two 1's, since beyond two 1's, that criteria will always fail. Reading in any 1 will move between these states, ending with the "more than two 1's" state in a dead end.



Since one criteria only cares about 0's being read in, and the other only cares about 1's being read in, we can combine the two into one machine with 8 states, each representing a state from the even/odd criteria *and* a state from the "how many 1's" criteria.



Q14.2


Draw a DFA that accepts the language consisting of the set of all strings of x 's and y 's that start with xy or yy followed by any string of x 's and y 's.

Q14.2 Answer

The given diagram represents a DFA comprising five states, s_0 to s_4 , with s_0 as the start state and s_2, s_4 as accept states. The transitions are defined by the input symbols 'a' and 'b':

- From s_0 , there is a transition to s_1 on input 'a' and a loop on 'b'.
- State s_1 transitions to s_2 on 'a' and to s_3 on 'b'.
- State s_2 has a loop on 'a'.
- State s_3 transitions to s_1 on 'a' and to s_4 on 'b'.
- State s_4 has a loop on 'b'.

This DFA accepts strings that end with 'a' leading to s_2 or strings with an even number of 'b's leading to s_4 .



Q14.3

Let L_1L_2 be the set of strings of 0's and 1's that both start and end with a 0.

Let $L_1 \cup L_2$ be the set of strings of 0's and 1's that start with a 0 or end with a 0 (or both).

Then the language $(L_1 \cup L_2)^*$ is the set of strings of 0's and 1's that start with a 0 or end with a 0 (or both) or that contain 00.

Q14.3 Answer

a. s_0, s_1, s_2, s_3 b. 0, 1 c. s_0 d. s_0

e.

		input	
		0	1
→ \odot state	s_0	s_0	s_1
	s_1	s_1	s_2
	s_2	s_2	s_3
	s_3	s_3	s_0

Q14.4

TODO: DFA Simplification question


Q14.4 Answer

Q14.5

Convert the NFA below into a DFA using the subset construction. This should be done by:

- Creating DFA states as subsets of the NFA states.
- Defining the start state of the DFA as the epsilon-closure of NFA state 0.
- Including any subset containing NFA state 3 as accept states in the DFA.
- Determining the DFA transitions for each symbol from each state subset.

Provide the transition table for the DFA.




Q	Σ_ϵ	Q
0	ϵ	3
0	a	1
0	a	2
1	b	3
2	ϵ	3
3	ϵ	1

Q14.5 Answer

Each state of the DFA is a set of states of the NFA. The DFA's initial state, S_0 , is the epsilon-closure of the NFA's initial state, which includes states $\{0, 1, 3\}$.


$$\varepsilon\text{-closure}\{0\} = \{0, 1, 3\} = S_0$$

$$\epsilon\text{-CLOSE}\{0\} = \{0, 1, 3\} = S_0$$




For input a , the transition from S_0 leads to a new DFA state, S_1 , determined by the epsilon-closure of the set $\{1, 2\}$, resulting in the set $\{1, 2, 3\}$. For input b , the transition from S_0 leads to state S_2 , which is the epsilon-closure of the set $\{3\}$, or $\{1, 3\}$. The DFA transitions from state S_1 on input a lead to a sinking state, as there are no corresponding NFA transitions. Conversely, on input b , S_1 transitions to the previously established state S_2 . The transitions from state S_2 on inputs a and b need to be determined following the same methodology, considering the epsilon-closures and transitions in the NFA. The final diagram is :

$$\begin{aligned} \delta(S_0, a) &= \epsilon\text{-CLOSE}\{1, 2\} = \{1, 2, 3\} = S_1 \\ \delta(S_0, b) &= \epsilon\text{-CLOSE}\{3\} = \{1, 3\} = S_2 \\ \delta(S_1, a) &= \epsilon\text{-CLOSE}\{\} = \emptyset = S_3 \\ \delta(S_1, b) &= \epsilon\text{-CLOSE}\{3\} = \{1, 3\} = S_2 \\ \delta(S_2, a) &= \epsilon\text{-CLOSE}\{\} = \emptyset = S_3 \\ \delta(S_2, b) &= \epsilon\text{-CLOSE}\{3\} = \{1, 3\} = S_2 \end{aligned}$$




Q14.6

Determining the transition from state S_2 on inputs a and b is straightforward; from the empty set of states, there are no transitions in the NFA. In the DFA, this is represented by a transition from the empty set back to itself.




Q14.6 Answer

All the edges of the DFA have been discovered.




Q	Σ	Q
{0, 1, 3}	a	{1, 2, 3}
{0, 1, 3}	b	{1, 3}
{1, 2, 3}	a	\emptyset
{1, 2, 3}	b	{1, 3}
{1, 3}	a	\emptyset
{1, 3}	b	{1, 3}
\emptyset	a	\emptyset
\emptyset	b	\emptyset

The final states of the DFA are determined from the final states of the NFA. State 3 was the only final state in the NFA. Any set of NFA states containing a final state is a final state in the DFA.



Q	Σ	Q
{0, 1, 3 }	a	{1, 2, 3 }
{0, 1, 3 }	b	{1, 3 }
{1, 2, 3 }	a	\emptyset
{1, 2, 3 }	b	{1, 3 }
{1, 3 }	a	\emptyset
{1, 3 }	b	{1, 3 }
\emptyset	a	\emptyset
\emptyset	b	\emptyset



Q14.7

Let L^R be any arbitrary regular language. Let the language $L^R = \{w^R : w \in L\}$ be the reversal of L . Show that L^R is regular.

Q14.7 Answer

Since we know that L is regular, we know that there exists some DFA or equivalent NFA which accepts L . Without loss of generality, we can assume that there exists only start or end state (If a DFA had multiple accepting states, you could eliminate them down to just one by redirecting all of the arrows going into other accepting states into a single accepting state, and eliminating the redundant accepting states). Then we can construct a DFA or equivalent NFA which accepts L^R by making the start state and accepting state, the accepting state a start state, and reversing every state transition (reversing the "arrows"). Then we have a machine that traverses backwards over every string in L , and have thus constructed a DFA for L^R . Since there exists a DFA / NFA for L^R , we know that L^R is regular.

Q14.8

For $\Sigma = \{a, b, c, d\}$, give a regular expression that captures all strings that use their letters in reverse alphabetical order, but use at most three of the four possible letters.

Note: The strings themselves can be longer than 3 letters long, since letters can repeat. Draw an NFA that captures the regular expression from above.

Q14.8 Answer


The regular expression that captures the above language is:

$$(d^*c^*b^*)|(d^*b^*a^*)|(d^*c^*a^*)|(c^*b^*a^*)$$


We will construct this NFA step by step. Initially, consider the NFA for a^* as follows:



The NFA for $d^*c^*b^*$ is constructed by chaining the NFAs for d^* , c^* , and b^* as shown:



Extra care is taken when connecting the NFAs sequentially; the final states of the first NFA are connected to the start state of the subsequent NFA, and the final states are made non-final to ensure the correctness of the overall NFA. Although it is safe to keep all states final in this specific case, it is generally prudent not to do so in order to avoid errors in the construction of the NFA. Now we can combine the components to construct the overall NFA.



Additional Resources for Finite Automata

1. [Myhill Nerode](#)
2. [DFA Minimization](#)
3. [Regular Expression to Automaton](#)
4. [Lambda/Epsilon NFA to NFA](#)

Chapter 15: Formal Language Theory

Q15.1

Let L be a language that is Turing Recognizable by some Turing Machine. Let L' be the language $\Sigma^* - L$, and let that language also be Turing Recognizable by some other Turing Machine. Prove that L is Turing Decidable. (Hint: Recall that a single-tape Turing Machine can be used to simulate a multi-tape Turing Machine)

Q15.1 Answer

We know this to be true by the TR / TD Theorem. But for a rigorous proof, recall that a multi-tape Turing Machine can be simulated by a single-tape Turing Machine. Then we will construct a multi-tape Turing Machine which runs the Turing Machines for L and L' in parallel. One of these is guaranteed to terminate, since $L + L' = \Sigma^*$. If, on any given string, the Turing Machine that recognizes L terminates, we know that string is in L . If the Turing Machine that recognizes L' terminates, we know that string is not in L . Then here we have constructed a Turing Machine that always decides whether or not a string is in L , and so L is Turing Decidable.

Additional Resources for Formal Language Theory

1. [Turing Machines Visualized](#)
2. [TR / TD](#)
3. [Halting Problem](#)
4. [Theory of Computation Playlist](#)
5. <https://www.geeksforgeeks.org/conversion-from-nfa-to-dfa/>
6. PS-1-SOL.PDF, accessed from <https://engineering.purdue.edu/~milind/ece468/2015fall/ps-1-sol.pdf>
7. Rabin-Scott subset construction, Loudon