



MMM11755280



CONFIDENTIAL

FOLR1/FR α

Immunohistochemistry Test

Histology Analysis for FOLR1

ORDERING HEALTHCARE PROVIDER

Karla Bowles MD

Myriad Women's Health Proficiency Testing Clinic
180 KIMBALL WAY
SOUTH SAN FRANCISCO, CA 94080

SPECIMEN

Specimen Type:	Tissue Block/Slide
Tumor Primary:	Ovary
Surgery/Biopsy Date:	Jun 11, 2024
TRF Received:	Jul 1, 2024
Sample Received:	Jul 1, 2024
Report Date:	Jul 18, 2024

PATIENT

Legal Name:	Pt Last Name Pt First Name
Date of Birth:	Jan 2, 1970
Patient ID:	Patient ID
Sex at Birth:	F
Accession #:	07253074-BLD
Requisition #:	90024548

PATHOLOGY

Josephia Pathologist MD

Specimen(s) Analyzed: Test Block

Results

FOLR1/FRA

POSITIVE

80%

REFERENCE RANGE: Positive: $\geq 75\%$ with 2+ and/or 3+ membrane staining, Negative: $< 75\%$

Stains reviewed and scored by a qualified pathologist using manual light microscopy. All controls reviewed and showed appropriate positive and negative reactivity.

FOLR1 immunohistochemical analysis:

VENTANA FOLR1 (FOLR1-2.1) RxDx Assay is a qualitative immunohistochemical assay using mouse monoclonal anti-FOLR1 clone FOLR1-2.1 intended for use in the assessment of folate receptor alpha (FOLR1) expression in formalin-fixed, paraffin-embedded epithelial ovarian, fallopian tube, or primary peritoneal cancer tissue specimens by light microscopy. This assay is for use with OptiView DAB IHC Detection Kit for staining on a BenchMark ULTRA instrument. FOLR1 expression clinical cut-off is $\geq 75\%$ viable tumor cells (TC) with membrane staining at moderate (2+) and/or strong (3+) intensity levels. This assay is indicated as an aid in identifying patients with epithelial ovarian, fallopian tube, or primary peritoneal cancer who may be eligible for treatment with ELAHERE™ (mirvetuximab soravtansine). See package insert for Ventana FOLR1 ELAHERE™ product label for additional information.

Methodology:

FOLR1 staining is performed on a BenchMark ULTRA instrument utilizing the Ventana FOLR1 FDA approved protocol in combination with OptiView DAB IC Detection Kit and ancillary reagents.

Disclaimer:

This test and its performance characteristics have been verified by and performed at the Intermountain Central Laboratory (5252 S Intermountain Dr. Murray UT 84107) for on-label use on epithelial ovarian, fallopian tube and/or peritoneal cancer tissue. This test is intended for in vitro diagnostic (IVD) use. This laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA-88) as qualified to perform high complexity clinical laboratory testing.

Sex assigned at birth refers to the classification of an individual as male or female, often based on physical characteristics at birth.

The lab director signature may reflect remote review of digital data.

This report was electronically reviewed and

signed out by:

Isaac E Lloyd

Sterling T. Bennett, MD

Laboratory Director

Dylan V. Miller, MD

Laboratory Director

Myriad Genetic Laboratories, Inc. | 320 Wakara Way, Salt Lake City, Utah 84108 | PH: 877-283-6709 FX: 801-883-8998

The format and contents of this report are proprietary and may not be copied or used without permission, except for purposes of diagnosis, counseling, and treating the patient identified in the report and members of his or her family. Myriad and the Myriad logo are either trademarks or registered trademarks of Myriad Genetics, Inc. in the United States and other jurisdictions. © 2024, Myriad Genetics, Inc.