




Introduction to Bioinformatics


S. Ravichandran, PhD, PMP
Hood College

Online servers that require user accounts

- NCBI
 - myNCBI (user account)
 - <https://www.ncbi.nlm.nih.gov/>
- Ensembl
- UCSC etc.
- Galaxy
 - <https://usegalaxy.org/> (user account)
- Create the accounts ahead of time




Human Cell As a Factory




Roughly 37.2 trillion cells in our body

Typical cell (across length) $10 \times 10^{-6}\text{m}$



Morbid Map of the Human mtDNA Genome


Inherited from
Mom to children



Fingerprints of Evolution
Circular Genome



Re-use with attribution permitted. www.mitomap.org.



Chromosome




autosomes

sex chromosomes

U.S. National Library of Medicine

Book of Life: 2 Multi-Volume Set



DAD






MOM


To understand the language of
DNA, we need to understand
some Chemistry/Biochemistry

Brief Introduction

Periodic Table



Ribose



Phosphate

Periodic Table of Elements

1																	18																																		
H																	He																																		
2																																																			
Li	Be											B	C	N	O	F	Ne																																		
Na	Mg	3	4	5	6	7	8	9	10	11	12	Al	Si	P	S	Cl	Ar																																		
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr																																		
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe																																		
Cs	Ba	*	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn																																		
Fr	Ra	**	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg																																									
		<table><tr><td>*</td><td>La</td><td>Ce</td><td>Pr</td><td>Nd</td><td>Pm</td><td>Sm</td><td>Eu</td><td>Gd</td><td>Tb</td><td>Dy</td><td>Ho</td><td>Er</td><td>Tm</td><td>Yb</td><td>Lu</td><td>D</td></tr><tr><td>**</td><td>Ac</td><td>Th</td><td>Pa</td><td>U</td><td>Np</td><td>Pu</td><td>Am</td><td>Cm</td><td>Bk</td><td>Cf</td><td>Es</td><td>Fm</td><td>Md</td><td>No</td><td>Lr</td><td>T</td></tr></table>																*	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	D	**	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	T
*	La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	D																																			
**	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	T																																			

General

NMR

Mass

Coloration

Characters :

Discoverer :

Name Origin :

Atomic Radius, A :

Electronegativity :

Ionization Potential, kJ/mol :

Electron Affinity, kJ/mol :

Density :

Melting Point, K :




Boiling Point, K :

OK

Cancel

Help

DNA has direction



Biology is the chemistry that crawls

What holds the molecules together?




COVALENT BONDS - BY WATERMELON56

WWW.TOONDOO.COM



Atomic level: Chemistry Rules

- Bonded interactions
 - Covalent bonds
- Non-bonded interactions
 - H-bonds
 - Holds DNA
 - Makes drug binding work
 - Ionic, VDW etc.



H-bonds

Which pair is easy to break?

A-T



Or

G-C

of H-bonds

Can we think of A-T being the site for
DNA actions such as
double-stranded → single-stranded

Replication and Transcription etc.




DNA/RNA 3D Structures

CGCGAATTCGGG
GCGCTTAAGCCC





GGGCUAUUAGCUCAGUUGGUUAGAGCGCACCC
CUGAUAAGGGUGAGGUCGCUGAUUCGAAUUC
AGCAUAGCCCA



RNAfold Server
structure drawing encoding base-pair
probabilities

DNA/RNA

- Different types of DNA
 - B, -Z etc
- Different RNAs
 - mRNA
 - Nucleus → Ribosomes
 - NR (non-coding; 95% of all RNAs)
 - tRNA
 - rRNA



- pre-mRNA
- 7-methylguanosime placed at 5' end (prevent RNA degradation)
- Poly A tail is added at the 3' end (200 bps)
- splicing
- Final product, mRNA

Theory published by **Crick** 1958 (Yes, the same Crick that worked with Watson)

Splicing

- Archaea and Bacteria
 - Usually have one chromosomes
 - Chromosomes are circular
 - some can be linear
- Eukaryotes
 - Multiple chromosomes
 - Linear
 - Packed into cell nucleus


Chromosome
(highlight: Gene)

No.	Exon / Intron	Start	End	Start Phase	End Phase	Length	Sequence
	5' upstream sequence					gcaggagccagggctgggcataaaagtcagggcagagccatctattgctt
1	ENSE00001829867	5,227,071	5,226,930	-	2	142	ACATTGCTTCTGACACAACCTGTGTTCACTAGCAACCTCAAACAGACACCATGGTGCATC TGA
	Intron 1-2	5,226,929	5,226,800			130	gttggtatcaaggttacaagacagg.....tattggtctattttccacccttag
2	ENSE00001057381	5,226,799	5,226,577	2	0	223	GCTGCTGGTGGTCTACCCCTGGACCCAGAGGTTCTTTGAGTCCTTTGGGGATCTGTCAC TCCTGATGCTGTTATGGGCAACCCTAAGGTGAAGGCTCATGGCAAGAAAGTGCTCGGTGC CTTTAGTGATGGCCTGGCTCACCTGGACAACCTCAAGGGCACCTTTGCCACACTGAGTGA GCTGCACTGTGACAAGCTGCACGTGGATCCTGAGAACTTCAGG
	Intron 2-3	5,226,576	5,225,727			850	gtgagtctatgggacgcttgatgtt.....catacctttatcttcctcccacag
3	ENSE00001600613	5,225,726	5,225,464	0	-	263	CTCCTGGGCAACGTGCTGGTCTGTGTGCTGGCCCATCACTTTGGCAAAGAATTCACCCCA CCAGTGCAGGCTGCCTATCAGAAAGTGGTGGCTGGTGTGGCTAATGCCCTGGCCACAAG TATCACTAAGCTCGCTTTCTTGCTGTCCAATTTCTATTAAAGGTTCCCTTTGTTCCCTAAG TCCAACTACTAAACTGGGGGATATTATGAAGGGCCTTGAGCATCTGGATTCTGCCTAATA AAAAACATTATTATTTTCATTGCAA
	3' downstream sequence						tgatgtattttaattattttctgaatattttactaaaaagggaatgtggga.....

Gene
sequence

mRNA (cDNA) and protein sequences


1	ACATTGCTTCTGACACAACCTGTGTTCACTAGCAACCTCAAACAGACACCATGGTGCATC	60
ATGGTGCATC	10
-M--V--H--	3
61	TGA	120
11	TGA	70
4	L--T--P--E--E--K--S--A--V--T--A--L--W--G--K--V--N--V--D--E--	23



Water molecules

Proteins in 3D

Drug binding
inhibition



Vitamin-H bound protein (183 aa protein)

Proteins vs DNA

Proteins

- Unstable
 - Seconds to months
 - Depends on protein & organism
 - <> life span of human proteins: 1 day
 - Destroyed after some time and recycled

DNA

- Stable
- DNA can be stable even for 100,000 years!!!

Related Dogmas

- Central Dogma of Genomics
 - Genome → Transcriptome → Proteome → Cellular Phenotype
- Study of organisms that inhabit the human body
 - microbiome

e-Genome

Genome → Computers

- Ecoli

a

0	1	1	0	0	0	1	1
---	---	---	---	---	---	---	---

- 4 million bases

- $4,000,000 * 1 \text{ Byte} = 4,000,000$

- ~4 MB Hard drive

- Human

- ~ **3,000,000,000** (3 Billion) bases (one copy)

- Each cell

- $3 \times 10^9 * 1 \text{ Byte} = 3 \times 10^9 = 3 \text{ GB}$

Learning Bioinformatics

- Software/Browser/OS
 - Version issues
 - Scripts sometimes fail
 - Web connectivity issues
 - NCBI/Ensembl might change their genomic browsers without notice; Genomic browsers behave sometimes differently with different browsers/OS
- Please be patient!

"it is better 100 guilty Persons should escape than that one innocent Person should suffer" Benjamin Franklin

Applications

- **Criminal Justice system** <https://www.innocenceproject.org/dna-exonerations-in-the-united-states/>
 - Genetic evidence and exoneration
 - First event happened in 1989
 - Since then many cases had been resolved using DNA exoneration
 - DNA evidence is admitted in criminal trials in almost all states in USA
- Disease, Therapy(?)
 - Breast cancer: Mutations in BRCA1/2 genes
 - Achondroplasia (Dwarfism): Mutations in FGFR3 gene
- **Evolving area**
 - Systems' view is lacking

Applications

Mail - sarangan.ravichan... Course Documents - Nu... Inbox (1,647) - saka.ravi... Bookshelf Online: Bioinfo... Influenza virus database... Flu DatasetExplorer - sec... Influenza A virus subtype... Revishendian

Secure | https://en.wikipedia.org/wiki/Influenza_A_virus_subtype_H1N1#Russian_flu

Apps v9.6.4 Unanet 9.6.4... http://www.google.co... IRF5 - Wikipedia, the ScienceProject Web Services provide Free Online Course M Google Inbox - Web Email A Cbl and human myel Add Site to Prism Fellowships & Posit

Tagalog
Українська
Tiếng Việt
粵語
中文
Edit links

The 1918 flu caused an unusual number of deaths, possibly due to it causing a **cytokine storm** in the body. (The current **H5N1 bird flu**, also an influenza A virus, has a similar effect.)^[8] The Spanish flu virus infected lung cells, leading to overstimulation of the **immune system** via release of **cytokines** into the **lung** tissue. This leads to extensive **leukocyte** migration towards the lungs, causing destruction of lung tissue and secretion of liquid into the organ. This makes it difficult for the patient to breathe. In contrast to other pandemics, which mostly kill the old and the very young, the 1918 pandemic killed unusual numbers of young adults, which may have been due to their healthy immune systems mounting a too-strong and damaging response to the infection.^[9]

The term "Spanish" flu was coined because **Spain** was at the time the only **European** country where the press were printing reports of the outbreak, which had killed thousands in the armies fighting **World War I**. Other countries suppressed the news in order to protect morale.^[10]

Fort Dix outbreak [edit]

Main article: 1976 swine flu outbreak

In 1976, a novel swine influenza A (H1N1) caused severe respiratory illness in 13 soldiers with 1 death at Fort Dix, New Jersey. The virus was detected only from January 19 to February 9 and did not spread beyond Fort Dix.^[11] Retrospective serologic testing subsequently demonstrated that up to 230 soldiers had been infected with the novel virus, which was an H1N1 strain. The cause of the outbreak is still unknown and no exposure to pigs was identified.^[12]

Russian flu [edit]

The 1977–1978 Russian flu **epidemic** was caused by strain *Influenza A/USSR/90/77 (H1N1)*. It infected mostly children and young adults under 23 because a similar strain was prevalent in 1947–57, causing most adults to have substantial immunity. Because of a striking similarity in the viral RNA of both strains – one which is unlikely to appear in nature due to **antigenic drift** – it was speculated that the later outbreak was due to a laboratory incident in Russia or Northern China, though this was denied by scientists in those countries.^{[13][14][15]} The virus was included in the 1978–1979 **influenza vaccine**.^{[16][17][18][19]}


See also 1889–1890 flu pandemic for the earlier Russian flu pandemic caused either by H3N8 or H2N2

2009 A(H1N1) pandemic [edit]

Main article: 2009 flu pandemic

In the **2009 flu pandemic**, the **virus** isolated from patients in the United States was found to be made up of genetic elements from four different flu viruses – North American swine influenza, North American avian influenza, human influenza, and swine influenza virus typically found in Asia and Europe – "an unusually **mongrelised** mix of genetic sequences."^[20] This new strain appears to be a result of **reassortment** of **human influenza** and **swine influenza** viruses, in all four different strains of subtype H1N1.

Preliminary genetic characterization found that the **hemagglutinin** (HA) gene was similar to that of swine flu viruses present in U.S. pigs since 1999, but the **neuraminidase** (NA) and **matrix protein** (M) genes resembled versions present in European swine flu isolates. The six genes from American swine flu are themselves mixtures of swine flu, bird flu, and human flu viruses.^[21] While viruses with this genetic makeup had



missing_misense_f....zip BIFX550-C13-112....pptx

Show all

Year 1977

Host Human

Protein HA

Subtype H1N1

5 protein sequences after collapsing (7 total)

<input checked="" type="checkbox"/>	Accession	Length	Host	Protein	Subtype	Country	Region	Date	Virus name
<input checked="" type="checkbox"/>	ABD60944	566	Human	HA	H1N1	Hong Kong	N	1977	Influenza A virus (A/Hong Kong/117/1977(H1N1))
<input checked="" type="checkbox"/>	ABO44134	566	Human	HA	H1N1	China	N	1977	Influenza A virus (A/Tientsin/78/1977(H1N1))
<input checked="" type="checkbox"/>	ABD95350	566	Human	HA	H1N1	Russia	N	1977	Influenza A virus (A/USSR/90/1977(H1N1))
<input checked="" type="checkbox"/>	APC57869	566	Human	HA	H1N1	USSR		1977	Influenza A virus (A/USSR/90/1977(H1N1))
<input checked="" type="checkbox"/>	ABD60933	566	Human	HA	H1N1	Russia	N	1977	Influenza A virus (A/USSR/92/1977(H1N1))

Influenza Virus Resource

Information, Search and Analysis

[HOME](#) | [SEARCH](#) | [SITE MAP](#) | [Flu home](#) | [Database](#) | [Genome Set](#) | [Alignment](#) | [Tree](#) | [BLAST](#) | [Annotation](#) | [FTP](#) | [Help](#) | [Contact us](#)

Multiple alignment for 5 protein sequences. Alignment length is 566.

Go to position

Position: 1 29 57 85 114 142 170 199 227 255 284 312 340 368 397 425 453 482 510 538 566

Consensus: MKAKLLVLLCALSDADTICIGYHANNSTDTVDIVLEKNVIVITHSVNLLDDSHNGKLCRLKGIAPLQLGKCSIAGWILGNPECESLFSKKSHSVIAETPNSENGTCYPGYFADYEELREQLSSVSSEFERFEIFPKERSWPKHNVTAGVTASCSHGKSSFYRNLLWLTEKNGSYPNLSKSYVNNKEKEVI


ABD60944:N.....

ABO44134:N.....

ABD95350:N.....


APC57869:N.....

ABD60933:V.....



What differentiates different
cells?

Gene Regulation






- Promoters, Transcriptor factor binding regions are identified by experiments.
 - Collecting samples and sequence alignments etc.

Same content (~25K Genes), different function

Active
NonActive

- **Brain Cells**
 - Amyloid, myosin, α -amylase
- **Muscle Cells**
 - Amyloid, myosin, α -amylase
- **Salivary Gland Cells**
 - Amyloid, myosin, α -amylase



Approximations in Bioinformatics

- 3 → 1 Letter (don't forget the chemistry)

- Protein (20 alphabet)

- Glu-Leu-Val-Ile-Ser-Thr-His-Glu-Lys-Ile-Gln-Gly

- ELVISTHEKING

- DNA/RNA (4 letter alphabet)

- Redundant information of DNA

- storage

ATGGAGCTGTCTTG
TACCTCGACAGAAG

What is the key issue in
Bioinformatics?

Release	Date	Bases (GB)	Seq (GB)	Bases(WGS)	Seq (WGS)
235	12/2019	388,417,258,009	215,333,020	6,277,551,200,690	1,127,023,870

Data Growth in numbers


Enormous data

Need for algorithms to model and carry out analysis

1 Billion
Need for Computers

Storage, Retrieval
and Analysis

1 Million



Whole Genome Shotgun (WGS)
High-throughput sequencing (Illumina etc.)

Image from NCBI
GenBank

Storage/Retrieval



What can we do with the information?
Can we compare/align them?
What do we learn from the alignments?

Comparing 3D is one way
Not all 3D information is available

So, sequence comparison is the common approach

How to find out whether two proteins are related?

Sequence Alignment

DNA can be compared instead of protein. But, for most cases proteins have more information than DNA

- Relatedness (homology) among proteins/DNAs
 - Common function?
 - Homology (common ancestor)
 - When two sequences (proteins/genes) are highly similar, they might be homologous
 - Converse is not true (lack of similarity != No Homology)
 - What is homology?

How can I compare two sequences?

- Not possible without the help of Math and Statistics
- Luckily for us the problem is addressed and some framework is available for us to use

Temple F. Smith



Michael S. Waterman



Creative Commons License

82.88% identity



Human	MVHLTPEEKSAVTALWGKVNVDEVGGEALGRLLVVYPWTQRFFESFGDLSTPDAVMGNPK
RT-Lemur	-TFLTPEENGHVTS LWGKVNVEKVGGEALGRLLVVYPWTQRFFESFGDLSSPDAIMG NPK ..*****;. **:*****::*****:*****:*****
Human	VKAHGKKVLGAFSDGLAHL DNLKGT FATLSELHCDKLHVDPENFRLLGNVLVCVLAHHFG
RT-Lemur	VKAHGKKVLSAFSEGLHHL DNLKGTFAQLSELHCVALHVDPENFKLLGNVLVIVLAHHFG *****.***:** ***** ***** *****:***** *****
Human	KEFTPPVQAAYQKVVAGVANALAHKYH
RT-Lemur	NDFSPQTQAAFQKVVTGVANALAHKYH ::*: * .***:*****:*****

51.37% identity



Human	MVHLTPEEKSAVTALWGKVNVDEVGGEALGRLLVVYPWTQRFFESFGDLSTPDAVMGNPK
Goldfish	-VEWTD AERSAI IGLWGKLN PDELGPQALARCLIVYPWTQRYFATFGNLSSPAAIMG NPK * . * *:***: .*****:* **: * :***. * *:*****:* **:***: * *:*****
Human	VKAHGKKVLGAFSDGLAHL DNLKGT FATLSELHCDKLHVDPENFRLLGNVLVCVLAHHFG
Goldfish	VAAHGRTVMGGLERAIKNMDNIKATYAPLSVMHSEKLHVDPDNFRLLADCITVCAAMKFG * ***:.*:*... .: :***:*. * :*. :*****:*****.: :. * : **
Human	-KEFTPPVQAAYQKVVAGVANALAHKYH
Goldfish	PSGFNADVQEAWQKFLSVVVSALCRQYH . * . ** *:***.: : *...*. : **

21.74% identity



Human	MVHLTPEEKSAVTALWGKVNVDEV---GGEALGRLLVVYPWTQRFFESFGDLSTPDAM
Bloodworm	-MGLSAAQRQVVASTWKDIAGSDNGAGVGKECFKFLSAHHD---IAAVFGFSG-----A
	: *: :...*: : * : : : * *: :*: : : : ** .
Human	GNPKVKAHGKKVLGAFSDGLAHLDNLKGTFATLS-----ELHCDKLHVDPENFRLLGNVL
Bloodworm	SDPGVADLGAKVLAQIGVAVSHLGDEGKMVAEMKAVGVRHKGYGKHIKAEYFEPLGASL
	.* * * ***. :. :*:*: : * :. . . *: . * * ** *
Human	VCVLAHHFGKEFTPPVQAAYQKVVAGVANALAHKYH-
Bloodworm	LSAMEHRIGGKMTAAAKDAWAAAYADISGALISGLQS
	:...: *:*: :*: : : *: . *...:** :

19.85% Identity



Human	MVHLTPEEKSAVTALWGKVN--VDEVGGEALGRLLVVYPWTQRFFESFGDLSTPDAMGN
Soybean	MVAFTEKQDALVSSSF EAFKANIPQYSVVFYTSILEKAPAAKDLFSFLA----NGVDPTN
	** :* :...: *: : : : : : * * : : *. :. . * *
Human	PKVKAHGKKVLGAFSDGLAHLDNLKG--TFATLSELHCDKLHVDPENFRLLGNVLVCVLA
Soybean	PKLTGHA EKLFALVRDSAGQLKASGTTVADAALGSVHAQKAVTDPQ-FVVVKEALLKTIK
	**:.*.*:*: : * . :.*. : *:*.*:*: * ** : * : : *. : :
Human	HHFGKEFT----PPVQAAYQKVVAGVANALAHKYH
Soybean	AAVGDKWSDELSRAWEVAYDELA AAIKKA-----
	.*. : : : ** : : *. : : *


Questions

- What sequences to use and why?
- What types of alignments
 - Global and Local
- Statistics of alignments
 - Scores and matrices

Two Cultures in Bioinformatics

- Two cultures
 - Web-based
 - Point-and-Click (no programming effort)
 - Command line
 - Sometimes steep learning curve (some programming)
- Which one is better?

Fig 1.5 from Bioinformatics and functional genomics / Jonathan Pevsner.— Third edition. Copyright Figure. Please do not distribute.



Validity of Predictions

- Can we use a software as a black-box?
- How do we know whether a software method is working properly?
- Each software team will in most cases do self evaluation
- Sensitivity(TPR) and Specificity(TNR)
 - Sensitivity: Detecting true cases
 - Specificity: Excluding those without disease

$$\text{TPR} = \frac{\text{TP}}{\text{TP} + \text{FN}}$$

$$\text{TNR} = \frac{\text{TN}}{\text{TN} + \text{FP}}$$

Evaluations

Name	Competition
Alignathon	Compare whole-genome sequence alignment methods
EGASP	ENCODE Genome Annotation Assessment Project
Assemblathon	Compare the performance of genome assemblers
GAGE	Genome Assembly Gold-standard Evaluations
ANRF	Assn. of Biomolecular Resource Facilities (ABRF) assessment of phosphorylation
CASP	Critical Assessment of Structure Prediction
CAFA	Critical Assessment of protein Function Annotation algorithms
CAGI	Critical Assessment of Genome Interpretation. Assess computational methods for predicting phenotypic impacts of genomic variation


New Paradigms for Learning Bioinformatics

- Online resources
 - Google
- Online Classes
 - MOOC
 - EdX, Stanford, Coursersa, Udacity etc.
- Programming
 - R, Python, Perl
 - Linux Shell scripting

Two perspectives in Bioinformatics


- Cell Perspective

- Contents of the cell
 - DNA/RNA/Protein
- Analysis of the sequences



- Organism Perspective

- How genes are expressed?
 - Age
 - Different tissues/cells
 - Race
 - Disease vs non-disease states



Command-line interface

- Command line or point-click or application focused
 - What OS?
 - Windows or Linux
 - Why?
 - Cost, ability to carry out tasks

Reproducibility

“More than 70% of researchers have tried and failed to reproduce another scientist’s experiments, and more than half have failed to reproduce their own experiments”

Is there a reproducibility crisis? M.Baker, Nature, 533, 452, 2016

Reproduce another scientist’s experiments (failed to reproduce their own experiment)

Chemistry: 90% (60%)

Biology: 80% (60%)

Physics & Engineering: 70% (50%)

Medicine: 70% (60%)

Earth and Env. Science: 60% (40%)

Reproducibility in Published Papers

- Script availability
 - Supplemental pages is a good place
 - Useful for checking the results
 - Useful for learning/teaching
 - Useful for reviewers
 - Etc.

Reproducibility using R

- R session
 - `sessionInfo()`
- What packages was used
 - `library(??)`
- Show the code
 - Use R command “`dput`” to make the user copy and use your code
- Show comments

Reproducible Code ; S. Ravichandran, Ph.D. 01/23/2017

load libraries at the top of the script

library(rafalib)

if not installed

install.packages("rafalib")

set seed for reproducibility

set.seed(100)

create 100 uniform normally distributed random numbers

x <- rnorm(100)

y <- rnorm(100)

use mypar from rafalib to plot 2 figs in 1 row

mypar(1,2)

hist(x,col="red"); hist(y,col="blue") # use two lines; for lack of space used 1 line

ttest <- t.test(x,y, alternative = "two.sided", conf.level = 0.95) # p-value = 0.94

add an outlier in y and call it my

m_y <- c(y, 150)

use dput(data) if you want to send data via email; ex. dput(m_y)

#use two lines; for lack of space used 1 line

hist(x,col="red"); hist(m_y,col="blue")

otest <- t.test(x, m_y, alternative = "two.sided", conf.level = 0.95) # p-value = 0.94


otest

sessionInfo() # provide sessionInfo()

Application-1

Sequence-based approach


Lactose Intolerance



"It has nothing to do with you, Bessie. It's just that I'm lactose intolerant."

"It has nothing to do with you, Bessie. It's just that I am Lactose intolerant"

Worldwide prevalence of lactose intolerance in recent populations (schematic)



Finland: 1/60K inborns have LCT intolerance


Very common in people of

- West African,
- Arabs,
- Jewish,
- Greek and
- Italian descent.

(ghr.nlm.nih.gov)

Lactose intolerance

- Lactase is the gene that produces Lactase (protein enzyme)
- Lactase digests Lactose → simple sugars



LCT Gene


- Intolerant
 - Gene turned off
- Tolerant (adult)
 - Persistence
- Remedy
 - Lactase pill
 - Which does not interfere with transcription but just provide a supply of Lactase enzyme
 - Need to be taken before the Lactose food

LCT Gene

- Lactase is active during childhood but slows or stops when child grows up for some people
- LCT (short name for the Lactase gene)



– Chr 2; 17 exons



Note the gene direction?

The -14010^*C variant associated with lactase persistence is located between an Oct-1 and HNF1 α binding site and increases lactase promoter activity

Tine G. K. Jensen · Anke Liebert · Rikke Lewinsky ·
Dallas M. Swallow · Jørgen Olsen · Jesper T. Troelsen






Figure 3

Locations of transcription factor-binding sites and predicted adaptive alleles upstream of *LCT*, the lactase gene. Three alleles were identified as potentially causal alleles in the African pastoral populations, whereas C/T-13910 was predicted to be the causal allele in Northern Europeans. Additionally, the T/G-13915 allele is correlated with lactase persistence in the Saudi Arabian population. The transcription factors and the sequence they bind in a supershift assay (48) are: HNF-4 α (–13854 to –13830), HNF-3 α and FOX (–13872 to –13848), Oct-1 and GAGA (–13933 to –13909), and Cdx-2 (–14040 to –14016).



Possible Mechanism

Protein (Activator) that bind in Enhancer regions far away from Promoter (non-binding) and bends and interacts with Promoter (TATA region) to positively affect transcription




<http://www.fda.gov/forconsumers/consumerupdates/ucm094550.htm>

Where is the variation (or lack of) that causes Lactose Persistence (Lactose intolerance)?



Application-2

Sickle Cell Disease (SCD): Structure-based approach



Data Source CDC

Malaria (2015)


214M (World-wide)
~1500 cases in US every year

Sickle-Cell Disease (SCD)

affects ~ 100,000 in US
occurs 1/365 black or African-American births
Occurs 1/16,300 Hispanic-American births



Figure taken from NCBI/NIH

Biology of RBC



S.E.M of
RBC
Wikipedia

- RBC (a.k.a. erythrocytes, haematids etc.)
- Nucleus lacking in human
- 2.4 M raw RBC are produced/second
- Produced in bone marrow and travel all over body carrying O_2 (& CO_2)
- Carrier protein complex: Hemoglobin
 - Not a single gene product ; 2 α and 2 β chains
 - HBA: α ; HBB: β




SCD



2 drops of
oxygenated/deoxygenated blood


https://en.wikipedia.org/wiki/Red_blood_cell

- Genetic disorder
- HBB:Glu7→Val
- Cell sickle & dies sooner




Picture taken from

<http://learn.genetics.utah.edu/content/disorders/singlegene/sicklecell/>





Related Disease



- SCD → Anemia
- Spleen the blood filter will be clogged by the sickle cells and damages → infections
- Malaria is caused by mosquito bites. Parasite invades the RBC and **destroys** it.
- In US disease most commonly affects African American

HBB: Where is it?

Ideogram



NCBI Database

Hemoglobin alpha/beta sequences

10	20	30	40	50	Alpha
MVLSPADKTN	VKAAWGKVGA	HAGEYGAEAL	ERMFLSFPTT	KTYFPHFDLS	
60	70	80	90	100	
HGSAQVKGHG	KKVADALTNA	VAHVDDMPNA	LSALSDLHAH	KLRVDPVNFK	
110	120	130	140		
LLSHCLLVTL	AAHLPAEFTP	AVHASLDKFL	ASVSTVLTSK	YR	

E→V → Sickle Cell Anemia

	10	20	30	40	50	60	
MVHLTP E EKS							Beta
	70	80	90	100	110	120	
VKAHGKKVLG							
	130	140					
KEFTPPVQAA							
YQKVVAGVAN							
ALAHKYH							

CLUSTAL multiple sequence alignment by MUSCLE (3.8)

sp P68871 HBB_HUMAN	VKAHGKKVLGAFSDGLAHLDNLKGTFATLSELHCDKLHVDPENFRLLGNVLVCVLAHHFG
sp P69905 HBA_HUMAN	-----KKVADALTNAVAHVDDMPNALSALSDLHAHKLRVDPVNFKLLSHCLLVTLAAHLPT *** .*:~::~*:~::~ ~::~*:~::~ **.*~::~ *.~::~ *: ~** *:
sp P68871 HBB_HUMAN	KEFTPPVQAAYQKVVAGVANALAHKYH
sp P69905 HBA_HUMAN	AETPAVHASLDKFLASVSTVLTSKYR ***** ~*:~::~ ~*:~::~ ~*:~::~ ~*:~::~ *

43.9% identity

Summary

- What is Bioinformatics?
- Basics of Bioinformatics
- Cell biology
 - DNA/Proteins/RNA
- Central Dogma
- Expression → Function
- Homology, sequence alignment (1D → 3D)
- Applications

Thanks

S. Ravichandran, Ph.D
ravichandran@hood.edu