


Non parametric Model

A **non-parametric algorithm** doesn't assume a **fixed shape for the data**, like a **line or curve**. Instead, **it adapts based on the actual data points available**.

This flexibility helps it work well even if the data has **complex patterns**.

K-Nearest Neighbour (KNN) - Classification

K-Nearest Neighbors (K-NN) is a simple, **non-parametric algorithm** used for **classification** and **regression**.



....

K-Nearest Neighbour (KNN)

Step1:

We have to initialize the K value

K can be **greater than 0** and to **any Number**

$K > 0$

$K = 1, 2, 3, 4, 5, \dots$


K is a **hyper parameter**, so that we can change the **K value** to different data set.

We have to **try** by changing **different K value** and we have to consider that particular **K value** which gives **highest accuracy** .

.....

Let us say we have chosen **K=5**

Step 2: Find the K Nearest neighbour for the New Test data



.....

Step 3:

From the **K=5** values ,How many **neighbor** belong to **0 category** and how many belongs to **1 category**.



Category 1 = 3


Category 0 = 2

Since **Category 1** has more **neighbours** , we can **predict** that the **new data point** belongs to **category1**

.....

To calculate the K-Nearest neighbour we will use **distance formula**.

1.Euclidian distance





$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

.....

2. Manhattan Distance

Also known "taxicab" distance

$$\text{Distance} = |x_2 - x_1| + |y_2 - y_1|$$



K-Nearest Neighbour (KNN) - Regression





If the **K value = 5**

Average of all these points to find out the **O/P** of the **New data point**



k-Nearest Neighbour-Classifier ☆

File Edit View Insert Runtime Tools Help [All changes saved](#)



+ Code + Text



✓
1s

[2] Suggested code may be subject to a license | tayyabmalik4/Data_preprocessing_And_Feature_Engineering_With_Tayyab

```
#importing libraries
```

```
import numpy as np
```

```
import pandas as pd
```

```
import matplotlib.pyplot as plt
```

```
import seaborn as sns
```



✓
0s



```
#import datasets
```

```
from sklearn.datasets import make_classification
```

```
x,y=make_classification(n_samples=1000,n_features=5,n_classes=2,random_state=42)
```



+ Code + Text



```
[9] #Train test split
from sklearn.model_selection import train_test_split
x_train,x_test,y_train,y_test=train_test_split(x,y,test_size=0.3,random_state=42)
```

```
#Import K-Nearest Neighbour classifier

from sklearn.neighbors import KNeighborsClassifier
knn=KNeighborsClassifier(n_neighbors=5,algorithm='auto')
knn.fit(x_train,y_train)
```



▼ KNeighborsClassifier ⓘ ?
KNeighborsClassifier()

Model will consider **the 5 nearest neighbors** of any new data point to predict its class label.

Allows the model to select the **most efficient algorithm** automatically (either **ball tree, kd_tree, or brute force**), based on the **data size and structure**.



+ Code + Text



```
[11] #prediction
y_pred=knn.predict(x_test)
```

```
Suggested code may be subject to a license | BHATNISAR/MARKETING-LEAD-CLASSIFIER
#perfomance metrics
from sklearn.metrics import accuracy_score,confusion_matrix,classification_report
print(accuracy_score(y_test,y_pred))
print(confusion_matrix(y_test,y_pred))
print(classification_report(y_test,y_pred))
```

```
⇒ 0.8966666666666666
[[137  9]
 [ 22 132]]
```

		precision	recall	f1-score	support
	0	0.86	0.94	0.90	146
	1	0.94	0.86	0.89	154
accuracy				0.90	300
macro avg		0.90	0.90	0.90	300
weighted avg		0.90	0.90	0.90	300

How to find the best K value?



Suggested code may be subject to a license | | Genedy-yonis/Step-by-Step-Diabetes-Classification-KNN-detailed

```
#grid search
from sklearn.model_selection import GridSearchCV
# Define the parameter grid with possible values for n_neighbors
param_grid = {'n_neighbors': range(1, 31)} # Testing values of K from 1 to 30

# Initialize the KNeighborsClassifier
knn = KNeighborsClassifier()

# Set up GridSearchCV without cross-validation (cv=1)
knn_cv = GridSearchCV(knn, param_grid, cv=5)

# Fit the model on the data
knn_cv.fit(x, y)

# Get the best K value
best_k = knn_cv.best_params_['n_neighbors']
print("Best K value:", best_k)
```

You define a **range** of possible values for each **hyperparameter** (like **n_neighbors** in **KNN**) in a **parameter grid**.



```
Best K value: 10
/usr/local/lib/python3.10/dist-packages/numpy/ma/core.py:2820: RuntimeWarning: invalid value encountered in cast
  _data = np.array(data, dtype=dtype, copy=copy,
```

Bayes Theorem

Probability: Is a measure of how likely an event is to happen

Take a Coin



S = {Head, Tail}

n(S) => Total No of Chances = **2**

A = Event of Getting Head

n(A) => Chance of getting head = **1**

$$\mathbf{P(A)} = \frac{\text{No of Chances for an Event}}{\text{Total No of Chances}}$$

$$\mathbf{P(A)} = \frac{1}{2} = \mathbf{0.5}$$

.....

Independent Events: The probability of one event occurring doesn't change based on the occurrence of the other.

For independent events A and B

Let A be the event of flipping a coin and getting heads.

Let B be the event of rolling a die and getting a 6.

These events are independent because the outcome of the coin flip doesn't influence the outcome of the die roll.

$$P(A \cap B) = P(A) \times P(B)$$

This means the probability of both events happening together is simply the product of their individual probabilities.

.....

Dependent Events: Where the occurrence of one event affects the probability of the other event happening.

For dependent events A and B , the probability of both occurring is:

$$P(A \cap B) = P(A) \times P(B|A)$$

$P(B|A)$: The conditional probability of B given that A has already occurred.

$P(A)$: Probability that event A occurs

$P(A \cap B)$: Probability that both A and B occur together.

.....

Conditional Probability: Is the probability of an event happening, given that another event has already occurred.

Event A: It's raining

Event B: You carry an umbrella.

Probability of **Event B** occurring, given that **Event A** has already occurred.

$$P(B | A) = \frac{P(B \cap A)}{P(A)}$$

P(B|A): Conditional probability of B given A.

P(B ∩ A): Probability that both A and B occur together.

P(A): Probability that event A occurs.

.....

Bayes Theorem Derivation:

$$P(B|A) = \frac{P(B \cap A)}{P(A)} \quad \text{Conditional probability of } B \text{ given } A.$$


$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad \text{Conditional probability of } A \text{ given } B.$$

$$P(A \cap B) = P(B \cap A) \quad \text{Both are same}$$

$$P(A|B) * P(B) = P(B|A) * P(A)$$

$$P(A|B) = \frac{P(B|A) * P(A)}{P(B)}$$

Bayes Theorem Formula:



The diagram shows the Bayes Theorem formula enclosed in an orange rectangular box. Four blue arrows point from labels outside the box to parts of the formula: 'Likely hood' points to $P(B|A)$, 'Prior Probability' points to $P(A)$, 'Posterior Probability' points to $P(A|B)$, and 'Marginal Probability' points to $P(B)$.

$$\mathbf{P(A|B)} = \frac{P(B|A) * P(A)}{P(B)}$$

Likely hood

Prior Probability

Posterior Probability

Marginal Probability

Naïve Bayes Classifier

The **Naive Bayes classifier** is a simple, **probabilistic classification algorithm** based on **Bayes' theorem**.

Types of Naive Bayes

- Gaussian Naive Bayes
- Multinomial Naive Bayes
- Bernoulli Naive Bayes

.....

Given Data set :

X = {**x1** , **x2** ,**x3**.....**xn**}

Independent Features

y

Dependent Variable

The formula for calculating the **probability** of a **class Y**

$$P(A|B) = \frac{P(B|A) * P(A)}{P(B)}$$

$$P(Y/(x1,x2,x3.....xn)) = \frac{P(x1|y)P(x2|y)P(x3|y).....P(xn|y) * P(Y)}{P(x1,x2,x3.....xn)}$$

Example- Naïve Bayes Classifier

Here's a sample dataset with features like **Outlook**, **Temperature**, **Humidity**, and **Windy** that are used to predict whether **Play Tennis** will be **Yes or No**.

Outlook	Temperature	Humidity	Windy	PlayTennis
Sunny	Hot	High	False	No
Sunny	Hot	High	True	No
Overcast	Hot	High	False	Yes
Rainy	Mild	High	False	Yes
Rainy	Cool	Normal	False	Yes
Rainy	Cool	Normal	True	No
Overcast	Cool	Normal	True	Yes
Sunny	Mild	High	False	No
Sunny	Cool	Normal	False	Yes
Rainy	Mild	Normal	False	Yes
Sunny	Mild	Normal	True	Yes
Overcast	Mild	High	True	Yes
Overcast	Hot	Normal	False	Yes
Rainy	Mild		True	No

.....

We want to **predict** Play Tennis for the following **conditions**?

Conditions=(Outlook = **Sunny**,Temperature = **Cool**,Humidity = **High**,Windy = **True**)

.....

- **Step-by-Step Naive Bayes Calculation**

Step 1: Calculate **Prior Probabilities**

calculate the probability of each class **$P(\text{Play Tennis}=\text{Yes})$** and

$P(\text{Play Tennis}=\text{No})$

Total Yes: 9 occurrences

$$P(\text{Yes}) = \frac{9}{14} = 0.64$$

Total No: 5 occurrences

$$P(\text{No}) = \frac{5}{14} = 0.35$$

Total records: 14

.....

Step 2: Calculate Likelihoods

Calculate the **likelihood** of each feature.

For Play Tennis = Yes:

$$P(\text{Outlook}=\text{Sunny}|\text{Yes}) = \frac{2}{9}$$

$$P(\text{Temperature}=\text{Cool}|\text{Yes}) = \frac{3}{9}$$

$$P(\text{Humidity}=\text{High}|\text{Yes}) = \frac{3}{9}$$

$$P(\text{Windy}=\text{True}|\text{Yes}) = \frac{3}{9}$$

.....

For Play Tennis = No:

$$P(\text{Outlook}=\text{Sunny} \mid \text{No})=\frac{3}{5}$$

$$P(\text{Temperature}=\text{Cool} \mid \text{No})=\frac{1}{5}$$

$$P(\text{Humidity}=\text{High} \mid \text{No})=\frac{4}{5}$$

$$P(\text{Windy}=\text{True} \mid \text{No})=\frac{2}{5}$$

.....

Step 3: Apply Bayes' Theorem

To classify, we find $P(\text{Yes}|\text{conditions})$ and $P(\text{No}|\text{conditions})$ and choose the one with the **higher probability**.

Conditions=(Outlook = **Sunny**,Temperature = **Cool**,Humidity = **High**,Windy = **True**)

$$\begin{aligned} P(\text{Yes}|\text{conditions}) &= \frac{P(\text{Sunny}|\text{yes})P(\text{cool}|\text{yes})P(\text{High}|\text{yes})P(\text{True}|\text{yes}) * P(\text{yes})}{P(\text{conditions})} \\ &= \frac{2}{9} \frac{3}{9} \frac{3}{9} \frac{3}{9} * 0.64 \\ &= \mathbf{0.0053} \end{aligned}$$

.....

$$\begin{aligned} \mathbf{P(\text{No} | \text{Conditions})} &= \frac{P(\text{Sunny}|\text{No})P(\text{cool}|\text{No})P(\text{High}|\text{No})P(\text{True}|\text{No})*P(\text{No})}{P(\text{conditions})} \\ &= \frac{3}{5} \frac{1}{5} \frac{4}{5} \frac{2}{5} * 0.36 \\ &= \mathbf{0.0115} \end{aligned}$$





Step 4: Compare and Classify

Since $P(\text{No}|\text{conditions}) > P(\text{Yes}|\text{conditions})$, the classifier predicts **Play Tennis = No** for these conditions.

Naive Bayes uses conditional probabilities to classify data.

Normal or Gaussian Distribution

In a normal distribution, the **mean (average)**, located at the **center** of the distribution.



Probability Density Function of the Normal distribution

PDF used to calculate the probability of the **Continuous Random variable** falling within a certain range.

For a **normal distribution** with mean μ and standard deviation σ , the PDF is given by

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right)$$

where:

- x is the value for which we want to find the probability density.
- μ is the mean (average) of the distribution.
- σ is the standard deviation, which measures the spread or "width" of the distribution.
- $\exp$ represents the exponential function, which can also be written as $e^{-\text{term}}$, where e is Euler's number (approximately 2.718).

Example - Gaussian Naïve Bayes Classification

Height	Weight	Foot Size	Gender
6.0	180	12	Male
5.92	190	11	Male
5.58	170	12	Male
5.92	165	10	Male
5.0	100	6	Female
5.5	150	8	Female
5.42	130	7	Female
5.75	150	9	Female

.....

Calculate Mean and Standard Deviation for Males and Females

- Mean (μ) for a feature:

$$\mu = \frac{\sum_{i=1}^N x_i}{N}$$

where x_i is each individual value, and N is the total number of values.

- Standard Deviation (σ) for a feature:

$$\sigma = \sqrt{\frac{\sum_{i=1}^N (x_i - \mu)^2}{N}}$$

Height	Weight	Foot Size	Gender
6.0	180	12	Male
5.92	190	11	Male
5.58	170	12	Male
5.92	165	10	Male
5.0	100	6	Female
5.5	150	8	Female
5.42	130	7	Female
5.75	150	9	Female

①

Step 1: Calculate Mean and Standard Deviation for Males

Height (Male)

1. Mean:

$$\begin{aligned}\mu_{\text{height, Male}} &= \frac{6.0 + 5.92 + 5.58 + 5.92}{4} \\ &= \frac{23.42}{4} = \boxed{5.855}\end{aligned}$$

2. Standard Deviation

$$\begin{aligned}\sigma_{\text{height, Male}} &= \sqrt{\frac{(6.0 - 5.855)^2 + (5.92 - 5.855)^2}{4} + \frac{(5.58 - 5.855)^2 + (5.92 - 5.855)^2}{4}} \\ &= \sqrt{\frac{(0.021025) + (0.004225) + (0.075625) + (0.004225)}{4}} \\ &= \sqrt{\frac{0.1051}{4}} = \boxed{0.162}\end{aligned}$$

Weight (Male)

1. Mean:

$$\begin{aligned}\mu_{\text{weight, Male}} &= \frac{180 + 190 + 170 + 165}{4} \\ &= \frac{705}{4} = \boxed{176.25}\end{aligned}$$

2. Standard Deviation:

$$\sigma_{\text{weight, Male}} = \sqrt{\frac{(180 - 176.25)^2 + (190 - 176.25)^2 + (170 - 176.25)^2 + (165 - 176.25)^2}{4}}$$

$$\begin{aligned}\sigma_{\text{weight, male}} &= \sqrt{\frac{14.0625 + 189.0625 + 39.0625 + 126.5625}{4}} \\ &= \sqrt{\frac{268.75}{4}} \\ &= \sqrt{67.1875} \\ &= \boxed{8.2}\end{aligned}$$

Foot size (Male)

③

1. Mean

$$\mu_{\text{foot size, male}} = \frac{12 + 11 + 12 + 10}{4} = \frac{45}{4} = 11.25$$

2. Standard Deviation

$$\begin{aligned}\sigma_{\text{foot size, male}} &= \sqrt{\frac{(12-11.25)^2 + (11-11.25)^2 + (12-11.25)^2 + (10-11.25)^2}{4}} \\ &= \sqrt{\frac{0.5625 + 0.0625 + 0.5625 + 1.5625}{4}} \\ &= \sqrt{\frac{2.75}{4}} = 0.83\end{aligned}$$

Step 2: Calculate Mean and Standard deviation for females

Height (Female)

1. Mean

$$\begin{aligned}\mu_{\text{height, female}} &= \frac{5.0 + 5.5 + 5.42 + 5.75}{4} \\ &= \frac{21.67}{4} = \boxed{5.4175}\end{aligned}$$

2. Standard Deviation:

$$\begin{aligned}\sigma_{\text{height, female}} &= \sqrt{\frac{(5.0 - 5.4175)^2 + (5.5 - 5.4175)^2}{4} + \frac{(5.42 - 5.4175)^2 + (5.75 - 5.4175)^2}{4}} \\ &= \boxed{0.278}\end{aligned}$$

(5)

Weight (female)1. Mean

$$\mu_{\text{weight, female}} = \frac{100 + 150 + 130 + 150}{4}$$

$$= \frac{530}{4} = \boxed{132.5}$$

2. Standard deviation

$$\sigma_{\text{weight, female}} = \sqrt{\frac{(100 - 132.5)^2 + (150 - 132.5)^2 + (130 - 132.5)^2 + (150 - 132.5)^2}{4}}$$

$$= \boxed{21.6}$$

Foot size (Female)

6

1. Mean

$$\begin{aligned}\mu_{\text{Foot size, Female}} &= \frac{6 + 8 + 7 + 9}{4} \\ &= \boxed{7.5}\end{aligned}$$

2. Standard Deviation

$$\begin{aligned}\sigma_{\text{Foot size, Female}} &= \sqrt{\frac{(6-7.5)^2 + (8-7.5)^2}{4} + (7-7.5)^2 + (9-7.5)^2} \\ &= \boxed{1.29}\end{aligned}$$

Male

7

Height		Weight		Foot size	
μ	σ	μ	σ	μ	σ
5.855	0.162	176.25	9.6	11.25	0.83

Female

Height		Weight		Foot size	
μ	σ	μ	σ	μ	σ
5.4175	0.278	132.5	21.6	7.5	1.29

Step-3 Define Bayes' Theorem

$$P(A|B) = \frac{P(B|A) * P(A)}{P(B)}$$

$$P(\text{Gender}|\text{Features}) = \frac{P(\text{Features}|\text{Gender}) \cdot P(\text{Gender})}{P(\text{Features})}$$

Here:

- $P(\text{Gender}|\text{Features})$ is the **posterior probability** (probability of gender given features).
- $P(\text{Features}|\text{Gender})$ is the **likelihood** (probability of features given gender).
- $P(\text{Gender})$ is the **prior probability** of each gender.
- $P(\text{Features})$ is the **evidence** (total probability of features).

Step-4 Calculate the Prior Probability

Assume we have 4 males and 4 females in our sample dataset, so:

$$P(\text{Male}) = \frac{4}{8} = 0.5, \quad P(\text{Female}) = \frac{4}{8} = 0.5$$

Step-5 Calculate the Likelihoods Using Gaussian PDF

The probability density function for a Gaussian distribution is:

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right)$$

where:

- x is the feature value (e.g., height, weight, foot size),
- μ is the mean of the feature for the gender,
- σ is the standard deviation of the feature for the gender.

Step-6 Calculate the Probability for Each Feature Given Each Gender

- Given a sample data, calculate the **likelihood** for each feature given each gender.

New Data

Height	Weight	Foot size	Gender
6.0	130	8	?

Gaussian Naive bayes Model should predict
the Gender for New data

Likelihood for each feature given each gender

for Male:

1. Height:

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$$

$$P(\text{height} = 6.0 | \text{Male}) = \frac{1}{\sqrt{2\pi(0.162)^2}} \exp\left(-\frac{(6.0 - 5.855)^2}{2(0.162)^2}\right)$$

$$P(\text{height} = 6.0 | \text{Male}) = \frac{1}{0.4064} \times 0.67 \approx$$

$$= 1.65$$

2. Weight

$$P(\text{Weight} = 130 | \text{Male}) = \frac{1}{\sqrt{2\pi(9.6)^2}} \exp\left(-\frac{(130 - 176.25)^2}{2 \times (9.6)^2}\right)$$

$$= \frac{1}{24.57} \times 6.6 \times 10^{-5} = \boxed{0.000276}$$

3. Foot size

$$P(\text{Foot size} = 8 | \text{Male}) = \frac{1}{\sqrt{2\pi(0.83)^2}} \exp\left(-\frac{(8 - 11.25)^2}{2 \times (0.83)^2}\right)$$

$$P(\text{Foot size} = 8 | \text{Male}) = \boxed{0.000251}$$

(11)

For female1. Height

$$P(\text{Height} = 6.0 | \text{female}) = \frac{1}{\sqrt{2\pi(0.278)^2}} \exp\left(-\frac{(6.0 - 5.417)^2}{2 \times (0.278)^2}\right)$$

$$P(\text{Height} = 6.0 | \text{female}) = \boxed{0.0041}$$

2. Weight

$$P(\text{Weight} = 130 | \text{female}) = \frac{1}{\sqrt{2\pi(21.6)^2}} \exp\left(-\frac{(130 - 132.5)^2}{2 \times (21.6)^2}\right)$$

$$P(\text{Weight} = 130 | \text{female}) = \boxed{0.0184}$$

(2)

3. Foot size

$$P(\text{Foot size} = 8 | \text{Female}) = \frac{1}{\sqrt{2\pi(1.29)^2}} \exp\left(-\frac{(8-7.5)^2}{2 \times (1.29)^2}\right)$$

$$P(\text{Foot size} = 8 | \text{Female}) = \boxed{0.303}$$

Step-7 Combine the Likelihoods

Multiply the likelihoods for each feature to get the combined likelihood for each gender:

$$P(\text{Features}|\text{Male}) = P(\text{height}|\text{Male}) \times P(\text{weight}|\text{Male}) \times P(\text{foot size}|\text{Male})$$

$$P(\text{Features}|\text{Female}) = P(\text{height}|\text{Female}) \times P(\text{weight}|\text{Female}) \times P(\text{foot size}|\text{Female})$$

Compute Posterior Probability

1. for male

$$P(\text{Male} | \text{Features}) = P(\text{height} | \text{Male}) \times P(\text{weight} | \text{Male}) \times P(\text{Foot size} | \text{Male}) \times P(\text{Male})$$

$$= 1.65 \times 0.000276 \times 0.000251 \times 0.5$$

$$= \boxed{5.72 \times 10^{-8}} \Rightarrow 0.000000572$$

2. for female

$$P(\text{Female} | \text{Features}) = P(\text{height} | \text{Female}) \times P(\text{weight} | \text{Female}) \times P(\text{Foot size} | \text{Female}) \times P(\text{Female})$$

$$= 0.0041 \times 0.0184 \times 0.303 \times 0.5$$

$$= \boxed{1.14 \times 10^{-5}} \Rightarrow 0.0000114$$

Step-7 Compute posterior probability using Bayes theorem

Finally, use Bayes' Theorem to get the posterior probabilities:

$$P(\text{Male}|\text{Features}) = \frac{P(\text{Features}|\text{Male}) \cdot P(\text{Male})}{P(\text{Features})}$$

$$P(\text{Female}|\text{Features}) = \frac{P(\text{Features}|\text{Female}) \cdot P(\text{Female})}{P(\text{Features})}$$

Since $P(\text{Features})$ is the same for both classes, we can ignore it for comparison. Whichever posterior probability is higher determines the predicted gender.

Step-8 Determine Classification

Since $P(\text{Female}|\text{Features}) > P(\text{Male}|\text{Features})$, the classifier would predict the gender as **Female** for this sample.

+ Code + Text

classify gender based on features like height, weight, and foot size

```
[1] import pandas as pd

# Sample dataset
data = {
    'height': [6.0, 5.92, 5.58, 5.92, 5.0, 5.5, 5.42, 5.75],
    'weight': [180, 190, 170, 165, 100, 150, 130, 150],
    'foot_size': [12, 11, 12, 10, 6, 8, 7, 9],
    'gender': ['male', 'male', 'male', 'male', 'female', 'female', 'female', 'female']
}
df = pd.DataFrame(data)
```

```
#Display the data frame
df
```

	height	weight	foot_size	gender
0	6.00	180	12	male
1	5.92	190	11	male
2	5.58	170	12	male
3	5.92	165	10	male
4	5.00	100	6	female
5	5.50	150	8	female

```
[4] #Encode Gender Labels
from sklearn.preprocessing import LabelEncoder
#convert the gender labels into numeric format, where 0 represents female and 1 represents male.
le = LabelEncoder()
df['gender'] = le.fit_transform(df['gender'])
```



df



	height	weight	foot_size	gender
0	6.00	180	12	1
1	5.92	190	11	1
2	5.58	170	12	1
3	5.92	165	10	1
4	5.00	100	6	0
5	5.50	150	8	0
6	5.42	130	7	0
7	5.75	150	9	0



✓
0s

```
[6] #Split Features and Labels  
#Separate the features (height, weight, foot_size) and the label (gender)  
X = df[['height', 'weight', 'foot_size']]  
y = df['gender']
```

✓
0s

▶ #Train the Gaussian Naive Bayes Classifier
#Using GaussianNB from scikit-learn, train the model on the data.

```
from sklearn.naive_bayes import GaussianNB  
  
# Initialize the Gaussian Naive Bayes classifier  
clf = GaussianNB()  
clf.fit(X, y)
```



▼ GaussianNB ⓘ ?
GaussianNB()



#Make Predictions

#After training, you can use the model to predict the gender based on new input data.

New data for prediction

```
new_data = [[6.0, 130, 8]] # height=6.0, weight=130, foot_size=8
```

Predict the gender

```
predicted_gender = clf.predict(new_data)
```

```
predicted_gender_label = le.inverse_transform(predicted_gender)
```

```
print("Predicted gender:", predicted_gender_label)
```



Predicted gender: ['female']

/usr/local/lib/python3.10/dist-packages/sklearn/base.py:493: UserWarning: X does not have valid feature names, but GaussianNB was fitted with feature names
warnings.warn(