

The Lund string: a linear potential, a world sheet that breaks, a logarithmic multiplicity (page 1)


(a) the string: energy grows linearly with length




(b) the break-up vertices of the 28 GeV u sbar string





(c) how many hadrons a string makes (gluons a



The break-up: flavour by tunnelling, transverse momentum by a Gaussian, species by spin counting (page 2)









The fragmentation function and the iteration from both ends (page 3)


(a) Lund symmetric $f(z) = \frac{1}{z}(1-z)^a \exp(-b\, m_T^2/z)$, $a = 0.68$, $b = 0.98$




(b) the fractions actually drawn




(c) the b quark's string (69 GeV) runs out: 16 hadrons




Gluons as kinks: regions of the world sheet, hadrons across the kinks (page 4)




(a) the *b* quark's fragmentation function at $m_T = m_{B^*}$




(b) the energy the B hadron keeps, same partons every time





(c) which B hadron: vector 2.2 : 1, strange 0.217, baryon 0



x_B in different settings: the hadron, the jet, the ensemble (page 6)









Our jet at this stage: the hadron-level family and the B hadron's share (page 7)

(a) the cone p_T before (dashed line) and after hadronisation



(b) more particles, more mass, same p_T



(c) the ledger: quark, hadron, jet

