

Two Strata Case

Ravjot Singh Kohli

Apr 2023

1 Kähler Reduction on \mathbb{C}^n

The simplest example of Kähler reduction with an isolated singular point is a linear action. Let (z^1, \dots, z^n) be holomorphic coordinates on \mathbb{C}^n . The standard Kähler form on \mathbb{C}^n is given by,

$$(\omega_{std})_z = \sum_{k=1}^n \frac{i}{2} dz^k \wedge d\bar{z}^k \quad (1)$$

The complex structure in these coordinates is given by

$$J_{std} \left(\frac{\partial}{\partial z^j} \right) = i \frac{\partial}{\partial z^j}, \quad J_{std} \left(\frac{\partial}{\partial \bar{z}^j} \right) = -i \frac{\partial}{\partial \bar{z}^j} \quad (2)$$

The standard metric on \mathbb{C}^n is given by

$$g_{std}(z) = \omega_{std}(-, J_{std}-) = \sum_{k=1}^n dz^k \otimes d\bar{z}^k \quad (3)$$

Recall that $\mathbb{C}^n \setminus \{0\} \simeq (0, \infty) \times S^{2n-1}$ with coordinates on the right given by polar coordinates (r, θ) . Here $r = |z|$ and θ denotes the coordinates on S^{2n-1} . The standard metric in these coordinates takes the form

$$g_{std}(r, \theta) = dr^2 + r^2 g_{S^{2n-1}}(\theta) \quad (4)$$

Now let $G \subset U(n)$ be a compact Lie group acting on \mathbb{C}^n . We assume that apart from the fixed point set, the action is free. Hence we have two strata given by the orbit types, $(\mathbb{C}^n)_G$ of orbit type (G) and $(\mathbb{C}^n)_e = (\mathbb{C}^n) \setminus (\mathbb{C}^n)_G$ of orbit type (e) where $e \in G$ is the identity element. Note that $(\mathbb{C}^n)_G$ is a linear **symplectic subspace**. Let $W := ((\mathbb{C}^n)_G)^\perp$ denote the perpendicular

subspace with respect to the standard metric on \mathbb{C}^n . Then W is a **symplectic subspace** as well. We can write the stratum $(\mathbb{C}^n)_e$ as the product

$$(\mathbb{C}^n)_e = (\mathbb{C}^n)_G \times (W \setminus \{0\}) \quad (5)$$

Note that the standard Riemannian metric and the symplectic form restricted to $(\mathbb{C}^n)_e$ is then compatible with respect to the above decomposition. Let (u_1, \dots, u_m) be the coordinates for $(\mathbb{C}^n)_G$ and let (w_1, \dots, w_l) the coordinates for W . Then for a point $z = (u, w) \in (\mathbb{C}^n)_e$,

$$g_{std}(u, w) = \sum_{k=1}^m du^k \otimes d\bar{u}^k + \sum_{k=1}^l dw^k \otimes d\bar{w}^k \quad (6)$$

Let $A \in \mathfrak{g} \subset \mathfrak{u}(n)$. The vector field generated on \mathbb{C}^n by the group action, denoted $A_{\mathbb{C}^n}$, is given by

$$A_{\mathbb{C}^n}(z) = \left. \frac{d}{dt} \right|_{t=0} \exp(tA) \cdot z = \begin{cases} 0 & z \in (\mathbb{C}^n)_G \\ A \cdot z & z \in (\mathbb{C}^n)_e \end{cases} \quad (7)$$

The moment map with respect to this action is given by

$$\Phi_{std}(z)(A) = (\omega_{std})_z(A_{\mathbb{C}^n}(z), z)$$

where $A \in \mathfrak{g} \subset \mathfrak{u}(n)$ is a skew-Hermitian matrix. From (7), we see that $(\mathbb{C}^n)_G \subset \Phi_{std}^{-1}(0)$.

Lemma 1.1. *The zero level set $Z_{std} := \Phi_{std}^{-1}(0)$ is a cone, i.e., $Z_{std} \simeq [0, \infty) \times L$ where $L = Z_{std} \cap S^{2n-1}$*

Proof. Let $p \in Z_{std}$. Consider the scalar multiplication of \mathbb{R}^+ on \mathbb{C}^n denoted by $t \cdot p$. Then we have

$$\begin{aligned} \Phi_{std}(t \cdot p)(A) &= (\omega_{std})_{t \cdot p}(A_{\mathbb{C}^n}(t \cdot p), t \cdot p) \\ &= (\omega_{std})_p(t \cdot A_{\mathbb{C}^n}(p), t \cdot p) \\ &= t^2 \Phi_{std}(p)(A) \\ &= 0 \end{aligned}$$

where we have used that $(\omega_{std})_p$ is independent of the point $p \in \mathbb{C}^n$ and that scalar multiplication on \mathbb{C}^n is linear and commutes with the action of $\mathfrak{u}(n)$. \square

Note that $0 \in Z_{std} \subset \mathbb{C}^n$ is the only singular point and $Z_{std} \setminus \{0\}$ is a smooth manifold. The reduced space defined as the quotient $\pi : Z_{std} \rightarrow Z_{std}/G =: (\mathbb{C}^n)_0$ has two strata, the point $\pi(0)$ and the rest. Hence outside the point $\pi(0)$, we can talk about the symplectic form $(\omega_{std})_0$ and the Riemannian metric $(g_{std})_0$ on the manifold $(\mathbb{C}^n)_0 \setminus \{\pi(0)\}$ coming from smooth Kähler reduction.

Note that the G -action on $Z_{std} \setminus \{0\} \simeq (0, \infty) \times L$ acts only on the link L and so

$$(\mathbb{C}^n)_0 \setminus \{\pi(0)\} \simeq (0, \infty) \times (L/G) \quad (8)$$

Now combining ??, ??, and the above decomposition, we get that the metric on the reduced space can be written as

$$(g_{std})_0(r, \phi) = dr^2 + r^2 g_{L/G}(\phi) \quad (9)$$

where $g_{L/G}$ is the quotient metric on manifold L/G (with coordinates (ϕ)).