

Measure Theory – 3

Independence of Events, Events Occurring Infinitely Often,
Measurable Functions and Piecewise Maps,
Induced Measures, and Composition of Random Variables

Ravleen Bajaj

Introduction

These notes collect the core definitions, results, and worked problems on independence of events, the probability of events occurring infinitely often, measurable functions (including piecewise-defined maps), induced (push-forward) measures, random variables defined by cases on an event, and the law of a composition $Y \circ X$. Each section states the theory first, then works the associated problem, and closes with remarks and common pitfalls.

Throughout, $(\Omega, \mathcal{F}, \mathbb{P})$ is a probability space, $\mathcal{B}_{\mathbb{R}}$ is the Borel σ -algebra on $\mathbb{R}$, and λ is Lebesgue measure.

1 Independence of Events

1.1 Definition and Complements

Definition 1.1 (Mutual independence). Events $A_1, \dots, A_m \in \mathcal{F}$ are **mutually independent** if for every non-empty $I \subseteq \{1, \dots, m\}$,

$$\mathbb{P}\left(\bigcap_{i \in I} A_i\right) = \prod_{i \in I} \mathbb{P}(A_i).$$

Lemma 1.2 (Complements of independent events). *If $A_1, \dots, A_m$ are mutually independent, then so are $A_1^c, \dots, A_m^c$.*

Proof. It suffices to replace one event at a time. Fix I containing 1, put $J = I \setminus \{1\}$ and $C = \bigcap_{i \in J} A_i$ (with $C = \Omega$ if $J = \emptyset$). Since $C = (A_1 \cap C) \sqcup (A_1^c \cap C)$,

$$\mathbb{P}(A_1^c \cap C) = \mathbb{P}(C) - \mathbb{P}(A_1) \mathbb{P}(C) = \mathbb{P}(A_1^c) \prod_{i \in J} \mathbb{P}(A_i).$$

So $A_1^c, A_2, \dots, A_m$ are mutually independent; repeat for each index. □

1.2 Problem: Probability of a Finite Union

Let $A_1, \dots, A_m$ be mutually independent events in $(\Omega, \mathcal{F}, \mathbb{P})$.

Key Result.

$$\mathbb{P}\left(\bigcup_{i=1}^m A_i\right) = 1 - \prod_{i=1}^m (1 - \mathbb{P}(A_i)).$$

Proof. 1. **De Morgan.** $(\bigcup_{i=1}^m A_i)^c = \bigcap_{i=1}^m A_i^c$.

2. **Complement rule.** Using $\mathbb{P}(E^c) = 1 - \mathbb{P}(E)$,

$$\mathbb{P}\left(\bigcup_{i=1}^m A_i\right) = 1 - \mathbb{P}\left(\bigcap_{i=1}^m A_i^c\right).$$

3. **Independence.** By Lemma 1.2,

$$\mathbb{P}\left(\bigcap_{i=1}^m A_i^c\right) = \prod_{i=1}^m \mathbb{P}(A_i^c) = \prod_{i=1}^m (1 - \mathbb{P}(A_i)).$$

4. **Substitute** Step 3 into Step 2. □

Example 1.3. Roll a fair die m times; let $A_i = \{\text{six on roll } i\}$. Then $\mathbb{P}(\text{at least one six}) = 1 - (5/6)^m$, which is $1 - 625/1296 \approx 0.518$ for $m = 4$. If $\mathbb{P}(A_i) \geq p > 0$ for all i , the union has probability at least $1 - (1 - p)^m \rightarrow 1$.

Common pitfalls.

- Pairwise independence is *not* enough: the product rule must hold for every subfamily.
- The formula $\mathbb{P}(\bigcup A_i) = \sum \mathbb{P}(A_i)$ requires *disjoint* events, not independent ones.

2 Events Occurring Infinitely Often

2.1 Definitions

For events $\{A_n\}_{n \geq 1}$ in $(\Omega, \mathcal{F}, \mathbb{P})$:

$$\begin{aligned} \{A_n \text{ i.o.}\} &= \limsup_{n \rightarrow \infty} A_n = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k, \\ \liminf_{n \rightarrow \infty} A_n &= \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k. \end{aligned}$$

- $\omega \in \{A_n \text{ i.o.}\}$ iff ω lies in **infinitely many** A_n .

- $\omega \in \liminf A_n$ iff ω lies in **all but finitely many** A_n .

Both sets are countable unions/intersections of events, hence belong to $\mathcal{F}$. Also $\liminf A_n \subseteq \limsup A_n$, and this inclusion can be strict.

2.2 Problem: A Lower Bound on $\mathbb{P}(A_n \text{ i.o.})$

Put $B_n = \bigcup_{k \geq n} A_k$. Then $B_1 \supseteq B_2 \supseteq \dots$ and $\limsup A_n = \bigcap_n B_n$.

Key Result.

If there exist $\epsilon > 0$ and N with $\mathbb{P}(A_n) \geq \epsilon$ for all $n \geq N$, then

$$\mathbb{P}(A_n \text{ i.o.}) \geq \epsilon.$$

Proof. For $n \geq N$, $A_n \subseteq B_n$, so $\mathbb{P}(B_n) \geq \mathbb{P}(A_n) \geq \epsilon$. Since $\mathbb{P}$ is finite and $B_n \downarrow \limsup A_n$, continuity from above gives

$$\mathbb{P}(A_n \text{ i.o.}) = \lim_{n \rightarrow \infty} \mathbb{P}(B_n) \geq \epsilon. \quad \square$$

The same argument gives the general inequality

$$\mathbb{P}(\liminf A_n) \leq \liminf_n \mathbb{P}(A_n) \leq \limsup_n \mathbb{P}(A_n) \leq \mathbb{P}(\limsup A_n).$$

2.3 Borel–Cantelli Lemmas

Theorem 2.1 (First Borel–Cantelli). *If $\sum_n \mathbb{P}(A_n) < \infty$, then $\mathbb{P}(A_n \text{ i.o.}) = 0$.*

Proof. For each n , $\mathbb{P}(A_n \text{ i.o.}) \leq \mathbb{P}(B_n) \leq \sum_{k \geq n} \mathbb{P}(A_k) \rightarrow 0$, the tail of a convergent series. $\square$

Theorem 2.2 (Second Borel–Cantelli). *If the A_n are independent and $\sum_n \mathbb{P}(A_n) = \infty$, then $\mathbb{P}(A_n \text{ i.o.}) = 1$.*

Proof. By De Morgan, $\{A_n \text{ i.o.}\}^c = \bigcup_n \bigcap_{k \geq n} A_k^c$. For $m \geq n$, independence and $1 - t \leq e^{-t}$ give

$$\mathbb{P}\left(\bigcap_{k=n}^m A_k^c\right) = \prod_{k=n}^m (1 - \mathbb{P}(A_k)) \leq \exp\left(-\sum_{k=n}^m \mathbb{P}(A_k)\right) \xrightarrow{m \rightarrow \infty} 0.$$

So each $\bigcap_{k \geq n} A_k^c$ is null, and so is the countable union. $\square$

Common pitfalls.

- The lower bound ϵ does *not* give probability 1; that needs independence and a divergent sum (second lemma).
- Independence cannot be dropped: if $A_n = A$ for all n with $\mathbb{P}(A) = \frac{1}{2}$, then $\sum \mathbb{P}(A_n) = \infty$ but $\mathbb{P}(A_n \text{ i.o.}) = \frac{1}{2}$.

3 Measurable Functions and Piecewise Maps

3.1 Definition and Tools

Definition 3.1 (Measurable function). Let $(X, \mathcal{F})$ and $(Y, \mathcal{G})$ be measurable spaces. A map $f : X \rightarrow Y$ is $(\mathcal{F}, \mathcal{G})$ -**measurable** if $f^{-1}(B) \in \mathcal{F}$ for every $B \in \mathcal{G}$.

Lemma 3.2 (Generating class). If $\mathcal{G} = \sigma(\mathcal{C})$, then f is measurable iff $f^{-1}(C) \in \mathcal{F}$ for all $C \in \mathcal{C}$.

Proof. $\{B : f^{-1}(B) \in \mathcal{F}\}$ is a σ -algebra, since preimages commute with complements and countable unions. If it contains $\mathcal{C}$, it contains $\sigma(\mathcal{C})$. $\square$

Consequently $f : X \rightarrow \mathbb{R}$ is measurable iff $\{f \leq a\} \in \mathcal{F}$ for all $a \in \mathbb{R}$, and every continuous $g : \mathbb{R} \rightarrow \mathbb{R}$ is Borel measurable.

Lemma 3.3 (Gluing on a measurable partition). Let $X = E_1 \sqcup \cdots \sqcup E_k$ with each $E_i \in \mathcal{F}$, and suppose each restriction $f|_{E_i}$ is measurable. Then f is measurable.

Proof. For $B \in \mathcal{G}$, $f^{-1}(B) = \bigcup_{i=1}^k (f|_{E_i})^{-1}(B)$. Each term lies in $\mathcal{F}$ (as $E_i \in \mathcal{F}$), and $\mathcal{F}$ is closed under finite unions. $\square$

3.2 Problem: A Piecewise Real-Valued Function

Consider $f : (\mathbb{R}, \mathcal{B}_{\mathbb{R}}) \rightarrow (\text{Range}(f), \mathcal{B}_{\text{Range}(f)})$,

$$f(x) = \begin{cases} -3x, & x < 0, \\ x, & 0 \leq x \leq 2, \\ 1 - x, & x > 2. \end{cases}$$

Range of each piece.

Piece	Domain	Range
1	$(-\infty, 0)$	$(0, \infty)$
2	$[0, 2]$	$[0, 2]$
3	$(2, \infty)$	$(-\infty, -1)$

Hence $\text{Range}(f) = (-\infty, -1) \cup [0, \infty)$. The first two ranges overlap on $(0, 2]$, so f is not injective.

Measurability. The intervals $(-\infty, 0)$, $[0, 2]$, $(2, \infty)$ are Borel and partition $\mathbb{R}$, and each piece is continuous, hence Borel. By Lemma 3.3, f is measurable. Explicitly, for Borel B ,

$$f^{-1}(B) = \left(-\frac{1}{3}(B \cap (0, \infty)) \right) \cup (B \cap [0, 2]) \cup \left(1 - (B \cap (-\infty, -1)) \right),$$

a union of three Borel sets (images of Borel sets under affine maps with non-zero slope).

Example 3.4. For $B = [1, 3]$: piece 1 gives $[-1, -\frac{1}{3}]$, piece 2 gives $[1, 2]$, and piece 3 gives $\emptyset$. So $f^{-1}([1, 3]) = [-1, -\frac{1}{3}] \cup [1, 2]$.

Common pitfalls.

- Continuity of each piece is only useful because the *domains* of the pieces are themselves Borel sets; check this.
- Invert each piece separately and intersect with its own domain; otherwise preimages from different pieces get mixed up.

4 Induced (Push-Forward) Measures

4.1 Definition

Definition 4.1 (Induced measure). Let $(X, \mathcal{M}, \mu)$ be a measure space and $f : X \rightarrow Y$ a measurable map into $(Y, \mathcal{N})$. The **induced (push-forward) measure** is

$$\mu_f(B) = \mu(f^{-1}(B)), \quad B \in \mathcal{N}.$$

Proposition 4.2. μ_f is a measure on $(Y, \mathcal{N})$; if μ is a probability measure, so is μ_f .

Proof. $\mu_f(\emptyset) = 0$. For disjoint $B_j \in \mathcal{N}$, the sets $f^{-1}(B_j)$ are disjoint and $f^{-1}(\bigcup_j B_j) = \bigcup_j f^{-1}(B_j)$, so countable additivity of μ transfers to μ_f . $\square$

Change of variables.

For measurable $g \geq 0$ (or integrable),

$$\int_Y g d\mu_f = \int_X g \circ f d\mu.$$

This holds for $g = \mathbf{1}_B$ by definition, extends to simple functions by linearity, and to general g by monotone convergence.

4.2 Problem: Push-Forward of Lebesgue Measure

Let $\mu = \lambda$ and

$$f(x) = \begin{cases} x, & x < 0, \\ 2x, & x \geq 0. \end{cases}$$

For $B \in \mathcal{B}_{\mathbb{R}}$, $f^{-1}(B) = (B \cap (-\infty, 0)) \cup \frac{1}{2}(B \cap [0, \infty))$. Since λ scales by $\frac{1}{2}$ under $x \mapsto x/2$,

$$\lambda_f(B) = \lambda(B \cap (-\infty, 0)) + \frac{1}{2} \lambda(B \cap [0, \infty)).$$

So λ_f has density 1 on $(-\infty, 0)$ and density $\frac{1}{2}$ on $[0, \infty)$. For instance, $\lambda_f((-1, 2)) = 1 + \frac{1}{2} \cdot 2 = 2$.

Common pitfalls.

- The push-forward is computed through the *preimage*, never the image: $\mu_f(B) \neq \mu(f(B))$ in general.
- Measurability of f is what ensures $f^{-1}(B) \in \mathcal{M}$ so that μ_f is defined.

5 Random Variables Defined by Cases

5.1 Problem: Gluing Two Random Variables on an Event

Let X, Y be random variables on $(\Omega, \mathcal{F}, \mathbb{P})$ and $A \in \mathcal{F}$. Define

$$Z(\omega) = X(\omega)\mathbf{1}_A(\omega) + Y(\omega)\mathbf{1}_{A^c}(\omega) = \begin{cases} X(\omega), & \omega \in A, \\ Y(\omega), & \omega \in A^c. \end{cases}$$

Key Result.

Z is a random variable.

Proof. For $B \in \mathcal{B}_{\mathbb{R}}$,

$$Z^{-1}(B) = (A \cap X^{-1}(B)) \cup (A^c \cap Y^{-1}(B)).$$

1. X, Y are random variables, so $X^{-1}(B), Y^{-1}(B) \in \mathcal{F}$.
2. $A \in \mathcal{F}$ implies $A^c \in \mathcal{F}$.
3. $\mathcal{F}$ is closed under finite intersections and unions, so $Z^{-1}(B) \in \mathcal{F}$. □

Common pitfalls.

- $A \in \mathcal{F}$ is essential. With $X \equiv 1, Y \equiv 0$ we get $Z = \mathbf{1}_A$, and $Z^{-1}(\{1\}) = A$, which is not an event if $A \notin \mathcal{F}$.
- This is Lemma 3.3 applied to the partition $\{A, A^c\}$ of Ω .

6 Composition of Random Variables and Equality of Law

6.1 Setup

Let $X : (\Omega, \mathcal{F}, \mathbb{P}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ be a random variable with law $\mathbb{P}_X(B) = \mathbb{P}(X^{-1}(B))$, and let $Y : (\mathbb{R}, \mathcal{B}_{\mathbb{R}}) \rightarrow (\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ be Borel measurable. Then $Y \circ X$ is a random variable, since

$$(Y \circ X)^{-1}(B) = X^{-1}(Y^{-1}(B)) \in \mathcal{F}.$$

6.2 Problem: Law of $Y \circ X$

Key Result.

Regarding Y as a random variable on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}}, \mathbb{P}_X)$, its law equals the law of $Y \circ X$ on $(\Omega, \mathcal{F}, \mathbb{P})$.

Proof. For $B \in \mathcal{B}_{\mathbb{R}}$,

$$\mathbb{P}_X(Y^{-1}(B)) = \mathbb{P}(X^{-1}(Y^{-1}(B))) = \mathbb{P}((Y \circ X)^{-1}(B)) = \mathbb{P}_{Y \circ X}(B).$$

Since B was arbitrary, the two laws coincide. □

Corollary 6.1. *For non-negative Borel h ,*

$$\mathbb{E}[h(Y(X))] = \int_{\mathbb{R}} h \circ Y \, d\mathbb{P}_X = \int_{\mathbb{R}} h \, d\mathbb{P}_{Y \circ X}.$$

Remark. Probabilities and expectations of functions of X can be computed entirely on $\mathbb{R}$ from the law $\mathbb{P}_X$, without reference to Ω . This is the change of variables formula of Section 4 applied to $f = X$.

Common pitfalls.

- The order of composition matters: $(Y \circ X)^{-1}(B) = X^{-1}(Y^{-1}(B))$, not $Y^{-1}(X^{-1}(B))$.
- Y must be Borel measurable, not merely defined on the range of X ; otherwise $Y^{-1}(B)$ may fail to lie in $\mathcal{B}_{\mathbb{R}}$.