2025 1.14

PHYSICS TEXTBOOK

David Griffiths

WILEY-VCH

Introduction to Elementary Particles

Second, Revised Edition



Contents

Introduction to Elementary Particles, Second Edition. David Griffiths

ISBN: 978-3-527-40601-2

High Energy Physics

Text: D. Griffiths: Introduction to Elementary Particles
John Wily & Sons , 2nd Edition(2008)

Reference:

F. Halzen and A.D. Martin: Quarks & Leptons John-Wiley & Sons (1984)

D.H. Perkins: Introduction to High Energy Physics
(4th Edition) Cambridge University Press (2000)

Fayyazuddin & Riazuddin: A Modern Introduction to Particle Physics
(2nd edition) World Scientific Publishing(2000)

Duncan Carlsmith: Particle Physics, Pearson Education (2013)

C H Oh

Physics Department

NUS National University of Singapore

General Reading:

(1) Brian Greene: The Elegant Universe (1999), QC794.6 Str. Gr.

The Fabric of the Cosmos (2003); The Hidden Reality (2011)

(1) M Veltman: Facts and Mysteries in Elementary Particle Physics (WSPC, 2003)

(2) Leo Lederman: The God Particle : If the Universe is the Answer, What is the question, Boston: Houghton Mifflin (1993), QC793.Bos.L

Websites:

Update of the Particle Listings available on the Web
PDG Berkeley website: http://pdg.lbl.gov/

The Berkeley website gives access to MIRROR sites in:
Brazil, CERN, Italy, Japan, Russia, and the United Kingdom.

Also see the Particle Adventure at: http://ParticleAdventure.org

http://www-ed.fnal.gov/lml/Leon_life.html (Leo Lederman)
http://www-ed.fnal.gov/trc/projects/index_all.html

Particle Physics Labs

- **Laboratories**BNL: The Department of Energy's Brookhaven National Laboratory in Upton, Long Island.
- CERN: Originally "Conseil Européenne pour Recherches Nucléaires," now the European Laboratory for Particle Physics, in Geneva, Switzerland.
- DESY: Deutches Elektronen SYnchrotron laboratory in Hamburg, Germany.
- FNAL: The Department of Energy's Fermi National Accelerator Laboratory in Batavia, Illinois.
- KEK: Koo Energy Ken. The High Energy Research Accelerator Organization in Tsukuba, Japan.
- SLAC: The Department of Energy's Stanford Linear Accelerator Center in Palo Alto, California.

Contents

Contents

Contents

Contents

Contents

1.1 Introduction

Elementary Particles = Basic constituents of matter.

A particle can be pointlike and wavelike.

To break matter into its smallest pieces, need high energy

$\therefore$ **Elementary particle physics = high energy physics**

Present energy achieved $\approx 1\ TeV \approx 1000\ GeV \approx 10^{12}\ eV$ (Fermilab)

LHC (2007) proton beams $7\ TeV + 7\ TeV = 14\ TeV$

Theoretical discussion on the unification of basic forces has reached the Planck energy scale

$$\left(\frac{\hbar c}{G_N}\right)^{1/2} = 10^{-5} gm = 10^{19} GeV = 10^{28} eV$$

Close to the energy scale at which the universe is created.

1 fm → Fermi $= 10^{-13}$ cm

$= 10^{-15}$ m size of nucleus

1 Å $= 10^{-8}$ cm size of atom

H atom

$|n, \ell, m\rangle$

$1, \frac{1}{2}, \frac{1}{2}$

History of Constituents of Matter



1869
Mendeleev

1.2 Particles

Leptons: Particles do not participate in strong interaction.

	Q	L_e	L_μ	L_τ
e	-1	1	0	0
ν_e	0	1	0	0
μ	-1	0	1	0
ν_μ	0	0	1	0
τ	-1	0	0	1
ν_τ	0	0	0	1

Electron pointlike up to

10^{-15} cm = 10^{-2} fm

Hadrons(strongly interacting particles)

Baryons: Half-integral spin particles (fermions) involve in all basic interactions, ***st*** (strong), ***wk*** (weak), ***em*** (electromagnetic), e g

$p, n, \Lambda, \Sigma^+, \Sigma^0, \Sigma^-, \Xi^0, \Xi^-, \Delta, \Omega^-$

Mesons: integer spin particles (bosons) involve in all basic interactions ***st, wk, em***

$\pi^+, \pi^\circ, \pi^-, k^\pm, k^\circ, \eta, \omega$

Baryons are made from three quarks q, q, q

Mesons are made from quark-antiquark $q, \bar{q}$

Three generations of quarks

	Q	U	D	C	S	T	B
u	2/3	1	0	0	0	0	0
d	-1/3	0	-1	0	0	0	0
c	2/3	0	0	1	0	0	0
s	-1/3	0	0	0	-1	0	0
t	2/3	0	0	0	0	1	0
b	-1/3	0	0	0	0	0	-1

each quark has a nonabelian charge, called colour (**source** of strong interaction); there are three different colours.

Classification symmetry group

The lepton number, like electric charge, is associated with the Abelian U(1) group.

$\{ e^{i\alpha_1}, e^{i\alpha_2}, \ldots \}$

α_i = phase

$e^{i\alpha_i}$ = phase factor

The lepton doublet and also quark doublet are associated with the non-Abelian SU(2), originally from the isospin symmetry of proton and neutron

Baryons and Mesons are bound states of quarks.

e.g.

proton = uud

antiproton = $\bar{u}\bar{u}\bar{d}$

neutron = udd

Pion $\pi^+ = u\bar{d}$

Pion $\pi^- = \bar{u}d$

Kaon $k^+ = u\bar{s}$

Kaon $k^- = \bar{u}s$

$J/\psi = c\bar{c}$

Gauge field particles (force field)

Photon γ	**electromagnetic interaction**
Graviton	**gravitation**
Gluons *g*	**strong interaction**
Intermediate Vector bosons $W^{\pm}$ Z	**weak interaction**

Mass: $m_{W^{\pm}} \approx 82GeV/c^2$, $m_Z \approx 92Gev/C^2$

1.3 Basic Interactions (forces)

Type of force:	Gravitational	Weak	Electro-magnetic	Strong
Range:	infinite	$\leq 10^{-16}$cm	infinite	$\leq 10^{-13}$cm
Strength relative to strong force at a distance 10^{-13} cm	10^{-38}	10^{-13}	10^{-2}	1
Decay time for a typical small mass hadron:		10^{-10}s	10^{-20}s	10^{-23}s
Mediator:	Graviton	W^+, W^-, Z^0	Photon γ	gluon
Mass of the mediator:	0	82 GeV/c^2 92 GeV/c^2	0	0

Theories: Strong interaction — **Quantum chromodynamics QCD**

***em* interaction** — **Quantum electrodynamics QED**

***Weak* interaction** — **Weinberg – Salam model (Flavour dynamics)**

Gravitation — **Quantum gravity (?)**
Einstein's general relativity

Standard Model in particle physics

(i) Electroweak unification 1967

So called Glashow-Salam-Weinberg Model unifying weak interaction with the electromagnetic interaction. Quantum flavor dynamics

The model is based on quantum field theory. Both the particle (matter lepton) and the interaction are represented by field operators and the interaction term is of the form of current(matter) × gauge field, or $J_\mu^a \times A_a^\mu$. The symmetry group is $U(1) \times SU(2)$

(ii) The strong interaction is described by quantum chromatic dynamics (QCD) ~1973. The symmetry group is $SU(3)$. Again Both the particle (matter quarks) and the interaction are represented by field operators and the interaction term is of the form of current(matter) × gauge field, or $J_\mu^a \times A_a^\mu$.

(iii) The standard model is based on the gauge group is $U(1) \times SU(2) \times SU(3)$. Strictly not a complete unification because it consists of 3 separate gauge group. Ideally unification should be based on one single gauge group.

THE EIGHTFOLD WAY

(1961)

classify hadrons (baryons, mesons) according to the multiplets (singlet, octet, decuplet) of SU(3) group, so called unitary symmetry.

Extension of isospin scheme SU(2)[17]

Alex

ALEXANDER JOHN GILL A0196530A

SHEN TIANWEI Daniel E1496743

XIE YUSHI A0294895A

MAJUMDAR SRINJOY A0246042N

Yang Meng Ting A0275638N

SUN BAISONG A0295059R

Jiang Peiyang A0292815J

LI XING YU A0275634X

MA XIANGYI A0298874B

Theo LEQUY A0313016X

Zhang Bohan A0295089L

Ramachandran Janani A0245979E

Ting Jun Rui [redacted] A0179506W

Zhang Yifeng A0298377E

2025. 1. 14

Attendance list

14 students