


IIT MATHEMATICS

COMPLEX NUMBER




COMPLEX NUMBER

1. DEFINITION :

Complex numbers are defined as expressions of the form $a + ib$ where $a, b \in \mathbb{R}$ & $i = \sqrt{-1}$. It is denoted by z i.e. $z = a + ib$. 'a' is called real part of z ($\text{Re } z$) and 'b' is called imaginary part of z ($\text{Im } z$).

Every Complex Number Can Be Regarded As



Note :

- (i) The set $\mathbb{R}$ of real numbers is a proper subset of the Complex Numbers. Hence the Complex Number system is $\mathbb{N} \subset \mathbb{W} \subset \mathbb{I} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$.
- (ii) Zero is both purely real as well as purely imaginary but not imaginary.
- (iii) $i = \sqrt{-1}$ is called the imaginary unit. Also $i^2 = -1$; $i^3 = -i$; $i^4 = 1$ etc.
In general $i^{4n} = 1$, $i^{4n+1} = i$, $i^{4n+2} = -1$, $i^{4n+3} = -i$, where $n \in \mathbb{I}$
- (iv) $\sqrt{a}\sqrt{b} = \sqrt{ab}$ only if atleast one of either a or b is non-negative

Illustration 1 : The value of $i^{57} + 1/i^{125}$ is :-

- (A) 0 (B) $-2i$ (C) $2i$ (D) 2


Solution :

$$\begin{aligned}
 i^{57} + 1/i^{125} &= i^{56} \cdot i + \frac{1}{i^{124} \cdot i} \\
 &= (i^4)^{14} i + \frac{1}{(i^4)^{31} i} \\
 &= i + \frac{1}{i} = i + \frac{i}{i^2} = i - i = 0
 \end{aligned}$$

Ans. (A)

2. ARGAND DIAGRAM :

Master Argand had done a systematic study on complex numbers and represented every complex number $z = x + iy$ as a set of ordered pair (x, y) on a plane called complex plane (Argand Diagram) containing two perpendicular axes. Horizontal axis is known as Real axis & vertical axis is known as Imaginary axis. All complex numbers lying on the real axis are called as purely real and those lying on imaginary axis as purely imaginary.



3. ALGEBRAIC OPERATIONS :

Fundamental operations with complex numbers :

- (a) Addition $(a + bi) + (c + di) = (a + c) + (b + d)i$
 (b) Subtraction $(a + bi) - (c + di) = (a - c) + (b - d)i$
 (c) Multiplication $(a + bi)(c + di) = (ac - bd) + (ad + bc)i$
 (d) Division $\frac{a + bi}{c + di} = \frac{a + bi}{c + di} \cdot \frac{c - di}{c - di} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i$

Note :

- (i) The algebraic operations on complex numbers are similar to those on real numbers treating i as a polynomial.
 (ii) Inequalities in complex numbers (non-real) are not defined. There is no validity if we say that complex number (non-real) is positive or negative.
 e.g. $z > 0$, $4 + 2i < 2 + 4i$ are meaningless.
 (iii) In real numbers, if $a^2 + b^2 = 0$, then $a = 0 = b$ but in complex numbers, $z_1^2 + z_2^2 = 0$ does not imply $z_1 = z_2 = 0$.

Illustration 2 : $\frac{3 + 2i \sin \theta}{1 - 2i \sin \theta}$ will be purely imaginary, if $\theta =$

- (A) $2n\pi \pm \frac{\pi}{3}, n \in I$ (B) $n\pi + \frac{\pi}{3}, n \in I$ (C) $n\pi \pm \frac{\pi}{3}, n \in I$ (D) none of these

Solution : $\frac{3 + 2i \sin \theta}{1 - 2i \sin \theta}$ will be purely imaginary, if the real part vanishes, i.e.,

$$\frac{(3 + 2i \sin \theta)}{(1 - 2i \sin \theta)} \times \frac{(1 + 2i \sin \theta)}{(1 + 2i \sin \theta)} = \frac{(3 - 4 \sin^2 \theta) + i(8 \sin \theta)}{(1 + 4 \sin^2 \theta)}$$

$$\frac{3 - 4 \sin^2 \theta}{1 + 4 \sin^2 \theta} = 0 \Rightarrow 3 - 4 \sin^2 \theta = 0 \text{ (only if } \theta \text{ be real)}$$

$$\Rightarrow \sin^2 \theta = \left(\frac{\sqrt{3}}{2} \right)^2 = \left(\sin \frac{\pi}{3} \right)^2$$

$$\Rightarrow \theta = n\pi \pm \frac{\pi}{3}, n \in I$$

Ans. (C)

Do yourself -1 :

- (i) Determine least positive value of n for which $\left(\frac{1+i}{1-i} \right)^n = 1$
 (ii) Find the value of the sum $\sum_{n=1}^5 (i^n + i^{n+2})$ where $i = \sqrt{-1}$.

4. EQUALITY IN COMPLEX NUMBER :

Two complex numbers $z_1 = a_1 + ib_1$ & $z_2 = a_2 + ib_2$ are equal if and only if their real & imaginary parts are respectively equal.

Illustration 3 : The values of x and y satisfying the equation $\frac{(1+i)x-2i}{3+i} + \frac{(2-3i)y+i}{3-i} = i$ are

- (A) $x = -1, y = 3$ (B) $x = 3, y = -1$ (C) $x = 0, y = 1$ (D) $x = 1, y = 0$

Solution : $\frac{(1+i)x-2i}{3+i} + \frac{(2-3i)y+i}{3-i} = i \Rightarrow (4+2i)x + (9-7i)y - 3i - 3 = 10i$

Equating real and imaginary parts, we get $2x - 7y = 13$ and $4x + 9y = 3$.

Hence $x = 3$ and $y = -1$.

Ans.(B)

Illustration 4 : Find the square root of $7 + 24i$.

Solution : Let $\sqrt{7+24i} = a + ib$

Squaring $a^2 - b^2 + 2iab = 7 + 24i$

Compare real & imaginary parts $a^2 - b^2 = 7$ & $2ab = 24$

By solving these two equations

We get $a = \pm 4, b = \pm 3$

$\sqrt{7+24i} = \pm (4 + 3i)$

Illustration 5 : If $x = -5 + 2\sqrt{-4}$, find the value of $x^4 + 9x^3 + 35x^2 - x + 4$.

Solution : We have, $x = -5 + 2\sqrt{-4}$

$$\Rightarrow x + 5 = 4i, \quad \Rightarrow (x + 5)^2 = 16i^2$$

$$\Rightarrow x^2 + 10x + 25 = -16 \Rightarrow x^2 + 10x + 41 = 0$$

Now,

$$x^4 + 9x^3 + 35x^2 - x + 4$$

$$\Rightarrow x^2(x^2 + 10x + 41) - x(x^2 + 10x + 41) + 4(x^2 + 10x + 41) - 160$$

$$\Rightarrow x^2(0) - x(0) + 4(0) - 160 = -160$$

Ans.

Do yourself -2 :

(i) Find the value of $x^3 + 7x^2 - x + 16$. Where $x = 1 + 2i$.

(ii) If $a + ib = \frac{c+i}{c-i}$, where c is a real number, then prove that : $a^2 + b^2 = 1$ and $\frac{b}{a} = \frac{2c}{c^2 - 1}$

(iii) Find square root of $-15 - 8i$

5. THREE IMPORTANT TERMS : CONJUGATE / MODULUS / ARGUMENT :

(a) CONJUGATE COMPLEX :

If $z = a + ib$ then its conjugate complex is obtained by changing the sign of its imaginary part & is denoted by $\bar{z}$. i.e. $\bar{z} = a - ib$.

Note that :

(i) $z + \bar{z} = 2 \operatorname{Re}(z)$


(ii) $z - \bar{z} = 2i \operatorname{Im}(z)$

(iii) $z \bar{z} = a^2 + b^2$, which is purely real

(iv) If z is purely real, then $z - \bar{z} = 0$

(v) If z is purely imaginary, then $z + \bar{z} = 0$

(vi) If z lies in the 1st quadrant, then $\bar{z}$ lies in the 4th quadrant and $-\bar{z}$ lies in the 2nd quadrant.



(b) Modulus :

If P denotes complex number $z = x + iy$, then the length OP is called modulus of complex number z . It is denoted by $|z|$.

$$OP = |z| = \sqrt{x^2 + y^2}$$


Geometrically $|z|$ represents the distance of point P from origin. ($|z| \geq 0$)

Note : Unlike real numbers, $|z| = \begin{cases} z & \text{if } z > 0 \\ -z & \text{if } z < 0 \end{cases}$ is not correct.

(c) Argument or Amplitude :

If P denotes complex number $z = x + iy$ and if OP makes an angle θ with real axis, then θ is called one of the arguments of z .

$$\theta = \tan^{-1} \frac{y}{x} \text{ (angle made by } OP \text{ with positive real axis)}$$




Note:

- (i) Argument of a complex number is a many valued function. If θ is the argument of a complex number, then $2n\pi + \theta$; $n \in \mathbb{I}$ will also be the argument of that complex number. Any two arguments of a complex number differ by $2n\pi$.
- (ii) The unique value of θ such that $-\pi < \theta \leq \pi$ is called **Amplitude (principal value of the argument)**.

(iii) Principal argument of a complex number $z = x + iy$ can be found out using method given below :

(a) Find $\theta = \tan^{-1} \left| \frac{y}{x} \right|$ such that $\theta \in \left(0, \frac{\pi}{2} \right)$

(b) Use given figure to find out the principal argument according as the point lies in respective quadrant.



(iv) Unless otherwise stated, $\arg z$ implies principal value of the argument.

(v) The unique value of $\theta = \tan^{-1} \frac{y}{x}$ such that $0 < \theta \leq 2\pi$ is called **least positive argument**.

(vi) If $z = 0$, $\arg(z)$ is not defined

(vii) If z is real & negative, $\arg(z) = \pi$.

(viii) If z is real & positive, $\arg(z) = 0$

(ix) If $\theta = \frac{\pi}{2}$, z lies on the positive side of imaginary axis.

(x) If $\theta = -\frac{\pi}{2}$, z lies on the negative side of imaginary axis.

By specifying the modulus & argument a complex number is defined completely. Argument impart direction & modulus impart distance from origin.

For the complex number $0 + 0i$ the argument is not defined and this is the only complex number which is given by its modulus only.

Illustration 6 : Find the modulus, argument, principal value of argument, least positive argument of complex numbers (a) $1 + i\sqrt{3}$ (b) $-1 + i\sqrt{3}$ (c) $1 - i\sqrt{3}$ (d) $-1 - i\sqrt{3}$

Solution : (a) For $z = 1 + i\sqrt{3}$


$$|z| = \sqrt{1^2 + (\sqrt{3})^2} = 2$$

$$\arg(z) = 2n\pi + \frac{\pi}{3}, n \in \mathbb{I}$$

Least positive argument is $\frac{\pi}{3}$

If the point is lying in first or second quadrant then $\arg(z)$ is taken in anticlockwise direction.

$$\text{In this case } \arg(z) = \frac{\pi}{3}$$




(b) For $z = -1 + i\sqrt{3}$

$$|z| = 2$$

$$\arg(z) = 2n\pi + \frac{2\pi}{3}, n \in \mathbb{I}$$

$$\text{Least positive argument} = \frac{2\pi}{3}$$

$$\text{amp}(z) = \frac{2\pi}{3}$$



(c) For $z = 1 - i\sqrt{3}$


$$|z| = 2$$

$$\arg(z) = 2n\pi - \frac{\pi}{3}, n \in \mathbb{I}$$

$$\text{Least positive argument} = \frac{5\pi}{3}$$

If the point lies in third or fourth quadrant then consider $\text{amp}(z)$ in clockwise direction.

$$\text{In this case } \text{amp}(z) = -\frac{\pi}{3}$$



(d) For $z = -1 - i\sqrt{3}$

$$|z| = 2$$

$$\arg(z) = 2n\pi - \frac{2\pi}{3}, n \in \mathbb{I}$$

$$\text{Least positive argument} = \frac{4\pi}{3}$$

$$\text{amp}(z) = -\frac{2\pi}{3}$$




Illustration 7 : Find modulus and argument for $z = 1 - \sin \alpha + i \cos \alpha$, $\alpha \in (0, 2\pi)$

Solution : $|z| = \sqrt{(1 - \sin \alpha)^2 + (\cos \alpha)^2} = \sqrt{2 - 2\sin \alpha} = \sqrt{2} \left| \cos \frac{\alpha}{2} - \sin \frac{\alpha}{2} \right|$

Case (i) For $\alpha \in \left(0, \frac{\pi}{2}\right)$, z will lie in I quadrant.

$$\text{amp}(z) = \tan^{-1} \frac{\cos \alpha}{1 - \sin \alpha} \Rightarrow \text{amp}(z) = \tan^{-1} \frac{\cos^2 \frac{\alpha}{2} - \sin^2 \frac{\alpha}{2}}{\left(\cos \frac{\alpha}{2} - \sin \frac{\alpha}{2}\right)^2} = \tan^{-1} \frac{\cos \frac{\alpha}{2} + \sin \frac{\alpha}{2}}{\cos \frac{\alpha}{2} - \sin \frac{\alpha}{2}}$$

$$\Rightarrow \arg z = \tan^{-1} \tan \left(\frac{\pi}{4} + \frac{\alpha}{2} \right)$$

$$\text{Since } \frac{\pi}{4} + \frac{\alpha}{2} \in \left(\frac{\pi}{4}, \frac{\pi}{2} \right)$$

$$\therefore \arg(z) = \left(\frac{\pi}{4} + \frac{\alpha}{2} \right), |z| = \sqrt{2} \left(\cos \frac{\alpha}{2} - \sin \frac{\alpha}{2} \right)$$

Case (ii) at $\alpha = \frac{\pi}{2}$: $z = 0 + 0i$

$$|z| = 0$$

amp (z) is not defined.

Case (iii) For $\alpha \in \left(\frac{\pi}{2}, \frac{3\pi}{2} \right)$, z will lie in IV quadrant

$$\text{so amp (z)} = -\tan^{-1} \tan \left(\frac{\alpha}{2} + \frac{\pi}{4} \right)$$

$$\text{Since } \frac{\alpha}{2} + \frac{\pi}{4} \in \left(\frac{\pi}{2}, \pi \right)$$

$$\therefore \arg(z) = -\left(\frac{\alpha}{2} + \frac{\pi}{4} - \pi \right) = \frac{3\pi}{4} - \frac{\alpha}{2}, |z| = \sqrt{2} \left(\sin \frac{\alpha}{2} - \cos \frac{\alpha}{2} \right)$$

Case (iv) at $\alpha = \frac{3\pi}{2}$: $z = 2 + 0i$

$$|z| = 2$$

$$\arg(z) = 0$$

Case (v) For $\alpha \in \left(\frac{3\pi}{2}, 2\pi \right)$

z will lie in I quadrant

$$\arg(z) = \tan^{-1} \tan \left(\frac{\alpha}{2} + \frac{\pi}{4} \right)$$

$$\text{Since } \frac{\alpha}{2} + \frac{\pi}{4} \in \left(\pi, \frac{5\pi}{4} \right)$$

$$\therefore \arg z = \frac{\alpha}{2} + \frac{\pi}{4} - \pi = \frac{\alpha}{2} - \frac{3\pi}{4}, |z| = \sqrt{2} \left(\sin \frac{\alpha}{2} - \cos \frac{\alpha}{2} \right)$$

Do yourself -3 :

Find the modulus and amplitude of following complex numbers :

(i) $-2 + 2\sqrt{3}i$

(ii) $-\sqrt{3} - i$

(iii) $-2i$

(iv) $\frac{1+2i}{1-3i}$

(v) $\frac{2+6\sqrt{3}i}{5+\sqrt{3}i}$

6. REPRESENTATION OF A COMPLEX NUMBER IN VARIOUS FORMS :

(a) Cartesian Form (Geometrical Representation) :

Every complex number $z = x + iy$ can be represented by a point on the cartesian plane known as complex plane by the ordered pair (x, y) . There exists a one-one correspondence between the points of the plane and the members of the set of complex numbers.

For $z = x + iy$; $|z| = \sqrt{x^2 + y^2}$; $\bar{z} = x - iy$ and $\theta = \tan^{-1} \frac{y}{x}$

Note :

(i) Distance between the two complex numbers z_1 & z_2 is given by $|z_1 - z_2|$.

(ii) $|z - z_0| = r$, represents a circle, whose centre is z_0 and radius is r .




Illustration 8 : Find the locus of :

(a) $|z - 1|^2 + |z + 1|^2 = 4$

(b) $\text{Re}(z^2) = 0$

Solution :

(a) Let $z = x + iy$

$$\Rightarrow (x + iy - 1)^2 + (x + iy + 1)^2 = 4$$

$$\Rightarrow (x - 1)^2 + y^2 + (x + 1)^2 + y^2 = 4$$

$$\Rightarrow x^2 - 2x + 1 + y^2 + x^2 + 2x + 1 + y^2 = 4 \Rightarrow x^2 + y^2 = 1$$

Above represents a circle on complex plane with center at origin and radius unity.

(b) Let $z = x + iy$

$$\Rightarrow z^2 = x^2 - y^2 + 2xyi$$

$$\therefore \text{Re}(z^2) = 0$$

$$\Rightarrow x^2 - y^2 = 0 \Rightarrow y = \pm x$$

Thus $\text{Re}(z^2) = 0$ represents a pair of straight lines passing through origin.

Illustration 9 : If z is a complex number such that $z^2 = (\bar{z})^2$, then

(A) z is purely real

(B) z is purely imaginary

(C) either z is purely real or purely imaginary

(D) none of these

Solution :

Let $z = x + iy$, then its conjugate $\bar{z} = x - iy$

$$\text{Given that } z^2 = (\bar{z})^2 \Rightarrow x^2 - y^2 + 2ixy = x^2 - y^2 - 2ixy \Rightarrow 4ixy = 0$$

If $x \neq 0$ then $y = 0$ and if $y \neq 0$ then $x = 0$.

Ans. (C)

Illustration 10 : Among the complex number z which satisfies $|z - 25i| \leq 15$, find the complex numbers z having

(a) least positive argument

(b) maximum positive argument

(c) least modulus

(d) maximum modulus

Solution :

The complex numbers z satisfying the condition

$$|z - 25i| \leq 15$$

are represented by the points inside and on the circle of radius 15 and centre at the point $C(0, 25)$.

The complex number having least positive argument and maximum positive arguments in this region are the points of contact of tangents drawn from origin to the circle


Here θ = least positive argument

and ϕ = maximum positive argument

$$\therefore \text{In } \triangle OCP, OP = \sqrt{(OC)^2 - (CP)^2} = \sqrt{(25)^2 - (15)^2} = 20$$

$$\text{and } \sin \theta = \frac{OP}{OC} = \frac{20}{25} = \frac{4}{5}$$

$$\therefore \tan \theta = \frac{4}{3} \Rightarrow \theta = \tan^{-1}\left(\frac{4}{3}\right)$$



Thus, complex number at P has modulus 20 and argument $\theta = \tan^{-1}\left(\frac{4}{3}\right)$

$$\therefore z_p = 20 (\cos \theta + i \sin \theta) = 20 \left(\frac{3}{5} + i \frac{4}{5} \right)$$

$$\therefore z_p = 12 + 16i$$

$$\text{Similarly } z_Q = -12 + 16i$$

From the figure, E is the point with least modulus and D is the point with maximum modulus.

$$\text{Hence, } z_E = \overrightarrow{OE} = \overrightarrow{OC} - \overrightarrow{EC} = 25i - 15i = 10i$$

$$\text{and } z_D = \overrightarrow{OD} = \overrightarrow{OC} + \overrightarrow{CD} = 25i + 15i = 40i$$

Do yourself -4 :

- (i) Find the distance between two complex numbers $z_1 = 2 + 3i$ & $z_2 = 7 - 9i$ on the complex plane.
- (ii) Find the locus of $|z - 2 - 3i| = 1$.
- (iii) If z is a complex number, then $z^2 + \bar{z}^2 = 2$ represents -
(A) a circle (B) a straight line (C) a hyperbola (D) an ellipse

(c) Trigonometric / Polar Representation :

$$z = r (\cos \theta + i \sin \theta) \text{ where } |z| = r ; \arg z = \theta ; \bar{z} = r (\cos \theta - i \sin \theta)$$

Note : $\cos \theta + i \sin \theta$ is also written as $\text{CiS } \theta$.

Euler's formula :

The formula $e^{ix} = \cos x + i \sin x$ is called Euler's formula.

It was introduced by Euler in 1748, and is used as a method of expressing complex numbers.

Also $\cos x = \frac{e^{ix} + e^{-ix}}{2}$ & $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$ are known as Euler's identities.

(d) Exponential Representation :

Let z be a complex number such that $|z| = r$ & $\arg z = \theta$, then $z = r.e^{i\theta}$

Illustration 11 : Express the following complex numbers in polar and exponential form :

(i) $\frac{1+3i}{1-2i}$ (ii) $\frac{i-1}{\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}}$

Solution :

(i) Let $z = \frac{1+3i}{1-2i} = \frac{1+3i}{1-2i} \times \frac{1+2i}{1+2i} = -1+i$

$|z| = \sqrt{(-1)^2 + 1^2} = \sqrt{2}$

$\tan \alpha = \frac{1}{-1} = -1 = \tan \frac{\pi}{4} \Rightarrow \alpha = \frac{\pi}{4}$

$\therefore \operatorname{Re}(z) < 0$ and $\operatorname{Im}(z) > 0 \Rightarrow z$ lies in second quadrant.

$\therefore \theta = \arg(z) = \pi - \alpha = \pi - \frac{\pi}{4} = \frac{3\pi}{4}$

Hence Polar form is $z = \sqrt{2} \left(\cos \frac{3\pi}{4} + i \sin \frac{3\pi}{4} \right)$

and exponential form is $z = \sqrt{2} e^{i \frac{3\pi}{4}}$

(ii) Let $z = \frac{i-1}{\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}} = \frac{i-1}{\frac{1}{2} + i \frac{\sqrt{3}}{2}} = \frac{2(i-1)}{(1+i\sqrt{3})}$

$\Rightarrow z = \frac{2(i-1)}{(1+i\sqrt{3})} \times \frac{(1-i\sqrt{3})}{(1-i\sqrt{3})} \Rightarrow z = \left(\frac{\sqrt{3}-1}{2} \right) + i \left(\frac{\sqrt{3}+1}{2} \right)$

$\therefore \operatorname{Re}(z) > 0$ and $\operatorname{Im}(z) > 0 \Rightarrow z$ lies in first quadrant.

$\therefore |z| = \sqrt{\left(\frac{\sqrt{3}-1}{2} \right)^2 + \left(\frac{\sqrt{3}+1}{2} \right)^2} = \sqrt{\frac{2(3+1)}{4}} = \sqrt{2}$

$\tan \theta = \frac{\sqrt{3}+1}{\sqrt{3}-1} = \tan \frac{5\pi}{12} \Rightarrow \alpha = \frac{5\pi}{12}$

Hence Polar form is $z = \sqrt{2} \left(\cos \frac{5\pi}{12} + i \sin \frac{5\pi}{12} \right)$

and exponential form is $z = \sqrt{2} e^{i \frac{5\pi}{12}}$

Illustration 12 : If $x_n = \cos\left(\frac{\pi}{2^n}\right) + i \sin\left(\frac{\pi}{2^n}\right)$, then $x_1 x_2 x_3 \dots \infty$ is equal to –

- (A) -1 (B) 1 (C) 0 (D) ∞

Solution :

$$x_n = \cos\left(\frac{\pi}{2^n}\right) + i \sin\left(\frac{\pi}{2^n}\right) = 1 \times e^{i \frac{\pi}{2^n}}$$

$$x_1 x_2 x_3 \dots \infty$$

$$= e^{i \frac{\pi}{2^1}} \cdot e^{i \frac{\pi}{2^2}} \dots e^{i \frac{\pi}{2^n}} = e^{i \left(\frac{\pi}{2} + \frac{\pi}{2^2} + \dots + \frac{\pi}{2^n} \right)}$$

$$= \cos\left(\frac{\pi}{2} + \frac{\pi}{2^2} + \frac{\pi}{2^3} + \dots\right) + i \sin\left(\frac{\pi}{2} + \frac{\pi}{2^2} + \frac{\pi}{2^3} + \dots\right) = -1$$

$$\left(\text{as } \frac{\pi}{2} + \frac{\pi}{2^2} + \frac{\pi}{2^3} + \dots = \frac{\pi/2}{1-1/2} = \pi \right)$$

Ans.

Do yourself -5 :

Express the following complex number in polar form and exponential form :

- (i) $-2 + 2i$ (ii) $-1 - \sqrt{3}i$ (iii) $\frac{(1+7i)}{(2-i)^2}$ (iv) $(1 - \cos\theta + i \sin\theta)$, $\theta \in (0, \pi)$

7. IMPORTANT PROPERTIES OF CONJUGATE :

- (a) $z + \bar{z} = 2 \operatorname{Re}(z)$ (b) $z - \bar{z} = 2i \operatorname{Im}(z)$ (c) $\overline{\bar{z}} = z$
- (d) $\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$ (e) $\overline{z_1 - z_2} = \bar{z}_1 - \bar{z}_2$
- (f) $\overline{z_1 z_2} = \bar{z}_1 \cdot \bar{z}_2$. In general $\overline{z_1 z_2 \dots z_n} = \bar{z}_1 \cdot \bar{z}_2 \dots \bar{z}_n$
- (g) $\overline{\left(\frac{z_1}{z_2}\right)} = \frac{\bar{z}_1}{\bar{z}_2}$; $z_2 \neq 0$ (h) If $f(\alpha + i\beta) = x + iy \Rightarrow f(\alpha - i\beta) = x - iy$

8. IMPORTANT PROPERTIES OF MODULUS:

- (a) $|z| \geq 0$ (b) $|z| \geq \operatorname{Re}(z)$ (c) $|z| \geq \operatorname{Im}(z)$
- (d) $|z| = |\bar{z}| = |-z| = |-\bar{z}|$ (e) $z\bar{z} = |z|^2$
- (f) $|z_1 z_2| = |z_1| \cdot |z_2|$. In general $|z_1 z_2 \dots z_n| = |z_1| \cdot |z_2| \dots |z_n|$
- (g) $\left|\frac{z_1}{z_2}\right| = \frac{|z_1|}{|z_2|}$, $z_2 \neq 0$
- (h) $|z^n| = |z|^n$, $n \in \mathbb{I}$
- (i) $|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2 + 2 \operatorname{Re}(z_1 \bar{z}_2)$
- (j) $|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2 + 2|z_1||z_2| \cos(\alpha - \beta)$, where α, β are $\arg(z_1), \arg(z_2)$ respectively.
- (k) $|z_1 + z_2|^2 + |z_1 - z_2|^2 = 2[|z_1|^2 + |z_2|^2]$
- (l) $||z_1| - |z_2|| \leq |z_1 + z_2| \leq |z_1| + |z_2|$ **[Triangle Inequality]**
- (m) $||z_1| - |z_2|| \leq |z_1 - z_2| \leq |z_1| + |z_2|$ **[Triangle Inequality]**

9. IMPORTANT PROPERTIES OF AMPLITUDE :

- (a) $\text{amp}(z_1, z_2) = \text{amp } z_1 + \text{amp } z_2 + 2k\pi ; k \in I$
 (b) $\text{amp}\left(\frac{z_1}{z_2}\right) = \text{amp } z_1 - \text{amp } z_2 + 2k\pi ; k \in I$
 (c) $\text{amp}(z^n) = n \text{amp}(z) + 2k\pi ; n, k \in I$
 where proper value of k must be chosen so that RHS lies in $(-\pi, \pi]$.

Illustration 13 : Find $\text{amp } z$ and $|z|$ if $z = \left[\frac{(3+4i)(1+i)(1+\sqrt{3}i)}{(1-i)(4-3i)(2i)} \right]^2$.

Solution : $\text{amp } z = 2[\text{amp}(3+4i) + \text{amp}(1+i) + \text{amp}(1+\sqrt{3}i) - \text{amp}(1-i) - \text{amp}(4-3i) - \text{amp}(2i)] + 2k\pi$
 where $k \in I$ and k chosen so that $\text{amp } z$ lies in $(-\pi, \pi]$.

$$\Rightarrow \text{amp } z = 2 \left[\tan^{-1} \frac{4}{3} + \frac{\pi}{4} + \frac{\pi}{3} - \left(-\frac{\pi}{4} \right) - \tan^{-1} \left(-\frac{3}{4} \right) - \frac{\pi}{2} \right] + 2k\pi$$

$$\Rightarrow \text{amp } z = 2 \left[\tan^{-1} \frac{4}{3} + \cot^{-1} \frac{4}{3} + \frac{\pi}{3} \right] + 2k\pi \Rightarrow \text{amp } z = 2 \left[\frac{\pi}{2} + \frac{\pi}{3} \right] + 2k\pi$$

$$\Rightarrow \text{amp } z = -\frac{\pi}{3} \quad [\text{at } k = -1]$$

Ans.

Also,

$$|z| = \left| \frac{(3+4i)(1+i)(1+\sqrt{3}i)}{(1-i)(4-3i)(2i)} \right|^2 \Rightarrow |z| = \left(\frac{|3+4i| |1+i| |1+\sqrt{3}i|}{|1-i| |4-3i| |2i|} \right)^2$$

$$\Rightarrow |z| = \left(\frac{5 \times \sqrt{2} \times 2}{\sqrt{2} \times 5 \times 2} \right)^2 = 1$$

Ans.

Aliter

$$z = \left[\frac{(3+4i)(1+i)(1+\sqrt{3}i)}{(1-i)(4-3i)(2i)} \right]^2 \Rightarrow z = \left[-\frac{\sqrt{3}+i}{2} \right]^2 \Rightarrow z = \frac{2-2\sqrt{3}i}{4} = \frac{1}{2} - \frac{\sqrt{3}i}{2}$$

$$\text{Hence } |z| = 1, \text{amp}(z) = -\frac{\pi}{3}.$$

Illustration 14 : If $\left| \frac{z-i}{z+i} \right| = 1$, then locus of z is –

- (A) x-axis (B) y-axis (C) x = 1 (D) y = 1

Solution :

$$\text{We have, } \left| \frac{z-i}{z+i} \right| = 1 \Rightarrow \left| \frac{x+i(y-1)}{x+i(y+1)} \right| = 1$$

$$\Rightarrow \frac{|x+i(y-1)|^2}{|x+i(y+1)|^2} = 1 \Rightarrow x^2 + (y-1)^2 = x^2 + (y+1)^2 \Rightarrow 4y = 0; y = 0$$

which is x-axis

Ans. (A)

Illustration 15 : If $|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2$ then $\left(\frac{z_1}{z_2}\right)$ is –

- (A) zero or purely imaginary (B) purely imaginary
(C) purely real (D) none of these

Solution : Here let $z_1 = r_1(\cos \theta_1 + i \sin \theta_1), |z_1| = r_1$

$$z_2 = r_2(\cos \theta_2 + i \sin \theta_2), |z_2| = r_2$$

$$\begin{aligned} \therefore |(z_1 + z_2)|^2 &= |(r_1 \cos \theta_1 + r_2 \cos \theta_2) + i(r_1 \sin \theta_1 + r_2 \sin \theta_2)|^2 \\ &= r_1^2 + r_2^2 + 2r_1 r_2 \cos(\theta_1 - \theta_2) = |z_1|^2 + |z_2|^2 \text{ if } \cos(\theta_1 - \theta_2) = 0 \end{aligned}$$

$$\therefore \theta_1 - \theta_2 = \pm \frac{\pi}{2}$$

$$\Rightarrow \text{amp}(z_1) - \text{amp}(z_2) = \pm \frac{\pi}{2} \Rightarrow \text{amp}\left(\frac{z_1}{z_2}\right) = \pm \frac{\pi}{2} \Rightarrow \frac{z_1}{z_2} \text{ is purely imaginary Ans. (B)}$$

Illustration 16 : z_1 and z_2 are two complex numbers such that $\frac{z_1 - 2z_2}{2 - z_1 z_2}$ is unimodular (whose modulus is one), while z_2 is not unimodular. Find $|z_1|$.

Solution : Here $\left|\frac{z_1 - 2z_2}{2 - z_1 z_2}\right| = 1 \Rightarrow \frac{|z_1 - 2z_2|}{|2 - z_1 z_2|} = 1$

$$\Rightarrow |z_1 - 2z_2| = |2 - z_1 \bar{z}_2| \Rightarrow |z_1 - 2z_2|^2 = |2 - z_1 \bar{z}_2|^2$$

$$\Rightarrow (z_1 - 2z_2)(\overline{z_1 - 2z_2}) = (2 - z_1 \bar{z}_2)(\overline{2 - z_1 \bar{z}_2})$$

$$\Rightarrow (z_1 - 2z_2)(\bar{z}_1 - 2\bar{z}_2) = (2 - z_1 \bar{z}_2)(2 - \bar{z}_1 z_2)$$

$$\Rightarrow z_1 \bar{z}_1 - 2z_1 \bar{z}_2 - 2z_2 \bar{z}_1 + 4z_2 \bar{z}_2 = 4 - 2\bar{z}_1 z_2 - 2z_1 \bar{z}_2 + z_1 \bar{z}_1 z_2 \bar{z}_2$$

$$\Rightarrow |z_1|^2 + 4|z_2|^2 = 4 + |z_1|^2 |z_2|^2 \Rightarrow |z_1|^2 - |z_1|^2 |z_2|^2 + 4|z_2|^2 - 4 = 0$$

$$\Rightarrow (|z_1|^2 - 4)(1 - |z_2|^2) = 0$$

But $|z_2| \neq 1$ (given)

$$\therefore |z_1|^2 = 4 \quad \text{Hence, } |z_1| = 2.$$

Illustration 17 : The locus of the complex number z in argand plane satisfying the inequality

$$\log_{1/2} \left(\frac{|z-1|+4}{3|z-1|-2} \right) > 1 \left(\text{where } |z-1| \neq \frac{2}{3} \right) \text{ is -}$$

- (A) a circle (B) interior of a circle
(C) exterior of a circle (D) none of these

Solution : We have, $\log_{1/2} \left(\frac{|z-1|+4}{3|z-1|-2} \right) > 1 = \log_{1/2} \left(\frac{1}{2} \right)$
 $\Rightarrow \frac{|z-1|+4}{3|z-1|-2} < \frac{1}{2}$ [$\because \log_a x$ is a decreasing function if $a < 1$]
 $\Rightarrow 2|z-1| + 8 < 3|z-1| - 2$ as $|z-1| > 2/3$
 $\Rightarrow |z-1| > 10$
 which is exterior of a circle. **Ans. (C)**


Illustration 18 : If $\left| z - \frac{4}{z} \right| = 2$, then the greatest value of $|z|$ is-

- (A) $1 + \sqrt{2}$ (B) $2 + \sqrt{2}$ (C) $\sqrt{3} + 1$ (D) $\sqrt{5} + 1$

Solution : We have $|z| = \left| z - \frac{4}{z} + \frac{4}{z} \right| \leq \left| z - \frac{4}{z} \right| + \frac{4}{|z|} = 2 + \frac{4}{|z|}$
 $\Rightarrow |z|^2 \leq 2|z| + 4 \Rightarrow (|z| - 1)^2 \leq 5$
 $\Rightarrow |z| - 1 \leq \sqrt{5} \Rightarrow |z| \leq \sqrt{5} + 1$
 Therefore, the greatest value of $|z|$ is $\sqrt{5} + 1$. **Ans. (D)**

Illustration 19 : Shaded region is given by -

- (A) $|z + 2| \geq 6, 0 \leq \arg(z) \leq \frac{\pi}{6}$
 (B) $|z + 2| \leq 6, 0 \leq \arg(z + 2) \leq \frac{\pi}{3}$
 (C) $|z + 2| \geq 6, 0 \leq \arg(z + 2) \leq \frac{\pi}{3}$
 (D) none of these



Solution : Note that $AB = 6$ and $1 + 3\sqrt{3}i = -2 + 3 + 3\sqrt{3}i$
 $= -2 + 6 \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i \right) = -2 + 6 \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right)$
 $\therefore \angle BAC = \frac{\pi}{3}$

Thus, shaded region is given by $|z + 2| \geq 6$ and $0 \leq \arg(z + 2) \leq \frac{\pi}{3}$ **Ans. (C)**

Do yourself -6 :

- (i) The inequality $|z - 4| < |z - 2|$ represents region given by -
 (A) $\operatorname{Re}(z) > 0$ (B) $\operatorname{Re}(z) < 0$ (C) $\operatorname{Re}(z) > 3$ (D) none
 (ii) If $z = re^{i\theta}$, then the value of $|e^{iz}|$ is equal to -
 (A) $e^{-r \cos \theta}$ (B) $e^{r \cos \theta}$ (C) $e^{r \sin \theta}$ (D) $e^{-r \sin \theta}$

10. SECTION FORMULA AND COORDINATES OF ORTHOCENTRE, CENTROID, CIRCUMCENTRE, INCENTRE OF A TRIANGLE :

If z_1 & z_2 are two complex numbers then the complex number $z = \frac{nz_1 + mz_2}{m+n}$ divides the join of z_1 & z_2 in the ratio $m : n$.

Note :

- (i) If a, b, c are three real numbers such that $az_1 + bz_2 + cz_3 = 0$; where $a + b + c = 0$ and a, b, c are not all simultaneously zero, then the complex numbers z_1, z_2 & z_3 are collinear.
- (ii) If the vertices A, B, C of a triangle represent the complex numbers z_1, z_2, z_3 respectively, then :



- Centroid of the $\Delta ABC = \frac{z_1 + z_2 + z_3}{3}$
- Orthocentre of the $\Delta ABC = \frac{(a \sec A)z_1 + (b \sec B)z_2 + (c \sec C)z_3}{a \sec A + b \sec B + c \sec C}$ or $\frac{z_1 \tan A + z_2 \tan B + z_3 \tan C}{\tan A + \tan B + \tan C}$
- Incentre of the $\Delta ABC = \frac{(az_1 + bz_2 + cz_3)}{(a + b + c)}$
- Circumcentre of the $\Delta ABC = \frac{(z_1 \sin 2A + z_2 \sin 2B + z_3 \sin 2C)}{(\sin 2A + \sin 2B + \sin 2C)}$

11. VECTORIAL REPRESENTATION OF A COMPLEX NUMBER :

- (a) In complex number every point can be represented in terms of position vector. If the point P represents the complex number z then, $\overrightarrow{OP} = z$ &

$$|\overrightarrow{OP}| = |z|$$


- (b) If $P(z_1)$ & $Q(z_2)$ be two complex numbers on argand plane then $\overrightarrow{PQ}$ represents complex number $z_2 - z_1$.



Note:


- (i) If $\overrightarrow{OP} = z = r e^{i\theta}$ then $\overrightarrow{OQ} = z_1 = r e^{i(\theta + \phi)} = z \cdot e^{i\phi}$. If $\overrightarrow{OP}$ and $\overrightarrow{OQ}$ are of unequal magnitude then $\hat{OQ} = \hat{OP} e^{i\phi}$ i.e.

$$\frac{z_1}{|z_1|} = \frac{z}{|z|} e^{i\phi}$$



(ii) In general, if z_1, z_2, z_3 be the three vertices of $\triangle ABC$


then $\frac{z_3 - z_1}{z_2 - z_1} = \frac{|z_3 - z_1|}{|z_2 - z_1|} e^{i\alpha}$. Here $\arg\left(\frac{z_3 - z_1}{z_2 - z_1}\right) = \alpha$



(iii) Note that the locus of z satisfying $\arg\left(\frac{z - z_1}{z - z_2}\right) = \alpha$ is :


Case (a) $0 < \alpha < \pi/2$

Locus is major arc of circle as shown
excluding z_1 & z_2



Case (b) $\frac{\pi}{2} < \alpha < \pi$

Locus is minor arc of circle as shown
excluding z_1 & z_2



(iv) If A, B, C & D are four points representing the complex numbers z_1, z_2, z_3 & z_4

then $AB \parallel CD$ if $\frac{z_4 - z_3}{z_2 - z_1}$ is purely real ;

$AB \perp CD$ if $\frac{z_4 - z_3}{z_2 - z_1}$ is purely imaginary.

(v) If z_1, z_2, z_3 are the vertices of an equilateral triangle where z_0 is its circumcentre then

$$(1) \quad z_1^2 + z_2^2 + z_3^2 - z_1 z_2 - z_2 z_3 - z_3 z_1 = 0$$

$$(2) \quad z_1^2 + z_2^2 + z_3^2 = 3z_0^2$$

Illustration 20 : Complex numbers z_1, z_2, z_3 are the vertices A, B, C respectively of an isosceles right angled triangle with right angle at C. Show that $(z_1 - z_2)^2 = 2(z_1 - z_3)(z_3 - z_2)$.

Solution : In the isosceles triangle ABC, $AC = BC$ and $BC \perp AC$. It means that AC is rotated through angle $\pi/2$ to occupy the position BC.

Hence we have, $\frac{z_2 - z_3}{z_1 - z_3} = e^{+i\pi/2} = +i \Rightarrow z_2 - z_3 = +i(z_1 - z_3)$

$$\Rightarrow z_2^2 + z_3^2 - 2z_2 z_3 = -(z_1^2 + z_3^2 - 2z_1 z_3)$$

$$\Rightarrow z_1^2 + z_2^2 - 2z_1 z_2 = 2z_1 z_3 + 2z_2 z_3 - 2z_1 z_2 - 2z_3^2 = 2(z_1 - z_3)(z_3 - z_2)$$

$$\Rightarrow (z_1 - z_2)^2 = 2(z_1 - z_3)(z_3 - z_2)$$




Illustration 21 : If the vertices of a square ABCD are z_1, z_2, z_3 & z_4 then find z_3 & z_4 in terms of z_1 & z_2 .

Solution : Using vector rotation at angle A

$$\frac{z_3 - z_1}{z_2 - z_1} = \frac{|z_3 - z_1|}{|z_2 - z_1|} e^{i\frac{\pi}{4}}$$

$$\because |z_3 - z_1| = AC \text{ and } |z_2 - z_1| = AB$$

$$\text{Also } AC = \sqrt{2} AB$$

$$\therefore |z_3 - z_1| = \sqrt{2} |z_2 - z_1|$$

$$\Rightarrow \frac{z_3 - z_1}{z_2 - z_1} = \sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$$

$$\Rightarrow z_3 - z_1 = (z_2 - z_1)(1 + i)$$

$$\Rightarrow z_3 = z_1 + (z_2 - z_1)(1 + i)$$

$$\text{Similarly } z_4 = z_2 + (1 + i)(z_1 - z_2)$$




Illustration 22 : Plot the region represented by $\frac{\pi}{3} \leq \arg\left(\frac{z+1}{z-1}\right) \leq \frac{2\pi}{3}$ in the Argand plane.


Solution : Let us take $\arg\left(\frac{z+1}{z-1}\right) = \frac{2\pi}{3}$, clearly z lies on the minor arc of the circle passing through $(1, 0)$ and $(-1, 0)$.

Similarly, $\arg\left(\frac{z+1}{z-1}\right) = \frac{\pi}{3}$ means that ' z ' is lying on the major arc of the circle passing through $(1, 0)$ and $(-1, 0)$.

Now if we take any point in the region included between two arcs say

$$P_1(z_1) \text{ we get } \frac{\pi}{3} \leq \arg\left(\frac{z+1}{z-1}\right) \leq \frac{2\pi}{3}$$

Thus $\frac{\pi}{3} \leq \arg\left(\frac{z+1}{z-1}\right) \leq \frac{2\pi}{3}$ represents the shaded region (excluding points $(1, 0)$ and $(-1, 0)$).



Do yourself -7 :

- (i) A complex number $z = 3 + 4i$ is rotated about another fixed complex number $z_1 = 1 + 2i$ in anticlockwise direction by 45° angle. Find the complex number represented by new position of z in argand plane.
- (ii) If A, B, C are three points in argand plane representing the complex number z_1, z_2, z_3 such that $z_1 = \frac{\lambda z_2 + z_3}{\lambda + 1}$, where $\lambda \in \mathbb{R}$, then find the distance of point A from the line joining points B and C.
- (iii) If $A(z_1), B(z_2), C(z_3)$ are vertices of ΔABC in which $\angle ABC = \frac{\pi}{4}$ and $\frac{AB}{BC} = \sqrt{2}$, then find z_2 in terms of z_1 and z_3 .
- (iv) If a & b are real numbers between 0 and 1 such that the points $z_1 = a + i, z_2 = 1 + bi$ and $z_3 = 0$ form an equilateral triangle then a and b are equal to :-
(A) $a = b = 1/2$ (B) $a = b = 2 - \sqrt{3}$ (C) $a = b = -2 + \sqrt{3}$ (D) $a = b = \sqrt{2} - 1$
- (v) If $\arg\left(\frac{z-1}{z+1}\right) = \frac{\pi}{4}$, find locus of z .

12. DE'MOIVRE'S THEOREM :

The value of $(\cos\theta + i\sin\theta)^n$ is $\cos n\theta + i\sin n\theta$ if 'n' is integer & it is one of the values of $(\cos\theta + i\sin\theta)^n$ if n is a rational number of the form p/q , where p & q are co-prime.

Note : Continued product of the roots of a complex quantity should be determined by using theory of equations.

Illustration 23 : If $\cos\alpha + \cos\beta + \cos\gamma = 0$ and also $\sin\alpha + \sin\beta + \sin\gamma = 0$, then prove that

(a) $\cos 2\alpha + \cos 2\beta + \cos 2\gamma = \sin 2\alpha + \sin 2\beta + \sin 2\gamma = 0$

(b) $\sin 3\alpha + \sin 3\beta + \sin 3\gamma = 3\sin(\alpha + \beta + \gamma)$

(c) $\cos 3\alpha + \cos 3\beta + \cos 3\gamma = 3\cos(\alpha + \beta + \gamma)$

Solution :

Let $z_1 = \cos\alpha + i\sin\alpha, z_2 = \cos\beta + i\sin\beta$ & $z_3 = \cos\gamma + i\sin\gamma$.

$\therefore z_1 + z_2 + z_3 = (\cos\alpha + \cos\beta + \cos\gamma) + i(\sin\alpha + \sin\beta + \sin\gamma)$
 $= 0 + i \cdot 0 = 0 \dots\dots\dots (i)$

(a) Also $\frac{1}{z_1} = (\cos\alpha + i\sin\alpha)^{-1} = \cos\alpha - i\sin\alpha$

$\frac{1}{z_2} = \cos\beta - i\sin\beta, \frac{1}{z_3} = \cos\gamma - i\sin\gamma$

$\therefore \frac{1}{z_1} + \frac{1}{z_2} + \frac{1}{z_3} = (\cos\alpha + \cos\beta + \cos\gamma) - i(\sin\alpha + \sin\beta + \sin\gamma) \dots\dots(ii)$
 $= 0 - i \cdot 0 = 0$

$$\text{Now } z_1^2 + z_2^2 + z_3^2 = (z_1 + z_2 + z_3)^2 - 2(z_1z_2 + z_2z_3 + z_3z_1)$$

$$= 0 - 2z_1z_2z_3 \left(\frac{1}{z_3} + \frac{1}{z_1} + \frac{1}{z_2} \right) = 0 - 2z_1z_2z_3 \cdot 0 = 0 \quad \{\text{using (i) and (ii)}\}$$

$$\text{or } (\cos\alpha + i\sin\alpha)^2 + (\cos\beta + i\sin\beta)^2 + (\cos\gamma + i\sin\gamma)^2 = 0$$

$$\text{or } \cos 2\alpha + i\sin 2\alpha + \cos 2\beta + i\sin 2\beta + \cos 2\gamma + i\sin 2\gamma = 0 + i \cdot 0$$

Equating real and imaginary parts on both sides,

$$\cos 2\alpha + \cos 2\beta + \cos 2\gamma = 0 \text{ and } \sin 2\alpha + \sin 2\beta + \sin 2\gamma = 0$$

(b) If $z_1 + z_2 + z_3 = 0$ then $z_1^3 + z_2^3 + z_3^3 = 3z_1z_2z_3$

$$\therefore (\cos\alpha + i\sin\alpha)^3 + (\cos\beta + i\sin\beta)^3 + (\cos\gamma + i\sin\gamma)^3 \\ = 3(\cos\alpha + i\sin\alpha)(\cos\beta + i\sin\beta)(\cos\gamma + i\sin\gamma)$$

$$\text{or } \cos 3\alpha + i\sin 3\alpha + \cos 3\beta + i\sin 3\beta + \cos 3\gamma + i\sin 3\gamma \\ = 3\{\cos(\alpha + \beta + \gamma) + i\sin(\alpha + \beta + \gamma)\}$$

Equating imaginary parts on both sides, $\sin 3\alpha + \sin 3\beta + \sin 3\gamma = 3\sin(\alpha + \beta + \gamma)$

(c) Equating real parts on both sides, $\cos 3\alpha + \cos 3\beta + \cos 3\gamma = 3\cos(\alpha + \beta + \gamma)$

Do yourself -8 :

- (i) If $z_r = \cos \frac{2r\pi}{5} + i\sin \frac{2r\pi}{5}$, $r = 0, 1, 3, 4, \dots$, then $z_1z_2z_3z_4z_5$ is equal to -
 (A) -1 (B) 0 (C) 1 (D) none of these
- (ii) If $(x-1)^4 - 16 = 0$, then the sum of nonreal complex values of x is -
 (A) 2 (B) 0 (C) 4 (D) none of these
- (iii) If $(\sqrt{3} - i)^n = 2^n$, $n \in \mathbb{Z}$, then n is a multiple of -
 (A) 6 (B) 10 (C) 9 (D) 12

13. CUBE ROOT OF UNITY :

(a) The cube roots of unity are $1, \frac{-1+i\sqrt{3}}{2}(\omega), \frac{-1-i\sqrt{3}}{2}(\omega^2)$.

(b) If ω is one of the imaginary cube roots of unity then $1 + \omega + \omega^2 = 0$. In general $1 + \omega^r + \omega^{2r} = 0$; where $r \in \mathbb{I}$ but is not the multiple of 3 & $1 + \omega^r + \omega^{2r} = 3$ if $r = 3\lambda$; $\lambda \in \mathbb{I}$

(c) In polar form the cube roots of unity are :

$$1 = \cos 0 + i\sin 0; \omega = \cos \frac{2\pi}{3} + i\sin \frac{2\pi}{3}, \omega^2 = \cos \frac{4\pi}{3} + i\sin \frac{4\pi}{3}$$

(d) The three cube roots of unity when plotted on the argand plane constitute the vertices of an equilateral triangle.

(e) The following factorisation should be remembered :

(a, b, c ∈ R & ω is the cube root of unity)

$$a^3 - b^3 = (a - b)(a - \omega b)(a - \omega^2 b); \quad x^2 + x + 1 = (x - \omega)(x - \omega^2);$$

$$a^3 + b^3 = (a + b)(a + \omega b)(a + \omega^2 b);$$

$$a^3 + b^3 + c^3 - 3abc = (a + b + c)(a + \omega b + \omega^2 c)(a + \omega^2 b + \omega c)$$




Illustration 24 : If α & β are imaginary cube roots of unity then αⁿ + βⁿ is equal to (where n ∈ I) –

(A) $2\cos\frac{2n\pi}{3}$ (B) $\cos\frac{2n\pi}{3}$ (C) $2i\sin\frac{2n\pi}{3}$ (D) $i\sin\frac{2n\pi}{3}$

Solution :

$$\alpha = \cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3};$$

$$\beta = \cos\frac{2\pi}{3} - i\sin\frac{2\pi}{3}$$

$$\begin{aligned} \alpha^n + \beta^n &= \left(\cos\frac{2\pi}{3} + i\sin\frac{2\pi}{3} \right)^n + \left(\cos\frac{2\pi}{3} - i\sin\frac{2\pi}{3} \right)^n \\ &= \left(\cos\frac{2n\pi}{3} + i\sin\frac{2n\pi}{3} \right) + \left(\cos\frac{2n\pi}{3} - i\sin\frac{2n\pi}{3} \right) = 2\cos\left(\frac{2n\pi}{3}\right) \end{aligned}$$

Ans.(A)

Illustration 25 : If α, β, γ are roots of x³ - 3x² + 3x + 7 = 0 (and ω is imaginary cube root of unity), then find the value of $\frac{\alpha-1}{\beta-1} + \frac{\beta-1}{\gamma-1} + \frac{\gamma-1}{\alpha-1}$.

Solution :

$$\text{We have } x^3 - 3x^2 + 3x + 7 = 0$$

$$\therefore (x-1)^3 + 8 = 0$$

$$\therefore (x-1)^3 = (-2)^3$$

$$\Rightarrow \left(\frac{x-1}{-2} \right)^3 = 1 \Rightarrow \frac{x-1}{-2} = (1)^{1/3} = 1, \omega, \omega^2 \quad (\text{cube roots of unity})$$

$$\therefore x = -1, 1 - 2\omega, 1 - 2\omega^2$$

$$\text{Here } \alpha = -1, \beta = 1 - 2\omega, \gamma = 1 - 2\omega^2$$

$$\therefore \alpha - 1 = -2, \beta - 1 = -2\omega, \gamma - 1 = -2\omega^2$$

$$\text{Then } \frac{\alpha-1}{\beta-1} + \frac{\beta-1}{\gamma-1} + \frac{\gamma-1}{\alpha-1} = \left(\frac{-2}{-2\omega} \right) + \left(\frac{-2\omega}{-2\omega^2} \right) + \left(\frac{-2\omega^2}{-2} \right) = \frac{1}{\omega} + \frac{1}{\omega} + \omega^2 = \omega^2 + \omega^2 + \omega^2$$

$$\text{Therefore } \frac{\alpha-1}{\beta-1} + \frac{\beta-1}{\gamma-1} + \frac{\gamma-1}{\alpha-1} = 3\omega^2.$$

Ans.

Do yourself -9 :

- (i) If ω is an imaginary cube root of unity, then $(1 + \omega - \omega^2)^2$ equals : -
 (A) ω (B) -4ω (C) ω^2 (D) 4ω
- (ii) If ω is a non real cube root of unity, then the expression $(1 - \omega)(1 - \omega^2)(1 + \omega^4)(1 + \omega^8)$ is equal to :-
 (A) 0 (B) 3 (C) 1 (D) 2

14. n^{th} ROOTS OF UNITY :

If $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$ are the n, n^{th} root of unity then :

- (a) They are in G.P. with common ratio $e^{i(2\pi/n)}$
- (b) Their arguments are in A.P. with common difference $\frac{2\pi}{n}$
- (c) The points represented by n, n^{th} roots of unity are located at the vertices of a regular polygon of n sides inscribed in a unit circle having center at origin, one vertex being on positive real axis.
- (d) $1^p + \alpha_1^p + \alpha_2^p + \dots + \alpha_{n-1}^p = 0$ if p is not an integral multiple of n
 $= n$ if p is an integral multiple of n
- (e) $(1 - \alpha_1)(1 - \alpha_2) \dots (1 - \alpha_{n-1}) = n$
- (f) $(1 + \alpha_1)(1 + \alpha_2) \dots (1 + \alpha_{n-1}) = 0$ if n is even and
 $= 1$ if n is odd.
- (g) $1, \alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1} = 1$ or -1 according as n is odd or even.




Illustration 26 : Find the value $\sum_{k=1}^6 \left(\sin \frac{2\pi k}{7} - \cos \frac{2\pi k}{7} \right)$

Solution :

$$\sum_{k=1}^6 \left(\sin \frac{2\pi k}{7} \right) - \sum_{k=1}^6 \left(\cos \frac{2\pi k}{7} \right) = \sum_{k=1}^6 \sin \frac{2\pi k}{7} - \sum_{k=0}^6 \cos \frac{2\pi k}{7} + 1$$

$$\sum_{k=0}^6 (\text{Sum of imaginary part of seven seventh roots of unity})$$

$$- \sum_{k=1}^6 (\text{Sum of real part of seven seventh roots of unity}) + 1 = 0 - 0 + 1 = 1$$

15. THE SUM OF THE FOLLOWING SERIES SHOULD BE REMEMBERED :


(a) $\cos\theta + \cos 2\theta + \cos 3\theta + \dots + \cos n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \cos\left(\frac{n+1}{2}\theta\right)$

(b) $\sin\theta + \sin 2\theta + \sin 3\theta + \dots + \sin n\theta = \frac{\sin(n\theta/2)}{\sin(\theta/2)} \sin\left(\frac{n+1}{2}\theta\right)$

Note : If $\theta = (2\pi/n)$ then the sum of the above series vanishes.


16. STRAIGHT LINES & CIRCLES IN TERMS OF COMPLEX NUMBERS :

(a) $\arg(z-\alpha) = \theta$ is a ray emanating from the complex




point α and inclined at an angle θ to the x-axis.


(b) $|z-a| = |z-b|$ is the perpendicular bisector of the segment joining a & b .



(c) The equation of a line joining z_1 & z_2 is given by; $z = z_1 + t(z_2 - z_1)$ where t is a parameter.



(d) $z = z_1(1 + it)$ where t is a real parameter, is a line through the point z_1 & perpendicular to z_1 .



- (e) The equation of a line passing through z_1 & z_2 can be expressed in the determinant form

as $\begin{vmatrix} z & \bar{z} & 1 \\ z_1 & \bar{z}_1 & 1 \\ z_2 & \bar{z}_2 & 1 \end{vmatrix} = 0$. This is also the condition for three complex numbers to be collinear.


- (f) Complex equation of a straight line through two given points z_1 & z_2 can be written as $z(\bar{z}_1 - \bar{z}_2) - \bar{z}(z_1 - z_2) + (z_1\bar{z}_2 - \bar{z}_1z_2) = 0$, which on manipulating takes the form as $\bar{\alpha}z + \alpha\bar{z} + r = 0$, where r is real and α is a non zero complex constant.

- (g) The equation of circle having centre z_0 & radius ρ is : $|z - z_0| = \rho$ or

$$z\bar{z} - z_0\bar{z} - \bar{z}_0z + \bar{z}_0z_0 - \rho^2 = 0 \text{ which is of the form}$$


$$z\bar{z} + \bar{\alpha}z + \alpha\bar{z} + r = 0 \text{ } r \text{ is real centre} = -\alpha \text{ \& radius} = \sqrt{\alpha\bar{\alpha} - r}. \text{ Circle}$$

will be real if $\alpha\bar{\alpha} - r \geq 0$.



- (h) $\arg\left(\frac{z - z_2}{z - z_1}\right) = \pm \frac{\pi}{2}$ or $(z - z_1)(\bar{z} - \bar{z}_2) + (z - z_2)(\bar{z} - \bar{z}_1) = 0$ this equation

represents the circle described on the line segment joining z_1 & z_2 as diameter.



- (i) Condition for four given points z_1, z_2, z_3 & z_4 to be concyclic is, the number $\frac{z_3 - z_1}{z_3 - z_2} \cdot \frac{z_4 - z_2}{z_4 - z_1}$ is real. Hence the equation of a circle through 3 non collinear points z_1, z_2 & z_3 can be taken as $\frac{(z - z_2)(z_3 - z_1)}{(z - z_1)(z_3 - z_2)}$ is real $\Rightarrow \frac{(z - z_2)(z_3 - z_1)}{(z - z_1)(z_3 - z_2)} = \frac{(\bar{z} - \bar{z}_2)(\bar{z}_3 - \bar{z}_1)}{(\bar{z} - \bar{z}_1)(\bar{z}_3 - \bar{z}_2)}$

Miscellaneous Illustration :

Illustration 27 : If z is a point on the Argand plane such that $|z - 1| = 1$, then $\frac{z - 2}{z}$ is equal to -

- (A) $\tan(\arg z)$ (B) $\cot(\arg z)$ (C) $i \tan(\arg z)$ (D) none of these

Solution :

Since $|z - 1| = 1$,

$\therefore$ let $z - 1 = \cos\theta + i \sin \theta$

Then, $z - 2 = \cos\theta + i \sin \theta - 1$

$$= -2\sin^2 \frac{\theta}{2} + 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2} = 2i \sin \frac{\theta}{2} \left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right) \dots\dots(i)$$

and $z = 1 + \cos \theta + i \sin \theta$

$$= 2\cos^2 \frac{\theta}{2} + 2i \sin \frac{\theta}{2} \cos \frac{\theta}{2} = 2\cos \frac{\theta}{2} \left(\cos \frac{\theta}{2} + i \sin \frac{\theta}{2} \right) \dots\dots(ii)$$

From (i) and (ii), we get $\frac{z - 2}{z} = i \tan \frac{\theta}{2} = i \tan(\arg z) \left(\because \arg z = \frac{\theta}{2} \text{ from (ii)} \right)$ **Ans.(C)**

Illustration 28 : Let a be a complex number such that $|a| < 1$ and $z_1, z_2, \dots, z_n$ be the vertices of a polygon such that $z_k = 1 + a + a^2 + \dots + a^k$, then show that vertices of the polygon lie within the circle $\left| z - \frac{1}{1-a} \right| = \frac{1}{|1-a|}$.

Solution : We have, $z_k = 1 + a + a^2 + \dots + a^k = \frac{1-a^{k+1}}{1-a}$
 $\Rightarrow z_k - \frac{1}{1-a} = \frac{-a^{k+1}}{1-a} \Rightarrow \left| z_k - \frac{1}{1-a} \right| = \frac{|a|^{k+1}}{|1-a|} < \frac{1}{|1-a|} \quad (\because |a| < 1)$
 $\therefore$ Vertices of the polygon $z_1, z_2, \dots, z_n$ lie within the circle $\left| z - \frac{1}{1-a} \right| = \frac{1}{|1-a|}$

Illustration 29 : If z_1 and z_2 are two complex numbers and $C > 0$, then prove that

$$|z_1 + z_2|^2 \leq (1+C)|z_1|^2 + (1+C^{-1})|z_2|^2$$

Solution : We have to prove that : $|z_1 + z_2|^2 \leq (1+C)|z_1|^2 + (1+C^{-1})|z_2|^2$
 i.e. $|z_1|^2 + |z_2|^2 + z_1 \bar{z}_2 + \bar{z}_1 z_2 \leq (1+C)|z_1|^2 + (1+C^{-1})|z_2|^2$
 or $z_1 \bar{z}_2 + \bar{z}_1 z_2 \leq C|z_1|^2 + C^{-1}|z_2|^2$
 or $C|z_1|^2 + \frac{1}{C}|z_2|^2 - z_1 \bar{z}_2 - \bar{z}_1 z_2 \geq 0$ (using $\text{Re}(z_1 \bar{z}_2) \leq |z_1 \bar{z}_2|$)
 or $\left(\sqrt{C}|z_1| - \frac{1}{\sqrt{C}}|z_2| \right)^2 \geq 0$ Which is always true.

Illustration 30 : If $\theta \in [\pi/6, \pi/3]$, $i = 1, 2, 3, 4, 5$ and $z^4 \cos \theta_1 + z^3 \cos \theta_2 + z^2 \cos \theta_3 + z \cos \theta_4 + \cos \theta_5 = 2\sqrt{3}$, then show that $|z| > \frac{3}{4}$

Solution : Given that $\cos \theta_1 z^4 + \cos \theta_2 z^3 + \cos \theta_3 z^2 + \cos \theta_4 z + \cos \theta_5 = 2\sqrt{3}$
 or $|\cos \theta_1 z^4 + \cos \theta_2 z^3 + \cos \theta_3 z^2 + \cos \theta_4 z + \cos \theta_5| = 2\sqrt{3}$
 $2\sqrt{3} \leq |\cos \theta_1 z^4| + |\cos \theta_2 z^3| + |\cos \theta_3 z^2| + |\cos \theta_4 z| + |\cos \theta_5|$
 $\therefore \theta_i \in [\pi/6, \pi/3]$
 $\therefore \frac{1}{2} \leq \cos \theta_i \leq \frac{\sqrt{3}}{2}$
 $2\sqrt{3} \leq \frac{\sqrt{3}}{2}|z|^4 + \frac{\sqrt{3}}{2}|z|^3 + \frac{\sqrt{3}}{2}|z|^2 + \frac{\sqrt{3}}{2}|z| + \frac{\sqrt{3}}{2}$
 $\Rightarrow 3 \leq |z|^4 + |z|^3 + |z|^2 + |z| \Rightarrow 3 < |z| + |z|^2 + |z|^3 + |z|^4 + |z|^5 + \dots \infty$
 $3 < \frac{|z|}{1-|z|} \Rightarrow 3-3|z| < |z| \Rightarrow 4|z| > 3$
 $\therefore |z| > \frac{3}{4}$

Illustration 31 : If z_1, z_2, z_3 are complex numbers such that $\frac{2}{z_1} = \frac{1}{z_2} + \frac{1}{z_3}$, show

that the points represented by z_1, z_2, z_3 lie on a circle passing through the origin.

Solution :

$$\text{We have, } \frac{2}{z_1} = \frac{1}{z_2} + \frac{1}{z_3} \Rightarrow \frac{1}{z_1} - \frac{1}{z_2} = \frac{1}{z_3} - \frac{1}{z_1} \Rightarrow \frac{z_2 - z_1}{z_1 z_2} = \frac{z_1 - z_3}{z_1 z_3}$$

$$\Rightarrow \frac{z_2 - z_1}{z_3 - z_1} = \frac{-z_2}{z_3} \Rightarrow \arg\left(\frac{z_2 - z_1}{z_3 - z_1}\right) = \arg\left(\frac{-z_2}{z_3}\right)$$

$$\arg\left(\frac{z_2 - z_1}{z_3 - z_1}\right) = \pi + \arg\left(\frac{z_2}{z_3}\right) \Rightarrow \text{or } \beta = \pi - \arg\frac{z_3}{z_2} = \pi - \alpha = \alpha + \beta = \pi$$

Thus the sum of a pair of opposite angle of a quadrilateral is 180° . Hence, the points O, z_1, z_2 and z_3 are the vertices of a cyclic quadrilateral i.e. lie on a circle.




Illustration 32 : Two given points P & Q are the reflection points w.r.t. a given straight line if the given line is the right bisector of the segment PQ. Prove that the two points denoted by the complex numbers z_1 & z_2 will be the reflection points for the straight line $\bar{\alpha}z + \alpha\bar{z} + r = 0$ if and only if ; $\bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0$, where r is real and α is non zero complex constant.

Solution :

Let P(z_1) is the reflection point of Q(z_2) then the perpendicular bisector of z_1 & z_2 must be the line

$$\bar{\alpha}z + \alpha\bar{z} + r = 0 \quad \dots\dots(i)$$

Now perpendicular bisector of z_1 & z_2 is, $|z - z_1| = |z - z_2|$

$$\text{or } (z - z_1)(\bar{z} - \bar{z}_1) = (z - z_2)(\bar{z} - \bar{z}_2)$$

$$-z\bar{z}_1 - z_1\bar{z} + z_1\bar{z}_1 = -z\bar{z}_2 - z_2\bar{z} + z_2\bar{z}_2 \quad (z\bar{z} \text{ cancels on either side})$$

$$\text{or } (\bar{z}_2 - \bar{z}_1)z + (z_2 - z_1)\bar{z} + z_1\bar{z}_1 - z_2\bar{z}_2 = 0 \quad \dots\dots(ii)$$

$$\text{Comparing (i) \& (ii) } \frac{\bar{\alpha}}{\bar{z}_2 - \bar{z}_1} = \frac{\alpha}{z_2 - z_1} = \frac{r}{z_1\bar{z}_1 - z_2\bar{z}_2} = \lambda$$

$$\therefore \bar{\alpha} = \lambda(\bar{z}_2 - \bar{z}_1) \quad \dots\dots(iii) \quad \alpha = \lambda(z_2 - z_1) \quad \dots\dots(iv)$$

$$r = \lambda(z_1\bar{z}_1 - z_2\bar{z}_2) \quad \dots\dots(v)$$

Multiplying (iii) by z_1 ; (iv) by $\bar{z}_2$ and adding

$$\bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0$$

Note that we could also multiply (iii) by z_2 & (iv) by $\bar{z}_1$ & add to get the same result.


$$\text{Hence } \bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0$$

Again, let $\bar{\alpha}z_1 + \alpha\bar{z}_2 + r = 0$ is true w.r.t. the line $\bar{\alpha}z + \alpha\bar{z} + r = 0$

$$\text{Subtracting } \bar{\alpha}(z - z_1) + \alpha(\bar{z} - \bar{z}_2) = 0$$


$$\text{or } |(z - z_1)| |\bar{\alpha}| = |\alpha| |(\bar{z} - \bar{z}_2)| \text{ or } |z - z_1| = \left| \frac{\alpha}{\bar{\alpha}} (\bar{z} - \bar{z}_2) \right| = |z - z_2|$$

Hence 'z' lies on the perpendicular bisector of joins of z_1 & z_2 .



ANSWERS FOR DO YOURSELF

- 1 : (i) $n = 4$ (ii) 0 2 : (i) $-17 + 24i$ (iii) $\pm(1 - 4i)$
- 3 : (i) $|z| = 4$; $\text{amp}(z) = \frac{2\pi}{3}$ (ii) $|z| = 2$; $\text{amp}(z) = -\frac{5\pi}{6}$ (iii) $|z| = 2$; $\text{amp}(z) = -\frac{\pi}{2}$
- (iv) $|z| = \frac{1}{\sqrt{2}}$; $\text{amp}(z) = \frac{3\pi}{4}$ (v) $|z| = 2$; $\text{amp}(z) = \frac{\pi}{3}$
- 4 : (i) 13 units
(ii) locus is a circle on complex plane with center at (2,3) and radius 1 unit. (iii) C
- 5 : (i) $2\sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right); 2\sqrt{2}e^{i\left(\frac{3\pi}{4}\right)}$ (ii) $2\left(\cos\frac{4\pi}{3} + i\sin\frac{4\pi}{3}\right); 2e^{i\left(\frac{4\pi}{3}\right)}$
- (iii) $\sqrt{2}\left(\cos\frac{3\pi}{4} + i\sin\frac{3\pi}{4}\right); \sqrt{2}e^{i\left(\frac{3\pi}{4}\right)}$
- (iv) $2\sin\left(\frac{\theta}{2}\right)\left(\cos\left(\frac{\pi}{2} - \frac{\theta}{2}\right) + i\sin\left(\frac{\pi}{2} - \frac{\theta}{2}\right)\right); 2\sin\left(\frac{\theta}{2}\right)e^{i\left(\frac{\pi}{2} - \frac{\theta}{2}\right)}$
- 6 : (i) C (ii) D
- 7 : (i) $1 + (2 + 2\sqrt{2})i$ (ii) 0 (iii) $z_2 = z_3 + i(z_1 - z_3)$ (iv) B
- (v) Locus is all the points on the major arc of circle as shown excluding points 1 & -1.



- 8 : (i) C (ii) A (iii) D
- 9 : (i) D (ii) B

EXERCISE # 1

PART-A

- Simplify and express the result in the form of $a + bi$
 (a) $\left(\frac{1+2i}{2+i}\right)^2$ (b) $-i(9+6i)(2-i)^{-1}$ (c) $\left(\frac{4i^3-i}{2i+1}\right)^2$ (d) $\frac{(2+i)^2}{2-i} - \frac{(2-i)^2}{2+i}$
 (e) A square $P_1P_2P_3P_4$ is drawn in the complex plane with P_1 at $(1, 0)$ and P_3 at $(3, 0)$. Let P_n denotes the point (x_n, y_n) $n = 1, 2, 3, 4$. Find the numerical value of the product of complex numbers $(x_1 + i y_1)(x_2 + i y_2)(x_3 + i y_3)(x_4 + i y_4)$.
- Given that $x, y \in \mathbb{R}$, solve :
 (a) $(x + 2y) + i(2x - 3y) = 5 - 4i$
 (b) $(x + iy) + (7 - 5i) = 9 + 4i$
 (c) $x^2 - y^2 - i(2x + y) = 2i$
- Find the square root of :
 (a) $9 + 40i$ (b) $-11 - 60i$ (c) $50i$
- (a) If $f(x) = x^4 + 9x^3 + 35x^2 - x + 4$, find $f(-5 + 4i)$
 (b) If $g(x) = x^4 - x^3 + x^2 + 3x - 5$, find $g(2 + 3i)$
- Solve the following equations over $\mathbb{C}$ and express the result in the form $a + ib$, $a, b \in \mathbb{R}$.
 (a) $ix^2 - 3x - 2i = 0$ (b) $2(1+i)x^2 - 4(2-i)x - 5 - 3i = 0$
- Locate the points representing the complex number z on the Argand plane :
 (a) $|z + 1 - 2i| = \sqrt{7}$ (b) $|z - 1|^2 + |z + 1|^2 = 4$
 (c) $\left|\frac{z-3}{z+3}\right| = 3$ (D) $|z - 3| = |z - 6|$
- If a & b are real numbers between 0 & 1 such that the points $z_1 = a + i$, $z_2 = 1 + bi$ & $z_3 = 0$ form an equilateral triangle, then find the values of 'a' and 'b'.
- Let $z_1 = 1 + i$ and $z_2 = -1 - i$. Find $z_3 \in \mathbb{C}$ such that triangle $z_1z_2z_3$ is equilateral.
- For what real values of x & y are the numbers $-3 + ix^2y$ & $x^2 + y + 4i$ conjugate complex ?
- If $(x + iy)^{1/3} = a + bi$, then prove that $4(a^2 - b^2) = \frac{x}{a} + \frac{y}{b}$.
- (a) Prove the identity, $|1 - z_1\bar{z}_2|^2 - |z_1 - z_2|^2 = (1 - |z_1|^2)(1 - |z_2|^2)$
 (b) Prove the identity, $|1 + z_1\bar{z}_2|^2 + |z_1 - z_2|^2 = (1 + |z_1|^2)(1 + |z_2|^2)$
 (c) For any two complex numbers, prove that $|z_1 + z_2|^2 + |z_1 - z_2|^2 = 2[|z_1|^2 + |z_2|^2]$. Also give the geometrical interpretation of this identity.

12. Find the Cartesian equation of the locus of 'z' in the complex plane satisfying, $|z - 4| + |z + 4| = 16$.

Paragraph for question nos. 13 to 15

Consider a complex number $w = \frac{z-i}{2z+1}$, where $z = x + iy$ and $x, y \in \mathbb{R}$.

13. If the complex number w is purely imaginary then locus of z is -
 (A) a straight line
 (B) a circle with centre $\left(-\frac{1}{4}, \frac{1}{2}\right)$ and radius $\frac{\sqrt{5}}{4}$.
 (C) a circle with centre $\left(\frac{1}{4}, -\frac{1}{2}\right)$ and passing through origin.
 (D) neither a circle nor a straight line.
14. If the complex number w is purely real then locus of z is
 (A) a straight line passing through origin
 (B) a straight line with gradient 3 and y intercept (-1)
 (C) a straight line with gradient 2 and y intercept 1.
 (D) none
15. If $|w| = 1$ then the locus of P is
 (A) a point circle (B) an imaginary circle
 (C) a real circle (D) not a circle.


PART-B

1. If $z + z^3 = 0$ then which of the following must be true on the complex plane?
 (A) $\text{Re}(z) < 0$ (B) $\text{Re}(z) = 0$ (C) $\text{Im}(z) = 0$ (D) $z^4 = 1$
2. Number of integral values of n for which the quantity $(n + i)^4$ where $i^2 = -1$, is an integer is
 (A) 1 (B) 2 (C) 3 (D) 4
3. Let $i = \sqrt{-1}$. The product of the real part of the roots of $z^2 - z = 5 - 5i$ is
 (A) -25 (B) -6 (C) -5 (D) 25
4. There is only one way to choose real numbers M and N such that when the polynomial $5x^4 + 4x^3 + 3x^2 + Mx + N$ is divided by the polynomial $x^2 + 1$, the remainder is 0. If M and N assume these unique values, then $M - N$ is
 (A) -6 (B) -2 (C) 6 (D) 2
5. The complex number z satisfying $z + |z| = 1 + 7i$ then the value of $|z|^2$ equals
 (A) 625 (B) 169 (C) 49 (D) 25
6. Number of values of z (real or complex) simultaneously satisfying the system of equations $1 + z + z^2 + z^3 + \dots + z^{17} = 0$ and $1 + z + z^2 + z^3 + \dots + z^{13} = 0$ is
 (A) 1 (B) 2 (C) 3 (D) 4

7. If $\frac{x-3}{3+i} + \frac{y-3}{3-i} = i$ where $x, y \in \mathbb{R}$ then
(A) $x = 2$ & $y = -8$ (B) $x = -2$ & $y = 8$ (C) $x = -2$ & $y = -6$ (D) $x = 2$ & $y = 8$
8. Number of complex numbers z satisfying $z^3 = \bar{z}$ is
(A) 1 (B) 2 (C) 4 (D) 5
9. Let $z = 9 + bi$ where b is non zero real and $i^2 = -1$. If the imaginary part of z^2 and z^3 are equal, then b^2 equals
(A) 261 (B) 225 (C) 125 (D) 361
10. The value of sum $\sum_{n=1}^{13} (i^n + i^{n+1})$, where $i = \sqrt{-1}$, equals
(A) i (B) $i - 1$ (C) -1 (D) 0

One or more than one is/are correct :

11. The diagram shows several numbers in the complex plane. The circle is the unit circle centered at the origin. One of these numbers is the reciprocal of F , which is



- (A) A (B) B (C) C (D) D
12. If $z = x + iy$ & $\omega = \frac{1-iz}{z-i}$ then $|\omega| = 1$ implies that, in the complex plane
(A) z lies on the imaginary axis (B) z lies on the real axis
(C) z lies on the unit circle (D) none
13. On the complex plane locus of a point z satisfying the inequality
 $2 \leq |z - 1| < 3$ denotes
(A) region between the concentric circles of radii 3 and 1 centered at $(1, 0)$
(B) region between the concentric circles of radii 3 and 2 centered at $(1, 0)$ excluding the inner and outer boundaries.
(C) region between the concentric circles of radii 3 and 2 centered at $(1, 0)$ including the inner and outer boundaries.
(D) region between the concentric circles of radii 3 and 2 centered at $(1, 0)$ including the inner boundary and excluding the outer boundary.
14. The complex number z satisfies $z + |z| = 2 + 8i$. The value of $|z|$ is
(A) 10 (B) 13 (C) 17 (D) 23

15. Let $Z_1 = (8 + i)\sin \theta + (7 + 4i)\cos \theta$ and $Z_2 = (1 + 8i)\sin \theta + (4 + 7i)\cos \theta$ are two complex numbers. If $Z_1 \cdot Z_2 = a + ib$ where $a, b \in \mathbb{R}$ then the largest value of $(a + b) \forall \theta \in \mathbb{R}$, is
(A) 75 (B) 100 (C) 125 (D) 130
16. The locus of z , for $\arg z = -\pi/3$ is
(A) same as the locus of z for $\arg z = 2\pi/3$
(B) same as the locus of z for $\arg z = \pi/3$
(C) the part of the straight line $\sqrt{3}x + y = 0$ with $(y < 0, x > 0)$
(D) the part of the straight line $\sqrt{3}x + y = 0$ with $(y > 0, x < 0)$
17. If z_1 & $\bar{z}_1$ represent adjacent vertices of a regular polygon of n sides with centre at the origin & if $\frac{\operatorname{Im} z_1}{\operatorname{Re} z_1} = \sqrt{2} - 1$ then the value of n is equal to :
(A) 8 (B) 12 (C) 16 (D) 24
18. All real numbers x which satisfy the inequality $|1 + 4i - 2^{-x}| \leq 5$ where $i = \sqrt{-1}$, $x \in \mathbb{R}$ are
(A) $[-2, \infty)$ (B) $(-\infty, 2]$ (C) $[0, \infty)$ (D) $[-2, 0]$
19. For $Z_1 = \sqrt[6]{\frac{1-i}{1+i\sqrt{3}}}$; $Z_2 = \sqrt[6]{\frac{1-i}{\sqrt{3}+i}}$; $Z_3 = \sqrt[6]{\frac{1+i}{\sqrt{3}-i}}$ which of the following holds good?
(A) $\sum |Z_1|^2 = \frac{3}{2}$ (B) $|Z_1|^4 + |Z_2|^4 = |Z_3|^8$
(C) $\sum |Z_1|^3 + |Z_2|^3 = |Z_3|^6$ (D) $|Z_1|^4 + |Z_2|^4 = |Z_3|^8$
20. Number of real or purely imaginary solution of the equation, $z^3 + iz - 1 = 0$ is :
(A) zero (B) one (C) two (D) three
21. A point ' z ' moves on the curve $|z - 4 - 3i| = 2$ in an argand plane. The maximum and minimum values of $|z|$ are
(A) 2, 1 (B) 6, 5 (C) 4, 3 (D) 7, 3
22. If z is a complex number satisfying the equation $|z + i| + |z - i| = 8$, on the complex plane then maximum value of $|z|$ is
(A) 2 (B) 4 (C) 6 (D) 8
23. Let z_r ($1 \leq r \leq 4$) be complex numbers such that $|z_r| = \sqrt{r+1}$ and $|30z_1 + 20z_2 + 15z_3 + 12z_4| = k |z_1z_2z_3 + z_2z_3z_4 + z_3z_4z_1 + z_4z_1z_2|$. Then the value of k equals
(A) $|z_1z_2z_3|$ (B) $|z_2z_3z_4|$ (C) $|z_3z_4z_1|$ (D) $|z_4z_1z_2|$

24. Let Z be a complex number satisfying the equation $(Z^3 + 3)^2 = -16$ then $|Z|$ has the value equal to
(A) $5^{1/2}$ (B) $5^{1/3}$ (C) $5^{2/3}$ (D) 5
25. If z_1, z_2, z_3 are 3 distinct complex numbers such that $\frac{3}{|z_2 - z_3|} = \frac{4}{|z_3 - z_1|} = \frac{5}{|z_1 - z_2|}$, then the value of $\frac{9}{z_2 - z_3} + \frac{16}{z_3 - z_1} + \frac{25}{z_1 - z_2}$ equals
(A) 0 (B) 3 (C) 4 (D) 5
26. The area of the triangle whose vertices are the roots $z^3 + iz^2 + 2i = 0$ is
(A) 2 (B) $\frac{3}{2}\sqrt{7}$ (C) $\frac{3}{4}\sqrt{7}$ (D) $\sqrt{7}$
27. Consider two complex numbers α and β as $\alpha = \left(\frac{a+bi}{a-bi}\right)^2 + \left(\frac{a-bi}{a+bi}\right)^2$, where $a, b \in \mathbb{R}$ and $\beta = \frac{z-1}{z+1}$, where $|z| = 1$, then
(A) Both α and β are purely real
(B) Both α and β are purely imaginary
(C) α is purely real and β is purely imaginary
(D) β is purely real and α is purely imaginary
28. Let Z is complex satisfying the equation $z^2 - (3+i)z + m + 2i = 0$, where $m \in \mathbb{R}$. Suppose the equation has a real root. The additive inverse of non real root, is
(A) $1-i$ (B) $1+i$ (C) $-1-i$ (D) -2
29. The minimum value of $|z-1+2i| + |4i-3-z|$ is
(A) $\sqrt{5}$ (B) 5 (C) $2\sqrt{13}$ (D) $\sqrt{15}$
30. If $i = \sqrt{-1}$, then $4 + 5\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^{334} + 3\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^{365}$ is equal to
(A) $1-i\sqrt{3}$ (B) $-1+i\sqrt{3}$ (C) $i\sqrt{3}$ (D) $-i\sqrt{3}$
31. Let C_1 and C_2 are concentric circles of radius 1 and $8/3$ respectively having centre at $(3, 0)$ on the argand plane. If the complex number z satisfies the inequality, $\log_{1/3}\left(\frac{|z-3|^2+2}{11|z-3|-2}\right) > 1$ then:
(A) z lies outside C_1 but inside C_2
(B) z lies inside of both C_1 and C_2
(C) z lies outside both of C_1 and C_2
(D) none of these

32. Identify the incorrect statement.
 (A) no non zero complex number z satisfies the equation, $\bar{z} = -4z$
 (B) $\bar{z} = z$ implies that z is purely real
 (C) $\bar{z} = -z$ implies that z is purely imaginary
 (D) if z_1, z_2 are the roots of the quadratic equation $az^2 + bz + c = 0$ such that $\text{Im}(z_1 z_2) \neq 0$ then a, b, c must be real numbers.
33. The equation of the radical axis of the two circles represented by the equations, $|z - 2| = 3$ and $|z - 2 - 3i| = 4$ on the complex plane is :
 (A) $3y + 1 = 0$ (B) $3y - 1 = 0$ (C) $2y - 1 = 0$ (D) none
34. $z_1 = \frac{a}{1-i}; z_2 = \frac{b}{2+i}; z_3 = a - bi$ for $a, b \in \mathbb{R}$
 if $z_1 - z_2 = 1$ then the centroid of the triangle formed by the points z_1, z_2, z_3 in the argand's plane is given by
 (A) $\frac{1}{9}(1+7i)$ (B) $\frac{1}{3}(1+7i)$ (C) $\frac{1}{3}(1-3i)$ (D) $\frac{1}{9}(1-3i)$
35. Consider the equation $10z^2 - 3iz - k = 0$, where z is a complex variable and $i^2 = -1$. Which of the following statements is True?
 (A) For all real positive numbers k , both roots are pure imaginary.
 (B) For negative real numbers k , both roots are pure imaginary.
 (C) For all pure imaginary numbers k , both roots are real and irrational.
 (D) For all complex numbers k , neither root is real.
36. Number of complex numbers z such that $|z| = 1$ and $\left| \frac{z}{\bar{z}} + \frac{\bar{z}}{z} \right| = 1$ is
 (A) 4 (B) 6 (C) 8 (D) more than 8
37. If z is a complex number satisfying the equation $|z - (1 + i)|^2 = 2$ and $\omega = \frac{2}{z}$, then the locus traced by ' ω ' in the complex plane is
 (A) $x - y - 1 = 0$ (B) $x + y - 1 = 0$ (C) $x - y + 1 = 0$ (D) $x + y + 1 = 0$
38. If P and Q are respectively by the complex numbers z_1 and z_2 such that $\left| \frac{1}{z_1} + \frac{1}{z_2} \right| = \left| \frac{1}{z_1} - \frac{1}{z_2} \right|$, then the circumcentre of ΔOPQ (where O is the origin) is
 (A) $\frac{z_1 - z_2}{2}$ (B) $\frac{z_1 + z_2}{2}$ (C) $\frac{z_1 + z_2}{3}$ (D) $z_1 + z_2$
39. If z_1 & z_2 are two non-zero complex numbers such that $|z_1 + z_2| = |z_1| + |z_2|$, then $\text{Arg } z_1 - \text{Arg } z_2$ is equal to
 (A) $-\pi$ (B) $-\pi/2$ (C) 0 (D) $\pi/2$

40. A particle starts from a point $z_0 = 1 + i$, where $i = \sqrt{-1}$. It moves horizontally away from origin by 2 units and then vertically away from origin by 3 units to reach a point z_1 . From z_1 particle moves $\sqrt{5}$ units in the direction of $2\hat{i} + \hat{j}$ and then it moves through an angle of $\operatorname{cosec}^{-1} \sqrt{2}$ in anticlockwise direction of a circle with centre at origin to reach a point z_2 . The $\arg z_2$ is given by
 (A) $\sec^{-1} 2$ (B) $\cot^{-1} 0$ (C) $\sin^{-1} \left(\frac{\sqrt{3}-1}{2\sqrt{2}} \right)$ (D) $\cos^{-1} \left(\frac{-1}{2} \right)$
41. Consider $az^2 + bz + c = 0$, where $a, b, c \in \mathbb{R}$ and $4ac > b^2$.
 (i) If z_1 and z_2 are the roots of the equation given above, then which one of the following complex numbers is purely real?
 (A) $z_1 \bar{z}_2$ (B) $\bar{z}_1 z_2$ (C) $z_1 - z_2$ (D) $(z_1 - z_2)i$
 (ii) In the argand's plane, if A is the point representing z_1 , B is the point representing z_2 & $z = \frac{\overrightarrow{OA}}{\overrightarrow{OB}}$ then
 (A) z is purely real (B) z is purely imaginary
 (C) $|z| = 1$ (D) ΔAOB is a scalene triangle.
42. If the complex number z satisfies the condition $|z| \geq 3$, then the least value of $\left| z + \frac{1}{z} \right|$ is equal to:
 (A) $5/3$ (B) $8/3$ (C) $11/3$ (D) none of these
43. Given $z_p = \cos \left(\frac{\pi}{2^p} \right) + i \sin \left(\frac{\pi}{2^p} \right)$, then $\lim_{n \rightarrow \infty} (z_1 z_2 z_3 \dots z_n) =$
 (A) 1 (B) -1 (C) i (D) $-i$
44. The maximum & minimum values of $|z + 1|$ when $|z + 3| \leq 3$ are :
 (A) (5, 0) (B) (6, 0) (C) (7, 1) (D) (5, 1)
45. If $|z| = 1$ and $|\omega - 1| = 1$ where $z, \omega \in \mathbb{C}$, then the largest set of values of $|2z - 1|^2 + |2\omega - 1|^2$ equals
 (A) [1, 9] (B) [2, 6] (C) [2, 12] (D) [2, 18]
46. If $\operatorname{Arg}(z + a) = \frac{\pi}{6}$ and $\operatorname{Arg}(z - a) = \frac{2\pi}{3}$; $a \in \mathbb{R}^+$, then
 (A) z is independent of a (B) $|a| = |z + a|$
 (C) $z = a$ (D) $z = a$ is $\frac{\pi}{6}$ is $\frac{\pi}{3}$
47. If z_1, z_2, z_3 are the vertices of the ΔABC on the complex plane which are also the roots of the equation, $z^3 - 3\alpha z^2 + 3\beta z + x = 0$, then the condition for the ΔABC to be equilateral triangle is
 (A) $\alpha^2 = \beta$ (B) $\alpha = \beta^2$ (C) $\alpha^2 = 3\beta$ (D) $\alpha = 3\beta^2$

48. The locus represented by the equation, $|z - 1| + |z + 1| = 2$ is :
 (A) an ellipse with focii $(1, 0)$; $(-1, 0)$
 (B) one of the family of circles passing through the points of intersection of the circles $|z - 1| = 1$ & $|z + 1| = 1$
 (C) the radical axis of the circles $|z - 1| = 1$ and $|z + 1| = 1$
 (D) the portion of the real axis between the points $(1, 0)$; $(-1, 0)$ including both.

Paragraph for question nos. 49 to 51

Let A, B, C be three sets of complex numbers as defined below.

$$A = \{z : |z + 1| \leq 2 + \operatorname{Re}(z)\}, B = \{z : |z - 1| \geq 1\} \text{ and } C = \left\{z : \left|\frac{z-1}{z+1}\right| \geq 1\right\}$$

49. The number of point(s) having integral coordinates in the region $A \cap B \cap C$ is
 (A) 4 (B) 5 (C) 6 (D) 10
50. The area of region bounded by $A \cap B \cap C$ is
 (A) $2\sqrt{3}$ (B) $\sqrt{3}$ (C) $4\sqrt{3}$ (D) 2
51. The real part of the complex number in the region $A \cap B \cap C$ and having maximum amplitude is
 (A) -1 (B) $-\frac{3}{2}$ (C) $\frac{1}{2}$ (D) -2

[MATCH THE COLUMN]

52. Match the equation in z, in Column-I with the corresponding values of $\arg(z)$ in Column-II.

Column-I

(equations in z)

(A) $z^2 - z + 1 = 0$

(B) $z^2 + z + 1 = 0$

(C) $2z^2 + 1 + i\sqrt{3} = 0$

(D) $2z^2 + 1 - i\sqrt{3} = 0$

Column-II

(principal value of $\arg(z)$)

(P) $-2\pi/3$

(Q) $-\pi/3$

(R) $\pi/3$

(S) $2\pi/3$

EXERCISE # 2

- Find the modulus, argument and the principal argument of the complex numbers.
(i) $6(\cos 310^\circ - i \sin 310^\circ)$ (ii) $-2(\cos 30^\circ + i \sin 30^\circ)$ (iii) $\frac{2+i}{4i+(1+i)^2}$
- Let Z is complex satisfying the equation, $z^2 - (3+i)z + m + 2i = 0$, where $m \in \mathbb{R}$. Suppose the equation has a real root, then find the value of m .
 - a, b, c are real numbers in the polynomial, $P(Z) = 2Z^4 + aZ^3 + bZ^2 + cZ + 3$. If two roots of the equation $P(Z) = 0$ are 2 and i , then find the value of ' a '.
- Find the real values of x & y for which $z_1 = 9y^2 - 4 - 10ix$ and $z_2 = 8y^2 - 20i$ are conjugate complex of each other.
- Solve the following equation $z^2 - (3-2i)z = (5i-5)$ expressing your answer in the form of $(a+ib)$.
 - If $(1-i)$ is a root of the equation $z^3 - 2(2-i)z^2 + (4-5i)z - 1 + 3i = 0$, then find the other two roots.
- If $iZ^3 + Z^2 - Z + i = 0$, then find $|Z|$.
 - Let z_1 and z_2 be two complex numbers such that $\left| \frac{z_1 - 2z_2}{2 - z_1\bar{z}_2} \right| = 1$ and $|z_2| \neq 1$, find $|z_1|$.
 - Find the minimum value of the expression $E = |z|^2 + |z-3|^2 + |z-6i|^2$
(where $z = x + iy, x, y \in \mathbb{R}$)
- Show that the product,

$$\left[1 + \left(\frac{1+i}{2} \right) \right] \left[1 + \left(\frac{1+i}{2} \right)^2 \right] \left[1 + \left(\frac{1+i}{2} \right)^{2^2} \right] \dots \left[1 + \left(\frac{1+i}{2} \right)^{2^n} \right]$$
is equal to $\left(1 - \frac{1}{2^{2^n}} \right) (1+i)$ where $n \geq 2$.
- Let z be a complex number such that $z \in \mathbb{C} \setminus \mathbb{R}$ and $\frac{1+z+z^2}{1-z+z^2} \in \mathbb{R}$, then prove that $|z| = 1$.
- Let $z = (0, 1) \in \mathbb{C}$. Express $\sum_{k=0}^n z^k$ in terms of the positive integer n .
- Among the complex numbers z satisfying the condition $|z + 3 - \sqrt{3}i| = \sqrt{3}$, find the number having the least positive argument.
- If A, B and C are the angles of a triangle $D = \begin{vmatrix} e^{-2iA} & e^{iC} & e^{iB} \\ e^{iC} & e^{-2iB} & e^{iA} \\ e^{iB} & e^{iA} & e^{-2iC} \end{vmatrix}$, where $i = \sqrt{-1}$, then find the value of D .

11. Dividing $f(z)$ by $z - i$, we get the remainder i and dividing it by $z + i$, we get the remainder $1 + i$. Find the remainder upon the division of $f(z)$ by $z^2 + 1$.
12. (a) Find all non-zero complex numbers Z satisfying $\bar{Z} = i Z^2$.
 (b) If the complex numbers $z_1, z_2, \dots, z_n$ lie on the unit circle $|z| = 1$ then show that $|z_1 + z_2 + \dots + z_n| = |z_1^{-1} + z_2^{-1} + \dots + z_n^{-1}|$.
13. Let $|z| = 2$ and $w = \frac{z+1}{z-1}$ where $z, w \in \mathbb{C}$ (where $\mathbb{C}$ is the set of complex numbers). If M and m respectively be the greatest and least modulus of w , then find the value of $(2010m + M)$.
14. If ω is the imaginary cube root of unity, then find the number of pairs of integers (a, b) such the $|a\omega + b| = 1$.



EXERCISE # 3 (JM)

1. If $|z + 4| \leq 3$, then the maximum value of $|z + 1|$ is- [AIEEE-2007]
 (1) 4 (2) 10 (3) 6 (4) 0
2. The conjugate of a complex number is $\frac{1}{i-1}$, then that complex number is- [AIEEE-2008]
 (1) $\frac{-1}{i-1}$ (2) $\frac{1}{i+1}$ (3) $\frac{-1}{i+1}$ (4) $\frac{1}{i-1}$
3. If $\left|Z - \frac{4}{Z}\right| = 2$, then the maximum value of $|Z|$ is equal to :- [AIEEE-2009]
 (1) 2 (2) $2 + \sqrt{2}$ (3) $\sqrt{3} + 1$ (4) $\sqrt{5} + 1$
4. The number of complex numbers z such that $|z - 1| = |z + 1| = |z - i|$ equals :- [AIEEE-2010]
 (1) 0 (2) 1 (3) 2 (4) ∞
5. If α and β are the roots of the equation $x^2 - x + 1 = 0$, then $\alpha^{2009} + \beta^{2009} =$ [AIEEE-2010]
 (1) -2 (2) -1 (3) 1 (4) 2
6. Let α, β be real and z be a complex number. If $z^2 + \alpha z + \beta = 0$ has two distinct roots on the line $\text{Re } z = 1$, then it is necessary that :- [AIEEE-2011]
 (1) $|\beta| = 1$ (2) $\beta \in (1, \infty)$ (3) $\beta \in (0, 1)$ (4) $\beta \in (-1, 0)$
7. If $\omega (\neq 1)$ is a cube root of unity, and $(1 + \omega)^7 = A + B\omega$. Then (A, B) equals :- [AIEEE-2011]
 (1) (1, 0) (2) (-1, 1) (3) (0, 1) (4) (1, 1)
8. If $z \neq 1$ and $\frac{z^2}{z-1}$ is real, then the point represented by the complex number z lies : [AIEEE-2012]
 (1) on the imaginary axis.
 (2) either on the real axis or on a circle passing through the origin.
 (3) on a circle with centre at the origin.
 (4) either on the real axis or on a circle not passing through the origin.
9. If z is a complex number of unit modulus and argument θ , then $\arg\left(\frac{1+z}{1+\bar{z}}\right)$ equals
 (1) $-\theta$ (2) $\frac{\pi}{2} - \theta$ (3) θ (4) $\pi - \theta$

[JEE (Main)-2013]

10. If z is a complex number such that $|z| \geq 2$, then the minimum value of $\left|z + \frac{1}{z}\right|$:
[JEE(Main)-2014]
- (1) is equal to $\frac{5}{2}$ (2) lies in the interval $(1, 2)$
(3) is strictly greater than $\frac{5}{2}$ (4) is strictly greater than $\frac{3}{2}$ but less than $\frac{5}{2}$
11. A complex number z is said to be unimodular if $|z| = 1$. Suppose z_1 and z_2 are complex numbers such that $\frac{z_1 - 2z_2}{2 - z_1\bar{z}_2}$ is unimodular and z_2 is not unimodular. Then the point z_1 lies on a :
[JEE(Main)-2015]
- (1) circle of radius 2 (2) circle of radius $\sqrt{2}$
(3) straight line parallel to x-axis (4) straight line parallel to y-axis
12. A value of θ for which $\frac{2 + 3i \sin \theta}{1 - 2i \sin \theta}$ is purely imaginary, is :
[JEE(Main)-2016]
- (1) $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$ (2) $\frac{\pi}{3}$ (3) $\frac{\pi}{6}$ (4) $\sin^{-1}\left(\frac{\sqrt{3}}{4}\right)$
13. Let ω be a complex number such that $2\omega + 1 = z$ where $z = \sqrt{-3}$. If $\begin{vmatrix} 1 & 1 & 1 \\ 1 & -\omega^2 - 1 & \omega^2 \\ 1 & \omega^2 & \omega^7 \end{vmatrix} = 3k$ then k is equal to :
[JEE(Main)-2017]
- (1) 1 (2) $-z$ (3) z (4) -1
14. If $\alpha, \beta \in \mathbb{C}$ are the distinct roots, of the equation $x^2 - x + 1 = 0$, then $\alpha^{101} + \beta^{107}$ is equal to :
[JEE(Main)-2018]
- (1) 2 (2) -1 (3) 0 (4) 1
15. Let α and β be two roots of the equation $x^2 + 2x + 2 = 0$, then $\alpha^{15} + \beta^{15}$ is equal to :
[JEE(Main)-2019]
- (1) 512 (2) -512 (3) -256 (4) 256
16. Let $A = \left\{ \theta \in \left(-\frac{\pi}{2}, \pi\right) : \frac{3 + 2i \sin \theta}{1 - 2i \sin \theta} \text{ is purely imaginary} \right\}$
Then the sum of the elements in A is :
[JEE(Main)-2019]
- (1) $\frac{5\pi}{6}$ (2) $\frac{2\pi}{3}$ (3) $\frac{3\pi}{4}$ (4) π
17. Let z_0 be a root of the quadratic equation, $x^2 + x + 1 = 0$. If $z = 3 + 6i z_0^{81} - 6i z_0^{93}$, then $\arg z$ is equal to :
[JEE(Main)-2019]
- (1) $\frac{\pi}{3}$ (2) 0 (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{4}$

18. Let z_1 and z_2 be any two non-zero complex numbers such that $3|z_1| = 4|z_2|$. If $z = \frac{3z_1}{2z_2} + \frac{2z_2}{3z_1}$ then:
[JEE(Main)-2019]
(1) $|z| = \frac{1}{2}\sqrt{\frac{17}{2}}$ (2) $\text{Im}(z) = 0$ (3) $\text{Re}(z) = 0$ (4) $|z| = \sqrt{\frac{5}{2}}$
19. Let $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^5 + \left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)^5$. If $R(z)$ and $I(z)$ respectively denote the real and imaginary parts of z , then :
[JEE(Main)-2019]
(1) $R(z) > 0$ and $I(z) > 0$ (2) $R(z) < 0$ and $I(z) > 0$
(3) $R(z) = -3$ (4) $I(z) = 0$
20. Let $\left(-2 - \frac{1}{3}i\right)^3 = \frac{x+iy}{27}$ ($i = \sqrt{-1}$), where x and y are real numbers, then $y - x$ equals ;
(1) 85 (2) -85 (3) 91 (4) -91
[JEE(Main)-2019]
21. Let z be a complex number such that $|z| + z = 3 + i$ (where $i = \sqrt{-1}$). Then $|z|$ is equal to:
(1) $\frac{\sqrt{41}}{4}$ (2) $\frac{5}{3}$ (3) $\frac{\sqrt{34}}{3}$ (4) $\frac{5}{4}$
[JEE(Main)-2019]
22. If $\frac{z-\alpha}{z+\alpha}$ ($\alpha \in \mathbb{R}$) is a purely imaginary number and $|z| = 2$, then a value of α is :
(1) $\sqrt{2}$ (2) 2 (3) 1 (4) $\frac{1}{2}$
[JEE(Main)-2019]
23. Let z_1 and z_2 be two complex numbers satisfying $|z_1| = 9$ and $|z_2 - 3 - 4i| = 4$. Then the minimum value of $|z_1 - z_2|$ is:
[JEE(Main)-2019]
(1) 0 (2) 1 (3) $\sqrt{2}$ (4) 2
24. If $z = \frac{\sqrt{3}}{2} + \frac{i}{2}$ ($i = \sqrt{-1}$), then $(1 + iz + z^5 + iz^8)^9$ is equal to :
[JEE(Main)-2019]
(1) -1 (2) $(-1 + 2i)^9$ (3) 0 (4) 1
25. All the points in the set $S = \left\{ \frac{\alpha+i}{\alpha-i}; \alpha \in \mathbb{R} \right\}$ ($i = \sqrt{-1}$) lie on a :
[JEE(Main)-2019]
(1) circle whose radius is 1 (2) straight line whose slope is 1
(3) circle whose radius is $\sqrt{2}$ (4) straight line whose slope is -1

26. Let $z \in \mathbb{C}$ be such that $|z| < 1$. If $\omega = \frac{5+3z}{5(1-z)}$, then : **[JEE(Main)-2019]**
 (1) $5 \operatorname{Re}(\omega) > 1$ (2) $4 \operatorname{Im}(\omega) > 5$ (3) $5 \operatorname{Re}(\omega) > 4$ (4) $5 \operatorname{Im}(\omega) < 1$
27. If $a > 0$ and $z = \frac{(1+i)^2}{a-i}$, has magnitude $\sqrt{\frac{2}{5}}$, then $\bar{z}$ is equal to : **[JEE(Main)-2019]**
 (1) $-\frac{3}{5} - \frac{1}{5}i$ (2) $\frac{1}{5} - \frac{3}{5}i$ (3) $-\frac{1}{5} - \frac{3}{5}i$ (4) $-\frac{1}{5} + \frac{3}{5}i$
28. If z and w are two complex numbers such that $|zw| = 1$ and $\arg(z) - \arg(w) = \frac{\pi}{2}$, then : **[JEE(Main)-2019]**
 (1) $z\bar{w} = \frac{1-i}{\sqrt{2}}$ (2) $z\bar{w} = \frac{-1+i}{\sqrt{2}}$ (3) $\bar{z}w = i$ (4) $\bar{z}w = -i$
29. The equation $|z-i| = |z-1|$, $i = \sqrt{-1}$, represents: **[JEE(Main)-2019]**
 (1) a circle of radius $\frac{1}{2}$. (2) the line through the origin with slope -1 .
 (3) the line through the origin with slope 1 . (4) a circle of radius 1 .
30. Let $z \in \mathbb{C}$ with $\operatorname{Im}(z) = 10$ and it satisfies $\frac{2z-n}{2z+n} = 2i - 1$ for some natural number n . Then : **[JEE(Main)-2019]**
 (1) $n = 20$ and $\operatorname{Re}(z) = -10$ (2) $n = 20$ and $\operatorname{Re}(z) = 10$
 (3) $n = 40$ and $\operatorname{Re}(z) = 10$ (4) $n = 40$ and $\operatorname{Re}(z) = -10$
31. If $\operatorname{Re}\left(\frac{z-1}{2z+i}\right) = 1$, where $z = x + iy$, then the point (x, y) lies on a : **[JEE(Main)-2020]**
 (1) circle whose diameter is $\frac{\sqrt{5}}{2}$. (2) straight line whose slope is $\frac{3}{2}$.
 (3) circle whose centre is at $\left(-\frac{1}{2}, -\frac{3}{2}\right)$. (4) straight line whose slope is $-\frac{2}{3}$.
32. If $\frac{3+i\sin\theta}{4-i\cos\theta}$, $\theta \in [0, 2\pi]$, is a real number, then an argument of $\sin\theta + i\cos\theta$ is : **[JEE(Main)-2020]**
 (1) $-\tan^{-1}\left(\frac{3}{4}\right)$ (2) $\tan^{-1}\left(\frac{4}{3}\right)$ (3) $\pi - \tan^{-1}\left(\frac{4}{3}\right)$ (4) $\pi - \tan^{-1}\left(\frac{3}{4}\right)$
33. If the equation, $x^2 + bx + 45 = 0$ ($b \in \mathbb{R}$) has conjugate complex roots and they satisfy $|z+1| = 2\sqrt{10}$, then : **[JEE(Main)-2020]**
 (1) $b^2 + b = 12$ (2) $b^2 - b = 30$ (3) $b^2 - b = 42$ (4) $b^2 + b = 72$

34. Let $\alpha = \frac{-1+i\sqrt{3}}{2}$, if $a = (1+\alpha)\sum_{k=0}^{100}\alpha^{2k}$ and $b = \sum_{k=0}^{100}\alpha^{3k}$ then [JEE(Main)-2020]
a and b are the roots of the quadratic equation
(1) $x^2 + 102x + 101 = 0$ (2) $x^2 + 101x + 100 = 0$
(3) $x^2 - 102x + 101 = 0$ (4) $x^2 - 101x + 100 = 0$
35. Let z be a complex number such that $\left|\frac{z-i}{z+2i}\right| = 1$ and $|z| = \frac{5}{2}$. Then value of $|z + 3i|$ is
(1) $\frac{7}{2}$ (2) $\sqrt{10}$ (3) $\frac{15}{4}$ (4) $2\sqrt{3}$
[JEE(Main)-2020]
36. If z be a complex number satisfying $|\operatorname{Re}(z)| + |\operatorname{Im}(z)| = 4$, then $|z|$ cannot be :
(1) $\sqrt{8}$ (2) $\sqrt{10}$ (3) $\sqrt{\frac{17}{2}}$ (4) $\sqrt{7}$
[JEE(Main)-2020]
37. The value of $\left(\frac{1+\sin\frac{2\pi}{9}+i\cos\frac{2\pi}{9}}{1+\sin\frac{2\pi}{9}-i\cos\frac{2\pi}{9}}\right)^3$ is : [JEE(Main)-2020]
(1) $\frac{1}{2}(1-i\sqrt{3})$ (2) $-\frac{1}{2}(1-i\sqrt{3})$ (3) $\frac{1}{2}(\sqrt{3}-i)$ (4) $-\frac{1}{2}(\sqrt{3}-i)$
38. The imaginary part of $(3+2\sqrt{-54})^{\frac{1}{2}} - (3-2\sqrt{-54})^{\frac{1}{2}}$ can be : [JEE(Main)-2020]
(1) $\sqrt{6}$ (2) $-2\sqrt{6}$ (3) $-\sqrt{6}$ (4) 6
39. If $\left(\frac{1+i}{1-i}\right)^{m/2} = \left(\frac{1+i}{i-1}\right)^{n/3} = 1$, ($m, n \in \mathbb{N}$) then the greatest common divisor of the least values of m and n is _____. [JEE(Main)-2020]
40. If z_1, z_2 are complex numbers such that $\operatorname{Re}(z_1) = |z_1 - 1|$, $\operatorname{Re}(z_2) = |z_2 - 1|$ and $\arg(z_1 - z_2) = \frac{\pi}{6}$, then $\operatorname{Im}(z_1 + z_2)$ is equal to : [JEE(Main)-2020]
(1) $\frac{\sqrt{3}}{2}$ (2) $\frac{2}{\sqrt{3}}$ (3) $\frac{1}{\sqrt{3}}$ (4) $2\sqrt{3}$
41. Let $u = \frac{2z+i}{z-ki}$, $z = x + iy$ and $k > 0$. If the curve represented by $\operatorname{Re}(u) + \operatorname{Im}(u) = 1$ intersects the y-axis at the points P and Q where $PQ = 5$, then the value of k is : [JEE(Main)-2020]
(1) $1/2$ (2) $3/2$ (3) 4 (4) 2

42. If a and b are real numbers such that $(2 + \alpha)^4 = a + b\alpha$, where $\alpha = \frac{-1+i\sqrt{3}}{2}$, then $a + b$ is equal to : [JEE(Main)-2020]
 (1) 33 (2) 9 (3) 57 (4) 24
43. If the four complex numbers z , $\bar{z}$, $\bar{z} - 2\operatorname{Re}(\bar{z})$ and $z - 2\operatorname{Re}(z)$ represent the vertices of a square of side 4 units in the Argand plane, then $|z|$ is equal to : [JEE(Main)-2020]
 (1) 2 (2) $4\sqrt{2}$ (3) $2\sqrt{2}$ (4) 4
44. The value of $\left(\frac{-1+i\sqrt{3}}{1-i}\right)^{30}$ is : [JEE(Main)-2020]
 (1) $-2^{15}i$ (2) $2^{15}i$ (3) 6^5 (4) -2^{15}
45. The region represented by $\{z = x + iy \in \mathbb{C} : |z| - \operatorname{Re}(z) \leq 1\}$ is also given by the inequality : [JEE(Main)-2020]
 (1) $y^2 \geq x + 1$ (2) $y^2 \geq 2(x + 1)$ (3) $y^2 \leq x + \frac{1}{2}$ (4) $y^2 \leq 2\left(x + \frac{1}{2}\right)$
46. Let $z = x + iy$ be a non-zero complex number such that $z^2 = i|z|^2$, where $i = \sqrt{-1}$, then z lies on the : [JEE(Main)-2020]
 (1) imaginary axis (2) real axis (3) line, $y = x$ (4) line, $y = -x$

EXERCISE # 4 (JA)

- A man walks a distance of 3 units from the origin towards the North-East ($N 45^\circ E$) direction. From there, he walks a distance of 4 units towards the North-West ($N 45^\circ W$) direction to reach a point P. Then the position of P in the Argand plane is **[JEE 2007, 3M]**

(A) $3e^{i\pi/4} + 4i$ (B) $(3 - 4i)e^{i\pi/4}$ (C) $(4 + 3i)e^{i\pi/4}$ (D) $(3 + 4i)e^{i\pi/4}$
- If $|z| = 1$ and $z \neq \pm 1$, then all the values of $\frac{z}{1-z^2}$ lie on **[JEE 2007, 3+3]**

(A) a line not passing through the origin (B) $|z| = \sqrt{2}$
(C) the x-axis (D) the y-axis
- A particle P starts from the point $z_0 = 1 + 2i$, where $i = \sqrt{-1}$. It moves first horizontally away from origin by 5 units and then vertically away from origin by 3 units to reach a point z_1 . From z_1 the particle moves $\sqrt{2}$ units in the direction of the vector $\hat{i} + \hat{j}$ and then it moves through an angle $\frac{\pi}{2}$ in anticlockwise direction on a circle with centre at origin, to reach a point z_2 . The point z_2 is given by **[JEE 2008, 3M, -1M]**

(A) $6 + 7i$ (B) $-7 + 6i$ (C) $7 + 6i$ (D) $-6 + 7i$
- Comprehension (3 questions together)**
Let A, B, C be three sets of complex numbers as defined below **[JEE 2008, 3 + 4 + 4 + 4]**
 $A = \{z : \operatorname{Im} z \geq 1\}$; $B = \{z : |z - 2 - i| = 3\}$; $C = \{z : \operatorname{Re}((1-i)z) = \sqrt{2}\}$

 - The number of elements in the set $A \cap B \cap C$ is
(A) 0 (B) 1 (C) 2 (D) ∞
 - Let z be any point in $A \cap B \cap C$. Then, $|z + 1 - i|^2 + |z - 5 - i|^2$ lies between
(A) 25 and 29 (B) 30 and 34 (C) 35 and 39 (D) 40 and 44
 - Let z be any point in $A \cap B \cap C$ and let w be any point satisfying $|w - 2 - i| < 3$. Then, $|z| - |w| + 3$ lies between
(A) -6 and 3 (B) -3 and 6 (C) -6 and 6 (D) -3 and 9
- Let $z = \cos \theta + i \sin \theta$. Then the value of $\sum_{m=1}^{15} \operatorname{Im}(z^{2m-1})$ at $\theta = 2^\circ$ is

(A) $\frac{1}{\sin 2^\circ}$ (B) $\frac{1}{3 \sin 2^\circ}$ (C) $\frac{1}{2 \sin 2^\circ}$ (D) $\frac{1}{4 \sin 2^\circ}$
- Let $z = x + iy$ be a complex number where x and y are integers. Then the area of the rectangle whose vertices are the roots of the equation $z^3 + \bar{z}z^3 = 350$ is **[JEE 2009, 3 + 3]**

(A) 48 (B) 32 (C) 40 (D) 80

7. Let z_1 and z_2 be two distinct complex numbers and let $z = (1 - t)z_1 + tz_2$ for some real number t with $0 < t < 1$. If $\text{Arg}(w)$ denotes the principal argument of a nonzero complex number w , then
- (A) $|z - z_1| + |z - z_2| = |z_1 - z_2|$
 (B) $\text{Arg}(z - z_1) = \text{Arg}(z - z_2)$ [JEE 2010, 3M]
 (C) $\begin{vmatrix} z - z_1 & \bar{z} - \bar{z}_1 \\ z_2 - z_1 & \bar{z}_2 - \bar{z}_1 \end{vmatrix} = 0$
 (D) $\text{Arg}(z - z_1) = \text{Arg}(z_2 - z_1)$

8. Let ω be the complex number $\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$. Then the number of distinct complex numbers z satisfying
- $$\begin{vmatrix} z+1 & \omega & \omega^2 \\ \omega & z+\omega^2 & 1 \\ \omega^2 & 1 & z+\omega \end{vmatrix} = 0$$
- is equal to [JEE 2010, 3M]

9. Match the statements in **Column-I** with those in **Column-II**.

[Note : Here z takes values in the complex plane and $\text{Im } z$ and $\text{Re } z$ denote, respectively, the imaginary part and the real part of z .] [JEE 10, 3+3+8]

Column I

Column II

- | | |
|--|--|
| (A) The set of points z satisfying $ z - i z = z + i z $ is contained in or equal to | (p) an ellipse with eccentricity $\frac{4}{5}$ |
| (B) The set of points z satisfying $ z + 4 + z - 4 = 10$ is contained in or equal to | (q) the set of points z satisfying $\text{Im } z = 0$ |
| (C) If $ w = 2$, then the set of points $z = w - \frac{1}{w}$ is contained in or equal to | (t) the set of points z satisfying $ \text{Im } z \leq 1$ |
| (D) If $ w = 1$, then the set of points $z = w + \frac{1}{w}$ is contained in or equal to | (s) the set of points z satisfying $ \text{Re } z \leq 2$ |
| | (r) the set of points z satisfying $ z \leq 3$ |

10. **Comprehension (3 questions together)**

Let a, b and c be three real numbers satisfying $[a \ b \ c] \begin{bmatrix} 1 & 9 & 7 \\ 8 & 2 & 7 \\ 7 & 3 & 7 \end{bmatrix} = [0 \ 0 \ 0]$... (E)

- (i) If the point $P(a, b, c)$, with reference to (E), lies on the plane $2x + y + z = 1$, then the value of $7a + b + c$ is
- (A) 0 (B) 12 (C) 7 (D) 6

(ii) Let ω be a solution of $x^3 - 1 = 0$ with $\text{Im}(\omega) > 0$. If $a = 2$ with b and c satisfying (E), then the value of $\frac{3}{\omega^a} + \frac{1}{\omega^b} + \frac{3}{\omega^c}$ is equal to –

- (A) –2 (B) 2 (C) 3 (D) –3

(iii) Let $b = 6$, with a and c satisfying (E). If α and β are the roots of the quadratic equation

$ax^2 + bx + c = 0$, then $\sum_{n=0}^{\infty} \left(\frac{1}{\alpha} + \frac{1}{\beta} \right)^n$ is - **[JEE 2011, 3+3+3]**

- (A) 6 (B) 7 (C) $\frac{6}{7}$ (D) ∞

11. If z is any complex number satisfying $|z - 3 - 2i| \leq 2$, then the minimum value of $|2z - 6 + 5i|$ is **[JEE 2011, 4M]**

12. Let $\omega = e^{i\pi/3}$, and a, b, c, x, y, z be non-zero complex numbers such that

$$a + b + c = x$$

$$a + b\omega + c\omega^2 = y$$

$$a + b\omega^2 + c\omega = z.$$

Then the value of $\frac{|x|^2 + |y|^2 + |z|^2}{|a|^2 + |b|^2 + |c|^2}$ is **[JEE 2011, 4M]**

13. Match the statements given in Column I with the values given in Column II

Column I

Column II

(A) If $\vec{a} = \hat{j} + \sqrt{3}\hat{k}$, $\vec{b} = -\hat{j} + \sqrt{3}\hat{k}$ and $\vec{c} = 2\sqrt{3}\hat{k}$ form a triangle,

(p) $\frac{\pi}{6}$

then the internal angle of the triangle between $\vec{a}$ and $\vec{b}$ is

(B) If $\int_a^b (f(x) - 3x) dx = a^2 - b^2$ then the value of $f\left(\frac{\pi}{6}\right)$ is

(q) $\frac{2\pi}{3}$

(C) The value of $\frac{\pi^2}{\ln 3} \int_{7/6}^{5/6} \sec(\pi x) dx$ is

(r) $\frac{\pi}{3}$

(D) The maximum value of $\left| \text{Arg} \left(\frac{1}{1-z} \right) \right|$ for

(s) π

$|z| = 1, z \neq 1$ is given by

(t) $\frac{\pi}{2}$

[JEE 2011, 2+2+2+2M]

14. Match the statements given in Column I with the intervals/union of intervals given in Column II

Column I

Column II

(A) The set $\left\{ \operatorname{Re} \left(\frac{2iz}{1-z^2} \right) : z \text{ is a complex number, } |z| = 1, z \neq \pm 1 \right\}$

(p) $(-\infty, -1) \cup (1, \infty)$

is

(B) The domain of the function $f(x) = \sin^{-1} \left(\frac{8(3)^{x-2}}{1-3^{2(x-1)}} \right)$ is

(q) $(-\infty, 0) \cup (0, \infty)$

(C) If $f(\theta) = \begin{vmatrix} 1 & \tan \theta & 1 \\ -\tan \theta & 1 & \tan \theta \\ -1 & -\tan \theta & 1 \end{vmatrix}$, then the set

(r) $[2, \infty)$

$\left\{ f(\theta) : 0 \leq \theta < \frac{\pi}{2} \right\}$

(s) $(-\infty, -1] \cup [1, \infty)$

(D) If $f(x) = x^{3/2}(3x - 10)$, $x \geq 0$, then $f(x)$ is increasing in

(t) $(-\infty, 0] \cup [2, \infty)$

[JEE 2011, 2+2+2+2M]

15. Let z be a complex number such that the imaginary part of z is nonzero and $a = z^2 + z + 1$ is real. Then a cannot take the value -

[JEE 2012, 3M, -1M]

(A) -1

(B) $\frac{1}{3}$

(C) $\frac{1}{2}$

(D) $\frac{3}{4}$

16. Let complex numbers α and $\frac{1}{\alpha}$ lie on circles $(x - x_0)^2 + (y - y_0)^2 = r^2$ and $(x - x_0)^2 + (y - y_0)^2 = 4r^2$ respectively. If $z_0 = x_0 + iy_0$ satisfies the equation $2|z_0|^2 = r^2 + 2$, then $|\alpha| =$

[JEE(Advanced) 2013, 2M]

(A) $\frac{1}{\sqrt{2}}$

(B) $\frac{1}{2}$

(C) $\frac{1}{\sqrt{7}}$

(D) $\frac{1}{3}$

17. Let ω be a complex cube root of unity with $\omega \neq 1$ and $P = [p_{ij}]$ be a $n \times n$ matrix with $p_{ij} = \omega^{i+j}$. Then $P^2 \neq 0$, when $n =$

[JEE(Advanced) 2013, 3, (-1)]

(A) 57

(B) 55

(C) 58

(D) 56

18. Let $w = \frac{\sqrt{3}+i}{2}$ and $P = \{w^n : n = 1, 2, 3, \dots\}$. Further $H_1 = \left\{ z \in \mathbb{C} : \operatorname{Re} z > \frac{1}{2} \right\}$ and $H_2 = \left\{ z \in \mathbb{C} : \operatorname{Re} z < \frac{-1}{2} \right\}$, where $\mathbb{C}$ is the set of all complex number. If $z_1 \in P \cap H_1$, $z_2 \in P \cap H_2$

and O represents the origin, then $\angle z_1 O z_2 =$

[JEE-Advanced 2013, 4, (-1)]

(A) $\frac{\pi}{2}$

(B) $\frac{\pi}{6}$

(C) $\frac{2\pi}{3}$

(D) $\frac{5\pi}{6}$

Paragraph for Question 19 and 20

Let $S = S_1 \cap S_2 \cap S_3$, where $S_1 = \{z \in \mathbb{C} : |z| < 4\}$, $S_2 = \left\{z \in \mathbb{C} : \operatorname{Im} \left[\frac{z-1+\sqrt{3}i}{1-\sqrt{3}i} \right] > 0 \right\}$ and $S_3 = \{z \in \mathbb{C} : \operatorname{Re} z > 0\}$.

19. $\min_{z \in S} |1-3i-z| =$ [JEE(Advanced) 2013, 3, (-1)]

(A) $\frac{2-\sqrt{3}}{2}$ (B) $\frac{2+\sqrt{3}}{2}$ (C) $\frac{3-\sqrt{3}}{2}$ (D) $\frac{3+\sqrt{3}}{2}$

20. Area of $S =$ [JEE(Advanced) 2013, 3, (-1)]

(A) $\frac{10\pi}{3}$ (B) $\frac{20\pi}{3}$ (C) $\frac{16\pi}{3}$ (D) $\frac{32\pi}{3}$

21. The quadratic equation $p(x) = 0$ with real coefficients has purely imaginary roots. Then the equation $p(p(x)) = 0$ has [JEE(Advanced) 2014, 3(-1)]

- (A) only purely imaginary roots
(B) all real roots
(C) two real and two purely imaginary roots
(D) neither real nor purely imaginary roots.

22. Let $z_k = \cos\left(\frac{2k\pi}{10}\right) + i\sin\left(\frac{2k\pi}{10}\right)$; $k=1, 2, \dots, 9$. [JEE(Advanced) 2014, 3(-1)]

List-I

- P. For each z_k there exists a z_j such that $z_k \cdot z_j = 1$
Q. There exists a $k \in \{1, 2, \dots, 9\}$ such that $z_1 \cdot z = z_k$ has no solution z in the set of complex numbers.

R. $\frac{|1-z_1| |1-z_2| \dots |1-z_9|}{10}$ equals

S. $1 - \sum_{k=1}^9 \cos\left(\frac{2k\pi}{10}\right)$ equals

List-II

1. True
2. False
3. 1
4. 2

Codes :

	P	Q	R	S
(A)	1	2	4	3
(B)	2	1	3	4
(C)	1	2	3	4
(D)	2	1	4	3

23. **Column-I**
- (A) In $\mathbb{R}^2$, if the magnitude of the projection vector of the vector $\alpha\hat{i} + \beta\hat{j}$ on $\sqrt{3}\hat{i} + \hat{j}$ is $\sqrt{3}$ and if $\alpha = 2 + \sqrt{3}\beta$, then possible value(s) of $|\alpha|$ is (are)
- (B) Let a and b be real numbers such that the function $f(x) = \begin{cases} -3ax^2 - 2, & x < 1 \\ bx + a^2, & x \geq 1 \end{cases}$ is differentiable for all $x \in \mathbb{R}$. Then possible value(s) of a is (are)
- (C) Let $\omega \neq 1$ be a complex cube root of unity. If $(3 - 3\omega + 2\omega^2)^{4n+3} + (2 + 3\omega - 3\omega^2)^{4n+3} + (-3 + 2\omega + 3\omega^2)^{4n+3} = 0$ then possible value(s) of n is (are)
- (D) Let the harmonic mean of two positive real number a and b be 4, If q is a positive real number such that a, 5, q, b is an arithmetic progression, then the value(s) of $|q - a|$ is (are)
- Column-II**
- (P) 1
- (Q) 2
- (R) 3
- (S) 4
- (T) 5
- [JEE 2015, 8(Each 2M, -1M)]
24. For any integer k, let $\alpha_k = \cos\left(\frac{k\pi}{7}\right) + i \sin\left(\frac{k\pi}{7}\right)$, where $i = \sqrt{-1}$. The value of the expression $\frac{\sum_{k=1}^{12} |\alpha_{k+1} - \alpha_k|}{\sum_{k=1}^3 |\alpha_{4k-1} - \alpha_{4k-2}|}$ is
- [JEE 2015, 4M, -0M]
25. Let $z = \frac{-1 + \sqrt{3}i}{2}$, where $i = \sqrt{-1}$, and $r, s \in \{1, 2, 3\}$. Let $P = \begin{bmatrix} (-z)^r & z^{2s} \\ z^{2s} & z^r \end{bmatrix}$ and I be the identity matrix of order 2. Then the total number of ordered pairs (r, s) for which $P^2 = -I$ is
- [JEE(Advanced)-2016, 3(0)]
26. Let $a, b \in \mathbb{R}$ and $a^2 + b^2 \neq 0$. Suppose $S = \left\{ z \in \mathbb{C} : z = \frac{1}{a + ibt}, t \in \mathbb{R}, t \neq 0 \right\}$, here $i = \sqrt{-1}$. If $z = x + iy$ and $z \in S$, then (x, y) lies on
- (A) the circle with radius $\frac{1}{2a}$ and centre $\left(\frac{1}{2a}, 0\right)$ for $a > 0, b \neq 0$
- (B) the circle with radius $-\frac{1}{2a}$ and centre $\left(-\frac{1}{2a}, 0\right)$ for $a < 0, b \neq 0$
- (C) the x-axis for $a \neq 0, b = 0$
- (D) the y-axis for $a = 0, b \neq 0$
- [JEE(Advanced)-2016, 4(-2)]

27. Let a, b, x and y be real numbers such that $a - b = 1$ and $y \neq 0$. If the complex number $z = x + iy$ satisfies $\operatorname{Im}\left(\frac{az+b}{z+1}\right) = y$, then which of the following is(are) possible value(s) of x ?
[JEE(Advanced)-2017, 4(-2)]
- (A) $-1 - \sqrt{1 - y^2}$ (B) $1 + \sqrt{1 + y^2}$ (C) $1 - \sqrt{1 + y^2}$ (D) $-1 + \sqrt{1 - y^2}$
28. For a non-zero complex number z , let $\arg(z)$ denote the principal argument with $-\pi < \arg(z) \leq \pi$. Then, which of the following statement(s) is (are) FALSE?
[JEE(Advanced)-2018, 4(-2)]
- (A) $\arg(-1 - i) = \frac{\pi}{4}$, Where $i = \sqrt{-1}$
- (B) The function $f : \mathbb{R} \rightarrow (-\pi, \pi]$, defined by $f(t) = \arg(-1 + it)$ for all $t \in \mathbb{R}$ is continuous at all points of $\mathbb{R}$, where $i = \sqrt{-1}$
- (C) For any two non-zero complex number z_1 and z_2 , $\operatorname{Arg}\left(\frac{z_1}{z_2}\right) - \arg(z_1) + \arg(z_2)$ is an integer multiple of 2π
- (D) For any three given distinct complex numbers z_1, z_2 and z_3 the locus of the point z satisfying the condition $\arg\left(\frac{(z - z_1)(z_2 - z_3)}{(z - z_3)(z_2 - z_1)}\right) = \pi$ lies on a straight line
29. Let s, t, r be non-zero complex numbers and L be the set of solutions $z = x + iy$ ($x, y \in \mathbb{R}, i = \sqrt{-1}$) of the equation $sz + t\bar{z} + r = 0$, where $\bar{z} = x - iy$. Then, which of the following statement(s) is (are) TRUE?
[JEE(Advanced)-2018, 4(0)]
- (A) If L has exactly one element, then $|s| \neq |t|$
- (B) If $|s| = |t|$, then L has infinitely many elements
- (C) The number of elements in $L \cap \{z : |z - 1 + i| = 5\}$ is at most 2
- (D) If L has more than one element, then L has infinitely many elements
30. Let S be the set of all complex numbers z satisfying $|z - 2 + i| \geq \sqrt{5}$. If the complex number z_0 is such that $\frac{1}{|z_0 - 1|}$ is the maximum of the set $\left\{\frac{1}{|z - 1|} : z \in S\right\}$, then the principal argument of $\frac{4 - z_0 - \bar{z}_0}{z_0 - \bar{z}_0 + 2i}$ is
[JEE(Advanced)-2019]
- (1) $-\frac{\pi}{2}$ (2) $\frac{3\pi}{4}$ (3) $\frac{\pi}{4}$ (4) $\frac{\pi}{2}$
31. Let $\omega \neq 1$ be a cube root of unity. Then the minimum of the set $\{|a + b\omega + c\omega^2|^2 : a, b, c \text{ distinct non-zero integers}\}$ equals ____
[JEE(Advanced)-2019]

32. Let S be the set of all complex numbers z satisfying $|z^2 + z + 1| = 1$. Then which of the following statements is/are TRUE ? **[JEE(Advanced)-2020]**
- (A) $\left|z + \frac{1}{2}\right| \leq \frac{1}{2}$ for all $z \in S$ (B) $|z| \leq 2$ for all $z \in S$
- (C) $\left|z + \frac{1}{2}\right| \geq \frac{1}{2}$ for all $z \in S$ (D) The set S has exactly four elements
33. For a complex number z , let $\text{Re}(z)$ denote the real part of z . Let S be the set of all complex numbers z satisfying $z^4 - |z|^4 = 4iz^2$, where $i = \sqrt{-1}$. Then the minimum possible value of $|z_1 - z_2|^2$, where $z_1, z_2 \in S$ with $\text{Re}(z_1) > 0$ and $\text{Re}(z_2) < 0$, is ____ **[JEE(Advanced)-2020]**



EXERCISE # 5

- Let z_1 & z_2 be non zero complex numbers satisfying the equation, $z_1^2 - 2z_1z_2 + 2z_2^2 = 0$. The geometrical nature of the triangle whose vertices are the origin and the points representing z_1 & z_2 is :
 (A) an isosceles right angled triangle
 (B) a right angled triangle which is not isosceles
 (C) an equilateral triangle
 (D) an isosceles triangle which is not right angled.
- Let P denotes a complex number z on the Argand's plane, and Q denotes a complex number $\sqrt{2|z|^2} \operatorname{cis}\left(\frac{\pi}{4} + \theta\right)$ where $\theta = \operatorname{amp} z$. If 'O' is the origin, then the ΔOPQ is :
 (A) isosceles but not right angled
 (B) right angled but not isosceles
 (C) right isosceles
 (D) equilateral.
- If $z = \frac{\pi}{4}(1+i)^4 \left(\frac{1-\sqrt{\pi}i}{\sqrt{\pi}+i} + \frac{\sqrt{\pi}-i}{1+\sqrt{\pi}i} \right)$, then $\left(\frac{|z|}{\operatorname{amp} z} \right)$ equals
 (A) 1
 (B) π
 (C) 3π
 (D) 4
- z is a complex number such that $z + \frac{1}{z} = 2 \cos 3^\circ$, then the value of $z^{2000} + \frac{1}{z^{2000}} + 1$ is equal to
 (A) 0
 (B) -1
 (C) $\sqrt{3}+1$
 (D) $1-\sqrt{3}$
- The complex number ω satisfying the equation $\omega^3 = 8i$ and lying in the second quadrant on the complex plane is
 (A) $-\sqrt{3}+i$
 (B) $-\frac{\sqrt{3}}{2}+\frac{1}{2}i$
 (C) $-2\sqrt{3}+i$
 (D) $-\sqrt{3}+2i$
- If $z^4 + 1 = \sqrt{3}i$
 (A) z^3 is purely real
 (B) z represents the vertices of a square of side $2^{1/4}$
 (C) z^9 is purely imaginary
 (D) z represents the vertices of a square of side $2^{3/4}$.
- Let z is a complex number satisfying the equation $Z^6 + Z^3 + 1 = 0$. If this equation has a root $re^{i\theta}$ with $90^\circ < \theta < 180^\circ$ then the value of ' θ ' is
 (A) 100°
 (B) 110°
 (C) 160°
 (D) 170°
- If A and B be two complex numbers satisfying $\frac{A}{B} + \frac{B}{A} = 1$. Then the two points represented by A and B and the origin form the vertices of
 (A) an equilateral triangle
 (B) an isosceles triangle which is not equilateral
 (C) an isosceles triangle which is not right angled
 (D) a right angled triangle

9. If $1, \alpha_1, \alpha_2, \dots, \alpha_{2008}$ are $(2009)^{\text{th}}$ roots of unity, then the value of $\sum_{r=1}^{2008} r(\alpha_r + \alpha_{2009-r})$ equals
(A) 2009 (B) 2008 (C) 0 (D) -2009
10. If $x = \frac{1+\sqrt{3}i}{2}$ then the value of the expression, $y = x^4 - x^2 + 6x - 4$, equals
(A) $-1 + 2\sqrt{3}i$ (B) $2 - 2\sqrt{3}i$ (C) $2 + 2\sqrt{3}i$ (D) none
11. (a) If $w(\neq 1)$ is a cube root of unity and $(1+w)^7 = A + Bw$, then A & B are respectively the numbers
(A) 0, 1 (B) 1, 1 (C) 1, 0 (D) $-1, 1$
- (b) If $(w \neq 1)$ is a cube root of unity then $\begin{vmatrix} 1 & 1+i+w^2 & w^2 \\ 1-i & -1 & w^2-1 \\ -i & -i+w-1 & -1 \end{vmatrix} =$
(A) 0 (B) 1 (C) i (D) w
12. If $z_n = \cos \frac{\pi}{(2n+1)(2n+3)} + i \sin \frac{\pi}{(2n+1)(2n+3)}$ then $\lim_{n \rightarrow \infty} (z_1 \cdot z_2 \cdot z_3 \dots z_n) =$
(A) $\cos \frac{\pi}{3} + i \sin \frac{\pi}{3}$ (B) $\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}$ (C) $\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6}$ (D) $\cos \frac{3\pi}{2} + i \sin \frac{3\pi}{2}$
13. The straight line $(1+2i)z + (2i-1)\bar{z} = 10i$ on the complex plane, has intercept on the imaginary axis equal to
(A) 5 (B) $5/2$ (C) $-5/2$ (D) -5
14. If $\cos\theta + i \sin\theta$ is a root of the equation $x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_{n-1}x + a_n = 0$ then the value of $\sum_{r=1}^n a_r \cos r\theta$ equals (where all coefficient are real)
(A) 0 (B) 1 (C) -1 (D) none
15. Intercept made by the circle $z\bar{z} + \bar{\alpha}z + \alpha\bar{z} + r = 0$ on the real axis on complex plane, is
(A) $\sqrt{(\alpha + \bar{\alpha}) - r}$ (B) $\sqrt{(\alpha + \bar{\alpha})^2 - 2r}$ (C) $\sqrt{(\alpha + \bar{\alpha})^2 + r}$ (D) $\sqrt{(\alpha + \bar{\alpha})^2 - 4r}$
16. If $Z_r ; r = 1, 2, 3, \dots, 50$ are the roots of the equation $\sum_{r=0}^{50} (Z)^r = 0$, then the value of $\sum_{r=1}^{50} \frac{1}{Z_r - 1}$ is
(A) -85 (B) -25 (C) 25 (D) 75
17. All roots of the equation, $(1+z)^6 + z^6 = 0$:
(A) lie on a unit circle with centre at the origin
(B) lie on a unit circle with centre at $(-1, 0)$
(C) lie on the vertices of a regular polygon with centre at the origin
(D) are collinear

18. It is given that complex numbers z_1 and z_2 satisfy $|z_1| = 2$ and $|z_2| = 3$. If the included angle of their corresponding vectors is 60° then $\left| \frac{z_1 + z_2}{z_1 - z_2} \right|$ can be expressed as $\frac{\sqrt{N}}{7}$ where N is natural number then N equals
(A) 126 (B) 119 (C) 133 (D) 19

One or more than one is/are correct :

19. If z is a complex number which simultaneously satisfies the equations $3|z - 12| = 5|z - 8i|$ and $|z - 4| = |z - 8|$ then the $\text{Im}(z)$ can be
(A) 15 (B) 16 (C) 17 (D) 8
20. Let z_1, z_2, z_3 be non-zero complex numbers satisfying the equation $z^4 = iz$. Which of the following statement(s) is/are correct?
(A) The complex number having least positive argument is $\left(\frac{\sqrt{3}}{2}, \frac{1}{2} \right)$.
(B) $\sum_{k=1}^3 \text{Amp}(z_k) = \frac{\pi}{2}$
(C) Centroid of the triangle formed by z_1, z_2 and z_3 is $\left(\frac{1}{\sqrt{3}}, \frac{-1}{3} \right)$
(D) Area of triangle formed by z_1, z_2 and z_3 is $\frac{3\sqrt{3}}{2}$
21. If $z \in \mathbb{C}$, which of the following relation(s) represents a circle on an Argand diagram?
(A) $|z - 1| + |z + 1| = 3$ (B) $(z - 3 + i)(\bar{z} - 3 - i) = 5$
(C) $3|z - 2 + i| = 7$ (D) $|z - 3| = 2$
22. The loci of a point $P(z)$ in the complex plane satisfying the $\left| z + \frac{1}{z} \right| = 2$ are two circles C_1 and C_2 . These circles
(A) have centres on real axis. (B) cut each other orthogonally.
(C) are congruent (D) have exactly two common tangents.
23. Let A and B be two distinct points denoting the complex numbers α and β respectively. A complex number z lies between A and B where $z \neq \alpha, z \neq \beta$. Which of the following relation(s) hold good?
(A) $|\alpha - z| + |z - \beta| = |\alpha - \beta|$
(B) $\exists$ a positive real number 't' such that $z = (1 - t)\alpha + t\beta$
(C) $\begin{vmatrix} z - \alpha & \bar{z} - \bar{\alpha} \\ \beta - \alpha & \bar{\beta} - \bar{\alpha} \end{vmatrix} = 0$ (D) $\begin{vmatrix} z & \bar{z} & 1 \\ \alpha & \bar{\alpha} & 1 \\ \beta & \bar{\beta} & 1 \end{vmatrix} = 0$

24. Let z_1, z_2, z_3 are the coordinates of the vertices of the triangle $A_1A_2A_3$. Which of the following statements are equivalent.
 (A) $A_1A_2A_3$ is an equilateral triangle.
 (B) $(z_1 + \omega z_2 + \omega^2 z_3)(z_1 + \omega^2 z_2 + \omega z_3) = 0$, where ω is the cube root of unity.
 (C) $\frac{z_2 - z_1}{z_3 - z_2} = \frac{z_3 - z_2}{z_1 - z_3}$ (D) $\begin{vmatrix} 1 & 1 & 1 \\ z_1 & z_2 & z_3 \\ \bar{z}_2 & \bar{z}_3 & \bar{z}_1 \end{vmatrix} = 0$
25. Let tangents at $A(z_1)$ and $B(z_2)$ are drawn to the circle $|z| = 2$. Then which of the following is/are **CORRECT** ?
 (A) The equation of tangent at A is given by $\frac{z}{z_1} + \frac{\bar{z}}{\bar{z}_1} = 2$.
 (B) If tangents at $A(z_1)$ and $B(z_2)$ intersect at $P(z_p)$, then $z_p = \frac{2z_1z_2}{z_1 + z_2}$.
 (C) Slope of tangent at $A(z_1)$ is $\frac{1}{i} \left(\frac{z_1 + \bar{z}_1}{z_1 - \bar{z}_1} \right)$
 (D) If points $A(z_1)$ and $B(z_2)$ on the circle $|z| = 2$ are such that $z_1 + z_2 = 0$, then tangents intersect at $\frac{\pi}{2}$.
26. Equation of a straight line on the complex plane passing through a point P denoting the complex number α and perpendicular to the vector $\overrightarrow{OP}$ where 'O' is the origin can be written as
 (A) $\text{Im} \left(\frac{z - \alpha}{\alpha} \right) = 0$ (B) $\text{Re} \left(\frac{z - \alpha}{\alpha} \right) = 0$ (C) $\text{Re}(\bar{\alpha} z) = 0$ (D) $\bar{\alpha} z + \alpha \bar{z} - 2|\alpha|^2 = 0$
27. If $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_{n-1}$ are the imaginary n^{th} roots of unity then the product $\prod_{r=1}^{n-1} (i - \alpha_r)$ (where $i = \sqrt{-1}$) can take the value equal to
 (A) 0 (B) 1 (C) i (D) $(1 + i)$
28. If the expression $(1 + ir)^3$ is of the form of $s(1 + i)$ for some real 's' where 'r' is also real and $i = \sqrt{-1}$, then the value of 'r' can be
 (A) $\cot \frac{\pi}{8}$ (B) $\sec \pi$ (C) $\tan \frac{\pi}{12}$ (D) $\tan \frac{5\pi}{12}$
29. Let point z moves on $|z - 1| = 1$ such that minimum & maximum value of $|z - 2\sqrt{6}i|$ are m & M respectively, then-
 (A) $m + M = 10$ (B) $m^2 + M^2 = 52$ (C) $m + M = 8$ (D) $m^2 + M^2 = 16$
30. Locus of all points z in argand plane which satisfy $|z^2 + 1| = |z^2 - 1|$ is -
 (A) same as $\text{Re}(z) = 0$ (B) same as $\text{Re}(z^2) = 0$
 (C) a pair of straight lines (D) a circle with unit radius

EXERCISE # 6

- Find the sum of the series $1(2 - \omega)(2 - \omega^2) + 2(3 - \omega)(3 - \omega^2) + \dots + (n-1)(n - \omega)(n - \omega^2)$ where ω is one of the imaginary cube root of unity.
- Resolve $Z^5 + 1$ into linear & quadratic factors with real coefficients. Deduce that:
 $4 \sin \frac{\pi}{10} \cdot \cos \frac{\pi}{5} = 1$.
- Prove that, with regard to the quadratic equation $z^2 + (p + ip')z + q + iq' = 0$ where p, p', q, q' are all real.
 - if the equation has one real root then $q'^2 - pp'q' + qp'^2 = 0$.
 - if the equation has two equal roots then $p^2 - p'^2 = 4q$ & $pp' = 2q'$.
 State whether these equal roots are real or complex.
- Let $z = x + iy$ be a complex number, where x and y are real numbers. Let A and B be the sets defined by
 $A = \{z : |z| \leq 2\}$ and $B = \{z : (1 - i)z + (1 + i)\bar{z} \geq 4\}$. Find the area of the region $A \cap B$.
 - If Z is a non-real complex number, then find the minimum value of $\frac{\text{Im } Z^5}{\text{Im } Z}$.
- Interpret the following locii in $z \in \mathbb{C}$.
 - $1 < |z - 2i| < 3$
 - $\text{Re}\left(\frac{z + 2i}{iz + 2}\right) \leq 4 \quad (z \neq 2i)$
 - $\text{Arg}(z + i) - \text{Arg}(z - i) = \pi/2$
 - $\text{Arg}(z - a) = \pi/3$ where $a = 3 + 4i$.
- If the equation $(z + 1)^7 + z^7 = 0$ has roots $z_1, z_2, \dots, z_7$, find the value of
 - $\sum_{r=1}^7 \text{Re}(Z_r)$
 - $\sum_{r=1}^7 \text{Im}(Z_r)$
- If the expression $z^5 - 32$ can be factorised into linear and quadratic factors over real coefficients as $(z^5 - 32) = (z - 2)(z^2 - pz + 4)(z^2 - qz + 4)$ then find the value of $(p^2 + 2p)$.
- Let $z_i (i = 1, 2, 3, 4)$ represent the vertices of a square all of which lie on the sides of the triangle with vertices $(0, 0)$, $(2, 1)$ and $(3, 0)$. If z_1 and z_2 are purely real, then area of triangle formed by z_3, z_4 and origin is $\frac{m}{n}$ (where m and n are in their lowest form). Find the value of $(m + n)$.
- Let $f(x) = ax^3 + bx^2 + cx + d$ be a cubic polynomial with real coefficients satisfying $f(i) = 0$ and $f(1 + i) = 5$. Find the value of $a^2 + b^2 + c^2 + d^2$. (where $i = \sqrt{-1}$)

10. A particle starts to travel from a point P on the curve $C_1 : |z - 3 - 4i| = 5$, where $|z|$ is maximum. From P, the particle moves through an angle $\tan^{-1} \frac{3}{4}$ in anticlockwise direction on $|z - 3 - 4i| = 5$ and reaches at point Q. From Q, it comes down parallel to imaginary axis by 2 units and reaches at point R. Find the complex number corresponding to point R in the Argand plane.
11. Evaluate : $\sum_{p=1}^{32} (3p+2) \left(\sum_{q=1}^{10} \left(\sin \frac{2q\pi}{11} - i \cos \frac{2q\pi}{11} \right) \right)^p$.
12. If the biquadratic $x^4 + ax^3 + bx^2 + cx + d = 0$ ($a, b, c, d \in \mathbb{R}$) has 4 non real roots, two with sum $3 + 4i$ and the other two with product $13 + i$. Find the value of 'b'.



ANSWER KEY

EXERCISE # 1

PART-A

1. (a) $\frac{7}{25} + \frac{24}{25}i$; (b) $\frac{21}{5} - \frac{12}{5}i$; (c) $3 + 4i$ (d) $\frac{22}{5}i$; (e) 15
2. (a) $x = 1, y = 2$; (b) (2,9); (c) $(-2, 2)$ or $\left(-\frac{2}{3}, -\frac{2}{3}\right)$
3. (a) $\pm(5 + 4i)$; (b) $\pm(5 - 6i)$; (c) $\pm 5(1 + i)$
4. (a) -160; (b) $-(77 + 108i)$ 5. (a) $-i, -2i$ (b) $\frac{3-5i}{2}$ or $-\frac{1+i}{2}$
6. (a) on a circle of radius $\sqrt{7}$ with centre $(-1, 2)$; (b) on a unit circle with centre at origin
(c) on a circle with centre $(-15/4, 0)$ & radius $9/4$; (d) a straight line.
7. $a = b = 2 - \sqrt{3}$ 8. $z_3 = \sqrt{3}(1-i)$ and $z'_3 = \sqrt{3}(-1+i)$
9. $x = 1, y = -4$ or $x = -1, y = -4$ 12. $\frac{x^2}{64} + \frac{y^2}{48} = 1$ 13. B 14. C 15. C

PART-B

- | | | | | | | |
|-------|-------|-------|--|-------|-------------------|-------|
| 1. B | 2. C | 3. B | 4. C | 5. A | 6. A | 7. B |
| 8. D | 9. B | 10. B | 11. C | 12. B | 13. D | 14. C |
| 15. C | 16. C | 17. A | 18. A | 19. B | 20. A | 21. D |
| 22. B | 23. D | 24. B | 25. A | 26. A | 27. C | 28. C |
| 29. C | 30. C | 31. A | 32. D | 33. B | 34. A | 35. B |
| 36. C | 37. A | 38. B | 39. C | 40. B | 41. (i) D; (ii) C | |
| 42. B | 43. B | 44. A | 45. D | 46. D | 47. A | 48. D |
| 49. B | 50. A | 51. B | 52. (A) Q,R; (B) P,S; (C) Q,S; (D) P,R | | | |

EXERCISE # 2

1. (i) Modulus = 6, Arg = $2k\pi + \frac{5\pi}{18}$ ($k \in I$), Principal Arg = $\frac{5\pi}{18}$
 (ii) Modulus = 2, Arg = $2k\pi + \frac{7\pi}{6}$ ($k \in I$), Principal Arg = $-\frac{5\pi}{6}$
 (iii) Modulus = $\frac{\sqrt{5}}{6}$, Arg = $2k\pi - \tan^{-1}2$ ($k \in I$), Principal Arg = $-\tan^{-1}2$
2. (a) 2, (b) $-11/2$ 3. $[(-2, 2); (-2, -2)]$ 4. (a) $z = (2 + i)$ or $(1 - 3i)$; (b) $z = 1$ or $2 - i$

5. (a) 1, (b) 2 (c) 30 8. $\begin{cases} (1,0) & \text{for } n = 4k \\ (1,1) & \text{for } n = 4k+1 \\ (0,1) & \text{for } n = 4k+2 \\ (0,0) & \text{for } n = 4k+3 \end{cases}$ 9. $-\frac{3}{2} + \frac{3\sqrt{3}}{2}i$ 10. -4
11. $\frac{iz}{2} + \frac{1}{2} + i$ 12. (a) $\frac{\sqrt{3}}{2} - \frac{i}{2}, -\frac{\sqrt{3}}{2} - \frac{i}{2}, i$ 13. 673 14. 6

EXERCISE # 3 (JM)

1. 3 2. 3 3. 4 4. 2 5. 3 6. 2 7. 4
8. 2 9. 3 10. 2 11. 1 12. 1 13. 2 14. 4
15. 3 16. 2 17. 4 18. Bonus 19. 4 20. 3 21. 2
22. 2 23. 1 24. 1 25. 1 26. 1 27. 3 28. 4
29. 3 30. 4 31. 1 32. 3 33. 2 34. 3 35. 1
36. 4 37. 4 38. 2 39. 4.00 40. 4 41. 4 42. 2
43. 3 44. 1 45. 4 46. 3

EXERCISE # 4 (JA)


1. D 2. D 3. D 4. (i) B; (ii) C; (iii) B or C or D 5. D
6. A 7. A,C,D 8. 1 9. $(A) \rightarrow (q,r), (B) \rightarrow (p), (C) \rightarrow (p,s,t), (D) \rightarrow (q,r,s,t)$
10. (i) D, (ii) A, (iii) B 11. 5 12. Bonus
13. $(A) \rightarrow (q); (B) \rightarrow (p) \text{ or } (p, q, r, s, t); (C) \rightarrow (s); (D) \rightarrow (t)$
14. $(A) \rightarrow (s); (B) \rightarrow (t); (C) \rightarrow (r); (D) \rightarrow (r)$ 15. D 16. C 17. B,C,D
18. C,D 19. C 20. B 21. D 22. C
23. $(A) \rightarrow (P,Q); (B) \rightarrow (P,Q); (C) \rightarrow (P,Q,S,T); (D) \rightarrow (Q,T)$ 24. 4 25. 1
26. A,C,D 27. A,D 28. A,B,D 29. A,C,D 30. (1)
31. (3.00) 32. B,C 33. (8)

EXERCISE # 5

1. A 2. C 3. D 4. A 5. A 6. D 7. C
8. A 9. D 10. A 11. (a) B; (b) A 12. B 13. A
14. C 15. D 16. B 17. D 18. C 19. C,D 20. A,B
21. B,C,D 22. B,C,D 23. A,B,C,D 24. A,B,C,D
25. A,B,C 26. B,D 27. A,B,C,D 28. B,C,D 29. A,B
30. B,C

EXERCISE # 6

1. $\left[\frac{n(n+1)}{2} \right]^2 - n$
2. $(Z + 1)(Z^2 - 2Z \cos 36^\circ + 1)(Z^2 - 2Z \cos 108^\circ + 1)$
4. (a) $\pi - 2$, (b) -4
5. (a) The region between the concentric circles with centre at $(0, 2)$ & radii 1 & 3 units
 (b) region outside or on the circle with centre $\frac{1}{2} + 2i$ and radius $\frac{1}{2}$.
 (c) semicircle (in the 1st & 4th quadrant) $x^2 + y^2 = 1$
 (d) a ray emanating from the point $(3 + 4i)$ directed away from the origin & having equation $\sqrt{3}x - y + 4 + 3\sqrt{3} = 0$
6. (a) $-\frac{7}{2}$, (b) zero
7. 4
8. 41
9. 26
10. $(3 + 7i)$
11. $48(1 - i)$
12. 51




ADDITIONAL EXERCISE ON COMPLEX NUMBER

[STRAIGHT OBJECTIVE TYPE]

- If z and w are complex number satisfying $\bar{z} + i\bar{w} = 0$ and $\text{Amp}(zw) = \pi$, then $\text{Amp}(z)$ is equal to
 (A) $\frac{\pi}{4}$ (B) $\frac{-\pi}{2}$ (C) $\frac{\pi}{2}$ (D) $\frac{3\pi}{4}$
- If $z_1 = -3 + 5i$; $z_2 = -5 - 3i$ and z is a complex number lying on the line segment joining z_1 & z_2 then $\arg z$ can be :
 (A) $-\frac{3\pi}{4}$ (B) $-\frac{\pi}{4}$ (C) $\frac{\pi}{6}$ (D) $\frac{5\pi}{6}$
- Given $z = f(x) + i g(x)$ where $f, g : (0, 1) \rightarrow (0, 1)$ are real valued functions then, which of the following holds good?
 (A) $z = \frac{1}{1-ix} + i\left(\frac{1}{1+ix}\right)$ (B) $z = \frac{1}{1+ix} + i\left(\frac{1}{1-ix}\right)$
 (C) $z = \frac{1}{1+ix} + i\left(\frac{1}{1+ix}\right)$ (D) $z = \frac{1}{1-ix} + i\left(\frac{1}{1-ix}\right)$
- Let z be a complex number having the argument θ , $0 < \theta < \pi/2$ and satisfying the equality $|z - 3i| = 3$. Then $\cot \theta - \frac{6}{z}$ is equal to :
 (A) 1 (B) -1 (C) i (D) $-i$
- The points $z_1 = 3 + \sqrt{3}i$ and $z_2 = 2\sqrt{3} + 6i$ are given on a complex plane. The complex number lying on the bisector of the angle formed by the vectors z_1 and z_2 is :
 (A) $z = \frac{(3+2\sqrt{3})}{2} + \frac{\sqrt{3}+2}{2}i$ (B) $z = 5 + 5i$
 (C) $z = -1 - i$ (D) none
- z_1 & z_2 are two distinct points in an argand plane. If $a|z_1| = b|z_2|$, (where $a, b \in \mathbb{R}$) then the point $\frac{az_1}{bz_2} + \frac{bz_2}{az_1}$ is a point on the :
 (A) line segment $[-2, 2]$ of the real axis
 (B) line segment $[-2, 2]$ of the imaginary axis
 (C) unit circle $|z| = 1$
 (D) the line with $\arg z = \tan^{-1} 2$.


7. A root of unity is a complex number that is a solution to the equation, $z^n = 1$ for some positive integer n . Number of roots of unity that are also the roots of the equation $z^2 + az + b = 0$, for some integer a and b is
(A) 6 (B) 8 (C) 9 (D) 10
8. If m and n are the smallest positive integers satisfying the relation $\left(2\text{Cis}\frac{\pi}{6}\right)^m = \left(4\text{Cis}\frac{\pi}{4}\right)^n$, then $(m + n)$ has the value equal to
(A) 120 (B) 96 (C) 72 (D) 60
9. Least positive argument of the 4th root of the complex number $2-i\sqrt{12}$ is
(A) $\pi/6$ (B) $5\pi/12$ (C) $7\pi/12$ (D) $11\pi/12$
10. Number of ordered pair(s) (z, ω) of the complex numbers z and ω satisfying the system of equations, $z^3 + \bar{\omega}^7 = 0$ and $z^5 \cdot \omega^{11} = 1$ is :
(A) 7 (B) 5 (C) 3 (D) 2
11. On the complex plane triangles OAP & OQR are similar and $l(OA) = 1$. If the points P and Q denotes the complex numbers z_1 & z_2 then the complex number 'z' denoted by the point R is given by :



- (A) $z_1 z_2$ (B) $\frac{z_1}{z_2}$ (C) $\frac{z_2}{z_1}$ (D) $\frac{z_1 + z_2}{z_2}$
12. Number of imaginary complex numbers satisfying the equation, $z^2 = \bar{z}2^{1-|z|}$ is
(A) 0 (B) 1 (C) 2 (D) 3
13. Let $A(z_1)$ and $B(z_2)$ represent two complex numbers on the complex plane. Suppose the complex slope of the line joining A and B is defined as $\frac{z_1 - z_2}{\bar{z}_1 - \bar{z}_2}$. Then the lines l_1 with complex slope ω_1 and l_2 with complex slope ω_2 on the complex plane will be perpendicular to each other if
(A) $\omega_1 + \omega_2 = 0$ (B) $\omega_1 - \omega_2 = 0$ (C) $\omega_1 \omega_2 = -1$ (D) $\omega_1 \omega_2 = 1$


14. If the equation, $z^4 + a_1z^3 + a_2z^2 + a_3z + a_4 = 0$, where a_1, a_2, a_3, a_4 are real coefficients different from zero has a pure imaginary root then the expression $\frac{a_3}{a_1a_2} + \frac{a_1a_4}{a_2a_3}$ has the value equal to:
(A) 0 (B) 1 (C) -2 (D) 2
15. Suppose A is a complex number & $n \in \mathbb{N}$, such that $A^n = (A + 1)^n = 1$, then the least value of n is
(A) 3 (B) 6 (C) 9 (D) 12
16. If z & w are two complex numbers simultaneously satisfying the equations, $z^3 + w^5 = 0$ and $z^2 \cdot \bar{w}^4 = 1$, then :
(A) z and w both are purely real
(B) z is purely real and w is purely imaginary
(C) w is purely real and z is purely imaginary
(D) z and w both are imaginary.
17. A function f is defined by $f(z) = (4 + i)z^2 + \alpha z + \gamma$ for all complex numbers z , where α and γ are complex numbers. If $f(1)$ and $f(i)$ are both real then the smallest possible value of $|\alpha| + |\gamma|$ equals
(A) 1 (B) $\sqrt{2}$ (C) 2 (D) $2\sqrt{2}$
18. Given $f(z)$ = the real part of a complex number z . For example, $f(3 - 4i) = 3$. If $a \in \mathbb{N}$, $n \in \mathbb{N}$ then the value of $\sum_{n=1}^{6a} \log_2 \left| f \left((1 + i\sqrt{3})^n \right) \right|$ has the value equal to
(A) $18a^2 + 9a$ (B) $18a^2 + 7a$ (C) $18a^2 - 3a$ (D) $18a^2 - a$
19. All complex numbers ' z ' which satisfy the relation $|z - |z + 1|| = |z + |z - 1||$ on the complex plane lie on the
(A) line $y = 0$ or an ellipse with foci $(-1, 0)$ and $(1, 0)$
(B) radical axis of the circles $|z - 1| = 1$ and $|z + 1| = 1$
(C) circle $x^2 + y^2 = 1$
(D) line $x = 0$ or on a line segment joining $(-1, 0)$ to $(1, 0)$
20. In the quadratic equation $x^2 + (p + iq)x + 3i = 0$, p & q are real. If the sum of the squares of the roots is 8 then
(A) $p = 3, q = -1$ (B) $p = -3, q = -1$ (C) $p = \pm 3, q = \pm 1$ (D) $p = -3, q = 1$
21. If $z = (3 + 7i)(p + iq)$ where $p, q \in \mathbb{I} - \{0\}$, is purely imaginary then minimum value of $|z|^2$ is
(A) 0 (B) 58 (C) $\frac{3364}{3}$ (D) 3364

22. If $x = 9^{1/3} 9^{1/9} 9^{1/27} \dots \infty$, $y = 4^{1/3} 4^{-1/9} 4^{1/27} \dots \infty$ and $z = \sum_{r=1}^{\infty} (1+i)^{-r}$ then, the argument of the complex number $w = x + yz$ is
 (A) 0 (B) $\pi - \tan^{-1}\left(\frac{\sqrt{2}}{3}\right)$ (C) $-\tan^{-1}\left(\frac{\sqrt{2}}{3}\right)$ (D) $-\tan^{-1}\left(\frac{2}{\sqrt{3}}\right)$
23. Which of the following represents a point on an argand's plane, equidistant from the roots of the equation $(z+1)^4 = 16z^4$?
 (A) (0, 0) (B) $\left(-\frac{1}{3}, 0\right)$ (C) $\left(\frac{1}{3}, 0\right)$ (D) $\left(0, \frac{2}{\sqrt{5}}\right)$
24. The value of $e(\text{CiS}(-i) - \text{CiS}(i))$ is equal to
 (A) 0 (B) $1 - e$ (C) $e - \frac{1}{e}$ (D) $e^2 - 1$
25. If $z_1 = 2 + 3i$, $z_2 = 3 - 2i$ and $z_3 = -1 - 2\sqrt{3}i$ then which of the following is true?
 (A) $\arg\left(\frac{z_3}{z_2}\right) = \arg\left(\frac{z_3 - z_1}{z_2 - z_1}\right)$ (B) $\arg\left(\frac{z_3}{z_2}\right) = \arg\left(\frac{z_2}{z_1}\right)$
 (C) $\arg\left(\frac{z_3}{z_2}\right) = 2\arg\left(\frac{z_3 - z_1}{z_2 - z_1}\right)$ (D) $\arg\left(\frac{z_3}{z_2}\right) = \frac{1}{2}\arg\left(\frac{z_3 - z_1}{z_2 - z_1}\right)$
26. On the Argand plane point 'A' denotes a complex number z_1 . A triangle OBQ is made directly similar to the triangle OAM, where $OM = 1$ and O is origin (as shown in the figure). If the point B denotes the complex number z_2 , then the complex number corresponding to the point 'Q' is



- (A) $z_1 z_2$ (B) $\frac{z_1}{z_2}$ (C) $\frac{z_2}{z_1}$ (D) $\frac{z_1 + z_2}{z_2}$
27. Number of ordered pairs(s) (a, b) of real numbers such that $(a + ib)^{2008} = a - ib$ holds good, is
 (A) 2008 (B) 2009 (C) 2010 (D) 1

28. If $z^3 + (3 + 2i)z + (-1 + ia) = 0$ has one real root, then the value of 'a' lies in the interval ($a \in \mathbb{R}$)
(A) $(-2, -1)$ (B) $(-1, 0)$ (C) $(0, 1)$ (D) $(1, 2)$
29. $P(z)$ is the point moving in the Argand's plane satisfying $\arg(z - 1) - \arg(z + i) = \pi$ then, P is
(A) a real number, hence lies on the real axis.
(B) an imaginary number, hence lies on the imaginary axis.
(C) a point on the hypotenuse of the right angled triangle OAB formed by $O \equiv (0, 0)$;
 $A \equiv (1, 0)$; $B \equiv (0, -1)$.
(D) a point on an arc of the circle passing through $A \equiv (1, 0)$; $B \equiv (0, -1)$.
30. If z_1 & z_2 are two complex numbers & if $\arg\left(\frac{z_1 + z_2}{z_1 - z_2}\right) = \frac{\pi}{2}$ but $|z_1 + z_3| \neq |z_1 - z_2|$, then the figure formed by the points represented by 0, z_1 , z_2 & $z_1 + z_2$ is
(A) a parallelogram but not a rectangle or a rhombus
(B) a rectangle but not a square
(C) a rhombus but not a square
(D) a square
31. If the six solutions of $x^6 = -64$ are written in the form $a + bi$, where a and b are real, then the product of those solutions with $a > 0$, is
(A) 4 (B) 8 (C) 16 (D) 64
32. Let z, w be complex numbers such that $\bar{z} + i\bar{w} = 0$ and $\arg zw = \pi$. Then $\arg z$ equals
[AIEEE-2004]
(A) $\pi/4$ (B) $\pi/2$ (C) $3\pi/4$ (D) $5\pi/4$
33. If $|z^2 - 1| = |z|^2 + 1$, then z lies on
[AIEEE-2004]
(A) the real axis (B) the imaginary axis
(C) a circle (D) an ellipse
34. If $z = x - iy$ and $z^{1/3} = p + iq$, then $\frac{\left(\frac{x}{p} + \frac{y}{q}\right)}{(p^2 + q^2)}$ is equal to-
[AIEEE-2004]
(A) 1 (B) -1 (C) 2 (D) -2
35. If z_1 and z_2 are two non zero complex numbers such that $|z_1 + z_2| = |z_1| + |z_2|$ then $\arg z_1 - \arg z_2$ is equal to-
[AIEEE-2005]
(A) $-\pi$ (B) $\frac{\pi}{2}$ (C) $-\frac{\pi}{2}$ (D) 0

36. If $w = \frac{z}{z - \frac{1}{3}i}$ and $|w| = 1$ then z lies on [AIEEE-2005]
 (A) a circle (B) an ellipse (C) a parabola (D) a straight line
37. ω is an imaginary cube root of unity. If $(1 + \omega^2)^m = (1 + \omega^4)^m$, then least positive integral value of m is [JEE 2004 (Scr)]
 (A) 6 (B) 5 (C) 4 (D) 3
38. The locus of z which lies in shaded region is best represented by
- 
- (A) $z : |z + 1| > 2, |\arg(z + 1)| < \pi/4$ (B) $z : |z - 1| > 2, |\arg(z - 1)| < \pi/4$
 (C) $z : |z + 1| < 2, |\arg(z + 1)| < \pi/2$ (D) $z : |z - 1| < 2, |\arg(z - 1)| < \pi/2$
39. If a, b, c are integers not all equal and w is a cube root of unity ($w \neq 1$), then the minimum value of $|a + bw + cw^2|$ is [JEE 2005 (Scr), 3 + 3]
 (A) 0 (B) 1 (C) $\frac{\sqrt{3}}{2}$ (D) $\frac{1}{2}$
40. If $w = \alpha + i\beta$ where $\beta \neq 0$ and $z \neq 1$, satisfies the condition that $\frac{w - \bar{w}z}{1 - z}$ is purely real, then the set of values of z is [JEE 2006, 3]
 (A) $\{z : |z| = 1\}$ (B) $\{z : z = \bar{z}\}$ (C) $\{z : z \neq 1\}$ (D) $\{z : |z| = 1, z \neq 1\}$
41. Let $f(x) = x^3 + ax^2 + bx + c$ be a cubic polynomial with real coefficients and all real roots. Also $|f(i)| = 1$ where $i = \sqrt{-1}$
Statement-1: All 3 roots of $f(x) = 0$ are zero
because
Statement-2: $a + b + c = 0$
 (A) Statement-1 is true, statement-2 is true and statement-2 is correct explanation for statement-1.
 (B) Statement-1 is true, statement-2 is true and statement-2 is NOT the correct explanation for statement-1.
 (C) Statement-1 is true, statement-2 is false.
 (D) Statement-1 is false, statement-2 is true.

[COMPREHENSION TYPE]

Paragraph for question number 42 to 44

For the complex number $w = \frac{4z - 5i}{2z + 1}$

42. The locus of z , when w is a real number other than 2, is

- (A) a point circle
- (B) a straight line with slope $-\frac{5}{2}$ and y-intercept $\frac{5}{4}$
- (C) a straight line with slope $\frac{5}{2}$ and y-intercept $\frac{5}{4}$
- (D) a straight line passing through the origin

43. The locus of z , when w is a purely imaginary number is

- (A) a circle with centre $\left(\frac{1}{2}, -\frac{5}{4}\right)$ passing through origin.
- (B) a circle with centre $\left(-\frac{1}{4}, \frac{5}{8}\right)$ passing through origin.
- (C) a circle with centre $\left(\frac{1}{4}, -\frac{5}{8}\right)$ and radius $\frac{\sqrt{29}}{8}$
- (D) any other circle

44. The locus of z , when $|w| = 1$ is

- (A) a circle with centre $\left(-\frac{5}{8}, \frac{1}{4}\right)$ and radius $\frac{1}{2}$
- (B) a circle with centre $\left(\frac{1}{4}, -\frac{5}{8}\right)$ and radius $\frac{1}{2}$
- (C) a circle with centre $\left(\frac{5}{8}, -\frac{1}{4}\right)$ and radius $\frac{1}{2}$
- (D) any other circle

[MATCHING COLUMN TYPE]

- | 45. Column-I | Column-II |
|---|----------------|
| (A) Let ω be a non real cube root of unity then the number of distinct elements in the set $\{(1 + \omega + \omega^2 + \dots + \omega^n)^m \mid m, n \in \mathbb{N}\}$ is | (P) 4 |
| (B) Let $1, \omega, \omega^2$ be the cube root of unity. The least possible degree of a polynomial with real coefficients having roots $2\omega, (2 + 3\omega), (2 + 3\omega^2), (2 - \omega - \omega^2)$, is | (Q) 5 |
| (C) $\alpha = 6 + 4i$ and $\beta = (2 + 4i)$ are two complex numbers on the complex plane. A complex number z satisfying $\arg\left(\frac{z - \alpha}{z - \beta}\right) = \frac{\pi}{6}$ moves on the major segment of a circle whose radius is | (R) 6
(S) 7 |

[SUBJECTIVE TYPE]

46. A polynomial $f(z)$ when divided by $(z - w)$ leaves remainder $2 + i\sqrt{3}$ and when divided by $(z - w^2)$ leaves remainder $2 - i\sqrt{3}$. If the remainder obtained when $f(z)$ is divided by $z^2 + z + 1$ is $az + b$ (where w is non-real cube root of unity and $a, b \in \mathbb{R}^+$), then find the value of $(a + b)$.
47. Let z_1, z_2, z_3 are three pair wise distinct complex numbers and t_1, t_2, t_3 are non-negative real numbers such that $t_1 + t_2 + t_3 = 1$. Prove that the complex number $z = t_1z_1 + t_2z_2 + t_3z_3$ lies inside a triangle with vertices z_1, z_2, z_3 or on its boundary.
48. Let $A \equiv z_1$; $B \equiv z_2$; $C \equiv z_3$ are three complex numbers denoting the vertices of an acute angled triangle. If the origin 'O' is the orthocentre of the triangle, then prove that
- $$z_1\bar{z}_2 + \bar{z}_1z_2 = z_2\bar{z}_3 + \bar{z}_2z_3 = z_3\bar{z}_1 + \bar{z}_3z_1$$
- hence show that the ΔABC is a right angled triangle
- $$\Leftrightarrow z_1\bar{z}_2 + \bar{z}_1z_2 = z_2\bar{z}_3 + \bar{z}_2z_3 = z_3\bar{z}_1 + \bar{z}_3z_1 = 0$$
49. Let $\alpha + i\beta$; $\alpha, \beta \in \mathbb{R}$, be a root of the equation $x^3 + qx + r = 0$; $q, r \in \mathbb{R}$. Find a real cubic equation, independent of α & β , whose one root is 2α .

50. Prove that $\left(\frac{1 + \sin \theta + i \cos \theta}{1 + \sin \theta - i \cos \theta}\right)^n = \cos\left(\frac{n\pi}{2} - n\theta\right) + i \sin\left(\frac{n\pi}{2} - n\theta\right)$
- Hence deduce that $\left(1 + \sin \frac{\pi}{5} + i \cos \frac{\pi}{5}\right)^5 + i \left(1 + \sin \frac{\pi}{5} - i \cos \frac{\pi}{5}\right)^5 = 0$

51. If $\cos(\alpha - \beta) + \cos(\beta - \gamma) + \cos(\gamma - \alpha) = -3/2$ then prove that :
- $\Sigma \cos 2\alpha = 0 = \Sigma \sin 2\alpha$
 - $\Sigma \sin(\alpha + \beta) = 0 = \Sigma \cos(\alpha + \beta)$
 - $\Sigma \sin^2 \alpha = \Sigma \cos^2 \alpha = 3/2$
 - $\Sigma \sin 3\alpha = 3 \sin(\alpha + \beta + \gamma)$
 - $\Sigma \cos 3\alpha = 3 \cos(\alpha + \beta + \gamma)$
 - $\cos^3(\theta + \alpha) + \cos^3(\theta + \beta) + \cos^3(\theta + \gamma) = 3 \cos(\theta + \alpha) \cdot \cos(\theta + \beta) \cdot \cos(\theta + \gamma)$ where $\theta \in \mathbb{R}$.
52. If $x = 1 + i\sqrt{3}$; $y = 1 - i\sqrt{3}$ & $z = 2$, then prove that $x^p + y^p = z^p$ for every prime $p > 3$.
53. Simplify and express the result in the form of $a + bi$:
- $\left(\frac{4i^3 - i}{2i + 1}\right)^2$
 - $\frac{3 + 2i}{2 - 5i} + \frac{3 - 2i}{2 + 5i}$
54. Find the modulus, argument and the principal argument of the complex numbers.
- $z = 1 + \cos\left(\frac{10\pi}{9}\right) + i \sin\left(\frac{10\pi}{9}\right)$
 - $(\tan 1 - i)^2$
 - $z = \frac{\sqrt{5+12i} + \sqrt{5-12i}}{\sqrt{5+12i} - \sqrt{5-12i}}$
 - $\frac{i - 1}{i\left(1 - \cos\frac{2\pi}{5}\right) + \sin\frac{2\pi}{5}}$
55. Given that $x, y \in \mathbb{R}$, solve :
- $\frac{x}{1+2i} + \frac{y}{3+2i} = \frac{5+6i}{8i-1}$
 - $(2+3i)x^2 - (3-2i)y = 2x - 3y + 5i$
56. Let $Z = 18 + 26i$ where $Z_0 = x_0 + iy_0$ ($x_0, y_0 \in \mathbb{R}$) is the cube root of Z having least positive argument. Find the value of $x_0 y_0 (x_0 + y_0)$.
57. Show that the locus formed by z in the equation $z^3 + iz = 1$ never crosses the co-ordinate axes in the Argand's plane. Further show that $|z| = \sqrt{\frac{-\operatorname{Im}(z)}{2 \operatorname{Re}(z) \operatorname{Im}(z) + 1}}$
58. Let $z_1, z_2 \in \mathbb{C}$ such that $z_1^2 + z_2^2 \in \mathbb{R}$. If $z_1(z_1^2 - 3z_2^2) = 10$ and $z_2(3z_1^2 - z_2^2) = 30$. Find the value of $(z_1^2 + z_2^2)$.

59. Let z_1 & z_2 be any two arbitrary complex numbers then prove that :

$$|z_1 + z_2| \geq \frac{1}{2} (|z_1| + |z_2|) \left| \frac{z_1}{|z_1|} + \frac{z_2}{|z_2|} \right|.$$

60. (i) Let C_r 's denotes the combinatorial coefficients in the expansion of $(1 + x)^n$, $n \in \mathbb{N}$. If the integers

$$a_n = C_0 + C_3 + C_6 + C_9 + \dots$$

$$b_n = C_1 + C_4 + C_7 + C_{10} + \dots$$

and $c_n = C_2 + C_5 + C_8 + C_{11} + \dots$, then

prove that (a) $a_n^3 + b_n^3 + c_n^3 - 3a_nb_nc_n = 2^n$, (b) $(a_n - b_n)^2 + (b_n - c_n)^2 + (c_n - a_n)^2 = 2$.

(ii) Prove the identity : $(C_0 - C_2 + C_4 - C_6 + \dots)^2 + (C_1 - C_3 + C_5 - C_7 + \dots)^2 = 2^n$

61. A function f is defined on the complex number by $f(z) = (a + bi)z$, where 'a' and 'b' are positive numbers. This function has the property that the image of each point in the complex plane is equidistant from that point and the origin. Given that $|a + bi| = 8$ and that $b^2 = \frac{u}{v}$ where u and v are coprimes. Find the value of $(u + v)$.

62. Prove that :

$$(a) \cos x + {}^nC_1 \cos 2x + {}^nC_2 \cos 3x + \dots + {}^nC_n \cos (n+1)x = 2^n \cdot \cos^n \frac{x}{2} \cdot \cos \left(\frac{n+2}{2} x \right)$$

$$(b) \sin x + {}^nC_1 \sin 2x + {}^nC_2 \sin 3x + \dots + {}^nC_n \sin (n+1)x = 2^n \cdot \cos^n x \cdot \sin \left(\frac{n+2}{2} x \right)$$

63. Let $\omega_1, \omega_2, \omega_3, \dots, \omega_n$ be the complex numbers. A line L on the complex plane is called a mean line for the points $\omega_1, \omega_2, \omega_3, \dots, \omega_n$ if L contains the points (complex numbers)

$z_1, z_2, z_3, \dots, z_n$ such that $\sum_{k=1}^n (z_k - \omega_k) = 0$. Now for the complex number $\omega_1 = 32 + 170i$,

$\omega_2 = -7 + 64i$, $\omega_3 = -9 + 200i$, $\omega_4 = 1 + 27i$ and $\omega_5 = -14 + 43i$, there is a unique mean line with y-intercept 3. Find the slope of the line.

64. Let a, b, c be distinct complex numbers such that $\frac{a}{1-b} = \frac{b}{1-c} = \frac{c}{1-a} = k$. Find the value of k.

65. Let α, β be fixed complex numbers and z is a variable complex number such that,

$$|z - \alpha|^2 + |z - \beta|^2 = k.$$

Find out the limits for 'k' such that the locus of z is a circle. Find also the centre and radius of the circle.

66. C is the complex number. $f : C \rightarrow R$ is defined by $f(z) = |z^3 - z + 2|$. Find the maximum value of $f(z)$ if $|z| = 1$.
67. If the area of the polygon whose vertices are the solutions (in the complex plane) of the equation $x^7 + x^6 + x^5 + x^4 + x^3 + x^2 + x + 1 = 0$, can be expressed in the simplest form as $\frac{a\sqrt{b} + c}{d}$, find the value $(a + b + c + d)$.
68. Let z_1, z_2 be complex numbers with $|z_1| = |z_2| = 1$, prove that $|z_1 + 1| + |z_2 + 1| + |z_1 z_2 + 1| \geq 2$.
69. One root of the cubic $2z^3 - (5 + 6i)z^2 + 9iz + 1 - 3i = 0$ is real. If all the three roots of this cubic are plotted on the complex plane, then find the area of the triangle formed by them.
70. If ω is the fifth root of 2 and $x = \omega + \omega^2$, prove that $x^5 = 10x^2 + 10x + 6$.
71. Let z_1, z_2, z_3, z_4 be the vertices A, B, C, D respectively of a square on the Argand diagram taken in anticlockwise direction then prove that :
(i) $2z_2 = (1 + i)z_1 + (1 - i)z_3$ & (ii) $2z_4 = (1 - i)z_1 + (1 + i)z_3$
72. Let $f(x) = \log_{\cos 3x}(\cos 2ix)$ if $x \neq 0$ and $f(0) = K$ (where $i = \sqrt{-1}$) is continuous at $x = 0$ then find the value of K .
73. Let $A = \{a \in R \mid \text{the equation } (1 + 2i)x^3 - 2(3 + i)x^2 + (5 - 4i)x + 2a^2 = 0\}$ has at least one real root. Find the value of $\sum_{a \in A} a^2$.
74. Find centre and radius of the circle determined by all complex numbers $z = x + iy$ satisfying $\left| \frac{(z - \alpha)}{(z - \beta)} \right| = k$, where $\alpha = \alpha_1 + i\alpha_2, \beta = \beta_1 + i\beta_2$ are fixed complex and $k \neq 1$.
- [JEE 2004, 2 out of 60]**
75. If one of the vertices of the square circumscribing the circle $|z - 1| = \sqrt{2}$ is $2 + \sqrt{3}i$. Find the other vertices of square.
- [JEE 2005 (Mains), 4]**

ANSWER KEY

1. D 2. D 3. B 4. C 5. B 6. A 7. B
8. C 9. B 10. D 11. A 12. C 13. A 14. B
15. B 16. A 17. B 18. D 19. D 20. C 21. D
22. C 23. C 24. D 25. C 26. C 27. C 28. B
29. C 30. C 31. A 32. C 33. B 34. D 35. D
36. D 37. D 38. A 39. B 40. D 41. B 42. C
43. B 44. D 45. (A) S; (B) Q; (C) P 46. 5
49. $x^3 + qx - r = 0$ 53. (a) $3 + 4i$ (b) $-\frac{8}{29} + 0i$
54. (i) Principal Arg $z = -\frac{4\pi}{9}$; $|z| = 2 \cos \frac{4\pi}{9}$; Arg $z = 2k\pi - \frac{4\pi}{9}$ $k \in \mathbb{I}$
(ii) Modulus = $\sec^2 1$, Arg = $2n\pi + (2 - \pi)$ ($n \in \mathbb{I}$); Principal Arg = $(2 - \pi)$
(iii) Principal value of Arg $z = -\frac{\pi}{2}$ & $|z| = \frac{3}{2}$; Principal value of Arg $z = \frac{\pi}{2}$ & $|z| = \frac{2}{3}$
(iv) Modulus = $\frac{1}{\sqrt{2}} \operatorname{cosec} \frac{\pi}{5}$, Arg $z = 2n\pi + \frac{11\pi}{20}$ ($n \in \mathbb{I}$), Principal Arg = $\frac{11\pi}{20}$
55. (a) $x = 1$ & $y = 2$ (b) $(1, 1)$ & $(0, \frac{5}{2})$ 56. 12 58. 10 61. 259
63. 163 64. $-\omega$ or $-\omega^2$ 65. $k > \frac{1}{2} |\alpha - \beta|^2$
66. $|f(z)|$ is maximum when $z = \omega$, where ω is the cube root unity and $|f(z)| = \sqrt{13}$
67. 8
69. $\frac{1}{4}$ 72. $K = -\frac{4}{9}$ 73. 18
74. Centre $\equiv \frac{k^2\beta - \alpha}{k^2 - 1}$, Radius = $\frac{1}{(k^2 - 1)} \sqrt{|\alpha - k^2\beta|^2 - (k^2 \cdot |\beta|^2 - |\alpha|^2) \cdot (k^2 - 1)}$
75. $z_2 = -\sqrt{3}i$; $z_3 = (1 - \sqrt{3}) + i$; $z_4 = (1 + \sqrt{3}) - i$

