


IIT MATHEMATICS

SEQUENCE & SERIES



INDEX

S.No.	Contents	Page No.
	SEQUENCE & SERIES	
1.	Theory	01-16
2.	Exercise # 1	17-18
3.	Exercise # 2	19-20
4.	Exercise # 3 (JM)	21-28
5.	Exercise # 4 (JA)	29-30
6.	Exercise # 5	31-33
7.	Exercise # 6	34-35
8.	Answer key	36-37

RESULTS OF BEST MENTORSHIP BY THE NUCLEUS TEAM



AIR-1
CHITRAANG MURDIA
Gen. Category - 2014



AIR-3
GOVIND LAHOTI
Gen. Category - 2014




AIR-6
NISHIT AGARWAL
Gen. Category - 2012




AIR-8
AMEY GUPTA
Gen. Category - 2014


JEE MAIN RESULTS 2020 (January Attempt) OF NUCLEUS EDUCATION




100 Percentile
(Physics)
DAKSH Khandelwal
2020



100 Percentile
(Maths & Physics)
VAIBHAV SAHA
2020




100 Percentile
(Physics)
ANISH MOHAN
2020



100 Percentile
(Maths)
ARCHIT PATNAIK
2020




100 Percentile
(Maths)
SWAPNIL YASASVI
2020




100 Percentile
(Maths)
PARSHANT ARORA
2020


JEE ADVANCED RESULTS OF NUCLEUS EDUCATION




AIR-10
LAKSHAY SHARMA
Gen. Category - 2017




AIR-12
YATEESH AGRAWAL
Gen. Category - 2017




AIR-23
ABHEY GOYAL
Gen. Category - 2017




AIR-24
TUSHAR GAUTAM
Gen. Category - 2017




AIR-37
PIYUSH TIBAREWAL
Gen. Category - 2017




AIR-42
SATVIK
Gen. Category - 2017



AIR-66
MAYANK DUBEY
Gen. Category - 2017



AIR-98
HRITHIK
Gen. Category - 2017



AIR-20
SHASHANK AGRAWAL
Gen. Category - 2018



AIR-27
RAAGHAV RAAJ
Gen. Category - 2018



AIR-32
SHREYA PATHAK
Gen. Category - 2018



AIR-61
SIDDHANT CHOUDHARY
Gen. Category - 2018



AIR-67
ANISWAR S KRISHNAN
Gen. Category - 2018
DLP



AIR-78
AAYUSH KADAM
Gen. Category - 2018



AIR-61
SARTHAK BEHERA
Gen. Category - 2018
*SDCCP




AIR-91
ANDREWS G. VARGHESE
Gen. Category - 2018
*SDCCP



AIR-2
HIMANSHU GAURAV SINGH
Gen. Category - 2019
*SDCCP



AIR-19
VIBHAV AGGARWAL
Gen. Category - 2019



AIR-33
S. PRAJEETH
Gen. Category - 2019
*SDCCP



AIR-48
SOHAM MISTRI
Gen. Category - 2019
DLP



AIR-51
SAYANTAN DHAR
Gen. Category - 2019
DLP



AIR-53
GAURAV KRISHAN GUPTA
Gen. Category - 2019
DLP



AIR-86
SATVIK JAIN
Gen. Category - 2019

SEQUENCE & SERIES

1. DEFINITION :

Sequence :

A succession of terms $a_1, a_2, a_3, a_4, \dots$ formed according to some rule or law.

Examples are : 1, 4, 9, 16, 25

$-1, 1, -1, 1, \dots$

$\frac{x}{1!}, \frac{x^2}{2!}, \frac{x^3}{3!}, \frac{x^4}{4!}, \dots$

A finite sequence has a finite (i.e. limited) number of terms, as in the first example above. An infinite sequence has an unlimited number of terms, i.e. there is no last term, as in the second and third examples.

Series :

The indicated sum of the terms of a sequence. In the case of a finite sequence $a_1, a_2, a_3, \dots, a_n$ the corresponding series is $a_1 + a_2 + a_3 + \dots + a_n = \sum_{k=1}^n a_k$. This series has a finite or limited number of terms and is called a finite series.

2. ARITHMETIC PROGRESSION (A.P.) :

A.P. is a sequence whose terms differ by a fixed number. This fixed number is called the common difference. If a is the first term & d the common difference, then A.P. can be written as

$a, a + d, a + 2d, \dots, a + (n - 1)d, \dots$

(a) n^{th} term of AP $T_n = a + (n - 1)d$, where $d = t_n - t_{n-1}$

(b) The sum of the first terms : $S_n = \frac{n}{2} [a + \ell] = \frac{n}{2} [2a + (n-1)d]$ where ℓ is n^{th} term.

Note :

- (i) n^{th} term of an A.P. is of the form $An + B$ i.e. a linear expression in 'n', in such a case the coefficient of n is the common difference of the A.P. i.e. A .
- (ii) Sum of first 'n' terms of an A.P. is of the form $An^2 + Bn$ i.e. a quadratic expression in 'n', in such case the common difference is twice the coefficient of n^2 . i.e. $2A$
- (iii) Also n^{th} term $T_n = S_n - S_{n-1}$

Illustration 1 : If $(x + 1)$, $3x$ and $(4x + 2)$ are first three terms of an A.P. then its 5^{th} term is -
(A) 14 (B) 19 (C) 24 (D) 28

Solution : $(x + 1)$, $3x$, $(4x + 2)$ are in AP
 $\Rightarrow 3x - (x + 1) = (4x + 2) - 3x \Rightarrow x = 3$
 $\therefore a = 4, d = 9 - 4 = 5$
 $\Rightarrow T_5 = 4 + (4)5 = 24$

Ans. (C)

Illustration 2 : The sum of first four terms of an A.P. is 56 and the sum of its last four terms is 112. If its first term is 11 then find the number of terms in the A.P.

Solution :
 $a + a + d + a + 2d + a + 3d = 56$
 $4a + 6d = 56$
 $44 + 6d = 56$ (as $a = 11$)
 $6d = 12$ hence $d = 2$
 Let total number of terms = n
 Now sum of last four terms.
 $a + (n-1)d + a + (n-2)d + a + (n-3)d + a + (n-4)d = 112$
 $\Rightarrow 4a + (4n-10)d = 112 \Rightarrow 44 + (4n-10)2 = 112$
 $\Rightarrow 4n - 10 = 34$
 $\Rightarrow n = 11$

Ans.

Illustration 3 : The sum of first n terms of two A.P.s. are in ratio $\frac{7n+1}{4n+27}$. Find the ratio of their 11th terms.

Solution : Let a_1 and a_2 be the first terms and d_1 and d_2 be the common differences of two A.P.s respectively then

$$\frac{\frac{n}{2}[2a_1 + (n-1)d_1]}{\frac{n}{2}[2a_2 + (n-1)d_2]} = \frac{7n+1}{4n+27} \Rightarrow \frac{a_1 + \left(\frac{n-1}{2}\right)d_1}{a_2 + \left(\frac{n-1}{2}\right)d_2} = \frac{7n+1}{4n+27}$$

For ratio of 11th terms

$$\frac{n-1}{2} = 10 \Rightarrow n = 21$$

$$\text{so ratio of 11th terms is } \frac{7(21)+1}{4(21)+27} = \frac{148}{111} = \frac{4}{3}$$

Ans.

Do yourself - 1 :

- (i) Write down the sequence whose n th terms is : (a) $\frac{2^n}{n}$ (b) $\frac{3+(-1)^n}{3^n}$
- (ii) For an A.P, show that $t_m + t_{2n+m} = 2t_{m+n}$
- (iii) If the sum of p terms of an A.P. is q and the sum of its q terms is p , then find the sum of its $(p+q)$ term.

3. PROPERTIES OF A.P. :

- (a) If each term of an A.P. is increased, decreased, multiplied or divided by the some nonzero number, then the resulting sequence is also an A.P.
- (b) Three numbers in A.P. : $a-d, a, a+d$
 Four numbers in A.P. : $a-3d, a-d, a+d, a+3d$
 Five numbers in A.P. : $a-2d, a-d, a, a+d, a+2d$
 Six numbers in A.P. : $a-5d, a-3d, a-d, a+d, a+3d, a+5d$ etc.

- (c) The common difference can be zero, positive or negative.
 (d) k^{th} term from the last = $(n - k + 1)^{\text{th}}$ term from the beginning (If total number of terms = n).
 (e) The sum of the two terms of an AP equidistant from the beginning & end is constant and equal to the sum of first & last terms. $\Rightarrow T_k + T_{n-k+1} = \text{constant} = a + \ell$.
 (f) Any term of an AP (except the first) is equal to half the sum of terms which are equidistant from it. $a_n = (1/2)(a_{n-k} + a_{n+k})$, $k < n$
 For $k = 1$, $a_n = (1/2)(a_{n-1} + a_{n+1})$; For $k = 2$, $a_n = (1/2)(a_{n-2} + a_{n+2})$ and so on.
 (g) If a, b, c are in AP, then $2b = a + c$.

Illustration 4 : Four numbers are in A.P. If their sum is 20 and the sum of their squares is 120, then the middle terms are -

- (A) 2, 4 (B) 4, 6 (C) 6, 8 (D) 8, 10

Solution :

Let the numbers are $a - 3d, a - d, a + d, a + 3d$

$$\text{given, } a - 3d + a - d + a + d + a + 3d = 20 \Rightarrow 4a = 20 \Rightarrow a = 5$$

$$\text{and } (a - 3d)^2 + (a - d)^2 + (a + d)^2 + (a + 3d)^2 = 120 \Rightarrow 4a^2 + 20d^2 = 120$$

$$\Rightarrow 4 \times 5^2 + 20d^2 = 120 \Rightarrow d^2 = 1 \Rightarrow d = \pm 1$$

Hence numbers are 2, 4, 6, 8 or 8, 6, 4, 2 **Ans. (B)**

Illustration 5 : If $a_1, a_2, a_3, \dots, a_n$ are in A.P. where $a_i > 0$ for all i , show that :

$$\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} = \frac{(n-1)}{\sqrt{a_1} + \sqrt{a_n}}$$

Solution :

$$\begin{aligned} \text{L.H.S.} &= \frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} \\ &= \frac{1}{\sqrt{a_2} + \sqrt{a_1}} + \frac{1}{\sqrt{a_3} + \sqrt{a_2}} + \dots + \frac{1}{\sqrt{a_n} + \sqrt{a_{n-1}}} \\ &= \frac{\sqrt{a_2} - \sqrt{a_1}}{(a_2 - a_1)} + \frac{\sqrt{a_3} - \sqrt{a_2}}{(a_3 - a_2)} + \dots + \frac{\sqrt{a_n} - \sqrt{a_{n-1}}}{a_n - a_{n-1}} \end{aligned}$$

Let 'd' is the common difference of this A.P.

$$\text{then } a_2 - a_1 = a_3 - a_2 = \dots = a_n - a_{n-1} = d$$

Now **L.H.S.**

$$\begin{aligned} &= \frac{1}{d} \{ \sqrt{a_2} - \sqrt{a_1} + \sqrt{a_3} - \sqrt{a_2} + \dots + \sqrt{a_{n-1}} - \sqrt{a_{n-2}} + \sqrt{a_n} - \sqrt{a_{n-1}} \} = \frac{1}{d} \{ \sqrt{a_n} - \sqrt{a_1} \} \\ &= \frac{a_n - a_1}{d(\sqrt{a_n} + \sqrt{a_1})} = \frac{a_1 + (n-1)d - a_1}{d(\sqrt{a_n} + \sqrt{a_1})} = \frac{1}{d} \frac{(n-1)d}{d(\sqrt{a_n} + \sqrt{a_1})} = \frac{n-1}{\sqrt{a_n} + \sqrt{a_1}} = \text{R.H.S.} \end{aligned}$$

Do yourself - 2 :

- (i) Find the sum of first 24 terms of the A.P. $a_1, a_2, a_3, \dots$, if it is known that $a_1 + a_5 + a_{10} + a_{15} + a_{20} + a_{24} = 225$.
- (ii) Find the number of terms common to the two A.P.'s 3, 7, 11, 407 and 2, 9, 16,, 709

4. GEOMETRIC PROGRESSION (G.P.) :

G.P. is a sequence of non zero numbers each of the succeeding term is equal to the preceding term multiplied by a constant. Thus in a GP the ratio of successive terms is constant. This constant factor is called the COMMON RATIO of the sequence & is obtained by dividing any term by the immediately previous term. Therefore $a, ar, ar^2, ar^3, ar^4, \dots$ is a GP with 'a' as the first term & 'r' as common ratio.

(a) n^{th} term ; $T_n = a r^{n-1}$

(b) Sum of the first n terms; $S_n = \frac{a(r^n - 1)}{r - 1}$, if $r \neq 1$

(c) Sum of infinite G.P. , $S_\infty = \frac{a}{1 - r}$; $0 < |r| < 1$

5. PROPERTIES OF GP :

- (a) If each term of a G.P. be multiplied or divided by the some non-zero quantity, then the resulting sequence is also a G.P.
- (b) Three consecutive terms of a GP : $a/r, a, ar$;
Four consecutive terms of a GP : $a/r^3, a/r, ar, ar^3$ & so on.
- (c) If a, b, c are in G.P. then $b^2 = ac$.
- (d) If in a G.P, the product of two terms which are equidistant from the first and the last term, is constant and is equal to the product of first and last term.
 $\Rightarrow T_k \cdot T_{n-k+1} = \text{constant} = a \cdot \ell$
- (e) If each term of a G.P. be raised to the same power, then resulting sequence is also a G.P.
- (f) In a G.P., $T_r^2 = T_{r-k} \cdot T_{r+k}$, $k < r$, $r \neq 1$
- (g) If the terms of a given G.P. are chosen at regular intervals, then the new sequence is also a G.P.
- (h) If $a_1, a_2, a_3, \dots, a_n$ is a G.P. of positive terms, then $\log a_1, \log a_2, \dots, \log a_n$ is an A.P. and vice-versa.
- (i) If $a_1, a_2, a_3, \dots$ and $b_1, b_2, b_3, \dots$ are two G.P.'s then $a_1 b_1, a_2 b_2, a_3 b_3, \dots$ & $\frac{a_1}{b_1}, \frac{a_2}{b_2}, \frac{a_3}{b_3}, \dots$ is also in G.P.

Illustration 6 : If a, b, c, d and p are distinct real numbers such that $(a^2 + b^2 + c^2)p^2 - 2p(ab + bc + cd) + (b^2 + c^2 + d^2) \leq 0$ then a, b, c, d are in
(A) A.P. (B) G.P. (C) H.P. (D) none of these

Solution : Here, the given condition $(a^2 + b^2 + c^2)p^2 - 2p(ab + bc + ca) + b^2 + c^2 + d^2 \leq 0$
 $\Rightarrow (ap - b)^2 + (bp - c)^2 + (cp - d)^2 \leq 0$
 $\therefore$ a square can not be negative
 $\therefore ap - b = 0, bp - c = 0, cp - d = 0 \Rightarrow p = \frac{b}{a} = \frac{c}{b} = \frac{d}{c} \Rightarrow a, b, c, d$ are in G.P.

Ans. (B)

Illustration 7 : If a, b, c are in G.P., then the equations $ax^2 + 2bx + c = 0$ and $dx^2 + 2ex + f = 0$ have a common root if $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in -
(A) A.P. (B) G.P. (C) H.P. (D) none of these

Solution : a, b, c are in G.P. $\Rightarrow b^2 = ac$

Now the equation $ax^2 + 2bx + c = 0$ can be rewritten as $ax^2 + 2\sqrt{ac}x + c = 0$
 $\Rightarrow (\sqrt{ax} + \sqrt{c})^2 = 0 \Rightarrow -\sqrt{\frac{c}{a}}, -\sqrt{\frac{c}{a}}$

If the two given equations have a common root, then this root must be $-\sqrt{\frac{c}{a}}$.

Thus $d\frac{c}{a} - 2e\sqrt{\frac{c}{a}} + f = 0 \Rightarrow \frac{d}{a} + \frac{f}{c} = \frac{2e}{c}\sqrt{\frac{c}{a}} = \frac{2e}{\sqrt{ac}} = \frac{2e}{b} \Rightarrow \frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in A.P. **Ans. (A)**

Illustration 8 : A number consists of three digits which are in G.P. the sum of the right hand and left hand digits exceeds twice the middle digit by 1 and the sum of the left hand and middle digits is two third of the sum of the middle and right hand digits. Find the numbers.

Solution : Let the three digits be a, ar and ar^2 then number is

$$100a + 10ar + ar^2 \quad \dots(i)$$

Given, $a + ar^2 = 2ar + 1$

or $a(r^2 - 2r + 1) = 1$

or $a(r - 1)^2 = 1 \quad \dots(ii)$

Also given $a + ar = \frac{2}{3}(ar + ar^2)$

$$\Rightarrow 3 + 3r = 2r + 2r^2 \Rightarrow 2r^2 - r - 3 = 0 \Rightarrow (r + 1)(2r - 3) = 0$$

$$\therefore r = -1, 3/2$$

for $r = -1, a = \frac{1}{(r-1)^2} = \frac{1}{4} \notin \mathbb{I} \quad \therefore r \neq -1$

for $r = 3/2, a = \frac{1}{\left(\frac{3}{2}-1\right)^2} = 4 \quad \{\text{from (ii)}\}$

From (i), number is $400 + 10 \cdot 4 \cdot \frac{3}{2} + 4 \cdot \frac{9}{4} = 469$

Ans.

Illustration 9 : Find the value of $0.32\overline{58}$

Solution : Let $R = 0.32\overline{58} \Rightarrow R = 0.32585858....$ (i)
Here number of figures which are not recurring is 2 and number of figures which are recurring is also 2.
then $100 R = 32.585858.....$ (ii)
and $10000 R = 3258.5858.....$ (iii)
Subtracting (ii) from (iii) , we get
 $9900 R = 3226 \Rightarrow R = \frac{1613}{4950}$

Aliter Method : $R = .32 + .0058 + .0058 + .000058 +$

$$= .32 + \frac{58}{10^4} \left(1 + \frac{1}{10} + \frac{1}{10^2} + \infty \right)$$

$$= .32 + \frac{58}{10^4} \left(\frac{1}{1 - \frac{1}{10}} \right)$$

$$= \frac{32}{100} + \frac{58}{9900} = \frac{3168 + 58}{9900} = \frac{3226}{9900} = \frac{1613}{4950}$$

Do yourself - 3 :

- (i) Find a three digit number whose consecutive digits form a G.P. If we subtract 792 from this number, we get a number consisting of the same digits written in the reverse order. Now, if we increase the second digit of the required number by 2, then the resulting digits will form an A.P.
- (ii) If the third term of G.P. is 4, then find the product of first five terms.
- (iii) If a, b, c are respectively the p^{th} , q^{th} and r^{th} terms of the given G.P., then show that $(q - r) \log a + (r - p) \log b + (p - q) \log c = 0$, where a, b, c > 0.
- (iv) Find three numbers in G.P., whose sum is 52 and the sum of whose products in pairs is 624.
- (v) The rational number which equals the number $2.\overline{357}$ with recurring decimal is -

(A) $\frac{2357}{999}$

(B) $\frac{2379}{997}$

(C) $\frac{785}{333}$

(D) $\frac{2355}{1001}$

6. HARMONIC PROGRESSION (H.P.) :

A sequence is said to be in H.P. if the reciprocal of its terms are in AP.

If the sequence $a_1, a_2, a_3,, a_n$ is an HP then $1/a_1, 1/a_2,, 1/a_n$ is an AP . Here we do not have the formula for the sum of the n terms of an HP. The general form of a harmonic

progression is $\frac{1}{a}, \frac{1}{a+d}, \frac{1}{a+2d}, \frac{1}{a+(n-1)d}$ Note : No term of any H.P. can be zero.

(i) If a, b, c are in HP, then $b = \frac{2ac}{a+c}$ or $\frac{a}{c} = \frac{a-b}{b-c}$

Illustration 10 : The sum of three numbers are in H.P. is 37 and the sum of their reciprocals is $1/4$. Find the numbers.

Solution : Three numbers are in H.P. can be taken as

$$\frac{1}{a-d}, \frac{1}{a}, \frac{1}{a+d}$$

$$\text{then } \frac{1}{a-d} + \frac{1}{a} + \frac{1}{a+d} = 37 \quad \dots\dots(i)$$

$$\text{and } a-d + a + a+d = \frac{1}{4} \Rightarrow a = \frac{1}{12}$$

$$\text{from (i), } \frac{12}{1-12d} + 12 + \frac{12}{1+12d} = 37 \Rightarrow \frac{12}{1-12d} + \frac{12}{1+12d} = 25$$

$$\Rightarrow \frac{24}{1-144d^2} = 25$$

$$\Rightarrow 1 - 144d^2 = \frac{24}{25} \Rightarrow d^2 = \frac{1}{25 \times 144}$$

$$\therefore d = \pm \frac{1}{60}$$

$$\therefore a-d, a, a+d \text{ are } \frac{1}{15}, \frac{1}{12}, \frac{1}{10} \text{ or } \frac{1}{10}, \frac{1}{12}, \frac{1}{15}$$

Hence, three numbers in H.P. are 15, 12, 10 or 10, 12, 15

Ans.

Illustration 11 : Suppose a is a fixed real number such that $\frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz}$

If p, q, r are in A.P., then prove that x, y, z are in H.P.

Solution : $\therefore$ p, q, r are in A.P.

$$\therefore q-p = r-q \quad \dots\dots(i)$$

$$\Rightarrow p-q = q-r = k \text{ (let)}$$

$$\text{given } \frac{a-x}{px} = \frac{a-y}{qy} = \frac{a-z}{rz} \Rightarrow \frac{\frac{a}{x}-1}{p} = \frac{\frac{a}{y}-1}{q} = \frac{\frac{a}{z}-1}{r}$$

$$\Rightarrow \frac{\left(\frac{a}{x}-1\right)-\left(\frac{a}{y}-1\right)}{p-q} = \frac{\left(\frac{a}{y}-1\right)-\left(\frac{a}{z}-1\right)}{q-r} \quad (\text{by law of proportion})$$

$$\Rightarrow \frac{\frac{a}{x}-\frac{a}{y}}{k} = \frac{\frac{a}{y}-\frac{a}{z}}{k} \quad \{\text{from (i)}\}$$

$$\Rightarrow a \left(\frac{1}{x} - \frac{1}{y} \right) = a \left(\frac{1}{y} - \frac{1}{z} \right) \Rightarrow \frac{1}{x} - \frac{1}{y} = \frac{1}{y} - \frac{1}{z}$$

$$\therefore \frac{2}{y} = \frac{1}{x} + \frac{1}{z} \quad \therefore \frac{1}{x}, \frac{1}{y}, \frac{1}{z} \text{ are in A.P.}$$

Hence x, y, z are in H.P.

Do yourself - 4 :

- (i) If the 7th term of a H.P. is 8 and the 8th term is 7. Then find the 28th term.
 (ii) In a H.P., if 5th term is 6 and 3rd term is 10. Find the 2nd term.
 (iii) If the pth, qth and rth terms of a H.P. are a,b,c respectively, then prove that $\frac{q-r}{a} + \frac{r-p}{b} + \frac{p-q}{c} = 0$

7. MEANS

(a) ARITHMETIC MEAN :

If three terms are in A.P. then the middle term is called the A.M. between the other two,

so if a, b, c are in A.P., b is A.M. of a & c. So A.M. of a and c = $\frac{a+c}{2} = b$.

n-ARITHMETIC MEANS BETWEEN TWO NUMBERS :

If a,b be any two given numbers & a, A₁, A₂,, A_n, b are in AP, then A₁, A₂,.....A_n are the 'n' A.M's between a & b then.

A₁ = a + d, A₂ = a + 2d,, A_n = a + nd or b - d, where

$$d = \frac{b-a}{n+1}$$

$$\Rightarrow A_1 = a + \frac{b-a}{n+1}, A_2 = a + \left(\frac{2b-a}{n+1}\right), \dots\dots\dots$$

Note : Sum of n A.M's inserted between a & b is equal to n times the single A.M. between a & b

$$\text{i.e. } \sum_{r=1}^n A_r = nA \text{ where } A \text{ is the single A.M. between } a \text{ \& } b.$$

(b) GEOMETRIC MEAN :

If a, b, c are in G.P., then b is the G.M. between a & c, $b^2 = ac$. So G.M. of a and c = $\sqrt{ac} = b$

n-GEOMETRIC MEANS BETWEEN TWO NUMBERS :

If a, b are two given positive numbers & a, G₁, G₂,, G_n, b are in G.P. Then G₁, G₂, G₃,.....G_n are 'n' G.Ms between a & b. where $b = ar^{n+1} \Rightarrow r = (b/a)^{1/n+1}$

$$G_1 = a(b/a)^{1/n+1}, \quad G_2 = a(b/a)^{2/n+1}, \dots\dots\dots, \quad G_n = a(b/a)^{1/n+1},$$

$$= ar, \quad = ar^2, \dots\dots\dots, \quad = ar^n = b/r$$

Note : The product of n G.Ms between a & b is equal to nth power of the single G.M.

$$\text{between } a \text{ \& } b \text{ i.e. } \prod_{r=1}^n G_r = (G)^n \text{ where } G \text{ is the single G.M. between } a \text{ \& } b$$

(c) **HARMONIC MEAN :**

If a, b, c are in H.P., then b is H.M. between a & c. So H.M. of a and c = $\frac{2ac}{a+c} = b$.

Insertion of 'n' HM's between a and b :

a, H₁, H₂, H₃,, H_n, b → H.P

$$\frac{1}{a}, \frac{1}{H_1}, \frac{1}{H_2}, \frac{1}{H_3}, \dots, \frac{1}{H_n}, \frac{1}{b} \rightarrow \text{A.P.}$$

$$\frac{1}{b} = \frac{1}{a} + (n+1)D \Rightarrow D = \frac{b-a}{n+1}$$

$$\frac{1}{H_n} = \frac{1}{a} + n \left(\frac{\frac{1}{b} - \frac{1}{a}}{n+1} \right)$$

Do yourself - 5 :

- (i) If $\frac{a^n + b^n}{a^{n-1} + b^{n-1}}$ is the G.M. between a & b then find the value of 'n'.
- (ii) If b is the harmonic mean between a and c, then prove that $\frac{1}{b-a} + \frac{1}{b-c} = \frac{1}{a} + \frac{1}{c}$.

8. ARITHMETICO - GEOMETRIC SERIES :

A series, each term of which is formed by multiplying the corresponding term of an A.P. & G.P. is called the Arithmetico-Geometric Series, e.g. $1 + 3x + 5x^2 + 7x^3 + \dots$

Here 1, 3, 5, are in A.P. & 1, x, x², x³ are in G.P.

(a) SUM OF N TERMS OF AN ARITHMETICO-GEOMETRIC SERIES :

$$\text{Let } S_n = a + (a+d)r + (a+2d)r^2 + \dots + [a + (n-1)d]r^{n-1}$$

$$\text{then } S_n = \frac{a}{1-r} + \frac{dr(1-r^{n-1})}{(1-r)^2} - \frac{[a + (n-1)d]r^n}{1-r}, r \neq 1$$

(b) SUM TO INFINITY :

$$\text{If } 0 < |r| < 1 \text{ \& } n \rightarrow \infty, \text{ then } \lim_{n \rightarrow \infty} r^n = 0, S_\infty = \frac{a}{1-r} + \frac{dr}{(1-r)^2}$$

Illustration 12 : Find the sum of series $4 - 9x + 16x^2 - 25x^3 + 36x^4 - 49x^5 + \dots \infty$.

Solution : Let $S = 4 - 9x + 16x^2 - 25x^3 + 36x^4 - 49x^5 + \dots \infty$

$$- Sx = -4x + 9x^2 - 16x^3 + 25x^4 - 36x^5 + \dots \infty$$

On subtraction, we get

$$S(1 + x) = 4 - 5x + 7x^2 - 9x^3 + 11x^4 - 13x^5 + \dots \infty$$

$$-S(1 + x)x = -4x + 5x^2 - 7x^3 + 9x^4 - 11x^5 + \dots \infty$$

On subtraction, we get

$$S(1 + x)^2 = 4 - x + 2x^2 - 2x^3 + 2x^4 - 2x^5 + \dots \infty$$

$$= 4 - x + 2x^2 (1 - x + x^2 - \dots \infty) = 4 - x + \frac{2x^2}{1 + x} = \frac{4 + 3x + x^2}{1 + x}$$

$$S = \frac{4 + 3x + x^2}{(1 + x)^3}$$

Ans.

Illustration 13 : Find the sum of series upto n terms $\left(\frac{2n+1}{2n-1}\right) + 3\left(\frac{2n+1}{2n-1}\right)^2 + 5\left(\frac{2n+1}{2n-1}\right)^3 + \dots$

Solution : For $x \neq 1$, let

$$S = x + 3x^2 + 5x^3 + \dots + (2n-3)x^{n-1} + (2n-1)x^n \quad \dots (i)$$

$$\Rightarrow xS = x^2 + 3x^3 + \dots + (2n-5)x^{n-1} + (2n-3)x^n + (2n-1)x^{n+1} \quad \dots (ii)$$

Subtracting (ii) from (i), we get

$$(1 - x)S = x + 2x^2 + 2x^3 + \dots + 2x^{n-1} + 2x^n - (2n-1)x^{n+1} = x + \frac{2x^2(1 - x^{n-1})}{1 - x} - (2n-1)x^{n+1}$$

$$= \frac{x}{1 - x} [1 - x + 2x - 2x^n - (2n-1)x^n + (2n-1)x^{n+1}]$$

$$\Rightarrow S = \frac{x}{(1 - x)^2} [(2n-1)x^{n+1} - (2n+1)x^n + 1 + x]$$

$$\text{Thus } \left(\frac{2n+1}{2n-1}\right) + 3\left(\frac{2n+1}{2n-1}\right)^2 + \dots + (2n-1)\left(\frac{2n+1}{2n-1}\right)^n$$

$$= \left(\frac{2n+1}{2n-1}\right)\left(\frac{2n-1}{2}\right)^2 \left[(2n-1)\left(\frac{2n+1}{2n-1}\right)^{n+1} - (2n+1)\left(\frac{2n+1}{2n-1}\right)^n + 1 + \frac{2n+1}{2n-1} \right]$$

$$= \frac{4n^2 - 1}{4} \cdot \frac{4n}{2n-1} = n(2n+1)$$

Ans.

Do yourself - 6 :

(i) Find sum to n terms of the series $3 + 5 \times \frac{1}{4} + 7 \times \frac{1}{4^2} + \dots$

(ii) If the sum to the infinity of the series $3 + 5r + 7r^2 + \dots$ is $\frac{44}{9}$, then find the value of r.

(iii) If the sum to infinity of the series $3 + (3 + d) \cdot \frac{1}{4} + (3 + 2d) \cdot \frac{1}{4^2} + \dots$ is $\frac{44}{9}$ then find d.

9. SIGMA NOTATIONS (Σ)

THEOREMS :

$$(a) \sum_{r=1}^n (a_r \pm b_r) = \sum_{r=1}^n a_r \pm \sum_{r=1}^n b_r \quad (b) \sum_{r=1}^n k a_r = k \sum_{r=1}^n a_r \quad (c) \sum_{r=1}^n k = nk ; \text{ where } k \text{ is a constant.}$$

10. RESULTS

$$(a) \sum_{r=1}^n r = \frac{n(n+1)}{2} \text{ (sum of the first } n \text{ natural numbers)}$$

$$(b) \sum_{r=1}^n r^2 = \frac{n(n+1)(2n+1)}{6} \text{ (sum of the squares of the first } n \text{ natural numbers)}$$

$$(c) \sum_{r=1}^n r^3 = \frac{n^2(n+1)^2}{4} = \left[\sum_{r=1}^n r \right]^2 \text{ (sum of the cubes of the first } n \text{ natural numbers)}$$

$$(d) \sum_{r=1}^n r^4 = \frac{n}{30} (n+1)(2n+1)(3n^2+3n-1)$$

$$(e) \sum_{r=1}^n (2r-1) = n^2 \text{ (sum of first } n \text{ odd natural numbers)}$$

$$(f) \sum_{r=1}^n 2r = n(n+1) \text{ (sum of first } n \text{ even natural numbers)}$$

Note :

If n^{th} term of a sequence is given by $T_n = an^3 + bn^2 + cn + d$ where a, b, c, d are constants, then sum of n terms $S_n = \Sigma T_n = a\Sigma n^3 + b\Sigma n^2 + c\Sigma n + \Sigma d$

This can be evaluated using the above results.

Illustration 14 : Sum up to 16 terms of the series $\frac{1^3}{1} + \frac{1^3+2^3}{1+3} + \frac{1^3+2^3+3^3}{1+3+5} + \dots$ is

- (A) 450 (B) 456 (C) 446 (D) none of these

Solution :

$$t_n = \frac{1^3+2^3+3^3+\dots+n^3}{1+3+5+\dots(2n-1)} = \frac{\left\{ \frac{n(n+1)}{2} \right\}^2}{\frac{n}{2} \{2+2(n-1)\}} = \frac{\frac{n^2(n+1)^2}{4}}{\frac{n}{2} \cdot n} = \frac{\frac{n^2(n+1)^2}{4}}{n^2} = \frac{(n+1)^2}{4} = \frac{n^2}{4} + \frac{n}{2} + \frac{1}{4}$$

$$\therefore S_n = \Sigma t_n = \frac{1}{4} \Sigma n^2 + \frac{1}{2} \Sigma n + \frac{1}{4} \Sigma 1 = \frac{1}{4} \cdot \frac{n(n-1)(2n-1)}{6} + \frac{1}{2} \cdot \frac{n(n+1)}{2} + \frac{1}{4} \cdot n$$

$$\therefore S_{16} = \frac{16 \cdot 17 \cdot 33}{24} + \frac{16 \cdot 17}{4} + \frac{16}{4} = 446 \quad \text{Ans.(C)}$$

11. METHOD OF DIFFERENCE :

Some times the n^{th} term of a sequence or a series can not be determined by the method, we have discussed earlier. So we compute the difference between the successive terms of given sequence for obtained the n^{th} terms.

If $T_1, T_2, T_3, \dots, T_n$ are the terms of a sequence then some times the terms $T_2 - T_1, T_3 - T_2, \dots$ constitute an AP/GP. n^{th} term of the series is determined & the sum to n terms of the sequence can easily be obtained.

- Case 1 :**
- (a) If difference series are in A.P., then
Let $T_n = an^2 + bn + c$, where a, b, c are constant
 - (b) If difference of difference series are in A.P.
Let $T_n = an^3 + bn^2 + cn + d$, where a, b, c, d are constant

- Case 2 :**
- (a) If difference are in G.P., then
Let $T_n = ar^n + b$, where r is common ratio & a, b are constant
 - (b) If difference of difference are in G.P., then
Let $T_n = ar^n + bn + c$, where r is common ratio & a, b, c are constant
Determine constant by putting $n = 1, 2, 3 \dots n$ and putting the value of $T_1, T_2, T_3 \dots$ and sum of series $(S_n) = \sum T_n$

Do yourself - 7 :

- (i) Find the sum of the series upto n terms $1 + \frac{1+2}{2} + \frac{1+2+3}{3} + \frac{1+2+3+4}{4} + \dots$
- (ii) Find the sum of ' n ' terms of the series whose n^{th} term is $t_n = 3n^2 + 2n$.

Miscellaneous Illustration :

Illustration 15 : If $\sum_{r=1}^n T_r = \frac{n}{8}(n+1)(n+2)(n+3)$, then find $\sum_{r=1}^n \frac{1}{T_r}$.

Solution : $\therefore T_n = S_n - S_{n-1}$

$$= \sum_{r=1}^n T_r - \sum_{r=1}^{n-1} T_r = \frac{n(n+1)(n+2)(n+3)}{8} - \frac{(n-1)(n+1)(n+2)}{8} = \frac{n(n+1)(n+2)}{8} [(n+3) - (n-1)]$$

$$T_n = \frac{n(n+1)(n+2)}{8} (4) = \frac{n(n+1)(n+2)}{2}$$

$$\Rightarrow \frac{1}{T_n} = \frac{2}{n(n+1)(n+2)} = \frac{(n+2) - n}{n(n+1)(n+2)} = \frac{1}{n(n+1)} - \frac{1}{(n+1)(n+2)} \dots (i)$$

$$\text{Let } V_n = \frac{1}{n(n+1)}$$

$$\therefore \frac{1}{T_n} = V_n - V_{n+1}$$

Putting $n = 1, 2, 3, \dots, n$

$$\Rightarrow \frac{1}{T_1} + \frac{1}{T_2} + \frac{1}{T_3} + \dots + \frac{1}{T_n} = (V_1 - V_{n+1}) \Rightarrow \sum_{r=1}^n \frac{1}{T_r} = \frac{n^2 + 3n}{2(n+1)(n+2)}$$

Illustration 16 : Find the sum of n terms of the series $1 \cdot 3 \cdot 5 + 3 \cdot 5 \cdot 7 + 5 \cdot 7 \cdot 9 + \dots$

Solution :

The n^{th} term is $(2n-1)(2n+1)(2n+3)$

$$T_n = (2n-1)(2n+1)(2n+3)$$

$$T_n = \frac{1}{8} (2n-1)(2n+1)(2n+3) \{(2n+5) - (2n-3)\}$$

$$= \frac{1}{8} (V_n - V_{n-1}) \text{ [Let } V_n = (2n-1)(2n+1)(2n+3)(2n+5)]$$

$$S_n = \sum T_n = \frac{1}{8} [V_n - V_0]$$

$$\therefore S_n = \frac{(2n-1)(2n+1)(2n+3)(2n+5)}{8} + \frac{15}{8} = n(2n^3 + 8n^2 + 7n - 2) \quad \text{Ans.}$$

Illustration 17 : Find the sum of n terms of the series $3 + 7 + 14 + 24 + 37 + \dots$

Solution :

Clearly here the differences between the successive terms are $7-3, 14-7, 24-14, \dots$ i.e. 4, 7, 10, 13,....., which are in A.P.

$$\text{Let } S = 3 + 7 + 14 + 24 + \dots + T_n$$

$$S = 3 + 7 + 14 + \dots + T_{n-1} + T_n$$

Subtracting, we get

$$0 = 3 + [4 + 7 + 10 + 13 + \dots (n-1) \text{ terms}] - T_n$$

$$\therefore T_n = 3 + S_{n-1} \text{ of an A.P. whose } a = 4 \text{ and } d = 3.$$

$$\therefore T_n = 3 + \left(\frac{n-1}{2} \right) (2 \cdot 4 + (n-2)3) = \frac{6 + (n-1)(3n+2)}{4} \text{ or, } T_n = \frac{1}{2} (3n^2 - n + 4)$$

Now putting $n = 1, 2, 3, \dots, n$ and adding

$$\therefore S_n = \frac{1}{2} [3\sum n^2 - \sum n + 4n]$$

$$= \frac{1}{2} \left[3 \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} + 4n \right] = \frac{n}{2} (n^2 + n + 4) \quad \text{Ans.}$$

Aliter Method :

$$\text{Let } T_n = an^2 + bn + c$$

$$\text{Now, } T_1 = 3 = a + b + c \quad \dots(i)$$

$$T_2 = 7 = 4a + 2b + c \quad \dots(ii)$$

$$T_3 = 14 = 9a + 3b + c \quad \dots(iii)$$

Solving (i), (ii) & (iii) we get

$$a = \frac{3}{2}, b = -\frac{1}{2} \text{ \& } c = 2$$

$$\therefore T_n = \frac{1}{2} (3n^2 - n + 4)$$

$$\Rightarrow s_n = \sum T_n = \frac{1}{2} [3\sum n^2 - \sum n + 4n] = \frac{1}{2} \left[3 \frac{n(n+1)(2n+1)}{6} - \frac{n(n+1)}{2} + 4n \right] = \frac{n}{2} (n^2 + n + 4)$$

Ans.

Illustration 18 : Find the sum of n-terms of the series $1 + 4 + 10 + 22 + \dots$

Solution : Let $S = 1 + 4 + 10 + 22 + \dots + T_n$ (i)

$S = 1 + 4 + 10 + \dots + T_{n-1} + T_n$ (ii)

(i) - (ii) $\Rightarrow T_n = 1 + (3 + 6 + 12 + \dots + T_n - T_{n-1})$

$$T_n = 1 + 3 \left(\frac{2^{n-1} - 1}{2 - 1} \right)$$

$$T_n = 3 \cdot 2^{n-1} - 2$$

$$\text{So } S_n = \sum T_n = 3 \sum 2^{n-1} - \sum 2$$

$$= 3 \left(\frac{2^n - 1}{2 - 1} \right) - 2n = 3 \cdot 2^n - 2n - 3 \text{ Ans.}$$

Aliter Method :

Let $T_n = ar^n + b$, where $r = 2$

Now $T_1 = 1 = ar + b$ (i)

$T_2 = 4 = ar^2 + b$ (ii)

Solving (i) & (ii), we get

$$a = \frac{3}{2}, b = -2$$

$$\therefore T_n = 3 \cdot 2^{n-1} - 2 \Rightarrow S_n = \sum T_n = 3 \sum 2^{n-1} - \sum 2$$

$$= 3 \left(\frac{2^n - 1}{2 - 1} \right) - 2n = 3 \cdot 2^n - 2n - 3$$

Ans.

Illustration 19 : The series of natural numbers is divided into groups (1), (2, 3, 4), (5, 6, 7, 8, 9) and so on. Show that the sum of the numbers in n^{th} group is $n^3 + (n-1)^3$

Solution : The groups are (1), (2, 3, 4), (5, 6, 7, 8, 9)

The number of terms in the groups are 1, 3, 5.....

$\therefore$ The number of terms in the n^{th} group $= (2n - 1)$

the last term of the n^{th} group is n^2

If we count from last term common difference should be -1

$$\text{So the sum of numbers in the } n^{\text{th}} \text{ group} = \left(\frac{2n-1}{2} \right) \{2n^2 + (2n-2)(-1)\}$$

$$= (2n-1)(n^2 - n + 1) = 2n^3 - 3n^2 + 3n - 1 = n^3 + (n-1)^3$$

Illustration 20 : Find the natural number 'a' for which $\sum_{k=1}^n f(a+k) = 16(2^n - 1)$, where the function f

satisfied $f(x+y) = f(x) \cdot f(y)$ for all natural number x, y and further $f(1) = 2$.

Solution : It is given that

$$f(x+y) = f(x) \cdot f(y) \text{ and } f(1) = 2$$

$$f(1+1) = f(1) \cdot f(1) \Rightarrow f(2) = 2^2, f(1+2) = f(1) \cdot f(2)$$

$$\Rightarrow f(3) = 2^3, f(2+2) = f(2) \cdot f(2)$$

$$\Rightarrow f(4) = 2^4$$

$$\text{Similarly } f(k) = 2^k \text{ and } f(a) = 2^a$$

$$\text{Hence, } \sum_{k=1}^n f(a+k) = \sum_{k=1}^n f(a) \cdot f(k) = f(a) \sum_{k=1}^n f(k) = 2^a \{ 2^1 + 2^2 + \dots + 2^n \}$$

$$= 2^a \left\{ \frac{2(2^n - 1)}{2 - 1} \right\} = 2^{a+1}(2^n - 1)$$

$$\text{But } \sum_{k=1}^n f(a + k) = 16(2^n - 1)$$

$$2_{a+1}(2^n - 1) = 16(2^n - 1)$$

$$\therefore 2^{a+1} = 2^4$$

$$\therefore a+1 = 4 \quad \Rightarrow \quad a = 3$$

Ans.

19. IMPORTANT RESULTS :

$$(ix) \quad \sin \alpha + \sin(\alpha + \beta) + \sin(\alpha + 2\beta) + \dots + \sin(\alpha + (n-1)\beta) = \frac{\sin \left\{ \alpha + \left(\frac{n-1}{2} \right) \beta \right\} \sin \left(\frac{n\beta}{2} \right)}{\sin \left(\frac{\beta}{2} \right)}$$

$$(x) \quad \cos \alpha + \cos(\alpha + \beta) + \cos(\alpha + 2\beta) + \dots + \cos(\alpha + (n-1)\beta) = \frac{\cos \left\{ \alpha + \left(\frac{n-1}{2} \right) \beta \right\} \sin \left(\frac{n\beta}{2} \right)}{\sin \left(\frac{\beta}{2} \right)}$$

Do yourself - 8 :

Evaluate $\sin \frac{\pi}{n} + \sin \frac{3\pi}{n} + \sin \frac{5\pi}{n} + \dots$ to n terms

Miscellaneous Illustration :

Illustration 21: Prove that

$$\tan \alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1} \tan 2^{n-1} \alpha + 2^n \cot 2^n \alpha = \cot \alpha$$

Solution :

We know $\tan \theta = \cot \theta - 2 \cot 2\theta$ (i)

Putting $\theta = \alpha, 2\alpha, 2^2\alpha, \dots$ in (i), we get

$$\tan \alpha = (\cot \alpha - 2 \cot 2\alpha)$$

$$2(\tan 2\alpha) = 2(\cot 2\alpha - 2 \cot 2^2\alpha)$$

$$2^2(\tan 2^2\alpha) = 2^2(\cot 2^2\alpha - 2 \cot 2^3\alpha)$$

.....

$$2^{n-1}(\tan 2^{n-1}\alpha) = 2^{n-1}(\cot 2^{n-1}\alpha - 2 \cot 2^n\alpha)$$

Adding,

$$\tan \alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1} \tan 2^{n-1} \alpha = \cot \alpha - 2^n \cot 2^n \alpha$$

$$\therefore \tan \alpha + 2 \tan 2\alpha + 2^2 \tan^2 \alpha + \dots + 2^{n-1} \tan 2^{n-1} \alpha + 2^n \cot 2^n \alpha = \cot \alpha$$

ANSWERS FOR DO YOURSELF

- 1: (i) (a) $\frac{2}{1}, \frac{4}{2}, \frac{8}{3}, \frac{16}{4}, \dots$ (b) $\frac{2}{3}, \frac{4}{9}, \frac{2}{27}, \frac{4}{81}, \dots$ (iii) $-(p + q)$
- 2: (i) 900 (ii) 14
- 3: (i) 931 (ii) 4^5 (iv) 4, 12, 36 (v) C
- 4: (i) 2 (ii) 15
- 5: (i) $\frac{1}{2}$
- 6: (i) $4 + \frac{8}{9} \left(1 - \frac{1}{4^{n-1}} \right) - \left(\frac{2n+1}{3 \times 4^{n-1}} \right)$ (ii) $\frac{1}{4}$ (iii) 2
- 7: (i) $\frac{n(n+3)}{4}$ (ii) $\frac{n(n+1)(2n+3)}{2}$
- 8: 0

EXERCISE # 1

- If $a_1, a_2, a_3, \dots, a_n, \dots$ are in A.P. such that $a_4 - a_7 + a_{10} = m$, then the sum of first 13 terms of this A.P., is :
[JEE-MAINS Online 2013]
(A) 15 m (B) 10 m (C) 12 m (D) 13 m
- Let $a_1, a_2, a_3, \dots$ be an A.P. such that $\frac{a_1 + a_2 + \dots + a_p}{a_1 + a_2 + a_3 + \dots + a_q} = \frac{p^3}{q^3}$; $p \neq q$. Then $\frac{a_6}{a_{21}}$ is equal to :
[JEE-MAINS Online 2013]
(A) $\frac{121}{1861}$ (B) $\frac{11}{41}$ (C) $\frac{121}{1681}$ (D) $\frac{41}{11}$
- Given sum of the first n terms of an A. P. is $2n + 3n^2$. Another A. P. is formed with the same first term and double of the common difference, the sum of n terms of the new A. P. is :-
[JEE-MAINS Online 2013]
(A) $n + 4n^2$ (B) $n^2 + 4n$ (C) $3n + 2n^2$ (D) $6n^2 - n$
- If a, b, c are in AP, then $(a - c)^2$ equals
(A) $4(b^2 - ac)$ (B) $4(b^2 + ac)$ (C) $4b^2 - ac$ (D) $b^2 - 4ac$
- If the sum of n terms of an AP is $Pn + Qn^2$, where P, Q are constants, then its common difference is
(A) $2Q$ (B) $P + Q$ (C) $2P$ (D) $P - Q$
- Given a sequence of 4 numbers, first three of which are in G.P. and the last three are in A.P. with common difference six. If first and last terms of this sequence are equal, then the last term is :
[JEE-MAINS Online 2013]
(A) 8 (B) 16 (C) 2 (D) 4
- The first term of an infinite G.P. is 1 and every term is equal to the sum of the successive terms, then its fourth term will be-
(A) $\frac{1}{2}$ (B) $\frac{1}{8}$ (C) $\frac{1}{4}$ (D) $\frac{1}{16}$
- If G be the GM between x and y , then the value of $\frac{1}{G^2 - x^2} + \frac{1}{G^2 - y^2}$ is equal to
(A) G^2 (B) $\frac{2}{G^2}$ (C) $\frac{1}{G^2}$ (D) $3G^2$
- $2 + 4 + 7 + 11 + 16 + \dots$ to n terms =
(A) $\frac{1}{6}(n^2 + 3n + 8)$ (B) $\frac{n}{6}(n^2 + 3n + 8)$ (C) $\frac{1}{6}(n^2 - 3n + 8)$ (D) $\frac{n}{6}(n^2 - 3n + 8)$

10. If a, b, c are in HP, then $\frac{a-b}{b-c}$ is equal to
 (A) $\frac{a}{b}$ (B) $\frac{b}{a}$ (C) $\frac{a}{c}$ (D) $\frac{c}{b}$
11. The value of $1^2 + 3^2 + 5^2 + \dots + 25^2$ is : [JEE-MAINS Online 2013]
 (A) 1728 (B) 1456 (C) 2925 (D) 1469
12. The sum of the series : $1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots$ up to 10 terms, is: [JEE-MAINS Online 2013]
 (A) $\frac{22}{13}$ (B) $\frac{18}{11}$ (C) $\frac{20}{11}$ (D) $\frac{16}{9}$
13. If $a = \sum_{n=0}^{\infty} x^n, b = \sum_{n=0}^{\infty} y^n, c = \sum_{n=0}^{\infty} (xy)^n$ where $|x|, |y| < 1$; then-
 (A) $abc = a + b + c$ (B) $ab + bc = ac + b$ (C) $ac + bc = ab + c$ (D) $ab + ac = bc + a$
14. If $r > 1$ and $x = a + \frac{a}{r} + \frac{a}{r^2} + \dots$ to ∞ , $y = b - \frac{b}{r} + \frac{b}{r^2} - \dots$ to ∞ and $z = c + \frac{c}{r^2} + \frac{c}{r^4} + \dots$ to ∞ ,
 then $\frac{xy}{z} =$
 (A) $\frac{ab}{c}$ (B) $\frac{ac}{b}$ (C) $\frac{bc}{a}$ (D) None of these
15. In a GP, first term is 1. If $4T_2 + 5T_3$ is minimum, then its common ratio is
 (A) $\frac{2}{5}$ (B) $-\frac{2}{5}$ (C) $\frac{3}{5}$ (D) $-\frac{3}{5}$
16. The sum $\frac{3}{1^2} + \frac{5}{1^2+2^2} + \frac{7}{1^2+2^2+3^2} + \dots$ upto 11-terms is :- [JEE-MAINS Online 2013]
 (A) $\frac{11}{4}$ (B) $\frac{60}{11}$ (C) $\frac{7}{2}$ (D) $\frac{11}{2}$
17. The sum of the series : $(2)^2 + 2(4)^2 + 3(6)^2 + \dots$ upto 10 terms is : [JEE-MAINS Online 2013]
 (A) 11300 (B) 12100 (C) 12300 (D) 11200

EXERCISE # 2

1. (i) The sum of n terms of two arithmetic series are in the ratio of $(7n + 1) : (4n + 27)$. Find the ratio of their n^{th} term.

 (ii) In an AP of which 'a' is the 1st term, if the sum of the 1st p terms is equal to zero, show that the sum of the next q terms is $-\left(\frac{aq(p+q)}{p-1}\right)$
2. The interior angles of a convex polygon form an arithmetic progression with a common difference of 4° . Determine the number of sides of the polygon if its largest interior angle is 172° .
3. There are n AM's between 1 & 31 such that 7th mean : $(n - 1)^{\text{th}}$ mean = 5 : 9, then find the value of n .
4. Find the value of the sum $\sum_{k=0}^{359} k \cdot \cos k^\circ$.
5. The first term of an arithmetic progression is 1 and the sum of the first nine terms equal to 369. The first and the ninth term of a geometric progression coincide with the first and the ninth term of the arithmetic progression. Find the seventh term of the geometric progression.
6. The sequence $a_1, a_2, a_3, \dots, a_{98}$ satisfies the relation $a_{n+1} = a_n + 1$ for $n = 1, 2, 3, \dots, 97$ and has the sum equal to 4949. Evaluate $\sum_{k=1}^{49} a_{2k}$.
7. For an increasing G.P. $a_1, a_2, a_3, \dots, a_n$, if $a_6 = 4a_4$, $a_9 - a_7 = 192$, then the value of $\sum_{i=1}^{\infty} \frac{1}{a_i}$ is
8. In a set of four numbers, the first three are in GP & the last three are in A.P. with common difference 6. If the first number is the same as the fourth, find the four numbers.
9. Find three numbers a, b, c between 2 & 18 such that ;
 - (i) their sum is 25
 - (ii) the numbers $2, a, b$ are consecutive terms of an AP &
 - (iii) the numbers $b, c, 18$ are consecutive terms of a G.P.

10. If the 10th term of an HP is 21 and 21st term of the same HP is 10, then find the 210th term.
11. The pth term T_p of H.P. is $q(p + q)$ and qth term T_q is $p(p + q)$ when $p > 2, q > 2$. Prove that
(a) $T_{p+q} = pq$; (b) $T_{pq} = p + q$; (c) $T_{p+q} > T_{pq}$
12. (a) The harmonic mean of two numbers is 4. The arithmetic mean A & the geometric mean G satisfy the relation $2A + G^2 = 27$. Find the two numbers.
(b) The AM of two numbers exceeds their GM by 15 & HM by 27. Find the numbers.
13. If $A_1, A_2, A_3, \dots, A_{51}$ are arithmetic means inserted between the numbers a and b, then find the value of $\left(\frac{b + A_{51}}{b - A_{51}} \right) - \left(\frac{A_1 + a}{A_1 - a} \right)$.
14. If number of coins earned in nth game is $n2^{n+2} - 2^n$ and total number of coins earned in first 10 games is $10(B \cdot 2^{10} + 1)$, where $B \in \mathbb{N}$, then the value of B is
15. Find the nth term and the sum to n terms of the sequence :
(i) $1 + 5 + 13 + 29 + 61 + \dots$ (ii) $6 + 13 + 22 + 33 + \dots$
16. Sum the following series to n terms and to infinity :
(i) $\frac{1}{1 \cdot 4 \cdot 7} + \frac{1}{4 \cdot 7 \cdot 10} + \frac{1}{7 \cdot 10 \cdot 13} + \dots$ (ii) $\sum_{r=1}^n r(r+1)(r+2)(r+3)$
(iii) $\sum_{r=1}^n \frac{1}{4r^2 - 1}$
17. Prove that : $\tan \alpha + 2 \tan 2\alpha + 4 \tan 4\alpha + 8 \cot 8\alpha = \cot \alpha$.

EXERCISE # 3 (JM)

- Let $a_1, a_2, a_3, \dots$ be terms of an A.P. If $\frac{a_1 + a_2 + \dots + a_p}{a_1 + a_2 + \dots + a_q} = \frac{p^2}{q^2}$, $p \neq q$ then $\frac{a_6}{a_{21}}$ equals-
 (1) $\frac{2}{7}$ (2) $\frac{11}{41}$ (3) $\frac{41}{11}$ (4) $\frac{7}{2}$ [AIEEE-2006]
- If $a_1, a_2, \dots, a_n$ are in H.P., then the expression $a_1a_2 + a_2a_3 + \dots + a_{n-1}a_n$ is equal to-
 (1) na_1a_n (2) $(n-1)a_1a_n$ (3) $n(a_1 - a_n)$ (4) $(n-1)(a_1 - a_n)$ [AIEEE-2006]
- In a geometric progression consisting of positive terms, each term equals the sum of the next two terms. Then the common ratio of this progression equals-
 (1) $\frac{1}{2}\sqrt{5}$ (2) $\sqrt{5}$ (3) $\frac{1}{2}(\sqrt{5}-1)$ (4) $\frac{1}{2}(1-\sqrt{5})$ [AIEEE-2007]
- The first two terms of a geometric progression add up to 12. The sum of the third and the fourth terms is 48. If the terms of the geometric progression are alternately positive and negative, then the first term is
 (1) -4 (2) -12 (3) 12 (4) 4 [AIEEE 2008]
- The sum to infinity of the series $1 + \frac{2}{3} + \frac{6}{3^2} + \frac{10}{3^3} + \frac{14}{3^4} + \dots$ is :-
 (1) 4 (2) 6 (3) 2 (4) 3 [AIEEE-2009]
- A person is to count 4500 currency notes. Let a_n denote the number of notes he counts in the n^{th} minute. If $a_1 = a_2 = \dots = a_{10} = 150$ and $a_{10}, a_{11}, \dots$ are in an AP with common difference -2 , then the time taken by him to count all notes is :-
 (1) 24 minutes (2) 34 minutes (3) 125 minutes (4) 135 minutes [AIEEE-2010]
- A man saves Rs. 200 in each of the first three months of his service. In each of the subsequent months his saving increases by Rs. 40 more than the saving of immediately previous month. His total saving from the start of service will be Rs. 11040 after :-
 (1) 20 months (2) 21 months (3) 18 months (4) 19 months [AIEEE-2011]
- Let a_n be the n^{th} term of an A.P. If $\sum_{r=1}^{100} a_{2r} = \alpha$ and $\sum_{r=1}^{100} a_{2r-1} = \beta$, then the common difference of the A.P. is :
 (1) $\frac{\alpha - \beta}{200}$ (2) $\alpha - \beta$ (3) $\frac{\alpha - \beta}{100}$ (4) $\beta - \alpha$ [AIEEE-2011]

9. **Statement-1** : The sum of the series
 $1 + (1 + 2 + 4) + (4 + 6 + 9) + (9 + 12 + 16) + \dots + (361 + 380 + 400)$ is 8000.
Statement-2 : $\sum_{k=1}^n (k^3 - (k-1)^3) = n^3$, for any natural number n. [AIEEE-2012]
 (1) Statement-1 is true, Statement-2 is false.
 (2) Statement-1 is false, Statement-2 is true.
 (3) Statement-1 is true, Statement-2 is true ; Statement-2 is a correct explanation for Statement-1.
 (4) Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for Statement-1.
10. If 100 times the 100th term of an A.P. with non-zero common difference equals the 50 times its 50th term, then the 150th term of this A.P. is : [AIEEE-2012]
 (1) zero (2) -150
 (3) 150 times its 50th term (4) 150
11. The sum of first 20 terms of the sequence 0.7, 0.77, 0.777,, is : [JEE(Main)-2013]
 (1) $\frac{7}{81}(179 - 10^{-20})$ (2) $\frac{7}{9}(99 - 10^{-20})$
 (3) $\frac{7}{81}(179 + 10^{-20})$ (4) $\frac{7}{9}(99 - 10^{-20})$
12. Let α and β be the roots of equation $px^2 + qx + r = 0$, $p \neq 0$. If p, q, r are in A.P. and $\frac{1}{\alpha} + \frac{1}{\beta} = 4$, then the value of $|\alpha - \beta|$ is: [JEE(Main)-2014]
 (1) $\frac{\sqrt{61}}{9}$ (2) $\frac{2\sqrt{17}}{9}$ (3) $\frac{\sqrt{34}}{9}$ (4) $\frac{2\sqrt{13}}{9}$
13. Three positive numbers form an increasing G.P. If the middle term in this G.P. is doubled, the new numbers are in A.P. Then the common ratio of the G.P. is : [JEE(Main)-2014]
 (1) $\sqrt{2} + \sqrt{3}$ (2) $3 + \sqrt{2}$ (3) $2 - \sqrt{3}$ (4) $2 + \sqrt{3}$
14. If $(10)^9 + 2(11)^1(10)^8 + 3(11)^2(10)^7 + \dots + 10(11)^9 = k(10)^9$, then k is equal to : [JEE(Main)-2014]
 (1) $\frac{121}{10}$ (2) $\frac{441}{100}$ (3) 100 (4) 110
15. If m is the A.M. of two distinct real numbers ℓ and n ($\ell, n > 1$) and G_1, G_2 and G_3 are three geometric means between ℓ and n, then $G_1^4 + 2G_2^4 + G_3^4$ equals- [JEE(Main)-2015]
 (1) $4\ell mn^2$ (2) $4\ell^2 m^2 n^2$ (3) $4\ell^2 mn$ (4) $4\ell m^2 n$

16. If the 2nd, 5th and 9th terms of a non-constant A.P. are in G.P., then the common ratio of this G.P. is:-
[JEE(Main)-2016]
(1) $\frac{7}{4}$ (2) $\frac{8}{5}$ (3) $\frac{4}{3}$ (4) 1
17. If the sum of the first ten terms of the series $\left(1\frac{3}{5}\right)^2 + \left(2\frac{2}{5}\right)^2 + \left(3\frac{1}{5}\right)^2 + 4^2 + \left(4\frac{4}{5}\right)^2 + \dots$, is $\frac{16}{5}m$, then m is equal to :-
[JEE(Main)-2016]
(1) 99 (2) 102 (3) 101 (4) 100
18. If, for a positive integer n, the quadratic equation,
 $x(x+1) + (x+1)(x+2) + \dots + (x+n-1)(x+n) = 10n$
has two consecutive integral solutions, then n is equal to :
[JEE (Main)2017]
(1) 11 (2) 12 (3) 9 (4) 10
19. For any three positive real numbers a, b and c, $9(25a^2 + b^2) + 25(c^2 - 3ac) = 15b(3a + c)$. Then :
[JEE (Main)2017]
(1) a, b and c are in G.P. (2) b, c and a are in G.P.
(3) b, c and a are in A.P. (4) a, b and c are in A.P.
20. Let $a, b, c \in \mathbb{R}$. If $f(x) = ax^2 + bx + c$ is such that $a + b + c = 3$ and $f(x+y) = f(x) + f(y) + xy, \forall x, y \in \mathbb{R}$, then $\sum_{n=1}^{10} f(n)$ is equal to :
[JEE (Main)2017]
(1) 255 (2) 330 (3) 165 (4) 190
21. The sum of all natural numbers 'n' such that $100 < n < 200$ and H.C.F. (91, n) > 1 is
[JEE-MAIN-2019]
(1) 3121 (2) 3303 (3) 3203 (4) 3221
22. The sum $\sum_{k=1}^{20} k \frac{1}{2^k}$ is equal to :
[JEE-MAIN-2019]
(1) $2 - \frac{3}{2^{17}}$ (2) $1 - \frac{11}{2^{20}}$ (3) $2 - \frac{11}{2^{19}}$ (4) $2 - \frac{21}{2^{20}}$
23. If three distinct numbers a, b, c are in G.P. and the equations $ax^2 + 2bx + c = 0$ and $dx^2 + 2ex + f = 0$ have a common root, then which one of the following statements is correct?
[JEE-MAIN-2019]
(1) d, e, f are in A.P. (2) $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in G.P.
(3) $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in A.P. (4) d, e, f are in G.P.

24. Let the sum of the first n terms of a non-constant A.P., $a_1, a_2, a_3, \dots$ be $50n + \frac{n(n-7)}{2}A$, where A is constant. If d is the common difference of this A.P., then the ordered pair (d, a_{50}) is equal to : **[JEE-MAIN-2019]**
 (1) $(A, 50 + 45A)$ (2) $(50, 50 + 46A)$ (3) $(A, 50 + 46A)$ (4) $(50, 50 + 45A)$
25. The sum of the series $1 + 2 \times 3 + 3 \times 5 + 4 \times 7 + \dots$ upto 11^{th} term is : **[JEE-MAIN-2019]**
 (1) 916 (2) 945 (3) 915 (4) 946
26. If the sum and product of the first three terms in an A.P. are 33 and 1155, respectively, then a value of its 11^{th} terms is: **[JEE-MAIN-2019]**
 (1) -36 (2) 25 (3) -35 (4) -25
27. The sum $\frac{3 \times 1^3}{1^2} + \frac{5 \times (1^3 + 2^3)}{1^2 + 2^2} + \frac{7 \times (1^3 + 2^3 + 3^3)}{1^2 + 2^2 + 3^2} + \dots$ upto 10^{th} term, is : **[JEE-MAIN-2019]**
 (1) 680 (2) 660 (3) 600 (4) 620
28. If $a_1, a_2, a_3, \dots, a_n$ are in A.P. and $a_1 + a_4 + a_7 + \dots + a_{16} = 114$, then $a_1 + a_6 + a_{11} + a_{16}$ is equal to : **[JEE-MAIN-2019]**
 (1) 76 (2) 98 (3) 64 (4) 38
29. Let $a_1, a_2, a_3, \dots$ be an A.P. with $a_6 = 2$. Then the common difference of this A.P, which maximises the product $a_1 a_4 a_5$, is : **[JEE-MAIN-2019]**
 (1) $\frac{3}{2}$ (2) $\frac{8}{5}$ (3) $\frac{2}{3}$ (4) $\frac{6}{5}$
30. Let a, b and c be in G.P. with common ratio r , where $a \neq 0$ and $0 < r \leq \frac{1}{2}$. If $3a, 7b$ and $15c$ are the first three terms of an A.P., then the 4^{th} term of this A.P. is : **[JEE-MAIN-2019]**
 (1) $\frac{2}{3}a$ (2) $\frac{7}{3}a$ (3) a (4) $5a$
31. The sum $1 + \frac{1^3 + 2^3}{1+2} + \frac{1^3 + 2^3 + 3^3}{1+2+3} + \dots + \frac{1^3 + 2^3 + 3^3 + \dots + 15^3}{1+2+3+\dots+15} - \frac{1}{2}(1+2+3+\dots+15)$ is equal to : **[JEE-MAIN-2019]**
 (1) 620 (2) 1240 (3) 1860 (4) 660

32. If α and β are the roots of the equation $375x^2 - 25x - 2 = 0$ then $\lim_{n \rightarrow \infty} \sum_{r=1}^n \alpha^r + \lim_{n \rightarrow \infty} \sum_{r=1}^n \beta^r$ is equal to
[JEE-MAIN-2019]
(1) $\frac{21}{346}$ (2) $\frac{1}{12}$ (3) $\frac{29}{358}$ (4) $\frac{7}{116}$
33. Let S_n denote the sum of the first n terms of an A.P. If $S_4 = 16$ and $S_6 = -48$, then S_{10} is equal to
[JEE-MAIN-2019]
(1) -280 (2) -260 (3) -320 (4) -410
34. If $a_1, a_2, a_3, \dots$ are in A.P. such that $a_1 + a_7 + a_{16} = 40$, then the sum of the first 15 terms of this A.P. is :
[JEE-MAIN-2019]
(1) 150 (2) 200 (3) 120 (4) 280
35. If a, b , and c be three distinct real numbers in G.P. and $a + b + c = xb$, then x cannot be :
[JEE-MAIN-2019]
(1) 4 (2) -3 (3) -2 (4) 2
36. Let $a_1, a_2, \dots, a_{30}$ be an A.P., $S = \sum_{i=1}^{30} a_i$ and $T = \sum_{i=1}^{15} a_{(2i-1)}$. If $a_5 = 27$ and $S - 2T = 75$, then a_{10} is equal to :
[JEE-MAIN-2019]
(1) 57 (2) 47 (3) 42 (4) 52
37. Let a, b and c be the 7th, 11th and 13th terms respectively of a non-constant A.P. If these are also the three consecutive terms of a G.P., then $\frac{a}{c}$ is equal to :
[JEE-MAIN-2019]
(1) $\frac{7}{13}$ (2) 4 (3) 2 (4) $\frac{1}{2}$
38. The sum of the following series
 $1 + 6 + \frac{9(1^2 + 2^2 + 3^2)}{7} + \frac{12(1^2 + 2^2 + 3^2 + 4^2)}{9} + \frac{15(1^2 + 2^2 + \dots + 5^2)}{11} + \dots$ up to 15 terms is :
[JEE-MAIN-2019]
(1) 7820 (2) 7510 (3) 7520 (4) 7830
39. The sum of all two digit positive numbers which when divided by 7 yield 2 or 5 as remainder is:
[JEE-MAIN-2019]
(1) 1365 (2) 1356 (3) 1256 (4) 1465

40. Let $a_1, a_2, \dots, a_{10}$ be a G.P. If $\frac{a_3}{a_1} = 25$. Then $\frac{a_9}{a_5}$ equals : [JEE-MAIN-2019]
 (1) 5^3 (2) $2(5^2)$ (3) $4(5^2)$ (4) 5^4
41. The sum of an infinite geometric series with positive terms is 3 and the sum of the cubes of its terms is $\frac{27}{19}$. Then the common ratio of this series is : [JEE-MAIN-2019]
 (1) $\frac{1}{3}$ (2) $\frac{2}{9}$ (3) $\frac{4}{9}$ (4) $\frac{2}{3}$
42. Let α and β be the roots of the quadratic equation $x^2 \sin \theta - x(\sin \theta \cos \theta + 1) + \cos \theta = 0$ ($0 < \theta < 45^\circ$), and $\alpha < \beta$. Then $\sum_{n=0}^{\infty} \left(\alpha^n + \frac{(-1)^n}{\beta^n} \right)$ is equal to: [JEE-MAIN-2019]
 (1) $\frac{1}{1+\cos \theta} - \frac{1}{1-\sin \theta}$ (2) $\frac{1}{1-\cos \theta} - \frac{1}{1+\sin \theta}$
 (3) $\frac{1}{1-\cos \theta} + \frac{1}{1+\sin \theta}$ (4) $\frac{1}{1+\cos \theta} + \frac{1}{1-\sin \theta}$
43. If 19th term of a non-zero A.P. is zero, then its (49th term) : (29th term) is: [JEE-MAIN-2019]
 (1) 2 : 1 (2) 3 : 1 (3) 4 : 1 (4) 1 : 3
44. Let $S_k = \frac{1+2+3+\dots+k}{k}$. If $S_1^2 + S_2^2 + \dots + S_{10}^2 = \frac{5}{12}A$, then A is equal to : [JEE-MAIN-2019]
 (1) 303 (2) 156 (3) 283 (4) 301
45. The product of three consecutive terms of a G.P. is 512. If 4 is added to each of the first and the second of these terms, the three terms now form an A.P. Then the sum of the original three terms of the given G.P. is : [JEE-MAIN-2019]
 (1) 36 (2) 32 (3) 24 (4) 28
46. If the sum of the first 15 terms of the series $\left(\frac{3}{4}\right)^4 + \left(1\frac{1}{2}\right)^3 + \left(2\frac{1}{4}\right)^3 + 3^3 + \left(3\frac{3}{4}\right)^3 + \dots$ is equal to 225 k, then k is equal to: [JEE-MAIN-2019]
 (1) 108 (2) 27 (3) 9 (4) 54

47. The greatest positive integer k , for which $49^k + 1$ is a factor of the sum $49^{125} + 49^{124} + \dots + 49^2 + 49 + 1$, is : **[JEE-MAIN-2020]**
(1) 32 (2) 63 (3) 60 (4) 65
48. Five numbers are in A.P., whose sum is 25 and product is 2520. If one of these five numbers is $-\frac{1}{2}$, then the greatest number amongst them is : **[JEE-MAIN-2020]**
(1) 27 (2) 7 (3) $\frac{21}{2}$ (4) 16
49. Let $a_1, a_2, a_3, \dots$ be a G.P. such that $a_1 < 0$, $a_1 + a_2 = 4$ and $a_3 + a_4 = 16$. If $\sum_{i=1}^9 a_i = 4\lambda$, then λ is equal to : **[JEE-MAIN-2020]**
(1) -171 (2) $\frac{511}{3}$ (3) -513 (4) 171
50. If the sum of the first 40 terms of the series $3 + 4 + 8 + 9 + 13 + 14 + 18 + 19 + \dots$ is $(102)m$, then m is equal to : **[JEE-MAIN-2020]**
(1) 5 (2) 10 (3) 20 (4) 25
51. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be such that for all $x \in \mathbb{R}$ ($2^{1+x} + 2^{1-x}$), $f(x)$ and $(3^x + 3^{-x})$ are in A.P., then the minimum value of $f(x)$ is : **[JEE-MAIN-2020]**
(1) 2 (2) 3 (3) 0 (4) 4
52. If the 10th term of an A.P. is $\frac{1}{20}$ and its 20th term is $\frac{1}{10}$ then the sum of its first 200 terms is **[JEE-MAIN-2020]**
(1) 100 (2) $100\frac{1}{2}$ (3) $50\frac{1}{4}$ (4) 50
53. The product $2^{\frac{1}{4}} \cdot 4^{\frac{1}{16}} \cdot 8^{\frac{1}{48}} \cdot 16^{\frac{1}{128}} \dots$ to ∞ is equal to : **[JEE-MAIN-2020]**
(1) 2 (2) $2^{\frac{1}{2}}$ (3) $2^{\frac{1}{4}}$ (4) 1
54. If $x = \sum_{n=0}^{\infty} (-1)^n \tan^{2n} \theta$ and $y = \sum_{n=0}^{\infty} \cos^{2n} \theta$, for $0 < \theta < \frac{\pi}{4}$, then : **[JEE-MAIN-2020]**
(1) $x(1+y) = 1$ (2) $x(1-y) = 1$ (3) $y(1-x) = 1$ (4) $y(1+x) = 1$

55. Let a_n be the n^{th} term of a G.P. of positive terms. If $\sum_{n=1}^{100} a_{2n+1} = 200$ and $\sum_{n=1}^{100} a_{2n} = 100$ then $\sum_{n=1}^{200} a_n$ is equal to [JEE-MAIN-2020]
(1) 150 (2) 300 (3) 225 (4) 175
56. The sum $\sum_{n=1}^7 \frac{n(n+1)(2n+1)}{4}$ is equal to [JEE-MAIN-2020]
57. The sum $\sum_{k=1}^{20} (1+2+3+\dots+k)$ is _____. [JEE-MAIN-2020]
58. The number of terms common to the two A.P.'s 3, 7, 11,, 407 and 2, 9, 16,, 709 is _____. [JEE-MAIN-2020]



EXERCISE # 4 (JA)

1. For $n = 1, 2, 3, \dots$, let $A_n = \left(\frac{3}{4}\right) - \left(\frac{3}{4}\right)^2 + \left(\frac{3}{4}\right)^3 + \dots + (-1)^{n-1} \left(\frac{3}{4}\right)^n$ and $B_n = 1 - A_n$. Find the smallest natural number n_0 such that $B_n > A_n$ for all $n \geq n_0$. [JEE 2006, 6]

Comprehension (3 questions)

Let V_r denote the sum of the first 'r' terms of an arithmetic progression (A.P.) whose first term is 'r' and the common difference is $(2r - 1)$. Let $T_r = V_{r+1} - V_r - 2$ and $Q_r = T_{r+1} - T_r$ for $r = 1, 2, \dots$

2. The sum $V_1 + V_2 + \dots + V_n$ is
 (A) $\frac{1}{12} n(n+1)(3n^2 - n + 1)$ (B) $\frac{1}{12} n(n+1)(3n^2 + n + 2)$
 (C) $\frac{1}{2} n(2n^2 - n + 1)$ (D) $\frac{1}{3} (2n^3 - 2n + 3)$ [JEE 2007, 4]
3. T_r is always
 (A) an odd number (B) an even number
 (C) a prime number (D) a composite number [JEE 2007, 4]
4. Which one of the following is a correct statement?
 (A) $Q_1, Q_2, Q_3, \dots$ are in A.P. with common difference 5.
 (B) $Q_1, Q_2, Q_3, \dots$ are in A.P. with common difference 6.
 (C) $Q_1, Q_2, Q_3, \dots$ are in A.P. with common difference 11.
 (D) $Q_1 = Q_2 = Q_3 = \dots$ [JEE 2007, 4]

Assertion & Reason :

5. Suppose four distinct positive numbers a_1, a_2, a_3, a_4 are in G.P.
 Let $b_1 = a_1, b_2 = b_1 + a_2, b_3 = b_2 + a_3$ and $b_4 = b_3 + a_4$.
Statement-1 : The numbers b_1, b_2, b_3, b_4 are neither in A.P. nor in G.P.
Statement-2 : The numbers b_1, b_2, b_3, b_4 are in H.P.
 (A) Statement-1 is True, Statement-2 is True; statement-2 is a correct explanation for statement-1
 (B) Statement-1 is True, Statement-2 is True; statement-2 is **NOT** a correct explanation for statement-1
 (C) Statement-1 is True, Statement-2 is False
 (D) Statement-1 is False, Statement-2 is True [JEE 2008, 3 (-1)]
6. If the sum of first n terms of an A.P. is cn^2 , then the sum of squares of these n terms is
 (A) $\frac{n(4n^2 - 1)c^2}{6}$ (B) $\frac{n(4n^2 + 1)c^2}{3}$ (C) $\frac{n(4n^2 - 1)c^2}{3}$ (D) $\frac{n(4n^2 + 1)c^2}{6}$ [JEE 2009, 3 (-1)]

7. For $0 < \theta < \frac{\pi}{2}$, the solution(s) of $\sum_{m=1}^6 \operatorname{cosec}\left(\theta + \frac{(m-1)\pi}{4}\right) \operatorname{cosec}\left(\theta + \frac{m\pi}{4}\right) = 4\sqrt{2}$ is (are) -
[JEE 2009, 4]
(A) $\frac{\pi}{4}$ (B) $\frac{\pi}{6}$ (C) $\frac{\pi}{12}$ (D) $\frac{5\pi}{12}$
8. Let $a_1, a_2, a_3, \dots, a_{11}$ be real numbers satisfying $a_1 = 15$, $27 - 2a_2 > 0$ and $a_k = 2a_{k-1} - a_{k-2}$ for $k = 3, 4, \dots, 11$. If $\frac{a_1^2 + a_2^2 + \dots + a_{11}^2}{11} = 90$, then the value of $\frac{a_1 + a_2 + \dots + a_{11}}{11}$ is equal to
[JEE 2010, 3+3]
9. Let $a_1, a_2, a_3, \dots, a_{100}$ be an arithmetic progression with $a_1 = 3$ and $S_p = \sum_{i=1}^p a_i, 1 \leq p \leq 100$. For any integer n with $1 \leq n \leq 20$, let $m = 5n$. If $\frac{S_m}{S_n}$ does not depend on n , then a_2 is
[JEE 2011, 4]
10. Let $a_1, a_2, a_3, \dots$ be in harmonic progression with $a_1 = 5$ and $a_{20} = 25$. The least positive integer n for which $a_n < 0$ is
[JEE 2012, 3 (-1)]
(A) 22 (B) 23 (C) 24 (D) 25
11. Let $S_n = \sum_{k=1}^{4n} (-1)^{\frac{k(k+1)}{2}} k^2$. Then S_n can take value(s)
[JEE-Advanced 2013, 4, (-1)]
(A) 1056 (B) 1088 (C) 1120 (D) 1332
12. A pack contains n cards numbered from 1 to n . Two consecutive numbered cards are removed from the pack and the sum of the numbers on the remaining cards is 1224. If the smaller of the numbers on the removed cards is k , then $k - 20 =$
[JEE-Advanced 2013, 4, (-1)]
13. Let a, b, c be positive integers such that $\frac{b}{a}$ is an integer. If a, b, c are in geometric progression and the arithmetic mean of a, b, c is $b + 2$, then the value of $\frac{a^2 + a - 14}{a + 1}$ is
[JEE(Advanced)-2014, 3]
14. Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first eleven terms is $6 : 11$ and the seventh term lies in between 130 and 140, then the common difference of this A.P. is
[JEE 2015, 4M, -0M]
15. The sides of a right angled triangle are in arithmetic progression. If the triangle has area 24, then what is the length of its smallest side?
[JEE 2017, 3M, -0M]
16. Let AP $(a; d)$ denote the set of all the terms of an infinite arithmetic progression with first term a and common difference $d > 0$. If
 $AP(1; 3) \cap AP(2; 5) \cap AP(3; 7) = AP(a; d)$
then $a + d$ equals____
[JEE(Advanced)-2019]

EXERCISE # 5

- If $x = \sum_{n=0}^{\infty} a^n$, $y = \sum_{n=0}^{\infty} b^n$, $z = \sum_{n=0}^{\infty} c^n$ where a, b, c are in A.P. and $|a| < 1$, $|b| < 1$, $|c| < 1$, then x, y, z are in- [AIEEE 2005]
 (A) HP (B) Arithmetic - Geometric Progression
 (C) AP (D) GP
- If a, b, c are distinct positive real in H.P., then the value of the expression, $\frac{b+a}{b-a} + \frac{b+c}{b-c}$ is equal to
 (A) 1 (B) 2 (C) 3 (D) 4
- Along a road lies an odd number of stones placed at intervals of 10 m. These stones have to be assembled around the middle stone. A person can carry only one stone at a time. A man carried out the job starting with the stone in the middle, carrying stones in succession, thereby covering a distance of 4.8 km. Then the number of stones is
 (A) 15 (B) 29 (C) 31 (D) 35
- If $S = 1^2 + 3^2 + 5^2 + \dots + (99)^2$ then the value of the sum $2^2 + 4^2 + 6^2 + \dots + (100)^2$ is
 (A) $S + 2550$ (B) $2S$ (C) $4S$ (D) $S + 5050$
- In an A.P. with first term 'a' and the common difference d ($a, d \neq 0$), the ratio 'p' of the sum of the first n terms to sum of n terms succeeding them does not depend on n . Then the ratio $\frac{a}{d}$ and the ratio 'p', respectively are
 (A) $\frac{1}{2}, \frac{1}{4}$ (B) $2, \frac{1}{3}$ (C) $\frac{1}{2}, \frac{1}{3}$ (D) $\frac{1}{2}, 2$
- The arithmetic mean of the nine numbers in the given set $\{9, 99, 999, \dots, 999999999\}$ is a 9 digit number N , all whose digits are distinct. The number N does not contain the digit
 (A) 0 (B) 2 (C) 5 (D) 9
- If for an A.P. $a_1, a_2, a_3, \dots, a_n, \dots, a_1 + a_3 + a_5 = -12$ and $a_1 a_2 a_3 = 8$ then the value of $a_2 + a_4 + a_6$ equals
 (A) -12 (B) -16 (C) -18 (D) -21
- An H.M. is inserted between the number $1/3$ and an unknown number. If we diminish the reciprocal of the inserted number by 6, it is the G.M. of the reciprocal of $1/3$ and that of the unknown number. If all the terms of the respective H.P. are distinct then
 (A) the unknown number is 27 (B) the unknown number is $1/27$
 (C) the H.M. is 15 (D) the G.M. is 21

9. If $x \in \mathbb{R}$, the numbers $(5^{1+x} + 5^{1-x})$, $a/2$, $(25^x + 25^{-x})$ form an A.P. then 'a' must lie in the interval
(A) $[1, 5]$ (B) $[2, 5]$ (C) $[5, 12]$ (D) $[12, \infty)$
10. If the sum of the first 11 terms of an arithmetic progression equals that of the first 19 terms, then the sum of its first 30 terms, is
(A) equal to 0 (B) equal to -1
(C) equal to 1 (D) non unique
11. Let $s_1, s_2, s_3, \dots$ and $t_1, t_2, t_3, \dots$ are two arithmetic sequences such that $s_1 = t_1 \neq 0$; $s_2 = 2t_2$ and $\sum_{i=1}^{10} s_i = \sum_{i=1}^{15} t_i$. Then the value of $\frac{s_2 - s_1}{t_2 - t_1}$ is
(A) $8/3$ (B) $3/2$ (C) $19/8$ (D) 2
12. If $\frac{1+3+5+\dots \text{upto } n \text{ terms}}{4+7+10+\dots \text{upto } n \text{ terms}} = \frac{20}{7 \log_{10} x}$
and $n = \log_{10} x + \log_{10} x^{\frac{1}{2}} + \log_{10} x^{\frac{1}{4}} + \log_{10} x^{\frac{1}{8}} + \dots + \infty$, then x is equal to
(A) 10^3 (B) 10^5 (C) 10^6 (D) 10^7
13. Let a_n , $n \in \mathbb{N}$ is an A.P. with common difference 'd' and all whose terms are non-zero. If n approaches infinity, then the sum $\frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \dots + \frac{1}{a_n a_{n+1}}$ will approach
(A) $\frac{1}{a_1 d}$ (B) $\frac{2}{a_1 d}$ (C) $\frac{1}{2a_1 d}$ (D) $a_1 d$
14. The sum of the first three terms of an increasing G.P. is 21 and the sum of their squares is 189. Then the sum of its first n terms is
(A) $3(2^n - 1)$ (B) $12\left(1 - \frac{1}{2^n}\right)$ (C) $6\left(1 - \frac{1}{2^n}\right)$ (D) $6(2^n - 1)$

15. If $a \neq 1$ and $\ln a^2 + (\ln a^2)^2 + (\ln a^2)^3 + \dots = 3(\ln a + (\ln a)^2 + (\ln a)^3 + (\ln a)^4 + \dots)$. then 'a' is equal to
(A) $e^{1/5}$ (B) $\sqrt{e}$ (C) $\sqrt[3]{e}$ (D) $\sqrt[4]{e}$
16. For which positive integers n is the ratio, $\frac{\sum_{k=1}^n k^2}{\sum_{k=1}^n k}$ an integer?
(A) odd n only
(B) even n only
(C) $n = 1 + 6k$ only, where $k \geq 0$ and $k \in \mathbb{I}$
(D) $n = 1 + 3k$, integer $k \geq 0$
17. Let $a_1, a_2, a_3, \dots$ and $b_1, b_2, b_3, \dots$ be arithmetic progressions such that $a_1 = 25$, $b_1 = 75$ and $a_{100} + b_{100} = 100$. Then which of the following is true.
(A) the difference between successive terms in progression 'a' is opposite of the difference in progression 'b'.
(B) $a_n + b_n = 100$ for any n.
(C) $(a_1 + b_1), (a_2 + b_2), (a_3 + b_3), \dots$ are in A.P.
(D) $\sum_{r=1}^{100} (a_r + b_r) = 10000$
18. If $\sin(x - y)$, $\sin x$ and $\sin(x + y)$ are in H.P., then $\left| \sin x \cdot \sec \frac{y}{2} \right| =$
(A) 2 (B) $\sqrt{2}$ (C) $-\sqrt{2}$ (D) -2
19. If $A = \sin \frac{2\pi}{7} + \sin \frac{4\pi}{7} + \sin \frac{8\pi}{7}$ and $B = \cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{8\pi}{7}$ then $\sqrt{A^2 + B^2}$ is equal to
(A) 1 (B) $\sqrt{2}$ (C) 2 (D) $\sqrt{3}$
20. If m and n are positive integers satisfying
 $1 + \cos 2\theta + \cos 4\theta + \cos 6\theta + \cos 8\theta + \cos 10\theta = \frac{\cos m\theta \cdot \sin n\theta}{\sin \theta}$ then (m + n) is equal to
(A) 9 (B) 10 (C) 11 (D) 12
21. Let $y = \frac{\cos x + \cos 2x + \cos 3x + \cos 4x + \cos 5x + \cos 6x + \cos 7x}{\sin x + \sin 2x + \sin 3x + \sin 4x + \sin 5x + \sin 6x + \sin 7x}$, then which of the following hold good?
(A) The value of y when $x = \pi/8$ is not defined.
(B) The value of y when $x = \pi/16$ is 1.
(C) The value of y when $x = \pi/32$ is $\sqrt{2} - 1$.
(D) The value of y when $x = \pi/48$ is $2 + \sqrt{3}$.

EXERCISE # 6

- If the first 3 consecutive terms of a geometrical progression are the real roots of the equation $2x^3 - 19x^2 + 57x - 54 = 0$ find the sum to infinite number of terms of G.P.
- Find the sum of the infinite series $\frac{1.3}{2} + \frac{3.5}{2^2} + \frac{5.7}{2^3} + \frac{7.9}{2^4} + \dots \infty$.
- Let $S = \sum_{n=1}^{99} \frac{5^{100}}{(25)^n + 5^{100}}$. Find $[S]$. Where $[y]$ denotes largest integer less than or equal to y .
- If $3^{2\sin 2x-1}$, 14 , $3^{4-2\sin 2x}$ form first three terms of an A.P., then find the sum $1 + \sin 2x + \sin^2 2x + \dots \infty$.
- If $\tan\left(\frac{\pi}{12} - x\right)$, $\tan \frac{\pi}{12}$, $\tan\left(\frac{\pi}{12} + x\right)$ in order are three consecutive terms of a G.P. then sum of all the solutions in $[0, 314]$ is $k\pi$. Find the value of k .
- If the roots of $10x^3 - cx^2 - 54x - 27 = 0$ are in harmonic progression, then find c and all the roots.
- In a GP the ratio of the sum of the first eleven terms to the sum of the last eleven terms is $1/8$ and the ratio of the sum of all the terms without the first nine to the sum of all the terms without the last nine is 2 . Find the number of terms in the GP.
- In the quadratic equation $A(\sqrt{3} - \sqrt{2})x^2 + \frac{B}{(\sqrt{3} + \sqrt{2})}x + C = 0$ with α, β as its roots
If $A = (49 + 20\sqrt{6})^{1/4}$; $B = \text{sum of the infinite G.P. as } 8\sqrt{3} + \frac{8\sqrt{6}}{\sqrt{3}} + \frac{16}{\sqrt{3}} + \dots \infty$
and $|\alpha - \beta| = (6\sqrt{6})^k$ where $k = \log_6 10 - 2 \log_6 \sqrt{5} + \log_6 \sqrt{(\log_6 18 + \log_6 72)}$,
then find the value of C .
- If a, b, c, d, e be 5 numbers such that a, b, c are in AP ; b, c, d are in GP & c, d, e are in HP then:
 - Prove that a, c, e are in GP.
 - Prove that $e = (2b - a)^2 / a$
 - If $a = 2$ & $e = 18$, find all possible values of b, c, d .

10. Prove that the average of the numbers $n \sin n^\circ$, $n = 2, 4, 6, \dots, 180$, is $\cot 1^\circ$.
11. If one AM 'a' and two GM's p and q be inserted between any two given numbers then show that $p^3 + q^3 = 2apq$.
12. Find the sum of the n terms of the sequence $\frac{1}{1+1^2+1^4} + \frac{2}{1+2^2+2^4} + \frac{3}{1+3^2+3^4} + \dots$
13. Let $k = 1^\circ$, then prove that $\sum_{n=0}^{88} \frac{1}{\cos nk \cdot \cos(n+1)k} = \frac{\cos k}{\sin^2 k}$
14. If $f(\theta) = \sum_{n=1}^6 \operatorname{cosec}\left(\theta + \frac{(n-1)\pi}{4}\right) \operatorname{cosec}\left(\theta + \frac{n\pi}{4}\right)$, where $0 < \theta < \frac{\pi}{2}$, then find the minimum value of $f(\theta)$.
15. Let $x_1 = \prod_{r=1}^5 \cos \frac{r\pi}{11}$ and $x_2 = \sum_{r=1}^5 \cos \frac{r\pi}{11}$, then show that $x_1 \cdot x_2 = \frac{1}{64} \left(\operatorname{cosec} \frac{\pi}{22} - 1 \right)$, where Π denotes the continued product.

ANSWER KEY

EXERCISE # 1

- | | | | | | | |
|-------|-------|-------|-------|-------|-------|-------|
| 1. D | 2. C | 3. D | 4. A | 5. A | 6. A | 7. B |
| 8. C | 9. B | 10. C | 11. C | 12. C | 13. C | 14. A |
| 15. B | 16. D | 17. B | | | | |

EXERCISE # 2

- | | | | | |
|--|---|--------------------|---------------------------|-------|
| 1. (i) $(14n - 6)/(8n + 23)$ | 2. 12 | 3. $n = 14$ | 4. -180 | 5. 27 |
| 6. 2499 | 7. 2 | 8. $(8, -4, 2, 8)$ | 9. $a = 5, b = 8, c = 12$ | 10. 1 |
| 12. (a) 6,3 (b) 120, 30 | 13. 102 | 14. 7 | | |
| 15. (i) $2^{n+1} - 3; 2^{n+2} - 4 - 3n$ | (ii) $n^2 + 4n + 1; (1/6)n(n+1)(2n+13) + n$ | | | |
| 16. (i) $S_n = (1/24) - [1/\{6(3n+1)(3n+4)\}]$; $S_\infty = 1/24$ | (ii) $(1/5)n(n+1)(n+2)(n+3)(n+4)$ | (iii) $n/(2n+1)$ | | |

EXERCISE # 3 (JM)

- | | | | | | | |
|------------|-------------|-----------|-------|-------|-------|-------|
| 1. 2 | 2. 2 | 3. 3 | 4. 2 | 5. 4 | 6. 2 | 7. 2 |
| 8. 3 | 9. 3 | 10. 1 | 11. 3 | 12. 4 | 13. 4 | 14. 3 |
| 15. 4 | 16. 3 | 17. 3 | 18. 1 | 19. 3 | 20. 2 | 21. 1 |
| 22. 3 | 23. 3 | 24. 3 | 25. 4 | 26. 4 | 27. 2 | 28. 1 |
| 29. 2 | 30. 3 | 31. 1 | 32. 2 | 33. 3 | 34. 2 | 35. 4 |
| 36. 4 | 37. 2 | 38. 1 | 39. 2 | 40. 4 | 41. 4 | 42. 3 |
| 43. 2 | 44. 1 | 45. 4 | 46. 2 | 47. 2 | 48. 4 | 49. 1 |
| 50. 3 | 51. 2 | 52. 2 | 53. 2 | 54. 3 | 55. 1 | |
| 56. 504.00 | 57. 1540.00 | 58. 14.00 | | | | |

EXERCISE # 4 (JA)


- | | | | | |
|--------------|--------|-------|-----------|-------|
| 1. $n_0 = 6$ | 2. B | 3. D | 4. B | 5. C |
| 6. C | 7. C,D | 8. 0 | 9. 9 or 3 | 10. D |
| 11. A,D | 12. 5 | 13. 4 | 14. 9 | 15. 6 |
| 16. 157.00 | | | | |

EXERCISE # 5

- | | | | | |
|---------|-------------|-------|-------|-------|
| 1. A | 2. B | 3. C | 4. D | 5. C |
| 6. A | 7. D | 8. B | 9. D | 10. A |
| 11. C | 12. B | 13. A | 14. A | 15. D |
| 16. D | 17. A,B,C,D | 18. B | 19. B | 20. C |
| 21. B,D | | | | |

EXERCISE # 6

- | | | | | |
|---|---------------------------------|-----------------|------|---------|
| 1. $\frac{27}{2}$ | 2. 23 | 3. 49 | 4. 2 | 5. 4950 |
| 6. $C = 9 ; (3, -3/2, -3/5)$ | 7. $n = 38$ | 8. 128 | | |
| 9. (iii) $b = 4, c = 6, d = 9$ or $b = -2, c = -6, d = -18$ | 12. $\frac{n(n+1)}{2(n^2+n+1)}$ | 14. $2\sqrt{2}$ | | |



Result JEE Main

Result Highlights 2019



100 Percentile
HIMANSHU GAURAV SINGH
2019 (*SDCCP)



99.99 Percentile
GAURAV KRISHAN GUPTA
2020 (DLP)



99.98 Percentile
SARTHAK ROUT
2020 (CCP)



99.98 Percentile
VIBHAV AGGARWAL
2019 (CCP)



99.97 Percentile
RITVIK GUPTA
2020 (DLP)



99.97 Percentile
BHAVYA JAIN
2020 (CCP)



99.96 Percentile
AYUSH PATTNAIK
2019 (CCP)



99.96 Percentile
SAYANTAN DHAR
2020 (DLP)

Result Highlights 2018

AIR
29



Sarthak Behera

AIR
36



Shashank Agrawal

AIR
46



Shreya Pathak

AIR
48



Raaghav Raaj

AIR
2
SC



Aayush Kadam

AIR
18
OBC



Gunjan

Result Highlights 2017

AIR
12



Abhay Goyal

AIR
14




Piyush Tibarwal

AIR
20



Abhinav Kumar

AIR
38



Tushar Gautam

AIR
49



Saurav Sahoo

AIR
60



Pratyush

Mentored by Team

AIR
4



Kushal Babel
2014

AIR
15



Gourika Bang
2015

AIR
21



Amit Malav
2013

AIR
24



Suraj Sanjay Jog
2012

AIR
38



Chitraang Murdia
2014

AIR
58



Pulkit Tandon
2012

AIR
68




Kapil Shobhnani
2016

AIR
70



Utkarsh Sinha
2015

AIR
80



S John S
2015