Table of 2D and 3D spectra of FUS-LC fibrils. Data files can be viewed with Sparky software, which is freely available at http://www.cgl.ucsf.edu/home/sparky.

Sample	Spectrum	Data file	NMR parameters	Total time	Processing
U-FUS-LC, 10 mg	2D CC	CC-2D-U-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 17.0 kHz; na = 240; τ_{pd} = 2.5 s; t_{1max} = 8.6 ms; t_{1inc} = 24 µs; τ_{dwell} = 15 µs; τ_{acq} = 15.4 ms; τ_{HC} =1.5 ms; τ_{DARR} = 25 ms; ν_{1H} = 85 kHz	120 h	GB = 75 Hz in t_1; GB = 75 Hz in t_2
U-FUS-LC, 10 mg	2D NCA	NCA-2D-U-FUS.ucsf	B_0 = 21.1 T; ν_{MAS} = 11 kHz; na = 64; τ_{pd} = 2.0 s; t_{1max} = 12.7 ms; t_{1inc} = 201.6 µs; τ_{dwell} = 10 µs; τ_{acq} = 20.5 ms; τ_{HC} = 1 ms; τ_{NC} = 4 ms; ν_{1H} = 85 kHz; ν_{1C} = 42 kHz, ν_{1N} = 31 kHz, and ν_{0C} = 53 ppm during τ_{NC}	5 h	GB = 10 Hz in t_1; GB = 50 Hz in t_2
U-FUS-LC, 10 mg	2D HC INEPT	HC-2D-U-FUS.ucsf	B_0 = 14.1 T; ν_{MAS}= 12 kHz; na = 32; τ_{pd} = 2.0 s; t_{1max} = 10 ms; t_{1inc} = 50 µs; τ_{dwell} = 15 µs; τ_{acq} = 15.4 ms; ν_{1H} = 13 kHz	7 h	GB = 10 Hz in t_1: GB = 50 Hz in t_2
U-FUS-LC, 10 mg	2D HN INEPT	HN-2D-U-FUS.ucsf	B_0 = 17.5 T; ν_{MAS}=17.0 kHz; na = 64; τ_{pd} = 2.0 s; t_{1max} = 14 ms; t_{1inc} = 200 µs; τ_{dwell} = 15 µs; τ_{acq} = 30.7 ms; ν_{1H} = 38 kHz	5 h	GB = 0 Hz in t_1: GB = 0 Hz in t_2
U-FUS-LC, 10 mg	3D NCACX	NCACX-3D-U-FUS.ucsf	B_0 = 21.1 T; ν_{MAS} = 13.8 kHz; na = 8; τ_{pd} = 2.0 s; t_{1max} = 12 ms; t_{1inc} = 222 µs; t_{2max} = 6.8 ms; t_{2inc} = 109 µs; τ_{dwell} = 5 µs; τ_{acq} = 10.2 ms; τ_{HN} = 1 ms; τ_{NC} = 4 ms; τ_{DARR} = 50 ms; ν_{1H} = 85 kHz; ν_{1C} = 22 kHz, ν_{1N} = 36 kHz, and ν_{0C} = 53 ppm during τ_{NC}	60 h	GB = 15 Hz in t_1; GB = 25 Hz in t_2; GB = 25 Hz in t_3
U-FUS-LC, 10 mg	3D NCOCX	NCOCX-3D-U-FUS.ucsf	B_0 = 21.1 T; ν_{MAS} = 13.8 kHz; na = 16; τ_{pd} = 2.0 s; t_{1max} = 12 ms; t_{1inc} = 222 µs; t_{2max} =9.1 ms; t_{2inc} = 222 µs; τ_{dwell} = 5 µs; τ_{acq} = 10.2 ms; τ_{HN} = 1 ms; τ_{NC} = 4 ms; τ_{DARR} = 50 ms; ν_{1H} = 83 kHz; ν_{1C} = 52 kHz, ν_{1N} = 39 kHz, and ν_{0C} = 175 ppm during τ_{NC}	79 h	GB = 15 Hz in t_1; GB = 25 Hz in t_2; GB = 25 Hz in t_3
U-FUS-LC, 10 mg	3D CONCA	CONCA-3D-U-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 12 kHz; na = 16; τ_{pd} = 2.0 s; t_{1max} = 5.9 ms; t_{1inc} = 196 µs; t_{2max} = 11.8 ms; t_{2inc} = 196 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{CN} = 4 ms; τ_{NC} = 4 ms; ν_{1H} = 70 kHz; ν_{1C} = 29 kHz, ν_{1N} = 17 kHz, and ν_{0C} = 168 ppm during τ_{CN}; ν_{1C} = 29 kHz, ν_{1N} = 17 kHz, and ν_{0C} = 60 ppm during τ_{NC}	68 h	GB = 25 Hz in t_1; GB = 15 Hz in t_2; GB = 25 Hz in t_3
U-FUS-LC, 10 mg	3D CANCX	CANCX-3D-U-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 12 kHz; na = 256; τ_{pd} = 2.0 s; t_{1max} = 4.0 ms; t_{1inc} = 165.2 µs; t_{2max}= 6.6 ms; t_{2inc} = 165.2 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{CN} = 4 ms; τ_{NC} = 4 ms; τ_{DARR} = 60 ms; ν_{1H} = 85 kHz; ν_{1C} = 26 kHz, ν_{1N} = 14 kHz, and ν_{0C} = 56 ppm during τ_{CN}; ν_{1C} = 16 kHz, ν_{1N} = 4 kHz, and ν_{0C} = 175 ppm during τ_{NC}	578 h	GB = 50 Hz in t_1; GB = 30 Hz in t_2; GB = 50 Hz in t_3
N112-FUS-LC, 6 mg	2D CC	CC-2D-N112-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 13.6 kHz; na = 32; τ_{pd} = 3.0 s; t_{1max} = 5.5 ms; t_{1inc} = 21.6 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{DARR} = 50 ms; ν_{1H} = 93 kHz	14 h	GB = 75 Hz in t_1: GB = 75 Hz in t_2
N112-FUS-LC, 6 mg	2D NCA	NCA-2D-N112-FUS.ucsf	B_0 = 14.1 T; ν_{MAS}=13.6 kHz; na = 384; τ_{pd} = 2.2 s; t_{1max} = 12.4 ms; t_{1inc} = 129.6 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HN} = 1.5 ms; τ_{NC} = 4 ms; ν_{1H} = 93 kHz; ν_{1C} = 34 kHz, ν_{1N} = 21 kHz, and ν_{0C} = 55 ppm during τ_{NC}	45 h	GB = 0 Hz in t_1: GB = 75 Hz in t_2
N112-FUS-LC, 6 mg	3D NCACX	NCACX-3D-N112-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 17 kHz; na = 32; τ_{pd} = 1.5 s; t_{1max} = 6.9 ms; t_{1inc} = 172.8 µs; t_{2max} = 5.2 ms; t_{2inc} = 64.8 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{NC} = 5 ms; ; τ_{DARR} = 50 ms; ν_{1H} = 93 kHz; ν_{1C} = 43 kHz, ν_{1N} = 26 kHz, and ν_{0C} = 51 ppm during τ_{NC};	170 h	GB = 10 Hz in t_1; GB = 10 Hz in t_2; GB = 100 Hz in t_3
N112-FUS-LC, 6 mg	3D CONCA	CONCA-3D-N112-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 17 kHz; na = 128; τ_{pd} = 2.0 s; t_{1max} = 4.4 ms; t_{1inc} = 259.2 µs; t_{2max} = 6.9 ms; t_{2inc} = 172.8 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{CN} = 4 ms; τ_{NC} = 5 ms; ν_{1H} = 93 kHz; ν_{1C} = 43 kHz, ν_{1N} = 26 kHz, and ν_{0C} = 175 ppm during τ_{CN}; ν_{1C} = 43 kHz, ν_{1N} = 26 kHz, and ν_{0C} = 51 ppm during τ_{NC}	192 h	GB = 10 Hz in t_1; GB = 10 Hz in t_2; GB = 100 Hz in t_3

C112-FUS-LC, 4 mg	2D CC	CC-2D-C112-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 13.6 kHz; na = 192; τ_{pd} = 2.2 s; t_{1max} = 4 ms; t_{1inc} = 22.4 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{DARR} = 50 ms; ν_{1H} = 78 kHz	42 h	GB = 75 Hz in t_1: GB = 75 Hz in t_2
C112-FUS-LC, 4 mg	2D NCA	NCA-2D-C112-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 13.6 kHz; na = 2880; τ_{pd} = 2.2 s; t_{1max} = 8.6 ms; t_{1inc} = 134.4 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HN} = 1.5 ms; τ_{NC} = 4 ms; ν_{1H} = 78 kHz; ν_{1C} = 34 kHz, ν_{1N} = 21 kHz, and ν_{0C} = 55 ppm during τ_{NC}	225 h	GB = 50 Hz in t_1: GB = 75 Hz in t_2
N60-FUS-LC, 4 mg	2D CC	CC-2D-N60-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 13.6 kHz; na = 128; τ_{pd} = 2.0 s; t_{1max} = 5.4 ms; t_{1inc} = 21.2 µs; τ_{dw} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{DARR} = 50 ms; ν_{1H} = 93 kHz	36 h	GB = 0 Hz in t_1: GB = 75 Hz in t_2
N60-FUS-LC, 4 mg	2D NCA	NCA-2D-N60-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 17.0 kHz; na = 512; τ_{pd} = 2.0 s; t_{1max} = 5.4 ms; t_{1inc} = 84 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HN} = 1.5 ms; τ_{NC} = 4 ms; ν_{1H} = 83 kHz; ν_{1C} = 24 kHz, ν_{1N} = 7 kHz, and ν_{0C} = 53 ppm during τ_{NC}	36 h	GB = 55 Hz in t_1: GB = 75 Hz in t_2
C60-FUS-LC, 3 mg	2D CC	CC-2D-C60-FUS.ucsf	B_0 = 14.1 T; ν_{MAS} = 13.6 kHz; na = 192; τ_{pd} = 2.0 s; t_{1max} = 5.4 ms; t_{1inc} = 21.2 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HC} = 1.5 ms; τ_{DARR} = 50 ms; ν_{1H} = 90 kHz	55 h	GB = 0 Hz in t_1: GB = 75 Hz in t_2
C60-FUS-LC, 3 mg	2D NCA	NCA-2D-C60-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 17.0 kHz; na = 896; τ_{pd} = 2.0 s; t_{1max} = 5.4 ms; t_{1inc} = 84 µs; τ_{dwell} = 15 µs; τ_{acq} = 7.7 ms; τ_{HN} = 1.5 ms; τ_{NC} = 4 ms; ν_{1H} = 83 kHz; ν_{1C} = 24 kHz, ν_{1N} = 7 kHz, and ν_{0C} = 53 ppm during τ_{NC}	64 h	GB = 55 Hz in t_1: GB = 75 Hz in t_2
U-FUS-LC, 10 mg	^{15}N-BARE	BAREn-2D-U-FUS.ucsf (n=1...6)	B_0 = 17.5 T; ν_{MAS} = 17 kHz; na = 384; τ_{pd} = 2.0 s; t_{1max} = 8.9 ms; t_{1inc} = 120 µs; τ_{dwell} = 15 µs; τ_{acq} = 5.5 ms; τ_{HN} = 1.5 ms; $\tau_{NCfsTEDOR}$ = 0.7 ms; ν_{1H} = 85 kHz during acquisition and t_1 evolution; ν_{0C} = 56 ppm during τ_{NC}; τ_{SD} = 30 ms; 28.2 ms constant-time recoupling period; ν_{NBARE} = 25.5 kHz; 6 periods of recoupling with 5.6 ms increment between blocks; ν_{1H} = 100 kHz during constant time recoupling block	32 h per 2D spectrum	GB = 125 Hz in t_1: GB = 60 Hz in t_2
2-Glyc-FUS-LC, 10 mg	^{15}N-BARE	BAREn-2D-2GLYC-FUS.ucsf (n=1...8)	B_0 = 17.5 T; ν_{MAS} = 12 kHz; na = 192; τ_{pd} = 2.3 s; t_{1max} = 7.0 ms; t_{1inc} = 100.8 µs; τ_{dwell} = 15 µs; τ_{acq} = 5.5 ms; τ_{HN} = 1.5 ms; τ_{NC} = 4 ms; ν_{1H} = 85 kHz; ν_{1C} = 28 kHz, ν_{1N} = 16 kHz; and ν_{0C} = 53 ppm during τ_{NC}; 28 ms constant-time recoupling period; ν_{NBARE} = 18 kHz; 8 periods of recoupling with 4 ms increment between blocks	17 h per 2D spectrum	GB = 0 Hz in t_1: GB = 0 Hz in t_2
2-Glyc-FUS-LC, 10 mg	3D NCACX	NCACX-3D-2GLYC-FUS.ucsf	B_0 = 21.1 T; ν_{MAS} = 13.95 kHz; na = 16; τ_{pd} = 2.0 s; t_{1max} = 8.8 ms; t_{1inc} = 180 µs; t_{2max} = 6.1 ms; t_{2inc} = 105 µs; τ_{dwell} = 5 µs; τ_{acq} = 12.0 ms; τ_{HN} = 0.8 ms; τ_{NC} = 6 ms; τ_{DARR} = 400 ms; ν_{1H} = 83.3 kHz; ν_{1C} = 22 kHz, ν_{1N} = 36 kHz, and ν_{0C} = 56 ppm during τ_{NC}	101 h	GB = 15 Hz in t_1; GB = 25 Hz in t_2; GB = 25 Hz in t_3
2-Glyc-FUS-LC, 10 mg	2D CC	CC-2D-2GLYC-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 11.7 kHz; na = 80; τ_{pd} = 1.5 s; t_{1max} = 7.0 ms; t_{1inc} = 22.4 µs; τ_{dwell} = 15 µs; τ_{acq} = 15.4 ms; τ_{HC} = 1.2 ms; τ_{DARR} = 300 ms; ν_{1H} = 85 kHz	21 h	GB = 50 Hz in t_1: GB = 50 Hz in t_2
1,3-Glyc-FUS-LC, 10 mg	3D NCOCX	NCOCX-3D-13GLYC-FUS.ucsf	B_0 = 21.1 T; ν_{MAS} = 13.8 kHz; na = 24; τ_{pd} = 2.0 s; t_{1max} = 9.1 ms; t_{1inc} = 190 µs; t_{2max} = 7.5 ms; t_{2inc} = 166 µs; τ_{dwell} = 5 µs; τ_{acq} = 12.0 ms; τ_{HN} = 0.6 ms; τ_{NC} = 5 ms; τ_{DARR} = 400 ms; ν_{1H} = 83.3 kHz; ν_{1C} = 35 kHz, ν_{1N} = 48 kHz, and ν_{0C} = 176 ppm during τ_{NC}	115 h	GB = 15 Hz in t_1; GB = 25 Hz in t_2; GB = 25 Hz in t_3
1,3-Glyc-FUS-LC, 10 mg	2D CC	CC-2D-13GLYC-FUS.ucsf	B_0 = 17.5 T; ν_{MAS} = 12.0 kHz; na = 48; τ_{pd} = 1.5 s; t_{1max} = 9.0 ms; t_{1inc} = 22.4 µs; τ_{dwell} = 15 µs; τ_{acq} = 15.4 ms; τ_{HC} = 1.2 ms; τ_{DARR} = 400 ms; ν_{1H} = 85 kHz	16 h	GB = 50 Hz in t_1: GB = 50 Hz in t_2

B_0 = magnetic field; ν_{MAS} = MAS frequency; ν_{1H} = ^{1}H radio-frequency field amplitude for decoupling; ν_{1X} = radio-frequency field amplitude for cross-polarization, X= N or C (^{15}N or ^{13}C); ν_{0C} = ^{13}C radio-frequency carrier frequency; na = number of scans per free-induction-decay; τ_{pd} = delay between scans; t_{1max} = maximum t_1 value; t_{1inc} = t_1 increment; t_{2max} = maximum t_2 value; t_{2inc} = t_2 increment; τ_{dwell} = digitization dwell time in free-induction-decay; τ_{acq} = free-induction-decay acquisition time; τ_{XY} = cross-polarization period, where X and Y are H, N, or C (^{1}H, ^{15}N, or ^{13}C); τ_{DARR} = DARR mixing time; GB = pure Gaussian line-broadening before Fourier transformation.