Lesson 1: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.

Lesson 2: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.

Lesson 3: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.

Lesson 4: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.

Lesson 5: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.

Lesson 6: A linear constraint

Let x + y = 2 be a constraint on two real variables.
We describe the solution by choosing one free coordinate.

$$x + y = 2$$

$$y = 2 - x$$

Every point on this line satisfies the original constraint.
The formula above stays unchanged when we translate the explanation.

For a quadratic example consider the following expression.

$$f(x) = x^{2}$$

Small raised symbols are part of the formula, not separate sentences.
A long translation must push later paragraphs down without overlap.