

# Linear Algebra and Differential Equations Project 1

Rayana Gottschall

December 2, 2024

## **Contents**

# 1 Part 1: Colley Method

## 1.1 Explanation

Colley's method is used to rank sports teams based on their win-loss ratio. The method was created by Wesley Colley to handle situations where teams may not have played each other an equal number of times. Colley's method takes advantage of Laplace's Rule of Succession.

$$\begin{aligned}total - wins &= \frac{total - wins}{2} + \frac{total - wins}{2} \\&= \left[ \frac{total - wins}{2} + \frac{total - wins}{2} \right] + \left[ \frac{total - loss}{2} - \frac{total - loss}{2} \right] \\&= \left[ \frac{total - wins}{2} - \frac{total - loss}{2} \right] + \left[ \frac{total - wins}{2} + \frac{total - loss}{2} \right] \\&= \left[ \frac{total - wins}{2} - \frac{total - loss}{2} \right] + \frac{1}{2} [total - wins + total - loss] \\&= \left[ \frac{total - wins}{2} - \frac{total - loss}{2} \right] + \frac{1}{2} [total - games]\end{aligned}$$

## 1.2 Laplace's Rule of Succession

Laplace's Rule of Succession provides a formula to relate observed instances to unobserved ones, formally referred to as "enumerative induction." The formula for probability is  $(k+1)/(n+2)$ . Where 'k' is the number of times an event has occurred, and 'n' is the total number of trials. In sports ranking, Laplace's rule provides a more accurate probability result for small data sets by accounting for future outcomes. This is achieved through the use of biases, +1 and +2. The bias also eases the jumps in ranking when the observed data, number of games played, is scarce.

# 2 Part 2: Massey Method

## 2.1 Introduction

Massey's method includes teams' differences in points, and assumes transitivity will hold true. The data involved includes a final score differential. And a matrix is constructed using the following:

$$M \cdot r = p$$


where:

- **M**: Massey matrix (based on the matchups).
- **r**: Vector of team ratings (to be solved for).
- **p**: Vector of point differentials.

## 2.2 Method of Least Squares

The method of least squares is used to find a line of best fit for a set of data. The goal is to minimize the distance, or error, between the line of best fit and the points of data.

### Least Square Method



## 3 Part 3: Application to Real Data

### 3.1 Data Collection

[Explain the source and characteristics of the data you used.]

### 3.2 Implementation

[Discuss how you applied the Colley and Massey methods to the real data.]

### 3.3 Comparison of Results

[Compare the outcomes of both methods when applied to the data.]

## 4 Part 4: Cutting Edge

### 4.1 Overview of New Methods

Additional methods to sport rankings include the Elo Rating System, the Bradley-Terry Model, and Bayesian Analysis. Each method listed has its own unique application. The Elo Rating System is primarily used for chess rankings, but it can be applied to any two player game. The Bradley-Terry Method can be used for various team sports and can even be applied to election outcomes, among other things. Lastly, the Bayesian Analysis approach is used to calculate rankings in motorsport racing.

### 4.2 Bayesian Analysis of Formula One Race Results

Unlike football and basketball, Formula One racing has an additional factor: the car. Each Formula One team has its own unique car, because of this, in what way can we compare driver skill? The Bayesian Analysis mathematics proposed by Kesteren and Bergkamp uses driver data from 2014-2021 and the attributes are as follows: driver id, constructor id, season, race number, finishing position, and status. Additional race information included weather conditions (ie wet or dry) and circuit type (street or permanent).

#### 4.2.1 Diving into the Math

For each race,  $r$ , we assume there is a set of competitors,  $C$ . The outcome is a vector which represents a ranking of competitors,  $y$ . The ranking follows a Rank-Ordered Logit model where finish position (ranking) is modeled as a function of driver skill, constructor advantage, and yearly variations in performance.

Rank-Ordered Logit:

$$\mathbf{y}_r \sim \text{RankOrderedLogit}(\boldsymbol{\theta}_r)$$

where  $\boldsymbol{\theta}_r$  represents the latent abilities of competitors in race  $r$ .

Latent Ability Decomposition: the ability of a competitor is decomposed into the following components


$$\theta_c = \theta_d + \theta_{ds} + \theta_t + \theta_{ts}$$

$\theta_d$ : Long-term driver skill,

$\theta_{ds}$ : Seasonal deviation of driver skill,

$\theta_t$ : Long-term constructor advantage,

$\theta_{ts}$ : Seasonal deviation of constructor advantage.



## References

The Elo System:

<https://compass.blogs.bristol.ac.uk/2020/12/17/the-elo-rating-system-from-chess-to-education/>

Bayesian Analysis:

<https://pmc.ncbi.nlm.nih.gov/articles/PMC10660124/>

Laplace's Rule of Sucession:

<https://jonathanweisberg.org/post/inductive-logic-2/>

Method of Least Squares:

<https://www.investopedia.com/terms/l/least-squares-method.asp: :text=The>