NOUVELLE
ANTI-AGING

NAD+ IV THERAPY — CONSENT FORM

NAD+ 静脉注射知情同意书

I request that Nouvelle Anti-Aging provide NAD+ IV Therapy as explained to me.

本人申请接受 Nouvelle Anti-Aging 向本人解释的 NAD+ 静脉注射项目。

A prescribed NAD+ preparation is infused intravenously. Evidence for routine anti-aging, improved cognition or energy is limited; benefit is uncertain and no rejuvenation outcome is promised.

静脉输注处方 NAD+制剂。常规抗衰、认知或精力改善的证据有限，获益不确定，不承诺年轻化结果。

Identify pharmacy/manufacturer, ingredients, dose, dilution, rate and regulatory basis. A compounded NAD+ preparation is not FDA-approved merely because supplied by a licensed pharmacy.

明确药房或厂家、成分、剂量、稀释液、速率及监管依据。由持牌药房提供不代表配制 NAD+获 FDA 核准。

RISKS / POSSIBLE COMPLICATIONS 风险及可能并发症

1. Needle/IV reactions: pain, warmth, bruising, bleeding, fainting, infiltration/extravasation, phlebitis and infection.

针刺或静脉反应：疼痛、温热、瘀青、出血、晕厥、渗漏或外渗、静脉炎及感染。

2. Possible symptoms include nausea, upset stomach, diarrhea, headache, dizziness, flushing, chills/fever, muscle cramps, weakness or palpitations; occurrence depends on the ingredients, dose and infusion rate.

可能出现恶心、胃部不适、腹泻、头痛、眩晕、潮红、寒战或发热、肌肉痉挛、无力或心悸，具体取决于成分、剂量及输注速率。

3. Allergy can cause itching, rash, hives, wheezing, mouth/throat swelling, breathing difficulty or anaphylaxis.

过敏可表现为瘙痒、皮疹、风团、喘鸣、口或喉肿胀、呼吸困难或过敏性休克。

4. Serious complications include fluid overload, electrolyte disturbance, abnormal heart rhythm, contamination, bloodstream infection/sepsis or endotoxin reactions.

严重并发症包括液体超负荷、电解质紊乱、心律失常、污染、血流感染或脓毒症、内毒素反应。

5. NAD+ infusion can cause abdominal or chest discomfort and changes in vital signs. Anti-aging benefit is uncertain; sterile formulation and lawful supply require verification.

NAD+输注可能引起腹部或胸部不适及生命体征变化。抗衰获益不确定，无菌配方及合法供应须核实。

NOUVELLE
ANTI-AGING

HEALTH DISCLOSURE 健康情况告知

Disclose kidney/heart/liver disease, fluid restrictions, pregnancy, allergies, prior infusion reactions and all medicines. Verify sterile injectable sourcing and the actual prescription before cannulation. Disclose diabetes/low glucose, seizure history, Leber disease if B12 is proposed, anemia/iron or folate deficiency, bone-marrow treatment and all medicines/supplements, including anticoagulants. Do not stop medication without medical advice.

告知肾心肝疾病、限液、孕期、过敏、既往输注反应及用药。穿刺前核实无菌注射制剂来源与实际处方。 另告知糖尿病或低血糖、癫痫史、拟用 B12 时的 Leber 病、贫血或铁叶酸缺乏、骨髓相关治疗及包括抗凝药在内的所有药物和补充剂。未经医生指导，不自行停药。

ALTERNATIVES / RESULTS 替代方案及效果

No infusion; assessment of fatigue or cognitive symptoms; sleep, nutrition and evidence-based treatment for identified causes.

不输注；检查疲劳或认知症状；睡眠营养管理及针对明确原因的规范治疗。

No result or duration is guaranteed; further sessions/correction may be needed. With no elective treatment, the original concern may remain. Fees and possible additional costs have been explained separately.

不保证效果或维持时间，可能需后续疗程或修复。不接受选择性治疗时，原有困扰可能持续。费用及可能的额外费用已另行说明。

AFTERCARE / WHEN TO SEEK HELP 护理及就医提示

Report all symptoms immediately; staff may stop or slow the infusion after assessment. Do not endure chest discomfort to complete it. Keep the IV site clean afterward and follow individualized hydration advice.

任何症状应立即报告，由医护评估后暂停或减速，不为完成输注而忍受胸部不适。之后保持针口清洁，并遵守个体化补水建议。

Chest pain, difficulty breathing, fainting or severe shaking/chills require immediate medical assessment; fever or worsening IV-site redness needs prompt care.

胸痛、呼吸困难、晕厥或严重寒战需立即医学评估；发热或针口红肿加重需及时就医。

Clinic: (425) 598-1111. For severe/life-threatening symptoms call 911; do not wait for a callback.

诊所：(425) 598-1111。严重或危及生命的症状请拨 911，不等待回电。

NOUVELLE
ANTI-AGING

ACKNOWLEDGMENT / CONSENT 确认与同意

I have read all pages and had my questions about the treatment, risks, benefits and alternatives answered. I consent to the treatment described in this form and repeat treatments within the agreed plan, with my willingness confirmed before each session.

本人已阅读全部页面，关于治疗、风险、获益及替代方案的问题已获解答。本人同意接受 NAD+ 静脉注射及约定方案内的后续重复治疗，每次治疗前确认本人愿意继续。

Patient / Responsible Party Name 顾客／合法代理人姓名

_____

Patient / Responsible Party Signature 顾客／合法代理人签字

_____

Date 日期

_____