SaddlepointExpansions for MATLAB

User manual

Simon A. Broda and Raymond Kan

Version 0.3.0, November 2026

Contents

1 Introduction **2**

2 A first example **3**

3 Conventions **3**

4 Cumulant generating functions **4**
4.1 The class CGF 4
4.2 The library 4
4.3 Transformations 5
4.4 Writing your own 5
4.5 Closed-form Taylor polynomials 6

5 Orthant probabilities **6**

6 Densities **7**

7 Ratios **7**
7.1 Usage 7
7.2 Examples 7
7.3 Cost 8

8 Studentized means **8**

9 Conditional distributions **10**

10 Lattice variables **11**

11 Quantiles **12**

12 Normalized densities **13**

13 Derivatives, saddlepoints and the Taylor class **13**
13.1 Double-double arithmetic 14

14 The expansions **14**
14.1 Whitened coefficients 14
14.2 Densities 15
14.3 Orthant probabilities 15
14.4 Ratios 15

14.5 Studentized means 16
14.6 Conditional expansions 16
14.7 Saddlepoint space 17

15 Accuracy and precision **17**

16 The engine **18**

17 Validation **18**

18 Function reference **20**

1 Introduction

SaddlepointExpansions for MATLAB computes saddlepoint expansions of any order for the distribution of a random vector from its cumulant generating function alone. It implements the expansions of Broda and Kan, "Uniform saddlepoint expansions for multivariate tail probabilities":

- the orthant probability $\mathrm{P}(X \geq x)$ of a random vector X of any dimension d, with lower and mixed corners, and with the inclusion–exclusion step for saddlepoints that have negative components;
- the density of X in any dimension;
- the joint density and the joint distribution function of a vector of ratios $R = (X_1/Y_1, \ldots, X_d/Y_d)$ with positive denominators, including ratios with a common denominator;
- in one dimension, the expansions of Daniels (density), Hauschildt and Robinson (distribution function) and Rice (density of a ratio).

It also implements the expansions of Bao, Broda and Kan, "Inversion formulas and saddlepoint expansions for Studentized means", for the density, the distribution function and the quantiles of a Studentized mean $Q = X/\sqrt{Y}$ with $Y > 0$, from the joint cumulant generating function of (X, Y) (§ 8), and those of Broda and Kan, "Conditional and lattice saddlepoint expansions of arbitrary order", for the conditional distribution function $\mathrm{P}(X_{\mathcal{A}} \geq x_{\mathcal{A}} \mid X_{\mathcal{B}} = x_{\mathcal{B}})$ and the conditional density of every order, for any numbers of tail and conditioning components and for continuous and lattice variables, with the expansions of $\mathrm{P}(S \geq k)$ and $\mathrm{P}(S = k)$ for sums of integer-valued vectors as the case without conditioning (§§ 9 and 10). Quantiles of the expansions and normalized densities are computed in saddlepoint space, with the point parametrized by its saddlepoint, so that no saddlepoint equation is solved on the way (§§ 11 and 12). The approximation of order M has relative error $O(n^{-M})$, where n is the large parameter (a sample size, a number of degrees of freedom).

The user writes the cumulant generating function K as an ordinary MATLAB function. Its derivatives of all orders are generated by Taylor-mode automatic differentiation (the class `tps`, truncated multivariate power series with overloaded arithmetic and elementary functions), directly in the whitened coordinates in which the expansions are evaluated, so that no derivative has to be written by hand and none is computed by finite differences. The expansions themselves are computed by the MATLAB programs that accompany the paper (folder `engine`), which can also be called directly with closed-form derivatives.

The package is the MATLAB counterpart of the Julia package `SaddlepointExpansions.jl` by Simon Broda (`https://github.com/s-broda/SaddlepointExpansions.jl`), which also has a Python version (`https://github.com/s-broda/saddlepoint-expansions`). The packages

have the same interface and share a set of reference values, on which they agree to about 10^{-13} relative (§ 17). Where double precision runs out, the Julia package recomputes in variable precision and this package in double-double arithmetic, about 32 digits (§ 13.1).

Theorem and corollary numbers refer to the version of the paper of November 2026: Theorem 1 (orthant probability), Theorem 2 (inclusion–exclusion), Theorem 3 (density), Theorem 4 and Corollary 3 (densities of ratios), Corollaries 1, 2, 4 and 5 (Daniels, Hauschildt–Robinson, the distribution function of a ratio, Rice). For the conditional paper they refer to its version of 9 November 2026: Theorem 3 (the conditional distribution function), Remark 1 (saddlepoints with negative tail components), Theorem 4 (the conditional density), Theorem 5 (lattice variables) and Corollary 1 (lattice variables without conditioning), Section 5.2 (the canonical coordinates).

Requirements. MATLAB (the package was tested with R2025b and R2026b) and the Statistics and Machine Learning Toolbox, whose `normcdf` and `normpdf` are used by the univariate engine. Nothing else is needed.

Installation. Copy the folder anywhere and run `spx_setup`, which adds the folder and its `engine` subfolder to the path. `help SaddlepointExpansions` lists the functions, `examples/demo_spx.m` runs the examples of this manual in a few seconds and `tests/test_spx.m` the tests of § 17 in a few minutes.

2 A first example

Let $X = (U_0 + U_1, U_0 + U_2)$ with U_0, U_1, U_2 independent unit exponential variables, the example of Kolassa and Li (2010), and let $\bar{X}$ be the mean of $n = 5$ independent copies of X. Then

```
spx_setup
K = CGF(@(t) -log(1 - t(1)) - log(1 - t(2)) - log(1 - t(1) - t(2)), 2, ...
        'domain', @(t) t(1) < 1 && t(2) < 1 && t(1) + t(2) < 1);
Kn = sample_mean(K, 5);                % the cgf 5 K(t/5) of the mean
r = tailprob(Kn, [3 3], 'order', 4);  % P(mean1 >= 3, mean2 >= 3)
r.F'                                   % 0.02142343 0.02214365 0.02215413 0.02215193
r.tau                                  % the saddlepoint
s = density(Kn, [3 3], 'order', 4);   % the joint density of the mean at (3, 3)
```

The exact probability is 0.02215196; the approximations of orders one to four have relative errors -3.3×10^{-2}, -3.8×10^{-4}, 9.8×10^{-5} and about -1.4×10^{-6}. The same cumulant generating function is in the library as `exponential_sums_cgf([1 0; 1 1; 0 1], 'n', 5)`.

3 Conventions

The large parameter. The large parameter n is absorbed in the cumulant generating function: the expansions are those of the vector whose cumulant generating function is supplied. For the mean of n copies one supplies $nK(t/n)$, which `sample_mean(K, n)` constructs; the density returned is then that of the mean.

Orders. Every expansion function takes `'order', M` (default three) and returns the approximations of orders $1, \dots, M$ in a column vector. The approximation of order M has relative error $O(n^{-M})$. For a density it keeps the terms $m = 0, \dots, M-1$ of the series in powers of n^{-1}, and for an orthant probability the terms $m = 0, \dots, 2M-1$ of the series in powers of $n^{-1/2}$. For densities, order one is Daniels's first-order approximation; for orthant probabilities, the leading term $m = 0$ is the approximation of Kolassa (2003).

Results. The expansion functions return a structure with the approximations (`F` or `f`), the saddlepoint (`tau`, or `shat` for ratios), a `status` string (`'ok'` or a warning about the numerical evaluation) and `info`, the diagnostics of the underlying engine program. The diagnostics measure the numerical quality of the evaluation, such as cancellation among the terms or the change with half the quadrature nodes; they do not measure the asymptotic error of the expansion, which is what the order controls.

Vectors. Points may be row or column vectors. The argument `t` of a cumulant generating function is always a $d \times 1$ column vector.

4 Cumulant generating functions

4.1 The class CGF

```
K = CGF(fun, d)
K = CGF(fun, d, 'domain', pred)
K = CGF(fun, d, 'Taylor', tfun)
```

`fun` is a function of a $d \times 1$ vector `t` returning $K(t)$, also for $d = 1$. `pred` is an optional predicate `@(t) logical` describing the domain, and `tfun` an optional function that returns Taylor polynomials in closed form (§ 4.5).

The function is evaluated on vectors of truncated power series to generate its derivatives, so it must be written with operations that the class `tps` supports:

- `+`, `-`, `.*`, `./`, `.^`, `*` and `/` by numbers, products `A*t` and `t.'*A` with constant matrices, `sum`, `prod`, `dot`, indexing `t(j)`, `t(1:k)`, concatenation, `reshape`, `diag`;
- `exp`, `expm1`, `log`, `log1p`, `sqrt`, `sin`, `cos`, `tan`, `sinh`, `cosh`, `tanh`, `atan`, `abs`, and `sinhc`, `sincu` of the package;
- comparisons (`<`, `<=`, ...), which compare the constant terms, so that code with tests on the argument can be differentiated.

Functions of `t` that call compiled numerical routines (`det`, `inv`, `chol`, `eig`, `expm`, ...) cannot be differentiated this way; § 4.4 shows how to write determinants and inverses.

The domain. Outside the domain, `fun` may raise an error with identifier `'tps:domain'` (the logarithm and the square root of a nonpositive number do so automatically) or return a nonfinite value; both are treated as "outside the domain", and the saddlepoint iteration steps back. When the formula is finite outside the true domain, as $1/(1-t)$ is for $t > 1$, the predicate `'domain'` is needed. It is also cheaper than an error.

4.2 The library

constructor	distribution
`normal_cgf(mu, Sigma)`	$N(\mu, \Sigma)$ (the expansions are exact at order one)
`gamma_cgf(a, b)`	gamma with shape a and scale b (default one); `gamma_cgf(1, b)` is the exponential distribution with mean b
`ncx2_cgf(m, lam)`	noncentral chi-squared $\chi^2_m(\lambda)$
`exponential_sums_cgf(A, 'n', n, 'shift', l)`	the mean of n independent copies of $A^{\mathsf{T}}U + l$, U a vector of independent unit exponential variables

constructor	distribution
`quadratic_forms_cgf(Mats, 'Sigma', S, 'mu', mu, 'scale', c)`	the quadratic forms $z^{\mathsf{T}}A_i z$, $i = 1, \ldots, d$, in $z \sim N(\mu, \Sigma)$, `Mats = {A_1,...,A_d}`, summed over c independent copies
`wishart_diag_cgf(V, n)`	the diagonal of a Wishart matrix $W_d(n, V)$
`common_component_cgf(m, lam)`	$Q_j = X_j + X_0$, $j = 1, \ldots, d$, with independent $X_i \sim \chi^2_{m_i}(\lambda_i)$, `m = [m_1 ... m_d m_0]`
`uniform_cgf(a, b)`	uniform on $[a, b]$ (default $[-1, 1]$)
`poisson_cgf(lam)`	Poisson with mean λ, $K(t) = \lambda(e^t - 1)$ (a lattice variable)
`binomial_cgf(m, p)`	binomial with m trials, $K(t) = m \log(1 - p + pe^t)$ (lattice)
`negative_binomial_cgf(r, p)`	the number of failures before the r-th success, $K(t) = r \log p - r \log\{1 - (1 - p)e^t\}$, $t < -\log(1 - p)$; r need not be an integer (lattice)
`logistic_regression_cgf(X, 'beta', b)`	the sufficient statistic $S = \sum_i y_i x_i$ of a logistic regression with design matrix X (rows x_i^{T}), $K(t) = \sum_i [\log\{1 + e^{x_i^{\mathsf{T}}(\beta+t)}\} - \log(1 + e^{x_i^{\mathsf{T}}\beta})]$, at $\beta =$ `b` (default zero); lattice for integer covariates

The cumulant generating function of quadratic forms is

$$K(t) = -\frac{1}{2}\log\det(I - 2C) + \nu^{\mathsf{T}} C (I - 2C)^{-1}\nu, \qquad C = \sum_{i=1}^{d} t_i \Sigma^{1/2} A_i \Sigma^{1/2}, \qquad \nu = \Sigma^{-1/2}\mu,$$

on the set where $I - 2C$ is positive definite. Its Taylor polynomials are computed in closed form by the trace recursion of the paper, which is much faster than automatic differentiation for matrices of large order.

4.3 Transformations

function	cumulant generating function of
`reflect(K, s)`	$(s_1 X_1, \ldots, s_d X_d)$, $s_j = \pm 1$; `reflect(K)` is that of $-X$
`marginal(K, idx)`	the components `idx` of X
`linear_combination(K, B)`	$B^{\mathsf{T}}X$ for a $d \times k$ matrix B, that is $K(Bu)$
`sample_mean(K, n)`	the mean of n independent copies, $nK(t/n)$
`shift(K, c)`	$X + c$
`K1 + K2`	the sum of two independent vectors of the same dimension

The transformations keep closed-form Taylor polynomials: a linear combination of quadratic forms, for example, is still differentiated by the trace recursion.

4.4 Writing your own

Removable singularities are the one trap: $\sinh(t)/t$ is $0/0$ at $t = 0$, where the saddlepoint iteration starts. Use `sinhc` and `sincu`, which switch to the power series there; the library defines the uniform distribution as `log(sinhc(t))`. Two patterns:

```
% independent parts, one per component, plus a common part
K = CGF(@(t) -sum(m(:)/2.*log(1 - 2*t)) - m0/2*log(1 - 2*sum(t)), d, ...
        'domain', @(t) all(t < 1/2) && sum(t) < 1/2);
```

```
% a log-determinant: a Cholesky factorization written in plain arithmetic
function v = logdet_cgf(t, B1, B2)
n = size(B1, 1); Q = eye(n) - 2*(t(1)*B1 + t(2)*B2);
L = Q(1, 1)*zeros(n);                 % a series array of the right class
for j = 1:n
   s = Q(j, j); for k = 1:j-1, s = s - L(j, k)^2; end
   L(j, j) = sqrt(s);
   for i = j+1:n
      w = Q(i, j); for k = 1:j-1, w = w - L(i, k)*L(j, k); end
      L(i, j) = w/L(j, j);
   end
end
v = -sum(log(diag(L)));
end
```

The test file `tests/test_spx.m` differentiates the complete cumulant generating function of a pair of quadratic forms written this way and checks it against the trace recursion. For quadratic forms in normal variables, `quadratic_forms_cgf` is faster.

4.5 Closed-form Taylor polynomials

The option `'Taylor', tfun` supplies a function `tfun(t, N, M)` that returns the Taylor polynomial of degree N in v of $K(t+Mv)$ as a scalar `tps` in `size(M, 2)` variables (`M = []` stands for the identity). It is then used instead of automatic differentiation; `fun` is still used for the values of K. The polynomial is built from its coefficients with `tps(C, [1 1], k, N)`, where `C` holds the coefficients of the monomials in the order of `tps.tables(k, N).mi` (§ 13). `quadratic_forms_cgf` is an example.

5 Orthant probabilities

```
r = tailprob(K, x, 'order', M)                  % P(X >= x)
r = tailprob(K, x, 'order', M, 'lower', true)   % P(X <= x)
r = tailprob(K, x, 'order', M, 'signs', s)      % e.g. s = [1 -1]: P(X1 >= x1, X2 <=
    x2)
```

`r.F` holds the approximations of orders $1, \dots, M$ and `r.tau` the saddlepoint, $K'(\hat\tau) = x$. Lower and mixed corners are computed as upper orthant probabilities of the reflected vector.

Saddlepoints with negative components. The expansion of Theorem 1 applies to saddlepoints in the closed positive orthant $\hat\tau \geq 0$. When some components are negative, `tailprob` writes the probability by the inclusion–exclusion identity of Theorem 2 as a signed sum of orthant probabilities of marginals with some components negated, each with a saddlepoint in its closed positive orthant, recursively, and expands every term (§ 14). This keeps the relative error $O(n^{-M})$; a direct evaluation of the formula at such a saddlepoint would have the wrong exponential order. The saddlepoint reported is that of the original problem.

Dimensions. In one dimension the expansion is that of Hauschildt and Robinson (Corollary 2), computed directly for the upper tail. In two dimensions the integrals of the expansion are computed by Gauss–Legendre quadrature (`spa_orthant2`, `spa_bi`), in three by Gauss quadrature on a conditioned grid (`spa_tri`), and in four or more by randomized quasi-Monte Carlo after Genz's sequential conditioning (`spa_mv`); `r.info.err` is then an error estimate (in four or more dimensions a standard error). A trivariate probability takes about a second at order three and tens of seconds at order four or five, mostly in the inclusion–exclusion terms.

Options. Name-value pairs other than `'order'`, `'lower'` and `'signs'` are passed to the engine: `'nG'` (Gauss–Legendre nodes per coordinate), `'qtol'` (quadrature tolerance), `'nGmax'`, `'method'` (`'quad'` or `'qmc'`), `'nqmc'`, `'nshift'`, `'seed'` and, in two dimensions, `'logscale'`, `c`, which returns the probability scaled by e^{-c} for probabilities below the range of double precision. See the help of `spa_bi`, `spa_tri` and `spa_mv`.

6 Densities

```
r = density(K, x, 'order', M)
```

`r.f` holds the approximations of orders $1, \dots, M$ of the density of X at x (Theorem 3; Daniels's expansion, Corollary 1, in one dimension). The expansion is a finite sum of Hermite polynomials at zero and needs no integration in any dimension. In three or more dimensions the terms of high orders can cancel by many orders of magnitude; `r.info.kappa_pdf` reports the cancellation (§ 15).

7 Ratios

7.1 Usage

Let $R = (X_1/Y_1, \dots, X_d/Y_d)$ with $Y_j > 0$. The cumulant generating function `K` is that of the vector $(X_1, \dots, X_d, Y_1, \dots, Y_d)$, of dimension $2d$, numerators first. For ratios with a common denominator Y it may instead be that of $(X_1, \dots, X_d, Y)$, of dimension $d + 1$. The number of ratios is the length of the point `r0`.

```
r = ratio_density(K, r0, 'order', M)                  % density of R at r0
r = ratio_tailprob(K, r0, 'order', M)                 % P(R >= r0)
r = ratio_tailprob(K, r0, 'order', M, 'lower', true)  % P(R <= r0)
r = ratio_tailprob(K, r0, 'order', M, 'signs', s)     % a mixed corner
```

`r.f` or `r.F` holds the approximations of orders $1, \dots, M$, and `r.shat` the saddlepoint of K_W (§ 14.4). For one ratio the density is Rice's expansion (Corollary 5), and for several the expansion of Theorem 4 and Corollary 3, whose term of order one is the approximation of Butler and Paolella (1998); the distribution function is the orthant probability of § 5 for $W = X - r_0 \circ Y$ (Corollary 4 for one ratio), with all its options.

7.2 Examples

Sample autocorrelations. With $z \sim N(0, I_N)$, $M = I_N - 1_N 1_N^{\mathsf{T}}/N$ and $(A_j)_{ts} = 1/2$ for $|t - s| = j$, the sample autocorrelations at lags one and two are $R_j = z^{\mathsf{T}} M A_j M z / z^{\mathsf{T}} M z$, ratios with the common denominator $z^{\mathsf{T}} M z$. For $N = 50$ at $r_0 = (-1, -1)/\sqrt{50}$:

```
N = 50; M = eye(N) - ones(N)/N;
A = @(j) (diag(ones(N-j, 1), j) + diag(ones(N-j, 1), -j))/2;
K = quadratic_forms_cgf({M*A(1)*M, M*A(2)*M, M});      % dimension d + 1 = 3
r0 = [-1 -1]/sqrt(N);
f = ratio_density(K, r0, 'order', 5).f
F = ratio_tailprob(K, r0, 'order', 5, 'lower', true).F
```

The relative errors of orders one to five, against the exact values of the paper, are

order	1	2	3	4	5
density	-1.4×10^{-3}	7.3×10^{-5}	-1.0×10^{-5}	2.2×10^{-6}	-6.4×10^{-7}
distribution function	-3.7×10^{-2}	1.5×10^{-4}	1.4×10^{-5}	5.8×10^{-7}	-1.6×10^{-7}

These are the values of the corresponding row of the table of sample autocorrelations in the paper, which were computed by a dedicated program.

Gamma ratios with a common denominator. If $X_1, \dots, X_d, Y$ are independent with $X_j \sim \text{Gamma}(a_j)$ and $Y \sim \text{Gamma}(b)$, the ratios $R_j = X_j/Y$ have the inverted Dirichlet density

$$f_R(r) = \frac{\Gamma(a_1 + \cdots + a_d + b)}{\Gamma(b)\prod_{j=1}^{d}\Gamma(a_j)} \, \frac{\prod_{j=1}^{d} r_j^{a_j - 1}}{(1 + r_1 + \cdots + r_d)^{a_1 + \cdots + a_d + b}}.$$

```
a = [3 4 5]; b = 6;
K = CGF(@(t) -sum(a(:).*log(1 - t(1:3))) - b*log(1 - t(4)), 4, 'domain', @(t) all(t <
    1));
f = ratio_density(K, [0.6 0.5 0.9], 'order', 5).f
```

The relative errors of orders one to five are 7.7×10^{-2}, -3.1×10^{-3}, -1.0×10^{-4}, 1.7×10^{-5} and 3.6×10^{-6}. Multiplying the shapes by n shows the rate $O(n^{-M})$; for $d = 2$ at $r_0 = (0.6, 0.5)$:

n	order 1	order 2	order 3	order 4
1	5.8×10^{-2}	-1.8×10^{-3}	-1.2×10^{-4}	1.3×10^{-5}
2	2.8×10^{-2}	-4.2×10^{-4}	-1.6×10^{-5}	6.6×10^{-7}
4	1.4×10^{-2}	-1.0×10^{-4}	-2.0×10^{-6}	3.7×10^{-8}
8	7.0×10^{-3}	-2.5×10^{-5}	-2.5×10^{-7}	2.2×10^{-9}

Each doubling of n divides the error of order M by about 2^M.

7.3 Cost

The density of several ratios needs one Taylor polynomial of $\mathcal{K}(s, -r \circ s + \tau)$ in the $2d$ variables (u, τ) of § 14.4, of degree $\max(2M, 2M - 2 + d)$. For two or three ratios every order up to five takes one or two seconds, also for quadratic forms in $N = 50$ variables; for four ratios order four takes several seconds. The distribution function costs what an orthant probability of dimension d costs (§ 5).

8 Studentized means

Let $Q = X/\sqrt{Y}$ with $Y > 0$, and let `K` be the joint cumulant generating function of (X, Y), of dimension two (numerator first). The expansions of Bao, Broda and Kan are

```
r = studentized_density(K, q, 'order', M)                % density of Q at q
r = studentized_tailprob(K, q, 'order', M)               % P(Q >= q)
r = studentized_tailprob(K, q, 'order', M, 'lower', true) % P(Q <= q)
r = studentized_quantile(K, p, 'order', M)               % the q with P(Q <= q) = p
sd = studentized_saddlepoint(K, q)                       % sd.s, sd.t, sd.v, sd.q0
loc = studentized_locus(K, s, tguess)                    % loc.t, loc.v, loc.q
```

Unlike a ratio, $X - q\sqrt{Y}$ is not linear in (X, Y), and the inversion formula is a double integral. Its saddlepoint $(\hat{s}, \hat{t}, \hat{v})$ solves

$$K_s(s, t) = qv, \qquad K_t(s, t) = v^2, \qquad qs + 2tv = 0,$$

with $\hat{v} > 0$; `studentized_saddlepoint` finds it by Newton's method with continuation in q from $q_0 = E(X)/E(Y)^{1/2}$, where $\hat{s} = \hat{t} = 0$. The density is the amplitude expansion of the double integral about the saddlepoint, and the distribution function an expansion of Hauschildt–Robinson type in s after the t-integral is expanded by Laplace's method (§ 14.5). `r.f(M)` keeps

the terms of the density of orders n^{-m}, $m \le M-1$, and `r.F(M)` those of the distribution function of orders $n^{-m/2}$, $m \le 2M-1$, both with relative error $O(n^{-M})$; `r.info.Fp(j+1)`, $j = 0, \dots, 2M-1$, holds the distribution function with the terms up to $n^{-j/2}$. `r.saddle` is the saddlepoint and `r.info.kappa` the cancellation indicator. The option `'saddle', sd` supplies a saddlepoint already computed. The expansions are not evaluated at $q = q_0$, where $\hat{t} = 0$.

The saddlepoints of all q lie on the curve $sK_s + 2tK_t = 0$ through the origin, on which $v = K_t^{1/2}$ and $q = K_s/v$ (`studentized_locus`). `studentized_quantile` searches along this curve: each trial value of $\hat{s}$ gives its q and its distribution function directly, so that no saddlepoint equation is solved for a trial q (a bracket in $\hat{s}$ from the origin, then the Illinois iteration; `'tol'`, default 10^{-10}, is the tolerance on the probability).

Example: the noncentral t distribution. With $X \sim N(\mu, 1)$ and $Y \sim \chi^2_\nu/\nu$ independent and n pairs, $\bar{X}/\sqrt{\bar{Y}} = T/\sqrt{n}$, T noncentral t with $n\nu$ degrees of freedom and noncentrality $\sqrt{n}\mu$:

```
mu = 1; nu = 3; n = 16;
K = sample_mean(CGF(@(t) mu*t(1) + t(1)^2/2 - nu/2*log(1 - 2*t(2)/nu), 2, ...
                    'domain', @(t) t(2) < nu/2), n);
studentized_tailprob(K, 6/sqrt(n), 'order', 3, 'lower', true).F'   % P(T <= 6)
%   0.9539837462   0.9539799842   0.9539798477    (exact 0.9539798478)
studentized_density(K, 6/sqrt(n), 'order', 3).f'/sqrt(n)           % density of T at 6
%   0.0757466061   0.0753761567   0.0753767005    (exact 0.0753767236)
studentized_quantile(K, 0.05, 'order', 3).q*sqrt(n)   % 2.32153317, the exact value
```

Example: the Dickey–Fuller t-ratio. `tests/dickey_fuller_cgf.m` gives the joint cumulant generating function of (U, V) whose ratio $U/\sqrt{V}$ is the limiting Dickey–Fuller t-ratio without deterministic terms (model 1), with a constant (model 2) or with a constant and a trend (model 3), under the unit root ($c = 0$) and local alternatives ($c < 0$), from Nabeya and Tanaka (1990) and Nabeya and Sørensen (1994). For model 2 at the 5% critical value,

```
K = dickey_fuller_cgf(2);
r = studentized_tailprob(K, -2.862, 'order', 3, 'lower', true);  r.info.Fp'
%   0.0475  0.0520  0.0484  0.0449  0.0453  0.0496
studentized_quantile(K, 0.05, 'order', 3).q                      % -2.85930
```

(the exact quantile of the limit is -2.861569). It is a test fixture, not part of the package.

Precision. Where $|\hat{t}|$ is large the terms of high order of the density cancel by many orders of magnitude (for the Dickey–Fuller model 2 above the indicator is about 10^8 at order three, 10^{13} at order four and 10^{17} at order five). When `r.info.kappa(end)` exceeds 10^6, `studentized_density` repeats the computation in double-double arithmetic (§ 13.1), as the Julia package repeats it in `BigFloat`: the saddlepoint is refined, and the Taylor polynomial of K, the whitening and all sums carry about 32 digits; `r.info.refined` is then `'double-double'`. The relative error of the refined values is about $\kappa \times 10^{-32}$: for the Dickey–Fuller model 2 the refined density agrees with the Julia package in `BigFloat` to the last digit of double precision up to order four and to 10^{-15} at order five; `r.status` remains `'coefficients inaccurate'` when $\kappa\, 2^{-104} > 10^{-10}$ (order six for that model). The options are

```
r = studentized_density(K, q, 'order', M, 'refine', false)    % double precision only
r = studentized_density(K, q, 'order', M, 'precision', 'dd')  % double-double from the
    start
r = studentized_tailprob(K, q, 'order', M, 'precision', 'dd')
```

without refinement `r.status` is `'coefficients inaccurate'` when the indicator exceeds 10^{10}, and `'refinement failed'` if K cannot be evaluated on double-double series. The distribution

function is much less affected; in double-double it costs a few seconds, mostly in the integrals I_m, which are computed in variable precision. A refined density of order six costs about ten seconds.

9 Conditional distributions

Let $\mathcal{A} \subseteq \{1, \dots, d\}$ hold the indices of the components whose tail is wanted and $\mathcal{B}$ the others, on which the distribution is conditioned. The expansions of Broda and Kan, "Conditional and lattice saddlepoint expansions of arbitrary order", are

```
r = conditional_tailprob(K, x, A, 'order', M)       % P(X_A >= x_A | X_B = x_B)
r = conditional_tailprob(K, x, A, 'order', M, 'lower', true)   % <= x_A
r = conditional_tailprob(K, x, A, 'order', M, 'signs', s)      % a mixed corner
r = conditional_density(K, x, A, 'order', M)        % f(x_A | x_B)
```

where `x` is the full point of length d and `A` an index or a vector of indices. The numerator of the conditional probability is the orthant-probability integral with the poles in the components of $\mathcal{A}$ only, expanded to the terms of orders $n^{-m/2}$, $m \le 2M-1$, and the denominator is the density expansion of $X_{\mathcal{B}}$ at its own saddlepoint with M terms, so that `r.F(M)` has relative error $O(n^{-M})$ (Theorem 3 of the paper). The conditional density `r.f(M)` is the ratio of the density expansions of X and of $X_{\mathcal{B}}$, both with M terms (Theorem 4); its leading term is the double saddlepoint approximation of Barndorff-Nielsen and Cox (1979), and unlike the distribution function it holds for saddlepoints in any orthant. With $\mathcal{B}$ empty the expansions are those of `tailprob` and `density`.

With one tail component and one conditioning variable the leading term, `r.info.Fp(1)`, is the conditional analogue $(\hat{u}/\hat{z})\, e^{(\hat{u}^2-\hat{w}^2)/2}\, \bar{\Phi}(\hat{u})$ of the approximation of Hauschildt and Robinson, in the notation of Skovgaard (1987), and the approximation of order one adds the terms of order $n^{-1/2}$ to it. The expansion has no singular terms, so that nothing special happens where the saddlepoint of a tail component vanishes, as it does for Skovgaard's formula at the median. When the saddlepoint has negative components in $\mathcal{A}$, the probability is written by the inclusion–exclusion identity of Remark 1 of the paper as a signed sum of conditional probabilities given the same $X_{\mathcal{B}}$, of the tail components that are kept, some of them negated, with the others marginalized, every term with its own saddlepoint (§ 14.6); `'lower'` and `'signs'` reflect tail components as in `tailprob`.

`r.tau` is the saddlepoint of K at x and `r.taut` that of the marginal of $X_{\mathcal{B}}$, embedded as $(0_{\mathcal{A}}, \tilde{s}_{\mathcal{B}})$. `r.info.Fp(j+1)`, $j = 0, \dots, 2M-1$, holds the expansion with the terms of the numerator up to $n^{-j/2}$ and $\lfloor j/2 \rfloor + 1$ terms of the denominator, so that `r.F(M)` is `r.info.Fp(2M)`; `r.info.kappa` and `r.info.kappa_den` are the cancellation indicators of the numerator, the terms of the inclusion–exclusion included, and of the denominator; `r.info.err` is the change of `r.F` with half the quadrature nodes of the orthant integrals, whose number per panel is the option `'nG'` (default 64). `r.status` is `'ok'`, or reports a large cancellation or an integration not resolved.

Example: a gamma sum given its total. Let U and V be independent Gamma(16) variables and $X = (U, U+V)$. Given $X_2 = 32$, $X_1/32$ has the Beta(16, 16) distribution, so that the conditional probabilities are known exactly:

```
K = CGF(@(t) -16*log(1 - t(1) - t(2)) - 16*log(1 - t(2)), 2, ...
        'domain', @(t) t(1) + t(2) < 1 && t(2) < 1);
x1 = 32*betaincinv(0.9, 16, 16);                    % the 90% point, 19.60597967
r = conditional_tailprob(K, [x1 32], 1, 'order', 5);
r.F'      % 0.1000254409 0.0999930422 0.0999999444 0.1000000055 0.1000000001
f = conditional_density(K, [x1 32], 1, 'order', 5);
```

The relative errors of orders one to five are

order	1	2	3	4	5
distribution function	2.5×10^{-4}	-7.0×10^{-5}	-5.6×10^{-7}	5.5×10^{-8}	1.3×10^{-9}
density	7.8×10^{-3}	-5.3×10^{-5}	-1.1×10^{-6}	1.5×10^{-8}	1.4×10^{-9}

These are cells of the first table of the paper, which the tests recompute in full (§ 17).

Computation and cost. The Taylor coefficients are generated by Taylor-mode differentiation directly in the canonical coordinates of Section 5.2 of the paper, in which the tail components are uncorrelated with the conditioning components under the tilted distribution and the latter are whitened (§ 14.6). The conditional probability does not depend on the coordinates, but the terms do: in the original coordinates they cancel by up to sixteen orders of magnitude in the examples of the paper, in the canonical ones they do not. With one tail component every integral of the expansion is one-dimensional, whatever the number of conditioning variables, and the cost is that of the Taylor coefficients: a conditional probability of order five in three dimensions takes a fraction of a second. With two or three tail components the orthant integrals are computed on a sequentially conditioned grid, which costs about a second at order three. The engine program is `spa_cond` (§ 16), the counterpart of the program of the same name that accompanies the paper.

10 Lattice variables

For an integer-valued vector S and an integer point k, the option `'lattice', true` gives the continuity-corrected expansions of $\mathrm{P}(S \geq k)$ and the expansions of $\mathrm{P}(S = k)$ (Corollary 1 of the conditional paper, its case without conditioning components), and, conditioned on other components, those of $\mathrm{P}(S_{\mathcal{A}} \geq k_{\mathcal{A}} \mid S_{\mathcal{B}} = k_{\mathcal{B}})$ and $\mathrm{P}(S_{\mathcal{A}} = k_{\mathcal{A}} \mid S_{\mathcal{B}} = k_{\mathcal{B}})$ (Theorem 5):

```
r = tailprob(K, k, 'order', M, 'lattice', true)                  % P(S >= k)
r = tailprob(K, k, 'order', M, 'lattice', true, 'lower', true)   % P(S <= k)
r = density(K, k, 'order', M, 'lattice', true)                   % P(S = k)
% P(S_A >= k_A | S_B = k_B) and P(S_A = k_A | S_B = k_B):
r = conditional_tailprob(K, k, A, 'order', M, 'lattice', true)
r = conditional_density(K, k, A, 'order', M, 'lattice', true)
```

In the distribution function the saddlepoint is taken at the continuity-corrected point $k_{\mathcal{A}} - 1/2$ in the tail components, and the integrand carries the amplitude $\prod_{j \in \mathcal{A}} s_j / \{2 \sinh(s_j/2)\}$, which is expanded about the saddlepoint (`'sinh', 'saddle'`, the default), so that the lattice factor is exact at the saddlepoint at every order. The option `'sinh', 'origin'` instead expands $s/\{2\sinh(s/2)\} = 1 - s^2/24 + 7s^4/5760 - \cdots$ about zero and keeps the terms up to s^{2M-2} in the approximation of order M, as Kolassa (2003, 2006) does; the two agree closely when the saddlepoint is small. The probability $\mathrm{P}(S = k)$ is the density expansion at the uncorrected point. The components in $\mathcal{B}$ may be lattice or continuous. In one dimension the approximation of order one, with the terms of orders n^0 and $n^{-1/2}$, has the relative error $O(n^{-1})$ of the continuity-corrected approximations of Daniels (1987). Lower tails and saddlepoints with negative components use the reflection $\mathrm{P}(S_j \leq k_j - 1) = \mathrm{P}(-S_j \geq -k_j + 1)$. Where the continuity-corrected point lies outside the interior of the convex hull of the support, as for $\mathrm{P}(S \geq 0)$ of a count, there is no saddlepoint (the probability is zero or one) and an error with identifier `'spa_cond:support'` is raised. `tailprob` and `conditional_tailprob` with $\mathcal{A} = \{1, \ldots, d\}$ make the same computation and return the same numbers; `r.info` holds `Fp`, `kappa`, `err` and `nG`.

Examples. For the Poisson distribution with mean five,

```
r = tailprob(poisson_cgf(5), 11, 'order', 4, 'lattice', true);
r.F'      % 0.01373079 0.01369706 0.01369526 0.01369527   (exact 0.01369527)
f = density(poisson_cgf(5), 11, 'order', 4, 'lattice', true);
f.f'      % 0.00830484 0.00824192 0.00824216 0.00824218   (exact 0.00824218)
```

with relative errors 2.6×10^{-3}, 1.3×10^{-4}, -7.3×10^{-7} and -1.3×10^{-7} for the tail. The standard conditional case is exact inference in a logistic regression: with y_i independent Bernoulli variables with $\log\{p_i/(1-p_i)\} = x_i^{\mathsf{T}}\beta$, the sufficient statistic $S = \sum_i y_i x_i$ has the cumulant generating function of `logistic_regression_cgf`, and the conditional distribution of one of its components given the others does not depend on the coefficients of the conditioning components. For thirty observations with an intercept and two integer covariates, under $\beta = 0$,

```
X = [ones(30, 1) mod((0:29)', 5) mod((0:29)', 3)];
K = logistic_regression_cgf(X);
r = conditional_tailprob(K, [15 37 15], 2, 'order', 4, 'lattice', true);
r.F'      % 0.05374981 0.05244006 0.05241372 0.05241465   (exact 0.05242119)
p = conditional_density(K, [15 37 15], 2, 'order', 4, 'lattice', true);
```

where the exact values come from the joint distribution of S by dynamic programming; the errors relative to $\min(P, 1-P)$ are 2.5×10^{-2}, 3.6×10^{-4}, -1.4×10^{-4} and -1.2×10^{-4}.

11 Quantiles

```
r = quantile(K, p, 'order', M)                    % P(X <= q) = p under the expansion
r = quantile(K, p, 'order', M, 'lower', false)    % P(X >= q) = p
r = ratio_quantile(K, p, 'order', M)              % P(X_1/X_2 <= r) = p, X_2 > 0
```

`quantile` returns the point at which the saddlepoint distribution function of order M of a univariate variable (the Hauschildt–Robinson expansion of `tailprob`) equals p; with `'lower', false` the upper tail is used, so that upper quantiles of small probability are computed directly. The search runs in saddlepoint space: the point is parametrized by its saddlepoint, $x = K'(s)$, so that each iterate comes with its saddlepoint and no saddlepoint equation is solved. Newton's method in s starts at $s = 0$ and uses the derivative $f_M(K'(s))K''(s)$, the density expansion of the same order times $K''(s)$; it is safeguarded by bracketing and bisection, its steps are limited to four standard deviations $K''(s)^{-1/2}$ and pulled back into the domain of K, and it stops when the distribution function is within `tol` $\min(p, 1-p)$ of p (`'tol'`, default 10^{-12}). `r.q` is the quantile, `r.tau` its saddlepoint, `r.F` the value of the expansion there, `r.iterations` the number of Newton steps and `r.status` `'ok'`, or `'bracket at rounding level'` when the bracket of the root has shrunk to rounding level first. The expansion of a lattice variable is a step function of k, and `quantile` is meant for continuous variables. It is a method of the class `CGF`, so that `quantile` of a numeric array is still MATLAB's own.

For a ratio $R = X_1/X_2$, `K` is the joint cumulant generating function of (X_1, X_2) and the parameter is the saddlepoint $\hat{s}$ of $W = X_1 - rX_2$. The saddlepoints of all r lie on the curve $sK_1(s,t) + tK_2(s,t) = 0$, $t = -r\hat{s}$, on which $r = K_1/K_2$ (§ 14.7); each trial $\hat{s}$ gives its t by a one-dimensional Newton iteration on the curve, continued from the previous iterate, and with it r and the distribution function of `ratio_tailprob`, without the saddlepoint equation of W for a trial r. The derivative in $\hat{s}$ is the density expansion of the ratio of the same order (`ratio_density`) times $dr/d\hat{s}$ along the curve, and the iteration is safeguarded as for `quantile`; `r.r` is the quantile and `r.shat` its saddlepoint. `studentized_quantile` (§ 8) is the counterpart for Studentized means.

```
q = quantile(gamma_cgf(3), 0.05, 'order', 4);
q.q               % 0.8176862582 (exact 0.8176914472)
q = quantile(gamma_cgf(3), 1e-6, 'order', 4, 'lower', false);
```

```
q.q                % 19.1291729204 (exact 19.1291681886)
```

12 Normalized densities

```
r = density(K, x, 'order', M, 'normalize', true)         % r.info.mass: c_1..c_M
r = density(K, x, 'order', M, 'normalize', r.info.mass)  % reuse them at another point
r = ratio_density(K, r0, 'order', M, 'normalize', true)  % one ratio
r = studentized_density(K, q, 'order', M, 'normalize', true)
```

The option divides the expansion of every order M by its total mass $c_M = \int f_M(x)\,dx$, the normalization that Daniels proposed for the leading term; the masses are returned in `r.info.mass` and do not depend on the point, so that they can be passed back as `'normalize', r.info.mass` (a numeric vector) to normalize at other points without recomputing them. The integral is computed in saddlepoint space,

$$c_M = \int f_M(x)\,dx = \int f_M(K'(s))\,|K''(s)|\,ds,$$

so that the saddlepoint is known at every quadrature node and no saddlepoint equation is solved there. The quadrature is adaptive Gauss–Kronrod with the 15-point Kronrod rule and its embedded 7-point Gauss rule, the segment with the largest error estimate being bisected until the sum of the estimates is below a relative tolerance (10^{-12} in one dimension and 10^{-9} in more), which is the algorithm and the tolerances of the Julia package, so that the masses agree with its values to rounding; the infinite ends are mapped to finite ones by the same changes of variable. The range is found from $s = 0$ by steps doubling in each direction until the leading term has fallen by e^{-50} from its largest value, with bisection toward the boundary of the domain and an infinite end where the integrand decays only algebraically. In more than one dimension the integral is nested over the coordinates, each inner line centred at the regression of the inner coordinates on the outer ones under $K''(0)$, and the expansion at a node is evaluated by `spa_cond` at the known saddlepoint. For the normal distribution $c_M = 1$; for the gamma distribution with shape a, $c_1 = \Gamma(a)e^a a^{1/2-a}/\sqrt{2\pi}$, and since the bracket of its expansion does not depend on x the normalized expansion of every order is exact:

```
r = density(gamma_cgf(3), 2, 'order', 4, 'normalize', true);
r.f'          % 0.270670566473 at every order, the exact density
r.info.mass'  % 1.0280645179 0.9995071702 0.9999038000 1.0000058955
```

For one ratio the integral over r is computed over the saddlepoint $\hat{s}$ along the curve of § 14.7, with $dr = (dr/d\hat{s})\,d\hat{s}$, and for a Studentized mean over $\hat{s}$ along the locus $sK_s + 2tK_t = 0$, on the two sides of $\hat{s} = 0$ separately (the expansion is not evaluated at $\hat{t} = 0$), with $dq = (dq/d\hat{s})\,d\hat{s}$; the expansions at the nodes are refined in double-double arithmetic where they cancel, as with `'refine', true`. A normalization costs a few hundred evaluations of the expansion in one dimension, under a second, and the square of that in two, about twenty seconds; in more dimensions it is expensive. `'normalize'` is for continuous variables; `density` raises an error with `'lattice', true`.

13 Derivatives, saddlepoints and the Taylor class

```
D = derivatives(K, t, N)       % all derivatives of orders <= N at t
[tau, info] = saddlepoint(K, x)
p = taylor(K, t, N, M)         % Taylor polynomial of K(t + M v) in v, degree N
```

`D.nu` holds the multi-indices ν, one per row in graded order, and `D.value` the derivatives $K^{(\nu)}(t)$, so that `D.value(1)` is $K(t)$. The saddlepoint is computed by Newton's method with backtracking on the convex function $K(\tau) - \tau^{\mathsf{T}} x$ (`spa_saddle`).

The class `tps` represents arrays of polynomials in k variables truncated at total degree N:

```
v = tps.vars(k, N)       % k x 1 array of the variables
p = exp(v(1)) * log(1 + v(2))
coef(p, [1 2])           % nu! times the coefficient of v^nu, the derivative at 0
double(p)                % the constant term
T = tps.tables(k, N);    % T.mi: the multi-indices in graded order
```

A function f of a series $p = p_0 + q$, q without constant term, is evaluated by composition, $f(p) = \sum_{j=0}^{N} f^{(j)}(p_0) q^j / j!$; where f is not analytic at p_0 an error with identifier `'tps:domain'` is raised. The coefficients are stored in `p.C`, one column per element of the array.

13.1 Double-double arithmetic

The class `dd` holds a number as the unevaluated sum $x_{\mathrm{hi}} + x_{\mathrm{lo}}$ of two doubles with $|x_{\mathrm{lo}}| \le \frac{1}{2}\mathrm{ulp}(x_{\mathrm{hi}})$, about 32 significant digits, with the error-free transformations of Knuth and Dekker (two-sum, two-product by splitting). Addition, multiplication, division and the square root are computed in this arithmetic, sums are compensated and pairwise; the other elementary functions (`exp`, `log`, `erfc`, ...) are evaluated through `vpa` with 40 digits and rounded back.

```
x = dd(sym(1)/3)             % from a symbolic number, a double, or (hi, lo)
y = sqrt(x)*x + 1;  vpa(y, 32)   % double(y) rounds to double
```

The class `tpsdd` is the Taylor class in double-double: the same interface as `tps` (`tpsdd.vars`, `tpsdd.const`, `coef`, arithmetic, `exp`, `log`, `sqrt`, `atan`, `sinhc`, ...), with the coefficients in `p.H` and `p.L`; `coef` returns a `dd`. The derivatives of a function at the expansion point, needed for the composition, are computed in `vpa`. `taylordd(K, t, N, M)` is the Taylor polynomial of K in double-double; it evaluates the function of `K` on `tpsdd` series, so it applies to the cumulant generating functions written with these operations (the library, the transformations, and `tests/dickey_fuller_cgf.m`), but not to a closed-form `'Taylor'` function, which works in double precision. A product of two bivariate series of degree 30 takes about ten milliseconds, several orders of magnitude less than the same product in `vpa`. The Symbolic Math Toolbox is needed for the constants and the elementary functions.

14 The expansions

This section states what the package computes. The paper gives the derivations, the conditions and the error bounds; here n is absorbed in K.

14.1 Whitened coefficients

Let $\hat{\tau}$ be the saddlepoint, $K'(\hat{\tau}) = x$, and $K''(\hat{\tau}) = LL^{\mathsf{T}}$ the Cholesky factorization. All terms are computed for the whitened vector $Y = L^{-1}X$, whose cumulant generating function $K_Y(u) = K(L^{-\mathsf{T}}u)$ has its saddlepoint at $\hat{u} = L^{\mathsf{T}}\hat{\tau}$ and Hessian I there. Let $\theta_\nu = K_Y^{(\nu)}(\hat{u})/\nu!$, $|\nu| \ge 3$, be its Taylor coefficients and define $c_{r,\nu}$ by

$$c_{1,\nu} = \theta_\nu, \qquad c_{r,\nu} = \frac{1}{r} \sum_{k=3}^{|\nu|-3(r-1)} \sum_{|\lambda|=k,\ \lambda \le \nu} \theta_\lambda \, c_{r-1,\nu-\lambda}, \qquad r \ge 2,$$

so that $c_{r,\nu}$ is the coefficient of u^ν in $\Theta(u)^r/r!$, where

$$\Theta(u) = \sum_{k=3}^{\infty} \sum_{|\nu|=k} \theta_\nu u^\nu.$$

The multivariate Hermite polynomials at zero factor into univariate ones,

$$\mathrm{He}_\nu(0) = \prod_{j=1}^{d} \mathrm{He}_{\nu_j}(0), \qquad \mathrm{He}_{2i}(0) = (-1)^i \prod_{j=1}^{i} (2j-1), \qquad \mathrm{He}_{2i+1}(0) = 0.$$

The package obtains the θ_ν by differentiating the composed function $K(\hat\tau + L^{-\mathsf{T}}u)$ directly. In the original coordinates the terms cancel much more when the components of X are strongly correlated under the tilted distribution, and the substitution of $L^{-\mathsf{T}}$ into the Taylor polynomial of K can lose many digits at high degree.

14.2 Densities

$$f(x) \approx \frac{\exp\{K(\hat\tau) - \hat\tau^{\mathsf{T}} x\}}{(2\pi)^{d/2} |K''(\hat\tau)|^{1/2}} \sum_{m=0}^{M-1} a_m, \qquad a_0 = 1, \qquad a_m = \sum_{r=1}^{2m} \sum_{|\nu|=2m+2r} c_{r,\nu} \mathrm{He}_\nu(0).$$

The coefficient a_m involves the θ_ν with $|\nu| \le 2m+2$.

14.3 Orthant probabilities

For $\hat\tau \ge 0$,

$$\mathrm{P}(X \ge x) \approx \exp\{K(\hat\tau) - \hat\tau^{\mathsf{T}} x + |\hat u|^2/2\} \sum_{m=0}^{2M-1} T_m, \qquad T_0 = \tilde I_0, \qquad T_m = \sum_{r=1}^{m} \sum_{|\nu|=m+2r} c_{r,\nu} \tilde I_\nu,$$

with

$$\tilde I_\nu = \int_{\{v \in \mathbb{R}^d:\ Lv \ge 0\}} \prod_{j=1}^{d} \mathrm{He}_{\nu_j}(v_j)\, \phi_d(v + \hat u)\, dv,$$

ϕ_d the standard normal density in d dimensions. The term T_m is of order $n^{-m/2}$. When $\hat\tau$ has negative components, let $\mathcal{N} = \{j : \hat\tau_j < 0\}$. Then

$$\mathrm{P}(X \ge x) = \sum_{T \subseteq \mathcal{N}} (-1)^{|T|}\, \mathrm{P}\big(\tilde X_{\mathcal{N}^c \cup T} \ge \tilde x_{\mathcal{N}^c \cup T}\big),$$

where $\tilde X$ and $\tilde x$ are X and x with the components in T negated and the subscript selects components; the term with $\mathcal{N}^c \cup T$ empty is one. Each term has a saddlepoint in its closed positive orthant, or is decomposed again.

14.4 Ratios

Let $\mathcal{K}(s,t)$, $s,t \in \mathbb{R}^d$, be the joint cumulant generating function of (X,Y), $W_j = X_j - r_j Y_j$, and $r \circ s = (r_1 s_1, \ldots, r_d s_d)$. The cumulant generating function of W is $K_W(s) = \mathcal{K}(s, -r \circ s)$, and $\mathrm{P}(R \le r) = \mathrm{P}(W \le 0)$ is the orthant probability of $-W$ at zero. For the density, let

$$\mathcal{K}_B = \frac{\partial^{|B|} \mathcal{K}}{\prod_{j \in B} \partial t_j}, \qquad B \subseteq \{1, \ldots, d\},$$

and let $\mathcal{P}_{d,j}$ be the set of partitions of $\{1, \ldots, d\}$ into j blocks and

$$g_k(s) = \sum_{\pi \in \mathcal{P}_{d,d-k}} \prod_{B \in \pi} \mathcal{K}_B(s, -r \circ s), \qquad k = 0, \ldots, d-1.$$

With $\hat s$ the saddlepoint of K_W, $K_W'(\hat s) = 0$, and $K_W''(\hat s) = LL^{\mathsf{T}}$,

$$f_R(r) \approx \frac{\exp\{K_W(\hat s)\}}{(2\pi)^{d/2} |K_W''(\hat s)|^{1/2}} \sum_{m=0}^{M-1} \tilde a_m, \qquad \tilde a_m = \sum_{k=0}^{\min(m, d-1)} b_{m-k}(g_k),$$

$$b_m(h) = \sum_{r=0}^{2m} \sum_{j=3r}^{2m+2r} \sum_{|\nu|=j} \sum_{|\mu|=2m+2r-j} c_{r,\nu}\, \gamma_\mu\, \mathrm{He}_{\nu+\mu}(0), \qquad c_{0,0} = 1,$$

where θ_ν, and hence $c_{r,\nu}$, are the whitened coefficients of K_W at $\hat s$ and γ_μ the Taylor coefficients of $h(\hat s + L^{-\mathsf T}u)$ in u. The leading coefficient is $\tilde a_0 = g_0(\hat s) = \prod_{j=1}^{d} \mathcal{K}_{\{j\}}(\hat s, -r \circ \hat s)$. All the coefficients come from one Taylor polynomial of

$$\mathcal{K}\big(\hat s + L^{-\mathsf T}u,\ -r \circ (\hat s + L^{-\mathsf T}u) + \tau\big)$$

in (u, τ): the coefficient of $\prod_{j\in B} \tau_j$ is the Taylor polynomial of $\mathcal{K}_B$ in u. The sums b_m are evaluated as He-weighted sums of the coefficients of the products $\Theta(u)^r\, h(u)/r!$.

14.5 Studentized means

For $Q = X/\sqrt{Y}$ the inversion formulas of Bao, Broda and Kan are double integrals,

$$f_Q(q) = \frac{1}{(2\pi i)^2} \iint e^{K(s,t)}\, 2J_2(qs, t)\, dt\, ds, \qquad \mathrm{P}(Q > q) = \frac{1}{(2\pi i)^2} \iint e^{K(s,t)}\, 2J_1(qs, t)\, dt\, \frac{ds}{s},$$

with the kernels $J_k(b, t) = \int_0^\infty v^k e^{-bv - tv^2}\, dv$, whose exponent is $q^2s^2/(4t)$ plus Gaussian terms. With $\psi(s, t) = K(s, t) + q^2s^2/(4t)$ the density is

$$f_Q(q) \approx \frac{\hat v^2 e^{\psi(\hat s, \hat t)}}{\sqrt{\pi}\, \{\hat t\, |\psi''(\hat s, \hat t)|\}^{1/2}} \Big\{1 + \sum_{m=1}^{M-1} \big(b_m(h_0) + b_{m-1}(h_1)\big)\Big\},$$

the amplitude expansion of § 14.4 in two variables with the amplitudes $h_0 = (s/\hat s)^2 (t/\hat t)^{-5/2}$ and $h_1 = (t/\hat t)^{-3/2}/(2\hat t \hat v^2)$; the Hessian ψ'' is indefinite where $\hat t < 0$, and the Hermite moments are evaluated in the coordinates of a complex factorization $\psi'' = LL^{\mathsf T}$ from the eigen-decomposition $\psi'' = V\Lambda V^{\mathsf T}$. The double-double computation uses the real whitening $W = V|\Lambda|^{-1/2}$, $W^{\mathsf T}\psi'' W = \mathrm{diag}(\sigma)$ with $\sigma_i = \pm 1$, and the Hermite moments of the indefinite form, $\prod_i (-1)^{n_i/2} \sigma_i^{n_i/2} (n_i - 1)!!$, which gives terms of the same magnitudes. For the distribution function the t-integral is expanded by Laplace's method, which leaves a univariate integral of Hauschildt–Robinson type in s with the exponent $\Psi(s) = \psi(s, t(s))$, $\psi_t(s, t(s)) = 0$, and amplitudes $H_k(s)$; the Taylor coefficients of Ψ and of the H_k at $\hat s$ are obtained from the Taylor polynomial of K at $(\hat s, \hat t)$ by Newton's method and composition on power series, exact up to rounding.

14.6 Conditional expansions

Let $\mathcal{A}$, with a elements, be the tail components and $\mathcal{B}$, with $b = d - a$ elements, the conditioning ones; let $\hat s$ be the saddlepoint of K at x, $K'(\hat s) = x$, and $\tilde s_{\mathcal{B}}$ that of $K_{\mathcal{B}}(s_{\mathcal{B}}) = K(0_{\mathcal{A}}, s_{\mathcal{B}})$ at $x_{\mathcal{B}}$. At $\hat s$ let

$$C = K''_{\mathcal{A}\mathcal{B}} (K''_{\mathcal{B}\mathcal{B}})^{-1}, \qquad L_{\mathcal{B}} L_{\mathcal{B}}^{\mathsf T} = K''_{\mathcal{B}\mathcal{B}}, \qquad S = K''_{\mathcal{A}\mathcal{A}\cdot\mathcal{B}} = K''_{\mathcal{A}\mathcal{A}} - C K''_{\mathcal{B}\mathcal{A}} = L_{\mathcal{A}} L_{\mathcal{A}}^{\mathsf T}.$$

The canonical coordinates of the paper are $Y_{\mathcal{A}} = X_{\mathcal{A}} - C X_{\mathcal{B}}$ and $Y_{\mathcal{B}} = L_{\mathcal{B}}^{-1} X_{\mathcal{B}}$, in which the tilted covariance matrix is $\mathrm{diag}(S, I)$; the package whitens the tail components as well, so that the dual variables are $s = Tu$ with $s_{\mathcal{A}} = L_{\mathcal{A}}^{-\mathsf T} u_{\mathcal{A}}$ and $s_{\mathcal{B}} = -C^{\mathsf T} L_{\mathcal{A}}^{-\mathsf T} u_{\mathcal{A}} + L_{\mathcal{B}}^{-\mathsf T} u_{\mathcal{B}}$. With θ_ν the Taylor coefficients of $K(\hat s + Tu)$ of degrees three and more and $c_{r,\nu}$ the coefficients of $\Theta(u)^r/r!$, $c_{0,0} = 1$, the numerator of the conditional probability with the terms of orders $n^{-m/2}$, $m \le p$, is

$$\mathcal{N}_p = \frac{\exp\{K(\hat s) - \hat s^{\mathsf T} x\}}{(2\pi)^{b/2} |L_{\mathcal{B}}|} \sum_{m=0}^{p} \sum_{r=0}^{m} \sum_{|\nu|=m+2r} c_{r,\nu}\, \mathrm{He}_{\nu_{\mathcal{B}}}(0)\, J(\nu_{\mathcal{A}}),$$

$$J(\mu) = \frac{1}{(2\pi)^{a/2}|S|^{1/2}} \int_{w\geq 0} \prod_{i=1}^{a} \mathrm{He}_{\mu_i}((L_{\mathcal{A}}^{-1}w)_i)\, \exp(-\hat{s}_{\mathcal{A}}^{\mathsf{T}}w - w^{\mathsf{T}}S^{-1}w/2)\, dw,$$

the Gaussian of the integrand factoring into the tail and the conditioning components in these coordinates. The approximation of order M is

$$\mathrm{P}(X_{\mathcal{A}} \geq x_{\mathcal{A}} \mid X_{\mathcal{B}} = x_{\mathcal{B}}) \approx \mathcal{N}_{2M-1}/\mathcal{D}_M,$$

$\mathcal{D}_M$ the density expansion of $X_{\mathcal{B}}$ at $x_{\mathcal{B}}$ with M terms, and the conditional density is $f_M(x)/\mathcal{D}_M$, both density expansions in whitened coordinates. For one tail component $J(\mu)$ is a univariate integral, $J(\mu) = \int_0^\infty \mathrm{He}_\mu(v)\exp(-\hat{u}v - v^2/2)\, dv/\sqrt{2\pi}$ with $\hat{u} = \hat{s}_{\mathcal{A}}S^{1/2}$, computed by three Gauss–Legendre panels split at $1/(1+\hat{u})$ and $4/(1+\hat{u})$ on $[0, v_{\max}]$, where $v_{\max}$ is the point beyond which $v^L\exp(-\hat{u}v - v^2/2)$, $L = 3(2M-1)$ the highest degree, has fallen by e^{-40} from its maximum. For more tail components the integral is computed on a grid in sequentially conditioned coordinates of $N(0, S)$, each coordinate with three panels placed by its own conditional truncation point.

When $\hat{s}_j < 0$ for j in $\mathcal{N} \subseteq \mathcal{A}$,

$$\mathrm{P}(X_{\mathcal{A}} \geq x_{\mathcal{A}} \mid X_{\mathcal{B}} = x_{\mathcal{B}}) = \sum_{T\subseteq\mathcal{N}} (-1)^{|T|}\, \mathrm{P}(\tilde{X}_{(\mathcal{A}\setminus\mathcal{N})\cup T} \geq \tilde{x}_{(\mathcal{A}\setminus\mathcal{N})\cup T} \mid X_{\mathcal{B}} = x_{\mathcal{B}}),$$

where $\tilde{X}$ and $\tilde{x}$ are X and x with the components in T negated, the components of $\mathcal{N}\setminus T$ are marginalized ($s_j = 0$, not conditioned on), the term without tail components is one, and the denominator is the same in every term.

For lattice variables the saddlepoint $\hat{s}$ is that of the continuity-corrected point, $x_{\mathcal{A}} - 1/2$ in the tail components, and the integrand of the numerator carries the amplitude $h(s) = \prod_{j\in\mathcal{A}} g(s_j)$, $g(s) = s/\{2\sinh(s/2)\}$, whose Taylor coefficients γ_μ at $\hat{s}$, in the whitened variables $u_{\mathcal{A}}$, enter through the products $\tilde{c}_{r,\nu} = \sum_{\mu\leq\nu} c_{r,\nu-\mu}\gamma_\mu$ in place of $c_{r,\nu}$, now with $r = 0$ and $\nu \neq 0$ allowed (the corrections for the lattice). With `'sinh', 'origin'` the coefficients are those of the polynomial $\sum_{l\leq M-1}\beta_l s^{2l}$, $\beta_0 = 1$, $\beta_1 = -1/24$, $\beta_2 = 7/5760$, the expansion of g about zero truncated at order M.

14.7 Saddlepoint space

Quantiles and masses are computed with the point parametrized by its saddlepoint. In one dimension $x = K'(s)$, and $dx = K''(s)\, ds$; in d dimensions $dx = |K''(s)|\, ds$. For a ratio $R = X_1/X_2$ the cumulant generating function of $W = X_1 - rX_2$ is $K_W(s) = K(s, -rs)$, and $K'_W(\hat{s}) = 0$ means $K_1 = rK_2$ at $(\hat{s}, -r\hat{s})$; the saddlepoints of all r therefore lie on the curve

$$G(s,t) = sK_1(s,t) + tK_2(s,t) = 0,$$

on which $r = K_1/K_2$, $t' = -G_s/G_t$ and $dr/ds = \{(K_{11} + K_{12}t')K_2 - K_1(K_{12} + K_{22}t')\}/K_2^2$. For a Studentized mean the curve is the locus $sK_s + 2tK_t = 0$ of § 8, with $v = K_t^{1/2}$ and $q = K_s/v$.

15 Accuracy and precision

The computations are in double precision, except the double-double refinement of the density of a Studentized mean (§ 13.1). Rounding errors are about 10^{-14} relative in the tests, far below the asymptotic errors of the expansions at the orders used in practice (10^{-3} to 10^{-9}), so double precision suffices for nearly all purposes. It matters in three situations:

- *Cancellation in three or more dimensions.* At high orders the terms can cancel by many orders of magnitude. The engine reports this (`kappa` in `info`) and, when the cancellation exceeds 10^6, tries to repeat the computation in variable precision. The Taylor class

`tps` works in double precision, and its double-double counterpart is used only for Studentized means, so in this package that refinement is not possible: the double-precision values are kept and `status` reports `'refinement failed'`. The Julia package repeats the computation in `BigFloat`.

- *Studentized means.* The density of a Studentized mean can cancel strongly where the saddlepoint has large $|\hat{t}|$; it is then recomputed in double-double arithmetic (§§ 8 and 13.1), and `status` reports what remains.
- *Probabilities below about* 10^{-308}. In two dimensions use `'logscale'` (§ 5).
- *Tables of very small relative errors*, at large n and high order, need exact values and expansions beyond 15 digits. The engine programs accept derivatives in variable precision (`vpa`, Symbolic Math Toolbox) when they are supplied in closed form (§ 16).

16 The engine

The folder `engine` contains the programs of the papers. They take the derivatives of K from a function `Kfun(nu, t)`, the derivative of order `nu` ($1 \times d$) at `t` ($1 \times d$), `NaN` outside the domain, and can be used directly when closed-form derivatives are available:

`spa_saddle`	saddlepoint by Newton's method with backtracking
`spa_uni`	Daniels and Hauschildt–Robinson expansions from $K^{(k)}(\hat{\tau})$
`spa_ratio_pdf`	Rice's expansion of the density of a ratio
`spa_bi`, `spa_orthant2`	bivariate density and orthant probability
`spa_tri`, `spa_mv`, `spa_orthant`	three and more dimensions
`spa_cond`	conditional and lattice expansions (§ 9), any dimension
`qf_Kmulti`	derivatives of the cgf of quadratic forms
`ncx2_spa`, `ncf_spa`, `dncf_spa`	noncentral χ^2, F and doubly noncentral F
`spa_uni_hp`, `spa_bi_hp`	variable precision

The class `CGF` builds `Kfun` and the whitened derivatives (`opts.Kwhite`) from the Taylor polynomials. See the help of each program. The conditional program is called as

```
[P, f, info] = spa_cond(Kfun, x, A, M, opts)
```

with the options `lattice`, `sinh`, `nG`, `tau0`, `Kwhite`, `tail`, `density` and `tauf` (a known saddlepoint for the density) in the structure `opts`; without `Kwhite` the coefficients in the canonical coordinates are obtained by substituting the linear maps into the Taylor polynomial of K at the saddlepoint, which loses digits when K'' is ill conditioned.

17 Validation

`tests/test_spx.m` runs the following checks and returns `true` if all pass (`test_spx(true)` prints every result):

- derivatives of the Taylor class against closed forms and finite differences, and the derivatives of a pair of quadratic forms by automatic differentiation through a generic Cholesky factorization against the trace recursion;
- the 559 reference values shared by the Julia and Python packages (`tests/julia_reference.csv`, generated by `test/reference_values.jl` of the Julia package; `tests/julia_reference.jl` copies them): derivatives, saddlepoints, orthant probabilities with lower and mixed corners and saddlepoints with negative

components, and densities in one to four dimensions (sums of exponentials including the examples of Wang and of Kolassa and Li, gamma, noncentral chi-squared, uniform, common-component, quadratic-form, Wishart and normal distributions), single ratios (the noncentral F) and several ratios. They agree to about 10^{-13} relative, at most 10^{-11}; the four-dimensional orthant probabilities, computed by quasi-Monte Carlo, agree within three standard errors;

- the expansion for several ratios at $d = 1$ against Rice's expansion (10^{-14}), and for two and three ratios against the inverted Dirichlet density and distribution function and the product of two independent beta-prime ratios, including the rate $O(n^{-4})$ of the density of order four;

- the tests of the Studentized-mean functions of the Julia and Python packages: the noncentral t distribution against exact values (distribution function to 2×10^{-9} at order three, density to 10^{-6}, the quantile and the saddlepoint locus), and the Dickey–Fuller t-ratio against the reference values of Bao, Broda and Kan (the distribution function with every half-order to 10^{-8}, the density to 2×10^{-6}, a saddlepoint and a probability under a local alternative, and the 5% quantile of model 2). The values agree with those of the Python package `saddlepoint-expansions` 0.2.2 to about 10^{-13} for the noncentral t and to 10^{-11}–10^{-9} for the Dickey–Fuller model, whose derivatives of high order carry rounding errors;

- double-double arithmetic: `dd` against `vpa` (about 10^{-32}), the derivatives of a `tpsdd` series against symbolic differentiation (about 10^{-32}), and the Studentized expansions in double-double against the Julia package in `BigFloat` with 256 bits (`tests/studentized_reference.csv`, generated by `tests/julia_studentized_reference.jl`): the density of orders one to six and the distribution function with every half-order up to order four, for the noncentral t at $q = 0.5$ and the Dickey–Fuller model under a local alternative, agree to the last digit of double precision, and the refined density of the Dickey–Fuller model 2 to 10^{-15} at order five. Over all the reference values (the noncentral t at $q = 0.5, 1.5, 2.5$ and the three Dickey–Fuller models), double-double agrees to the last digit except the density of model 2 at order five (10^{-15}) and order six (4×10^{-10}, $\kappa \approx 10^{23}$); for that model double precision has errors of 2×10^{-9} at order three and gives no correct digit at order six;

- the conditional expansions, lattice sums, quantiles and normalized densities against the Julia package 0.3.0: the 150 rows of the third part of `julia_reference.csv` and the 247 rows of `tests/conditional_reference.csv`, generated by `tests/julia_conditional_reference.jl` from the Julia package (masses in two dimensions and of a Studentized mean, one, two and three tail components of a Gaussian vector with saddlepoints of every sign, `spa_cond` with closed-form derivatives, quantiles, and conditional expansions in `BigFloat` with 256 bits). They agree to about 10^{-15}, at most 8×10^{-14}, except `spa_cond` with closed-form derivatives, whose coefficients are obtained by substitution into the Taylor polynomial (6×10^{-13}), and the mass of a Studentized density (7×10^{-11}, its quadrature having the relative tolerance 10^{-9}); the double-precision values agree with those in `BigFloat` to 4×10^{-15} absolute for the gamma sum of order five and to 10^{-14} relative otherwise;

- every printed cell of the tables of the conditional paper, recomputed from the cumulant generating function alone against exact values (the rows in `tests/conditional_rows.txt`, shared with the Julia and Python tests): the gamma sum given its total (450 cells, against the beta distribution), a quadratic form given another (300, against the exact conditional distribution by quadrature over the common component), the means of five and ten in-

dependent ratios of quadratic forms (150 of 155 cells, conditional on five and ten variables, against the exact quantiles of the paper; Skovgaard's formula at the median, where $1/\hat{w} - 1/\hat{z}$ cancels in double precision, is left out), the logistic regression (189, against the exact distribution by dynamic programming) and the lattice sums (88), each relative error to its two printed digits;

- the consistency checks of the Julia tests: the expansions with $\mathcal{B}$ empty against `tailprob` and `density`; Gaussian vectors, for which the conditional expansions are exact at every order, with one, two and three tail components, saddlepoints of every sign, lower tails and mixed corners (10^{-13}); the Dirichlet distribution; `spa_cond` with closed-form derivatives and with the derivatives of `CGF`; the leading term against the conditional Hauschildt–Robinson formula (20) of the paper (10^{-13}); the complement identities of the lattice tails (10^{-14}); binomial, negative binomial and Poisson tails and probabilities against exact values; the quantiles of the gamma and the noncentral F (the expansion at the quantile equals p to $10^{-12}\min(p, 1-p)$, and the quantile of the normal distribution is exact); and the normalized densities, which integrate to one (10^{-9} to 10^{-8}), with the masses of the gamma distribution in one and two dimensions against their closed forms.

In addition, for all 24 points of the table of sample autocorrelations of the paper, the density and distribution function of two ratios computed by the package agree with those of the dedicated program of the paper (`acf_spa`) to 1.5×10^{-12} at every order from one to five.

18 Function reference

call	result
`K = CGF(fun, d, 'domain', pred, 'Taylor', tfun)`	a cumulant generating function
`r = tailprob(K, x, 'order', M, 'lower', tf, 'signs', s, ...)`	orthant probabilities; `r.F`, `r.tau`, `r.status`, `r.info`
`r = density(K, x, 'order', M, ...)`	densities; `r.f`, `r.tau`, `r.status`, `r.info`
`r = ratio_density(K, r0, 'order', M)`	densities of $d =$ `numel(r0)` ratios; `r.f`, `r.shat`, `r.info` (for $d \geq 2$: the terms `a`, the log prefactor `logC`)
`r = ratio_tailprob(K, r0, 'order', M, 'lower', tf, 'signs', s, ...)`	distribution functions of ratios; `r.F`, `r.shat`, `r.status`, `r.info`
`r = studentized_density(K, q, 'order', M, 'saddle', sd, 'refine', tf, 'precision', p)`	density of $X/\sqrt{Y}$; `r.f`, `r.saddle`, `r.status`, `r.info` (`'refine'`, default `true`: double-double when $\kappa > 10^6$; `p`: `'double'` or `'dd'`)
`r = studentized_tailprob(K, q, 'order', M, 'lower', tf, 'saddle', sd, 'precision', p)`	distribution function of $X/\sqrt{Y}$; `r.F`, `r.saddle`, `r.status`, `r.info.Fp`, `r.info.kappa`
`r = studentized_quantile(K, p, 'order', M, 'tol', tol)`	quantile of the expansion; `r.q`, `r.saddle`
`r = conditional_tailprob(K, x, A, 'order', M, 'lower', tf, 'signs', s, 'lattice', tf, 'sinh', m, 'nG', n)`	$\mathrm{P}(X_{\mathcal{A}} \geq x_{\mathcal{A}} \mid X_{\mathcal{B}} = x_{\mathcal{B}})$; `r.F`, `r.tau`, `r.taut`, `r.status`, `r.info` (`Fp`, `kappa`, `kappa_den`, `err`, `nG`); `m`: `'saddle'` or `'origin'`

call	result
`r = conditional_density(K, x, A, 'order', M, 'lattice', tf)`	conditional density (probability for lattice variables); `r.f`, `r.tau`, `r.taut`, `r.status`, `r.info` (`kappa`, `kappa_den`)
`r = tailprob(K, k, 'lattice', true, 'sinh', m)`, `density(K, k, 'lattice', true)`	$\mathrm{P}(S \geq k)$ and $\mathrm{P}(S = k)$ of a lattice variable
`r = quantile(K, p, 'order', M, 'lower', tf, 'tol', tol)`	quantile of the expansion; `r.q`, `r.tau`, `r.F`, `r.iterations`, `r.status`
`r = ratio_quantile(K, p, 'order', M, 'lower', tf, 'tol', tol)`	quantile of the expansion of a ratio; `r.r`, `r.shat`, `r.F`, `r.iterations`, `r.status`
`'normalize', true` or `'normalize', c` in `density`, `ratio_density`, `studentized_density`	normalized expansions; `r.info.mass`
`sd = studentized_saddlepoint(K, q)`, `studentized_locus(K, s, tguess)`	the saddlepoint and the saddlepoint locus
`[tau, info] = saddlepoint(K, x)`	the saddlepoint
`D = derivatives(K, t, N)`	`D.nu`, `D.value`
`p = taylor(K, t, N, M)`	Taylor polynomial of $K(t + Mv)$
`v = eval(K, t)`	$K(t)$
`reflect`, `marginal`, `linear_combination`, `sample_mean`, `shift`, `+`	transformations (§ 4)
`tps.vars(k, N)`, `tps.const(c, k, N)`, `coef(p, nu)`, `double(p)`, `tps.tables(k, N)`	the Taylor class
`x = dd(a)`, `dd(hi, lo)`, `vpa(x, n)`, `double(x)`	double-double numbers (§ 13.1)
`tpsdd.vars(k, N)`, `tpsdd.const(c, k, N)`, `tpsdd.fromtps(p)`, `coef(p, nu)`, `ddconst(p)`	the Taylor class in double-double
`p = taylordd(K, t, N, M)`	Taylor polynomial of $K(t + Mv)$ in double-double
`sinhc(x)`, `sincu(x)`	$\sinh(x)/x$ and $\sin(x)/x$
`spx_setup`	adds the package to the path

References

Bao, Y., Broda, S. A. and Kan, R. Inversion formulas and saddlepoint expansions for Studentized means. Working paper, 2026.

Barndorff-Nielsen, O. E. and Cox, D. R. (1979). Edgeworth and saddle-point approximations with statistical applications. *J. R. Statist. Soc. B* 41, 279–312.

Broda, S. A. and Kan, R. Conditional and lattice saddlepoint expansions of arbitrary order. Working paper, 2026.

Broda, S. A. and Kan, R. Uniform saddlepoint expansions for multivariate tail probabilities. Working paper, 2026.

Butler, R. W. (2007). *Saddlepoint Approximations with Applications.* Cambridge University Press.

Butler, R. W. and Paolella, M. S. (1998). Approximate distributions for the various serial correlograms. *Bernoulli* 4, 497–518.

Daniels, H. E. (1954). Saddlepoint approximations in statistics. *Ann. Math. Statist.* 25, 631–650.

Daniels, H. E. (1987). Tail probability approximations. *Int. Statist. Rev.* 55, 37–48.

Hauschildt, R. (1969). Saddlepoint approximations to distribution functions. Master's thesis, University of British Columbia, Vancouver.

Kolassa, J. E. (2003). Multivariate saddlepoint tail probability approximations. *Ann. Statist.* 31, 274–286.

Kolassa, J. E. (2006). *Series Approximation Methods in Statistics*, 3rd edition. Springer, New York.

Kolassa, J. E. and Li, J. (2010). Multivariate saddlepoint approximations in tail probability and conditional inference. *Bernoulli* 16, 1191–1207.

Nabeya, S. and Sørensen, B. E. (1994). Asymptotic distributions of the least-squares estimators and test statistics in the near unit root model with non-zero initial value and local drift and trend. *Econometric Theory* 10, 937–966.

Nabeya, S. and Tanaka, K. (1990). A general approach to the limiting distribution for estimators in time series regression with nonstable autoregressive errors. *Econometrica* 58, 145–163.

Skovgaard, I. M. (1987). Saddlepoint expansions for conditional distributions. *J. Appl. Probab.* 24, 875–887.