

A method for generating triangulated 2-spheres of arbitrary geometry

Joshua H. Cooperman

Department of Physics, University of California, Davis, CA 95616

Jonah M. Miller

Department of Physics, University of Colorado, Boulder, CO 80309

August 24, 2012

Consider a 2-sphere \mathcal{S}^2 triangulated by N_2 regular 2-simplices. Let $\{|\Upsilon_\lambda^\mu\rangle\}$ for $\lambda \in \{0, \dots, l_{N_2}\}$ and $\mu \in \{-\lambda, \dots, \lambda\}$ be the set of discrete spherical harmonics required to characterize the geometry of the 2-sphere \mathcal{S}^2 composed of the finite number N_2 of regular 2-simplices. Denoting by $|\mathcal{G}_{\mathcal{S}^2}\rangle$ the geometry of the 2-sphere \mathcal{S}^2 , we suggestively express the geometry as a linear combination of the discrete spherical harmonics,

$$|\mathcal{G}_{\mathcal{S}^2}\rangle = \sum_{\lambda=0}^{l_{N_2}} \sum_{\mu=-\lambda}^{\lambda} c_\mu^\lambda |\Upsilon_\lambda^\mu\rangle. \quad (1)$$

The set of coefficients $\{c_\mu^\lambda\}$ thus completely specifies the geometry $|\mathcal{G}_{\mathcal{S}^2}\rangle$. We define an inner product $A[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ on the space of geometries of 2-spheres as the projection of one geometry $|\mathcal{G}_{\mathcal{S}^2}\rangle$ onto another geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$:

$$A[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}] = \langle \mathcal{G}_{\tilde{\mathcal{S}}^2} | \mathcal{G}_{\mathcal{S}^2} \rangle. \quad (2)$$

Assuming that the set $\{|\Upsilon_\lambda^\mu\rangle\}$ of discrete spherical harmonics is orthonormal for any number N_2 of regular 2-simplices, the inner product $A[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ assumes the form

$$A[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}] = \sum_{\lambda=0}^{\min[l_{N_2}, l_{\tilde{N}_2}]} \sum_{\mu=-\lambda}^{\lambda} \tilde{c}_\mu^\lambda c_\mu^\lambda \quad (3)$$

Given the inner product $A[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$, we further define a measure $\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ of the closeness of the geometry $|\mathcal{G}_{\mathcal{S}^2}\rangle$ to the geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$ as

$$\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}] = A[\mathcal{G}_{\tilde{\mathcal{S}}^2} - \mathcal{G}_{\mathcal{S}^2} | \mathcal{G}_{\tilde{\mathcal{S}}^2}]. \quad (4)$$

The measure $\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ readily simplifies to

$$\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}] = 1 - \sum_{\lambda=0}^{\min[l_{N_2}, l_{\tilde{N}_2}]} \sum_{\mu=-\lambda}^{\lambda} \tilde{c}_\mu^\lambda c_\mu^\lambda. \quad (5)$$

Conceiving of the measure $\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ as an action functional on the space of geometries of 2-spheres, we construct a partition function

$$Z(\tilde{\mathcal{S}}^2) = \sum_{\mathcal{G}_{\mathcal{S}^2}} e^{-\beta \mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]} \quad (6)$$

for the chosen fixed 2-sphere $\tilde{\mathcal{S}}^2$ triangulated by \tilde{N}_2 regular 2-simplices of geometry $\mathcal{G}_{\tilde{\mathcal{S}}^2}$. β is a tunable coupling constant. The partition function $Z(\tilde{\mathcal{S}}^2)$ is related by Legendre transform to the partition function $Z_{\tilde{N}_2}(\tilde{\mathcal{S}}^2)$ for fixed number \tilde{N}_2 of regular 2-simplices:

$$Z(\tilde{\mathcal{S}}^2) = \sum_{N_2} Z_{\tilde{N}_2}(\tilde{\mathcal{S}}^2) e^{-N_2}. \quad (7)$$

Suppose now that we wish to generate a 2-sphere $\tilde{\mathcal{S}}^2$ triangulated by \tilde{N}_2 regular 2-simplices of geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$ specified by the set of coefficients $\{\tilde{c}_\mu^\lambda\}$. We propose the following method: design a Monte Carlo algorithm based on the partition function $Z_{\tilde{N}_2}(\tilde{\mathcal{S}}^2)$. Interpreting the weighting factor $e^{-\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]}$ as the conditional probability of the geometry $|\mathcal{G}_{\mathcal{S}^2}\rangle$ given the geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$, those geometries $|\mathcal{G}_{\mathcal{S}^2}\rangle$ closest to the geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$ are most probable. Since a Monte Carlo algorithm is designed to produce an ensemble of geometries representative of the partition function $Z_{\tilde{N}_2}(\tilde{\mathcal{S}}^2)$, the algorithm should converge on geometries close to the given geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$.

Our Monte Carlo algorithm functions as follows. First, we choose the fixed geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$ —in particular, the set of coefficients $\{\tilde{c}_\mu^\lambda\}$ —to define the measure $\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$. Next, we initialize our simulation by generating the 2-sphere \mathcal{S}_{min}^2 minimally triangulated by regular 2-simplices, namely the regular tetrahedron. Then, by applying the known ergodic moves on the space of 2-surfaces triangulated by regular 2-simplices, we increase the number N_2 of regular 2-simplices of the 2-sphere \mathcal{S}_{min}^2 to the value \tilde{N}_2 of regular 2-simplices of the fixed geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$. Next, we run a standard Metropolis algorithm based on the partition function $Z_{\tilde{N}_2}(\tilde{\mathcal{S}}^2)$. To determine whether particular ergodic moves are accepted or rejected as the Metropolis algorithm runs, we must compute the change in the measure $\mu[\mathcal{G}_{\mathcal{S}^2}|\mathcal{G}_{\tilde{\mathcal{S}}^2}]$ upon implementation of the ergodic move. This requires that we compute the sets of coefficients $\{c_\mu^\lambda\}$ characterizing the intermediate geometries $|\mathcal{G}_{\mathcal{S}^2}\rangle$; we employ the technique of Sachs to perform these computations [1]. After the Monte Carlo algorithm has run for a sufficient computational time, the geometry $|\mathcal{G}_{\mathcal{S}^2}\rangle$ should be sufficiently close to the fixed geometry $|\mathcal{G}_{\tilde{\mathcal{S}}^2}\rangle$; we validate this closeness by employing again the technique of Sachs.

References

- [1] M. K. Sachs. “Testing Lattice Quantum Gravity in 2 + 1 Dimensions.” arXiv: gr-qc/1110.6880.