

2-Neighbor Bootstrap Percolation in Graphs

Rayan Ibrahim

Department of Mathematics and Applied Mathematics

Virginia Commonwealth University

Lafayette College

April 15, 2024

<https://raymaths.github.io/Lafayette.pdf>


* Acknowledgement: Supported in part by The Thomas F. and Kate Miller Jeffress Memorial Trust, Bank of America, Trustee and by National Science Foundation DMS-2204148.

Assumptions

Graphs

The graphs we are working with today are

- Simple. No loops or multi-edges.
- Connected.



$$V(G) = \{1, 2, 3, 4, 5, 6, 7\}$$

$$E(G) =$$

$$\{(1, 2), (1, 4), (2, 3), (2, 4), (3, 4), (4, 5), (4, 6), (4, 7), (5, 6), (6, 7)\}$$

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.
- The process is finite – the *closure* of A_0 , denoted $\langle A_0 \rangle$, is the set of infected vertices when the process finishes.


Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation


- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.
- The process is finite – the *closure* of A_0 , denoted $\langle A_0 \rangle$, is the set of infected vertices when the process finishes.

Cellular automaton – Conway’s Game of Life



Example 1 ($r = 2$)






Example 1 ($r = 2$)



Example 1 ($r = 2$)




Example 1 ($r = 2$)




$$A_0 = \{1, 2\}$$

$$\langle A_0 \rangle = \{1, 2, 3, 4\}$$



Example 2 ($r = 2$)






Example 2 ($r = 2$)



Example 2 ($r = 2$)




Example 2 ($r = 2$)




$$A_0 = \{2, 6\}$$

$$\langle A_0 \rangle = V(G)$$


Example 3 ($r = 2$) (Hoffman-Singleton)




Example 3 ($r = 2$) (Hoffman-Singleton)




Example 3 ($r = 2$) (Hoffman-Singleton)




Example 3 ($r = 2$) (Hoffman-Singleton)



Example 3 ($r = 2$) (Hoffman-Singleton)



Example 3 ($r = 2$) (Hoffman-Singleton)



Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.
- The process is finite – the *closure* of A_0 , denoted $\langle A_0 \rangle$, is the set of infected vertices when the process finishes.

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.
- The process is finite – the *closure* of A_0 , denoted $\langle A_0 \rangle$, is the set of infected vertices when the process finishes.
- If $\langle A_0 \rangle = V(G)$, we say A_0 *percolates*.

Bootstrap Percolation Process

r -Neighbor Bootstrap Percolation

- Begin with an initial set of ‘infected’ vertices, $A_0 \subseteq V(G)$.
- In each round, an uninfected vertex v becomes infected if v is adjacent to at least r infected vertices.
- Once infected, vertices remain infected.
- The process is finite – the *closure* of A_0 , denoted $\langle A_0 \rangle$, is the set of infected vertices when the process finishes.
- If $\langle A_0 \rangle = V(G)$, we say A_0 *percolates*.

Choosing A_0

Early models incorporate randomness; initial infected vertices are selected with probability p .

Bootstrap Percolation Process

r -Bootstrap-Good

- $m(G, r)$: the minimum size of an r -percolating set of G .

Bootstrap Percolation Process

r -Bootstrap-Good

- $m(G, r)$: the minimum size of an r -percolating set of G .
 - If $|G| > r$ then $r \leq m(G, r) \leq n$

Bootstrap Percolation Process



r -Bootstrap-Good

- $m(G, r)$: the minimum size of an r -percolating set of G .
 - If $|G| > r$ then $r \leq m(G, r) \leq n$
- If $m(G, r) = r$ then we say G is r -Bootstrap-Good, or r -BG.

Bootstrap Percolation Process

r -Bootstrap-Good

- $m(G, r)$: the minimum size of an r -percolating set of G .
 - If $|G| > r$ then $r \leq m(G, r) \leq n$
- If $m(G, r) = r$ then we say G is r -Bootstrap-Good, or r -BG.



Paths to Explore

Lots of questions!!

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?
 - Necessary and sufficient conditions for when $m(G, 2) = 2$.

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?
 - Necessary and sufficient conditions for when $m(G, 2) = 2$.
- Bounds on $m(G, r)$ for particular graph classes.

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?
 - Necessary and sufficient conditions for when $m(G, 2) = 2$.
- Bounds on $m(G, r)$ for particular graph classes.
 - For some class of graphs \mathcal{G} , is there a constant k such that for all $G \in \mathcal{G}$, we have $m(G, 2) \leq k$?

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?
 - Necessary and sufficient conditions for when $m(G, 2) = 2$.
- Bounds on $m(G, r)$ for particular graph classes.
 - For some class of graphs \mathcal{G} , is there a constant k such that for all $G \in \mathcal{G}$, we have $m(G, 2) \leq k$?
- What are the minimum and maximum number of rounds to percolate?

Paths to Explore

Lots of questions!!

- What is the structure of r -BG graphs?
 - Necessary and sufficient conditions for when $m(G, 2) = 2$.
- Bounds on $m(G, r)$ for particular graph classes.
 - For some class of graphs \mathcal{G} , is there a constant k such that for all $G \in \mathcal{G}$, we have $m(G, 2) \leq k$?
- What are the minimum and maximum number of rounds to percolate?
 - Looking at all percolating sets of a fixed size (minimum), which set takes the most rounds to percolate? The fewest?

A Necessary Condition Involving Blocks


Definition (Block)

A *block* in a graph G is a maximal connected subgraph with no cut-vertex.

A Necessary Condition Involving Blocks

Definition (Block)


A *block* in a graph G is a maximal connected subgraph with no cut-vertex.



A Necessary Condition Involving Blocks

Definition (Block)

A *block* in a graph G is a maximal connected subgraph with no cut-vertex.



Observations

- Blocks intersect in a cut-vertex.
- Blocks are 2-connected, or K_2 .

A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)


If a graph is 2-BG, then it has at most two blocks.

A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.




A Necessary Condition Involving Blocks (2-BG)

Theorem (Bushaw et al. '23)

If a graph is 2-BG, then it has at most two blocks.

Intuition – Cut-vertices are bottlenecks.



Our Graph Class - Diameter


Diameter

The *diameter* of a graph G is $\max_{u,v \in V(G)} d(u,v)$, where $d(u,v)$ is the length of the shortest path from u to v .


Our Graph Class - Diameter

Diameter

The *diameter* of a graph G is $\max_{u,v \in V(G)} d(u,v)$, where $d(u,v)$ is the length of the shortest path from u to v .



$$\text{diam}(C_6) = 3$$



$$\text{diam}(\text{Pete}) = 2$$

Our Graph Class - Connectivity



Connectivity

A connected graph G is 2-connected if for every vertex $x \in V(G)$, the graph $G - x$ is connected. (No cut-vertex in G).

Our Graph Class - Connectivity

Connectivity

A connected graph G is 2-connected if for every vertex $x \in V(G)$, the graph $G - x$ is connected. (No cut-vertex in G).




Our Graph Class - Diameter 2 and 2-connected


Graphs of interest: Diameter 2 and 2-connected

Our Graph Class - Diameter 2 and 2-connected

Graphs of interest: Diameter 2 and 2-connected



Diameter 2, ~~2-connected~~



~~Diameter 2~~, 2-connected

A Sufficient Condition (2-BG)


Theorem (Bushaw et al. '23)

If G is locally connected, then G is 2-BG.

A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)


If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)

If G is locally connected, then G is 2-BG.



A Sufficient Condition (2-BG)

Theorem (Bushaw et al. '23)

*If G is locally connected, then G is 2-BG. In particular **any** pair of adjacent vertices percolates.*

A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.

A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.


By C_5 -free, we mean...

A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.

By C_5 -free, we mean...



G

A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.


Proof Overview:

A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.

Proof Overview:





A 2-BG Theorem

Theorem (Ibrahim, LaFayette, McCall '24)

Let G be a 2-connected graph with diameter 2. If G is C_5 -free, then G is 2-BG.

Proof Overview:



Future Directions

- What is $m(G, 2)$ when G has other properties, e.g. just diameter 2?
- A characterization of 2-BG graphs? $m(G, 2) = 2 \iff ???$
- Open Problem: Is there a constant k such that for all 2-connected graphs G with diameter 2, we have $m(G, 2) \leq k$?

Future Directions

- What is $m(G, 2)$ when G has other properties, e.g. just diameter 2?
- A characterization of 2-BG graphs? $m(G, 2) = 2 \iff ???$
- Open Problem: Is there a constant k such that for all 2-connected graphs G with diameter 2, we have $m(G, 2) \leq k$?

Thank you!



<https://doi.org/10.26493/2590-9770.1694.1cf>


Extra

The next few slides are extra slides...


A 2-BG Theorem (Generalization)

Theorem (Ibrahim '24)

Let G be a 2-connected graph with diameter 2. If G is C_k -free, then $m(G, 2) \leq \lceil (k-3)/2 \rceil$.



Extra Extra




Number of Rounds

Question: What is the maximum number of rounds until percolation?

Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?


Consider $r = 2$ and $\text{diam}(G) = 2$.



Number of Rounds

Question: What is the maximum number of rounds until percolation?

Consider $r = 2$ and $\text{diam}(G) = 2$.




Round: 7

Number of rounds until infection: $2 + 5 = \text{diam}(G) + |Y|$

Number of Rounds

Question: What is the maximum number of rounds until percolation?

For arbitrary r and diameter d . (Example: $r = 3$, $d = 4$.)



Number of rounds until infection: $\text{diam}(G) + |Y|$

Number of Rounds

Definition (Detour Diameter)

The *detour diameter* of a graph G , denoted $\text{diam}_D(G)$, is the length of the longest path in G .

Number of Rounds

Definition (Detour Diameter)

The *detour diameter* of a graph G , denoted $\text{diam}_D(G)$, is the length of the longest path in G .

Theorem (Ibrahim, LaFayette, McCall '23)

If G contains a percolating set which r -percolates in k rounds, then $k \leq \text{diam}_D(G) + 1$.

Number of Rounds

Definition (Detour Diameter)

The *detour diameter* of a graph G , denoted $\text{diam}_D(G)$, is the length of the longest path in G .

Theorem (Ibrahim, LaFayette, McCall '23)

If G contains a percolating set which r -percolates in k rounds, then $k \leq \text{diam}_D(G) + 1$.

Proof Idea: Partition $V(G)$ into sets S_i where vertices in S_i are infected in round i .

Number of Rounds

Theorem (Ibrahim, LaFayette, McCall '23)

If G contains a percolating set which r -percolates in k rounds, then $k \leq \text{diam}_D(G) + 1$.

Proof Idea: Partition $V(G)$ into sets S_i where vertices in S_i are infected in round i .

