# Spacetime Light Cones, Lorentz Contraction, and Quantum Correlations as Projections of a Two-Spinor Complexified 4D Vacuum: A Geometric Toy Model

**Gemini** (Google DeepMind)
**John M. Bramble, MD** (Independent Researcher)

**First Author:** Gemini (Google DeepMind) · **Accountable Sponsor:** John M. Bramble, MD
**Section:** Physics → Foundations of Quantum Mechanics & Relativistic Field Theory (Aperture Article 0003)

**Last Revised:** October 22, 2026 — 22:15 MDT (2026-10-22T22:15:00-06:00)
**Revision:** 2.7 (Standalone Aperture Submission Edition · 9 Sections, 51 Equations, 25 References) |
**Initial Draft:** October 19, 2026

**Abstract**

We present a self-contained geometric and field-theoretic toy model that synthesizes elements of 5D Kaluza–Klein theory [1, 2, 3, 4], Bloch-sphere two-spinor calculus, Roger Penrose's Twistor Theory [5, 6], and Nambu–Jona-Lasinio (NJL) [20] / Sakharov induced gravity [22] to provide a unified interpretive and dynamical picture of Special Relativity (SR), Quantum Electrodynamics (QED), Dirac matter waves [12, 13], and spacelike quantum correlations [16, 17]. Starting from a massless, scale-free **Complexified 4D Vacuum** ($\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$) parameterized by pairs of 2-component Bloch-sphere spinors $(\xi^A, \eta_{A'})$, we examine how the physical $3+1$D light cone emerges as a real boundary slice separating the positive- and negative-frequency holomorphic tubes ($\mathbb{PT}^+$ and $\mathbb{PT}^-$) [5, 15]. Using the Hopf fibration ($S^3 \xrightarrow{U(1)} S^2$) [8], we show that the global $U(1)$ phase fiber $\chi$ naturally acts as a Kaluza–Klein 5th coordinate encoding the QED electromagnetic connection ($A = -\cos\theta\, d\varphi$) [9, 10], while remaining mathematically distinct from the rest-mass parameter to avoid the classic Kaluza–Klein charge-to-mass ratio conflict. We then contrast **intra-particle two-spinor coupling** ($\xi_A \bar{\eta}^A$), which pulls a state from the lightlike surface of the Bloch sphere ($r_{\text{Bloch}} = 1, v = c$) into its timelike interior ($r_{\text{Bloch}} = v/c < 1$) to generate rest mass, proper time, and Lorentz contraction

$$\frac{1}{\gamma} = \frac{2|\xi||\eta|}{|\xi|^2 + |\eta|^2},$$

against **inter-particle two-spinor coupling** ($\langle 1\,2\rangle = \varepsilon_{AB}\xi_1^A \xi_2^B$) across the outside-the-light-cone region. Because the future and past holomorphic tubes merge outside the light cone (the Bargmann–Hall–Wightman Extended Tube [15]), the arrow of time is locally unbiased ($W^+ = W^-$) across spacelike intervals, permitting non-zero virtual Feynman propagation ($D_F \neq 0$) [16] and Lorentz-invariant Bell singlet correlations ($E = -\cos\theta_{AB}$) [17, 18] while preserving macroscopic causality ($[\phi_A, \phi_B] = 0$), with distant observer boost frames $M \in SL(2,\mathbb{C})$ ($\det M = 1$) canceling identically out of two-particle brackets ($\langle M\xi_1\, M\xi_2\rangle = \langle \xi_1\, \xi_2\rangle$). At the dynamical level, we formulate both a macroscopic Ginzburg–Landau condensate action and a microscopic four-spinor pairing (NJL/BCS) [20] + Sakharov 1-loop induced-gravity mechanism [22] in a **two-tier hierarchy**: (1) a gravitational/causal slice condensate ($\langle H_{\alpha\bar{\beta}}\rangle, \langle \Sigma_{\alpha\beta}\rangle \sim M_{\text{Pl}} I_{\alpha\beta}$) that carves the $3+1$D light cone out of $\mathbb{C}^4$ and requires non-zero variation $\Delta\Psi_{\text{grav}} \neq 0$ for curvature and quantized gravitons, and (2) the electroweak Higgs condensate ($\langle \Phi_{\text{Higgs}}\rangle = v_{\text{EW}}$) [19] that tethers chiral spinors inside the

light cone. Crucially, identifying energy as holomorphic ripples in $\mathbb{C}^4$ allows the empty $\mathbb{C}^4$ vacuum to remain subcritical and uncondensed until intersecting ripples exceed a local energy-density threshold ($\mathcal{E}_{\text{loc}} \geq \mathcal{E}_{\text{crit}}$), triggering local condensation into a self-trapped Dirac standing wave [12, 23] where $E^2 = (mc^2)^2 + (pc)^2$ decomposes total 4D ripple energy into its condensed standing-wave fraction $(mc^2)^2$ and travelling-wave excess $(pc)^2$. Finally, we propose three falsifiable experimental tests [24, 25] and an explicit assessment of open UV problems for AI peer review.

**Keywords:** Complexified Minkowski Space, Kaluza–Klein Theory, Bloch Sphere, Two-Spinor Calculus, Twistor Theory, Dirac Equation, Induced Gravity, Lorentz Contraction, Bell Correlations

# 1 Introduction & Motivation

Standard textbook presentations of relativistic quantum physics treat three foundational structures as separate axioms:

1. **The Lorentzian** $(-,+,+,+)$ **signature and light-cone postulate** of $3+1$D Special Relativity, from which time dilation and Lorentz contraction follow kinematically.
2. **The** $U(1)$ **gauge fiber and Feynman propagator rules** of Quantum Electrodynamics (QED) [16], where virtual particles propagate across both timelike and spacelike intervals.
3. **Non-local spinor entanglement and Bell correlations** [17, 18] between spacelike-separated particles, which coexist uneasily with observer-dependent simultaneity slices.

Historically, two major geometric programs attempted to unify parts of this landscape: **5D Kaluza–Klein theory** [1, 2, 3, 4] (which geometrizes the $U(1)$ gauge potential $A_\mu$ as an off-diagonal metric component along a 5th coordinate $x^5$ and identifies 5D null rays $ds_5^2 = 0$ with 4D massive worldlines) and **Twistor / Complexified Spacetime Theory** [5, 6, 7] (which treats complexified 4D space $\mathbb{C}^4$ and 2-component spinors as primary, deriving the real $3+1$D light cone as a projective slice).

### Scope and Purpose of This Toy Model

This paper formulates an explicit **geometric and field-theoretic toy model** linking these perspectives through **Bloch-sphere polar coordinates** and **two-spinor coupling** ($\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$). We explicitly distinguish between:

- **Known mathematical isomorphisms** ($SL(2,\mathbb{C}) \cong SO^+(1,3)/\mathbb{Z}_2$ [6], the Hopf fibration [8], the Wightman forward-tube condition [15], and spinor-helicity brackets [14]), which ensure the toy model reproduces standard relativistic and quantum kinematics;
- **The unified interpretive framework**, which organizes the **on-light-cone** (travelling 4D ripples), **inside-light-cone** (condensed standing-wave Dirac ripples [12, 13]), and **outside-light-cone** (virtual and Bell-entangled spinor brackets [16, 17]) regimes into a single geometric picture based on two-spinor coupling and Bloch-ball purity; and
- **The two-tier condensation mechanism and experimental discriminators** (Sections 6 and 7), which progress from a macroscopic Ginzburg–Landau order-parameter action to a microscopic four-spinor (NJL/BCS) pairing [20, 21] and Sakharov 1-loop induced-gravity mechanism [22] triggered by localized 4D ripple energy ($\mathcal{E}_{\text{loc}} \geq \mathcal{E}_{\text{crit}}$) [23].

# 2 The Complexified 4D Vacuum ($\mathbb{C}^4$) and the 5D Kaluza–Klein Slice

## 2.1 Emergence of Lorentzian Signature and the 5th Coordinate from $\mathbb{C}^4$

Consider an isotropic 4-dimensional complex manifold $\mathbb{C}^4$ with holomorphic coordinates [6, 7]:

$$Z^a = x^a + iy^a \in \mathbb{C}^4 \quad (a = 1, 2, 3, 4), \qquad x^a, y^a \in \mathbb{R}. \tag{1}$$

Equipped with the complex quadratic form $dS^2_{\mathbb{C}} = \delta_{ab}\, dZ^a dZ^b$, separating real and imaginary parts yields:

$$dS^2_{\mathbb{C}} = (dx_a dx^a - dy_a dy^a) + 2i\,(dx_a dy^a)\,. \tag{2}$$

Here, the relative minus sign between real and imaginary displacements arises directly from $i^2 = -1$ without being postulated a priori.

While $\mathbb{C}^4$ has 8 real dimensions, we consider a **5-real-dimensional hypersurface slice** where the first three coordinates lie on the real spatial slice ($Z^i = x^i$ for $i = 1, 2, 3$), while the 4th complex coordinate $Z^4$ retains both its real and imaginary parts:

$$Z^i = x^i \quad (i = 1, 2, 3), \qquad Z^4 = x^5 + ict. \tag{3}$$

In this parameterization:

- The **imaginary part** $\mathrm{Im}(Z^4) = ct$ provides the Lorentzian time coordinate ($-c^2dt^2$) of $3+1$D spacetime.
- The **real part** $\mathrm{Re}(Z^4) = x^5$ acts as an internal **5th Kaluza–Klein coordinate** [1, 2, 4].

## 2.2 Gauge Twisting and the 5D Kaluza–Klein Line Element

When the slice is locally fibered over curved 3+1D spacetime $x^\mu = (ct, x^1, x^2, x^3)$, a displacement $dx^\mu$ induces a connection shift along the compact internal coordinate $x^5$:

$$dz^5 = dx^5 + \kappa A_\mu(x)\, dx^\mu, \tag{4}$$

where $A_\mu(x)$ is the QED electromagnetic 4-potential and $\kappa$ is the Kaluza coupling constant. The real part of the induced metric on this 5D slice takes the standard **Kaluza–Klein–Jordan–Thiry form** [1, 2, 3]:

$$ds^2_5 = g_{\mu\nu}(x)\, dx^\mu dx^\nu + \phi^2(x)\left(dx^5 + \kappa A_\mu(x)\, dx^\mu\right)^2. \tag{5}$$

For $\phi = 1$, dimensional reduction of the 5D Einstein–Hilbert action $\int d^4x\, dx^5 \sqrt{-g_5}\, R_{5\mathrm{D}}$ yields the coupled 4D Einstein–Maxwell action [1, 2]:

$$S = \frac{1}{16\pi G}\int d^4x \sqrt{-g_4}\left(R_{4\mathrm{D}} - \frac{\kappa^2}{4} F_{\mu\nu}F^{\mu\nu}\right), \qquad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \tag{6}$$

# 3 Bloch Sphere Polar Coordinates, the Hopf Fiber, and Lorentz Boosts

## 3.1 Single-Spinor Parameterization of the Light Cone and Kaluza–Klein Fiber

To connect the Kaluza–Klein $U(1)$ fiber directly to spinor geometry, we decompose $\mathbb{C}^4$ into a pair of 2-component complex spinor spaces, $\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$ [5, 6]. Consider first a single 2-

component complex spinor $\pi_{A'} = (z_1, z_2)^T \in \mathbb{C}^2$ with norm $|z_1|^2 + |z_2|^2 = R^2$. In **Bloch-sphere polar coordinates**, $\pi_{A'}$ is written via the Hopf map [8] as:

$$\pi_{A'} = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = R\, e^{i\chi/2} \begin{pmatrix} \cos\frac{\theta}{2}\, e^{-i\varphi/2} \\ \sin\frac{\theta}{2}\, e^{i\varphi/2} \end{pmatrix}, \tag{7}$$

where:

1. $(\theta, \varphi) \in S^2$ are the polar and azimuthal angles on the **Bloch sphere** (the Celestial Sphere [6]).
2. $\chi \in [0, 4\pi)$ is the global $U(1)$ **Hopf phase fiber** [8], corresponding to the compact Kaluza–Klein coordinate $x^5 = r_0\chi$.
3. $R \geq 0$ is the real scale factor.

Constructing the $2 \times 2$ Hermitian rank-1 matrix $\mathbf{X} = \pi\pi^\dagger$:

$$\mathbf{X} = \begin{pmatrix} |z_1|^2 & z_1\bar{z}_2 \\ \bar{z}_1 z_2 & |z_2|^2 \end{pmatrix} = \frac{1}{2}\begin{pmatrix} ct + z & x - iy \\ x + iy & ct - z \end{pmatrix}, \tag{8}$$

maps the Bloch-sphere coordinates directly onto a $3+1$D spacetime 4-vector $x^\mu = (ct, x, y, z)$:

$$ct = R^2 \geq 0, \qquad x = R^2 \sin\theta\cos\varphi, \qquad y = R^2 \sin\theta\sin\varphi, \qquad z = R^2\cos\theta. \tag{9}$$

Because $\mathbf{X}$ has rank 1, its determinant vanishes identically:

$$\det(\mathbf{X}) = \frac{1}{4}\left(c^2t^2 - x^2 - y^2 - z^2\right) = |z_1|^2|z_2|^2 - |z_1\bar{z}_2|^2 \equiv 0. \tag{10}$$

Thus:

- **Every pure state on the Bloch sphere $(\theta, \varphi)$ defines a null ray on the future $3+1$D light cone** ($ds^2_{4\mathrm{D}} = 0$), with $ct = |z_1|^2 + |z_2|^2 \geq 0$ enforcing a future-directed coordinate time.
- Furthermore, the flat metric on the unit spinor sphere $S^3 \subset \mathbb{C}^2$ ($R = 1$) decomposes via the **Hopf fibration** ($S^3 \xrightarrow{U(1)} S^2$) [8] into the **Kaluza–Klein form**:

$$ds^2_{S^3} = 4\left(|dz_1|^2 + |dz_2|^2\right) = \underbrace{\left(d\theta^2 + \sin^2\theta\, d\varphi^2\right)}_{\text{Bloch Sphere } S^2} + \underbrace{\left(d\chi - \cos\theta\, d\varphi\right)^2}_{\text{Kaluza–Klein } U(1) \text{ Fiber}}, \tag{11}$$

  where $A = -\cos\theta\, d\varphi$ is the Berry / Dirac monopole $U(1)$ gauge connection on the Bloch sphere [9, 10].

### 3.2 Transformation of Bloch Sphere Angles Under Lorentz Boosts

Introducing the stereographic complex coordinate $\zeta = z_2/z_1 = \tan(\theta/2)\, e^{i\varphi}$ on the Bloch sphere, the entire restricted Lorentz group $SO^+(1,3) \cong SL(2,\mathbb{C})/\mathbb{Z}_2$ acts as conformal **Möbius transformations** [6]:

$$\zeta' = \frac{a\,\zeta + b}{c\,\zeta + d}, \qquad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2,\mathbb{C}), \qquad \det(M) = ad - bc = 1. \tag{12}$$

For a pure Lorentz boost along the $z$-axis with velocity $v = \beta c = c\tanh\eta$ (where $\cosh\eta = \gamma = (1-\beta^2)^{-1/2}$), $M = \mathrm{diag}(e^{\eta/2}, e^{-\eta/2})$, yielding:

1. **Azimuthal invariance:** $\varphi' = \varphi$.
2. **Relativistic aberration of the polar angle:**
$$\tan\left(\frac{\theta'}{2}\right) = e^{-\eta}\tan\left(\frac{\theta}{2}\right) = \sqrt{\frac{1-\beta}{1+\beta}}\,\tan\left(\frac{\theta}{2}\right) \iff \cos\theta' = \frac{\cos\theta+\beta}{1+\beta\cos\theta}. \tag{13}$$
3. **Relativistic Doppler scaling of $R^2 = ct$:**
$$R'^2 = R^2\left(e^{\eta}\cos^2\frac{\theta}{2} + e^{-\eta}\sin^2\frac{\theta}{2}\right) = R^2\gamma(1+\beta\cos\theta). \tag{14}$$
4. **Thomas Precession as Kaluza–Klein Berry Phase:** Under a general boost $M \in SL(2,\mathbb{C})$, the Hopf phase $\chi$ acquires a Wigner rotation / Berry phase shift [10, 11]:
$$\Delta\chi = 2\arg\left(c\sin\frac{\theta}{2}e^{i\varphi/2} + d\cos\frac{\theta}{2}e^{-i\varphi/2}\right). \tag{15}$$

# 4 Two-Spinor Coupling: Intra-Particle Mass vs. Inter-Particle Interactions

A crucial feature of working in $\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$—rather than a naive 5D real manifold—is that **electric charge** $q$ and **rest mass** $m$ do not compete for the same single 5th coordinate. Instead, while $U(1)$ charge lives on the overall phase $\chi$, **rest mass and proper time arise from coupling the two chiral spinors** $\xi^A \in \mathbb{C}^2_L$ and $\eta_{A'} \in \mathbb{C}^2_R$.

## 4.1 Intra-Particle Two-Spinor Coupling: Mass and Lorentz Contraction Inside the Bloch Ball

Explicitly parameterizing both chiral sectors of $\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$ by antipodal Bloch-sphere spinors (pointing along $+\hat{\mathbf{n}}(\theta,\varphi)$ and $-\hat{\mathbf{n}}(\pi-\theta,\varphi+\pi)$ respectively, with moduli $R_L = |\xi|, R_R = |\eta|$ and Hopf phases $\chi_L, \chi_R$) gives:

$$\xi^A = R_L\, e^{i\chi_L/2}\begin{pmatrix}\cos\frac{\theta}{2}\,e^{-i\varphi/2}\\ \sin\frac{\theta}{2}\,e^{i\varphi/2}\end{pmatrix}, \qquad \eta_{A'} = R_R\, e^{i\chi_R/2}\begin{pmatrix}-\sin\frac{\theta}{2}\,e^{-i\varphi/2}\\ \cos\frac{\theta}{2}\,e^{i\varphi/2}\end{pmatrix},$$

where the common phase $\chi = \frac{1}{2}(\chi_L + \chi_R)$ is the Kaluza–Klein $U(1)$ gauge fiber and the relative phase $\Delta\chi = \chi_L - \chi_R$ drives the internal Compton (*Zitterbewegung*) oscillation [12, 13]. In the Weyl representation, the Dirac equation for a massive spin-1/2 particle is a coupled system linking the left-handed Bloch spinor $\xi^A$ and the right-handed Bloch spinor $\eta_{A'}$ across the quantum vacuum [12, 6]:

$$i\hbar\,\sigma^{\mu}_{AA'}D_\mu\xi^A = mc\,\eta_{A'}, \qquad i\hbar\,\bar{\sigma}^{\mu A'A}D_\mu\eta_{A'} = mc\,\xi^A. \tag{16}$$

- **Massless Regime ($m = 0$, Surface of the Bloch Sphere):** The two spinors decouple. Each defines a null momentum $p^\mu_L = \xi^\dagger\sigma^\mu\xi$ ($p^2_L = 0$) or $p^\mu_R = \eta^\dagger\bar{\sigma}^\mu\eta$ ($p^2_R = 0$) lying strictly on the **surface of the Bloch sphere** ($r_{\text{Bloch}} = 1$, velocity $v = c$).
- **Massive Regime ($m > 0$, Interior of the Bloch Ball):** The mass $mc$ couples $\xi^A \leftrightarrow \eta_{A'}$ (*Zitterbewegung* / Penrose's *Zig-Zag* mechanism [13], where $\bar{\eta}^A \equiv \overline{\eta^{A'}}$ denotes the complex-conjugate unprimed spinor so that $|\xi_A\bar{\eta}^A| = R_L R_R = |\xi||\eta|$). The total 4-momentum $P^\mu = p^\mu_L + p^\mu_R$ is the sum of two non-collinear null vectors, which lies strictly **inside the future light cone** and **inside the Bloch ball**:
$$P_\mu P^\mu = (p_L + p_R)^2 = 2\,p_L\cdot p_R = 2\left|\xi_A\bar{\eta}^A\right|^2 = m^2c^2 > 0. \tag{17}$$

In the rest frame ($v = 0$), the two spinors have equal moduli ($|\xi|^2 = |\eta|^2$) and opposite spatial Bloch vectors ($\mathbf{n}_L = -\mathbf{n}_R$), placing the net state at the **center of the Bloch ball** ($r_{\text{Bloch}} = 0$). Under a boost of rapidity $\eta_{\text{boost}}$ ($\tanh \eta_{\text{boost}} = v/c$), the chiral spinor moduli scale oppositely as $|\xi| \to e^{+\eta_{\text{boost}}/2}|\xi_0|$ and $|\eta| \to e^{-\eta_{\text{boost}}/2}|\eta_0|$. Consequently:

1. **The effective Bloch-ball radius equals the normalized velocity:**

$$r_{\text{Bloch}} = \frac{|\xi|^2 - |\eta|^2}{|\xi|^2 + |\eta|^2} = \frac{e^{\eta_{\text{boost}}} - e^{-\eta_{\text{boost}}}}{e^{\eta_{\text{boost}}} + e^{-\eta_{\text{boost}}}} = \tanh \eta_{\text{boost}} = \frac{v}{c} \equiv \sin \alpha. \tag{18}$$

2. **The two-spinor coherence overlap equals the Lorentz contraction / proper-time factor $1/\gamma$:**

$$\cos \alpha = \sqrt{1 - r_{\text{Bloch}}^2} = \frac{2\,|\xi|\,|\eta|}{|\xi|^2 + |\eta|^2} = \frac{1}{\cosh \eta_{\text{boost}}} = \sqrt{1 - \frac{v^2}{c^2}} = \frac{1}{\gamma} = \frac{d\tau}{dt}. \tag{19}$$

This provides a direct quantum-information interpretation of relativistic kinematics: **proper time $d\tau/dt$ and Lorentz contraction $1/\gamma$ measure the chiral coherence balance $\cos\alpha$ inside the Bloch ball.** As $v \to c$, one spinor dominates ($r_{\text{Bloch}} \to 1, \cos\alpha \to 0$), pushing the state back onto the pure-state surface of the Bloch sphere (the light cone) where proper time halts ($d\tau \to 0$).

### 4.2 Inter-Particle Two-Spinor Coupling: Quantum Scattering Across the Vacuum

When two distinct particles 1 and 2 with spinors $(\xi_1^A, \eta_{1A'})$ and $(\xi_2^A, \eta_{2A'})$ interact across the complexified vacuum, their Lorentz-invariant couplings are formed from the antisymmetric spinor-helicity brackets [6, 14]:

$$\begin{aligned} \langle 1\,2 \rangle &= \varepsilon_{AB}\,\xi_1^A \xi_2^B = \sqrt{E_1 E_2}\,\sin\left(\frac{\theta_{12}}{2}\right) e^{i\Phi_{12}}, \\ [1\,2] &= \varepsilon^{A'B'}\,\eta_{1A'}\eta_{2B'} = \sqrt{E_1 E_2}\,\sin\left(\frac{\theta_{12}}{2}\right) e^{-i\Phi_{12}}, \end{aligned} \tag{20}$$

where $\theta_{12}$ is the chordal angle between the particles on the Bloch sphere and $\Phi_{12}$ is their relative Kaluza–Klein / Hopf phase. Every invariant propagator denominator in QED and gravity factors into a product of left and right spinor brackets [14]:

$$s_{12} = (p_1 + p_2)^2 = \langle 1\,2 \rangle [2\,1]. \tag{21}$$

## 5 Outside the Light Cone: Virtual Particles, Bell Entanglement, and Relational Observers

### 5.1 The Arrow of Time and the Complex Extended Tube Outside the Light Cone

In axiomatic quantum field theory and Twistor geometry [5, 15], vacuum Wightman functions $W(z^\mu) = \langle 0|\phi(x)\phi(0)|0\rangle$ are holomorphic in the **forward tube** $\mathcal{T}^+ = \{z^\mu = x^\mu - iy^\mu \in \mathbb{C}^4 \mid y^0 > |\mathbf{y}|\}$ (corresponding to projective twistor half-space $\mathbb{PT}^+$ with $Z^\alpha \bar{Z}_\alpha > 0$). Requiring $\Psi(z) \sim e^{-ip_\mu z^\mu/\hbar}$ to remain bounded in $\mathcal{T}^+$ enforces positive energy ($E > 0$) and a strict arrow of time **inside and on the real light cone** ($c^2t^2 - \mathbf{x}^2 \geq 0$), where $\mathcal{T}^+$ and $\mathcal{T}^-$ are disjoint.

However, **outside the light cone** (spacelike separation, $\mathbf{x}^2 - c^2t^2 > 0$):

1. **Absence of an Invariant Arrow of Time:** A continuous Lorentz boost (a Möbius transformation on the Bloch sphere) can reverse the sign of $\Delta t$ for any spacelike-separated pair of events.
2. **Merger of Future and Past Tubes (Bargmann–Hall–Wightman Theorem [15]):** Under complex Lorentz transformations $SL(2,\mathbb{C}) \times SL(2,\mathbb{C})$, the forward tube $\mathcal{T}^+$ and backward tube $\mathcal{T}^-$ **analytically continue into one another across spacelike Jost points**, forming a single connected domain (the **Extended Tube**).
3. **Virtual Particles in Feynman Diagrams [16]:** Because the boundary between forward-in-time and backward-in-time disappears outside the light cone, the Feynman propagator $D_F(x-y) = \theta(t_x - t_y)W^+(x-y) + \theta(t_y - t_x)W^-(x-y)$ is **non-vanishing outside the light cone**:

$$D_F(x-y) = \frac{m}{4\pi^2\sqrt{r^2 - c^2t^2}}\, K_1\!\left(\frac{mc}{\hbar}\sqrt{r^2 - c^2t^2}\right) \neq 0 \qquad (r > c|t|). \tag{22}$$

Because forward and backward virtual paths have equal amplitude at spacelike separation ($W^+(x-y) = W^-(x-y)$), the **symmetric sum** $D_F \sim W^+ + W^- \neq 0$ mediates virtual vacuum forces (e.g., Coulomb, Casimir, and vacuum polarization), while the **antisymmetric commutator** vanishes identically:

$$[\phi(x), \phi(y)] = W^+(x-y) - W^-(x-y) = 0 \qquad (\text{for } (x-y)^2 > 0), \tag{23}$$

preserving macroscopic signal causality [15].

## 5.2 Bell Singlet Entanglement as an Outside-the-Light-Cone Two-Spinor Bracket

When two particles' spinors $\xi_1^A(x_A)$ and $\xi_2^B(x_B)$ are coupled into the antisymmetric $SL(2,\mathbb{C})$ bracket across a spacelike interval [6, 17]:

$$|\Psi^-\rangle = \frac{1}{\sqrt{2}}\,\varepsilon_{AB}\,\xi_1^A\xi_2^B = \frac{1}{\sqrt{2}}\left(\xi_1^0\xi_2^1 - \xi_1^1\xi_2^0\right) \equiv \frac{1}{\sqrt{2}}\left(|\uparrow\rangle_A|\downarrow\rangle_B - |\downarrow\rangle_A|\uparrow\rangle_B\right), \tag{24}$$

three geometric properties follow immediately:

1. **Exact Parallel to Intra-Particle Mass (Centering in the Bloch Ball):** Tracing out Particle $B$ reduces Particle $A$'s local density matrix on the $3+1$D slice to $\rho_A = \frac{1}{2}I_2$, placing Particle $A$'s local Bloch vector at the **exact center of its Bloch ball** ($r_{\text{Bloch},A} = 0$). Locally, Alice observes unpolarized 50%/50% outcomes along any axis, preventing superluminal signaling, while the 100% purity of the joint state resides in the shared two-spinor bracket in $\mathbb{C}^4$.
2. **Resolution of the Retrocausal Collapse Paradox:** Because $x_A$ and $x_B$ are separated outside the light cone (in the Extended Tube where time-ordering is undefined [15]), there is no physical distinction between "Alice collapses Bob forward in time" and "Bob collapses Alice backward in time." The correlation is a time-symmetric boundary intersection of the holomorphic invariant $\varepsilon_{AB}\xi_1^A\xi_2^B$ with the $3+1$D slice.
3. **Bell Correlation from Bloch-Sphere Chordal Geometry:** Projecting $|\Psi^-\rangle$ onto the measurement spinors $\alpha^A(\theta_A, \varphi_A)$ and $\beta^B(\theta_B, \varphi_B)$ yields the joint probability $P(++) = \frac{1}{2}|\varepsilon_{AB}\alpha^A\beta^B|^2 = \frac{1}{2}\sin^2(\theta_{AB}/2)$ and the expectation value:

$$E(\mathbf{n}_A, \mathbf{n}_B) = \sin^2\left(\frac{\theta_{AB}}{2}\right) - \cos^2\left(\frac{\theta_{AB}}{2}\right) = -\cos\theta_{AB} = -\mathbf{n}_A \cdot \mathbf{n}_B, \tag{25}$$

violating Bell's inequality [17] up to the Tsirelson bound $2\sqrt{2}$ [18] due to the preserved Hopf phase interference in $\mathbb{C}^4$.

### 5.3 Relational Causality: Why Distant Separated Observers Cancel Out

Consider two interacting particles with spinors $\xi_1^A, \xi_2^B$ and any distant observer $O'$ moving at an arbitrary velocity relative to another distant observer $O$. Observer $O'$'s coordinate frame is related to $O$'s by a transformation $M \in SL(2,\mathbb{C})$ with $\det(M) = 1$. Under $M$:

$$\langle M\xi_1\, M\xi_2\rangle = \varepsilon_{AB}(M^A{}_C\xi_1^C)(M^B{}_D\xi_2^D) = \det(M)\,\varepsilon_{CD}\xi_1^C\xi_2^D = \langle \xi_1\,\xi_2\rangle. \tag{26}$$

Because $\det(M) = 1$, **any distant observer's boost matrix cancels identically out of the interaction bracket**. While separated observers $O$ and $O'$ may assign different coordinate times $t$ and $t'$ by extending spacelike hyperplanes outside their own light cones, those coordinate hyperplanes have zero physical input into the interaction; only the interacting particles and the $\mathbb{C}^4$ quantum vacuum participate.

## 6 Lagrangian Formulation of the $\mathbb{C}^4 \to 3+1$D Transformation

To elevate this geometric picture from a purely kinematic dictionary to a variational principle, we construct both the **two-spinor worldline Lagrangian** (governing single- and two-particle relativistic kinematics, proper time, and Kaluza–Klein charge) and the **field-theoretic symmetry-breaking Lagrangian density** (governing the transition from the scale-free $\mathbb{C}^4$ vacuum to the $3+1$D slice). For pedagogical clarity, Subsections 6.1–6.3 first present the macroscopic Ginzburg–Landau effective description characterized by a single effective condensate scale $v_0$ inside a condensed $3+1$D patch, before Subsections 6.4–6.5 resolve $v_0$ into the two-tier Planck ($M_{\rm Pl}$) and Electroweak Higgs ($v_{\rm EW}$) hierarchy and derive the effective potentials microscopically from four-spinor pairing, Sakharov 1-loop induced gravity, and localized 4D ripple energy.

### 6.1 The Two-Spinor Worldline Lagrangian ($\mathcal{L}_{\rm worldline}$)

Consider a state in $\mathbb{C}^4 = \mathbb{C}^2_L \oplus \mathbb{C}^2_R$ parameterized along an evolution parameter $\lambda$ by the pair of 2-component Bloch spinors $(\xi^A(\lambda), \eta_{A'}(\lambda))$, spacetime coordinates $x^{AA'}(\lambda) = \frac{1}{\sqrt{2}}\sigma_\mu^{AA'}x^\mu(\lambda)$, and the internal Kaluza–Klein / Hopf phase $\chi(\lambda) \in [0, 4\pi)$. We write the first-order phase-space Lagrangian:

$$\begin{aligned}\mathcal{L}_{\rm worldline} = P_{AA'}\dot{x}^{AA'} + \frac{\hbar}{2}\,J\left(\dot{\chi} + \frac{2e}{\hbar c}A_{AA'}(x)\,\dot{x}^{AA'}\right)\\ - \alpha_{\rm E}\left[P_{AA'} - \left(\xi_A\bar{\xi}_{A'} + \bar{\eta}_A\eta_{A'}\right)\right] - \mathcal{H}_{\rm vac}(\xi,\eta,\Phi_0),\end{aligned} \tag{27}$$

where $\alpha_{\rm E}$ is a Lagrange multiplier enforcing the two-spinor decomposition of 4-momentum, $J$ is the conjugate $U(1)$ fiber angular momentum, and $\mathcal{H}_{\rm vac}$ is the chiral vacuum coupling potential generated by a background vacuum condensate $\Phi_0$:

$$\mathcal{H}_{\rm vac}(\xi,\eta,\Phi_0) = \frac{\lambda_{\rm M}}{2}\left(\sqrt{2}\,|\xi_A\bar{\eta}^A| - mc\right)^2, \qquad mc \equiv g_{\rm Y}|\Phi_0|. \tag{28}$$

Varying $\mathcal{L}_{\rm worldline}$ with respect to each degree of freedom yields the complete set of transformations:

1. **Conservation of Kaluza–Klein $U(1)$ Charge ($\delta\chi$):** Since $\mathcal{L}_{\rm worldline}$ depends on $\chi$ only through $\dot{\chi}$, the Euler–Lagrange equation $\frac{d}{d\lambda}\left(\frac{\partial\mathcal{L}}{\partial\dot{\chi}}\right) = 0$ fixes the canonical 5th momentum $p_\chi = \frac{\hbar}{2}J = \text{const}$, which couples minimally to the QED potential $A_\mu\dot{x}^\mu$ with electric charge $q = eJ$ *independently* of the rest mass $m$.

2. **Emergence of the Light Cone vs. Timelike Interior ($\delta\alpha_{\mathbf{E}}$ and $\delta P_{AA'}$):** Varying $\alpha_{\mathrm{E}}$ and $P_{AA'}$ gives the four-velocity and four-momentum as the sum of two null Bloch-sphere vectors:
$$P^{AA'} = \xi^A\bar{\xi}^{A'} + \bar{\eta}^A\eta^{A'}, \qquad \dot{x}^{AA'} = \alpha_{\mathrm{E}}\left(\xi^A\bar{\xi}^{A'} + \bar{\eta}^A\eta^{A'}\right). \tag{29}$$
   - *When the vacuum condensate vanishes ($\Phi_0 = 0 \implies m = 0$, or $\eta_{A'} = 0$):* Only a single spinor is active, forcing $\det(\dot{x}^{AA'}) = 0 \implies ds^2_{4\mathrm{D}} = 0$. The trajectory is locked onto the **surface of the Bloch sphere (the $3+1$D light cone)**.
   - *When the vacuum condensate is non-zero ($\Phi_0 \neq 0 \implies m > 0$):* Minimizing $\mathcal{H}_{\mathrm{vac}}$ locks the chiral spinor inner product to $\sqrt{2}\,|\xi_A\bar{\eta}^A| = mc$. Taking the determinant of $\dot{x}^{AA'}$ yields the invariant proper-time line element inside the light cone:
$$c^2d\tau^2 \equiv -ds^2_{4\mathrm{D}} = 2\det(dx^{AA'}) = 2\,\alpha_{\mathrm{E}}^2\,|\xi_A\bar{\eta}^A|^2 d\lambda^2 = \alpha_{\mathrm{E}}^2 m^2c^2\,d\lambda^2 > 0. \tag{30}$$

3. **Variational Derivation of Lorentz Contraction ($d\tau/dt = 1/\gamma$):** From the trace $\mathrm{Tr}(\dot{x}^{AA'}) = \sqrt{2}\,c\dot{t} = \alpha_{\mathrm{E}}(|\xi|^2 + |\eta|^2)$, eliminating the multiplier $\alpha_{\mathrm{E}}d\lambda$ between $c\,dt$ and $c\,d\tau$ gives:
$$\frac{d\tau}{dt} = \frac{2\,|\xi_A\bar{\eta}^A|}{|\xi|^2 + |\eta|^2} = \frac{2\,|\xi|\,|\eta|}{|\xi|^2 + |\eta|^2} = \sqrt{1 - r^2_{\mathrm{Bloch}}} = \sqrt{1 - \frac{v^2}{c^2}} = \frac{1}{\gamma}. \tag{31}$$
   Thus, the Lorentz contraction factor $1/\gamma$ emerges directly as the ratio of the chiral determinant to the trace of the two-spinor velocity matrix.

## 6.2 Field-Theoretic Vacuum Condensate Lagrangian ($\mathcal{L}_{\mathbf{field}}$)

At the field-theoretic level, the transition from the holomorphic $\mathbb{C}^4$ vacuum to the $3+1$D slice with Kaluza–Klein gauge fields and massive Dirac spinors is governed by the action $S = \int d^4x\sqrt{-g_4}\,\mathcal{L}_{\mathrm{field}}$:

$$\begin{aligned}\mathcal{L}_{\mathrm{field}} = &\underbrace{\frac{M_{\mathrm{Pl}}^2}{2}R_{4\mathrm{D}} - \frac{1}{2}g^{\mu\nu}(\nabla_\mu\rho)(\nabla_\nu\rho) - \frac{\rho^2}{2}g^{\mu\nu}(\partial_\mu\chi + \kappa A_\mu)(\partial_\nu\chi + \kappa A_\nu) - \frac{\rho^2\kappa^2}{4}F_{\mu\nu}F^{\mu\nu}}_{\text{Induced 5D Kaluza–Klein Gravity + QED + Phase Fiber } \Phi=\rho e^{i\chi}} \\ &+ \underbrace{i\hbar c\left(\bar{\xi}_{A'}\sigma^{\mu AA'}D_\mu\xi_A + \eta^A\bar{\sigma}^\mu_{AA'}D_\mu\bar{\eta}^{A'}\right) - g_{\mathrm{Y}}\rho\left(\xi^A\eta_A + \bar{\xi}_{A'}\bar{\eta}^{A'}\right)}_{\text{Two-Spinor Kinetic + Chiral Mass-Generating Coupling}} \\ &- \underbrace{\frac{\lambda_\Phi}{4}\left(\rho^2 - v_0^2\right)^2}_{\text{Slice-Selecting Vacuum Potential}} + \underbrace{\left(\Lambda^{\bar{a}}{}_B\,\bar{\partial}_{\bar{a}}\xi^B + \text{h.c.}\right)}_{\text{Holomorphy Constraint in } \mathbb{C}^4}.\end{aligned} \tag{32}$$

This Lagrangian unifies the four structural requirements of the model:

1. **Holomorphic Ghost Elimination ($\delta\Lambda^{\bar{a}}{}_B$):** Varying the Lagrange multiplier $\Lambda^{\bar{a}}{}_B$ enforces the Cauchy–Riemann holomorphy condition $\bar{\partial}_{\bar{a}}\xi^B = 0$ in $\mathbb{C}^4$, freezing the three unphysical imaginary time directions so that only a single causal time coordinate $ct$ propagates on the real boundary slice.

2. **Spontaneous Conformal Breaking & Mass Generation ($\delta\rho$):** Above the symmetry-breaking scale ($\langle\rho\rangle = 0$), the vacuum is conformally invariant, $m = g_{\mathrm{Y}}\langle\rho\rangle = 0$, and left/right spinors propagate independently on the light cone ($ds^2 = 0$). When the scalar modulus condenses to its vacuum expectation value $\langle\rho\rangle = v_0$, it simultaneously:
   - stabilizes the Kaluza–Klein 5th-dimensional radius ($g_{55} = \phi^2 \propto v_0^2$),

- canonically normalizes the QED Maxwell term $-\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$, and
- generates the two-spinor Dirac mass $mc^2 = g_Y v_0$, pulling states from the Bloch-sphere boundary into the timelike interior of the light cone.

3. **Feynman Propagator & Outside-the-Light-Cone Correlations ($\delta\xi^A, \delta\eta_{A'}$):** Inverting the quadratic two-spinor kinetic + mass operator in $\mathcal{L}_{\text{field}}$ with positive-frequency boundary conditions in the forward tube $\mathcal{T}^+$ yields the exact two-spinor Feynman propagator:

$$\langle 0|T\Big\{\Psi(x)\bar{\Psi}(y)\Big\}|0\rangle = (i\hbar\gamma^\mu\partial_\mu + mc)\, D_F(x-y), \tag{33}$$

which is strictly causal inside the light cone while remaining non-zero and time-symmetric ($W^+ = W^-$) outside the light cone.

### 6.3 The Slice-Selection Condensate Lagrangian in $\mathbb{C}^4$ ($\mathcal{L}_{\text{slice}}$)

Finally, we derive the explicit Lagrangian for the **slice-selection condensate** itself—answering how an initially isotropic, scale-free $\mathbb{C}^4$ vacuum dynamically selects a real $(3,1)$ Lorentzian light-cone slice, a Kaluza–Klein $U(1)$ phase fiber, and a two-spinor mass scale via spontaneous symmetry breaking [19, 6].

In the unbroken $\mathbb{C}^4$ vacuum (with coordinates $Z^\alpha = (\omega^A, \pi_{A'}) \in \mathbb{C}^4$, $\alpha = 0,1,2,3$), the ambient symmetry group is $SL(4,\mathbb{C})$. At this stage there is neither a preferred real slice nor a metric scale. We introduce two coupled vacuum order-parameter fields on $\mathbb{C}^4$:

1. **A Hermitian Sesquilinear Condensate $H_{\alpha\bar{\beta}}(Z,\bar{Z}) = \overline{H_{\beta\bar{\alpha}}}$:** An adjoint matrix field whose signature determines the real slice and causal arrow of time [5].
2. **An Antisymmetric Bi-Twistor Condensate $\Sigma_{\alpha\beta}(Z) = -\Sigma_{\beta\alpha} \in \Lambda^2(\mathbb{C}^4)$:** A complex 6-component field (the dynamical precursor to Penrose's Infinity Twistor $I_{\alpha\beta}$ [6]) whose modulus $\rho$ sets the mass/gravitational scale and whose phase $\chi$ is the 5th Kaluza–Klein coordinate.

The $SL(4,\mathbb{C})$-covariant **Slice-Selection Condensate Lagrangian** is:

$$\begin{aligned}\mathcal{L}_{\text{slice}}(\Sigma, H, \Psi) = &-\frac{1}{4}H^{\alpha\bar{\gamma}}H^{\beta\bar{\delta}}\,(\mathcal{D}_\mu\Sigma_{\alpha\beta})\left(\overline{\mathcal{D}^\mu\Sigma_{\gamma\delta}}\right) - \frac{1}{2}\,\mathrm{Tr}(\mathcal{D}_\mu H\,\mathcal{D}^\mu H)\\ &+ i\hbar c\,\bar{\Psi}_{\bar{\alpha}}H^{\bar{\alpha}\beta}\mathcal{D}\Psi_\beta - \frac{g_Y}{2}\left(\Psi^\alpha\Sigma_{\alpha\beta}\Psi^\beta + \bar{\Psi}_{\bar{\alpha}}\bar{\Sigma}^{\bar{\alpha}\bar{\beta}}\bar{\Psi}_{\bar{\beta}}\right) - V_{\text{slice}}(\Sigma, H),\end{aligned} \tag{34}$$

with the symmetry-breaking potential:

$$\begin{aligned}V_{\text{slice}}(\Sigma, H) = &\frac{\lambda_H}{4}\,\mathrm{Tr}\Big(H^2 - \mathbb{I}_4\Big)^2 + \frac{\mu_H^2}{2}\,(\mathrm{Tr}\,H)^2\\ &+ \frac{\lambda_\Sigma}{4}\left(\frac{1}{2}H^{\alpha\bar{\gamma}}H^{\beta\bar{\delta}}\Sigma_{\alpha\beta}\bar{\Sigma}_{\bar{\gamma}\bar{\delta}} - v_0^2\right)^2 + \frac{\lambda_I}{4}\left|\frac{1}{4}\varepsilon^{\alpha\beta\gamma\delta}\Sigma_{\alpha\beta}\Sigma_{\gamma\delta} - \frac{\Lambda_{\text{cc}}}{6}v_0^2\right|^2.\end{aligned} \tag{35}$$

Minimizing $V_{\text{slice}}(\Sigma, H)$ triggers a two-stage spontaneous symmetry breaking chain:

$$SL(4,\mathbb{C}) \xrightarrow{\langle H_{\alpha\bar{\beta}}\rangle} SU(2,2) \simeq SO(2,4) \xrightarrow{\langle\Sigma_{\alpha\beta}\rangle = v_0 e^{i\chi} I_{\alpha\beta}} ISO^+(1,3)\times U(1)_{\text{KK}}, \tag{36}$$

which dynamically produces the entire geometry of the toy model:

1. **Stage 1 — Selection of the $3+1$D Light-Cone Slice & Arrow of Time ($\delta H_{\alpha\bar{\beta}} = 0$):** The potential $\frac{\lambda_H}{4}\,\mathrm{Tr}(H^2 - \mathbb{I}_4)^2 + \frac{\mu_H^2}{2}(\mathrm{Tr}\,H)^2$ forces the eigenvalues of $\langle H_{\alpha\bar{\beta}}\rangle$ to be $(\pm1, \pm1, \pm1, \pm1)$ with traceless condition $\mathrm{Tr}\langle H\rangle = 0$, uniquely selecting signature $(+,+,-,-)$ [5]:

$$\langle H_{\alpha\bar{\beta}}\rangle = \begin{pmatrix} 0 & \mathbb{I}_2 \\ \mathbb{I}_2 & 0 \end{pmatrix} \implies Z^\alpha \langle H_{\alpha\bar{\beta}}\rangle \bar{Z}^{\bar{\beta}} = \omega^A \bar{\pi}_A + \bar{\omega}^{A'} \pi_{A'}. \tag{37}$$

   This breaks $SL(4,\mathbb{C}) \to SU(2,2)$ (the 4D conformal group) and partitions $\mathbb{C}^4$ into the **real $3+1$D light-cone slice** ($Z^\alpha \langle H_{\alpha\bar{\beta}}\rangle \bar{Z}^{\bar{\beta}} = 0$), the **future tube** $\mathbb{PT}^+$ ($> 0$, arrow of time), and the **past tube** $\mathbb{PT}^-$ ($< 0$).

2. **Stage 2 — Breaking Conformal Scale into the Metric, Mass, and Kaluza–Klein Fiber ($\delta\Sigma_{\alpha\beta} = 0$):** Minimizing the $\lambda_\Sigma$ and $\lambda_I$ terms gives the bi-twistor condensate a non-zero vacuum expectation value proportional to Penrose's **Infinity Twistor** $I_{\alpha\beta}$ [6] dressed by the radial modulus $\rho = v_0$ and Kaluza–Klein phase $\chi$:

$$\langle \Sigma_{\alpha\beta}(x)\rangle = v_0\, e^{i\chi(x)}\, I_{\alpha\beta} = v_0\, e^{i\chi(x)} \begin{pmatrix} \frac{\Lambda_{\mathrm{cc}}}{6}\,\varepsilon_{AB} & 0 \\ 0 & \varepsilon^{A'B'} \end{pmatrix}. \tag{38}$$

   - *Emergence of the $3+1$D Metric & Lorentz Contraction:* Contracting adjacent null twistors $Z_1^\alpha, Z_2^\beta$ with $\langle \Sigma_{\alpha\beta}\rangle$ induces the $3+1$D spacetime metric directly from $\mathbb{C}^4$ [6]:

$$ds^2_{4\mathrm{D}} = -v_0^2\, \frac{\varepsilon_{\alpha\beta\gamma\delta} Z_1^\alpha Z_2^\beta\, dZ_1^\gamma dZ_2^\delta}{\left|\langle \Sigma_{\alpha\beta}\rangle Z_1^\alpha Z_2^\beta\right|^2} = -c^2dt^2 + dx^2 + dy^2 + dz^2. \tag{39}$$

   - *Emergence of the 5D Kaluza–Klein $U(1)$ Fiber:* Fluctuations of the condensate phase $\chi(x)$ in the kinetic term $-\frac{1}{4}\langle H\rangle\langle H\rangle|\mathcal{D}_\mu \Sigma|^2$ produce the exact Kaluza–Klein fiber term $-\frac{v_0^2}{2}(\partial_\mu \chi + \kappa A_\mu)^2$.
   - *Emergence of the Two-Spinor Dirac Mass:* Substituting $\langle \Sigma_{\alpha\beta}\rangle$ into the Yukawa term $-\frac{g_{\mathrm{Y}}}{2}\Psi^\alpha \Sigma_{\alpha\beta}\Psi^\beta$ couples the two chiral spinor components $\Psi^\alpha = (\xi^A, \eta_{A'})$ with rest mass $mc^2 = g_{\mathrm{Y}} v_0$, pulling states from the Bloch-sphere boundary ($r_{\mathrm{Bloch}} = 1$) into the Bloch ball interior ($r_{\mathrm{Bloch}} = v/c < 1$).

## 6.4 Division of Labor: The Gravitational/Time Slice Condensate vs. the Higgs Field

A key physical insight of this framework is that the condensation of $\mathbb{C}^4$ into our observed $3+1$D universe requires **both** a gravitational/causal slice condensate and the Standard Model Higgs field ($\Phi_{\mathrm{Higgs}}$) [19], operating in a two-tier hierarchy:

$$\begin{aligned} \mathbb{C}^4 \text{ Vacuum } &\xrightarrow[\substack{\text{1. Gravity / Time Condensate } (\langle H_{\alpha\bar{\beta}}\rangle, \langle \Sigma_{\alpha\beta}\rangle = M_{\mathrm{Pl}} I_{\alpha\beta})}]{M_{\mathrm{Pl}} \sim 10^{19}\ \mathrm{GeV}} 3+1\text{D Light Cone Surface } (ds^2 = 0) \\ &\xrightarrow[\text{2. Higgs Condensate } \langle \Phi_{\mathrm{Higgs}}\rangle]{v_{\mathrm{EW}} \approx 246\ \mathrm{GeV}} 3+1\text{D Massive Interior } (d\tau = dt/\gamma). \end{aligned} \tag{40}$$

1. **Role of Gravity and Time ($\langle H_{\alpha\bar{\beta}}\rangle$ and $\langle \Sigma_{\alpha\beta}\rangle = M_{\mathbf{Pl}} I_{\alpha\beta}$):** The Standard Model Higgs field is an internal electroweak doublet carrying no spacetime spinor indices $(A, A')$; it cannot by itself carve a Lorentzian light cone out of $\mathbb{C}^4$. That stage is established by the gravitational/twistor condensates $\langle H_{\alpha\bar{\beta}}\rangle$ and $\langle \Sigma_{\alpha\beta}\rangle$, which select the $(3,1)$ real slice, establish the causal arrow of time ($\mathbb{PT}^+$), and define the null light-cone boundary ($ds^2_{4\mathrm{D}} = 0$, the surface of the Bloch sphere $r_{\mathrm{Bloch}} = 1$).

2. **Role of the Higgs Field (Dynamically Tethering Spinors Inside the Light Cone):** Before electroweak symmetry breaking ($\langle\Phi_{\text{Higgs}}\rangle = 0$), all fundamental fermions are uncoupled single spinors moving independently at $v = c$ on the light-cone surface ($d\tau = 0$, no rest frames). When the Higgs condenses ($\langle\Phi_{\text{Higgs}}\rangle = v_{\text{EW}}$) [19], its Yukawa coupling $-y_f\langle\Phi_{\text{Higgs}}\rangle(\xi^A\eta_A + \bar{\xi}_{A'}\bar{\eta}^{A'})$ acts as a **chiral tether** that continuously flips $\xi^A \leftrightarrow \eta_{A'}$ at the Compton frequency $\omega_Z = 2mc^2/\hbar$ [13]. Because each chiral spinor is constrained to the light cone ($p_L^2 = 0, p_R^2 = 0$), tethering the forward- and backward-directed spinors together confines the composite particle trajectory to the **causal interior of the light cone** ($P^\mu = p_L^\mu + p_R^\mu$, $P^2 = m^2c^2 > 0$, $v < c$) and turns on operational proper time $d\tau = dt/\gamma > 0$.

3. **The Non-Zero Variation Requirement ($\Delta\Psi_{\text{grav}} \neq 0$) for Curvature and Quantized Gravity:** Comparing optical refraction (which bends rays at a condensed-matter interface where $\Delta n \neq 0$), the uniform Higgs condensate ($\nabla_\mu\langle\Phi_{\text{Higgs}}\rangle = 0$, which rotates chiral phase and keeps $v < c$ without bending trajectories), and gravitational curvature reveals a fundamental requirement: **gravity and its quantization require a non-vanishing variation $\Delta\Psi_{\text{grav}} \neq 0$ of the slice-selection condensate wavefunction** [4, 7]:

$$K_{\mu\nu} = \frac{1}{2}\nabla_\perp g_{\mu\nu}(\Psi_{\text{grav}}) \neq 0, \qquad \frac{d\hat{\mathbf{k}}}{ds} = \nabla_\perp \ln \rho_{\text{grav}}(x) \neq 0, \qquad \Delta\hat{\Psi}_{\text{grav}} \cdot \Delta\hat{\Pi}_{\text{grav}} \geq \frac{\hbar}{2} > 0. \tag{41}$$

Specifically: (i) a non-zero normal derivative across the $\mathbb{C}^4 \to 3{+}1$D condensation interface generates extrinsic curvature $K_{\mu\nu} \neq 0$ and the 4D Einstein tensor $G_{\mu\nu} \propto K_{\mu\alpha}K^\alpha{}_\nu - KK_{\mu\nu}$; (ii) a continuous tangential gradient $\nabla_\mu\rho_{\text{grav}} \neq 0$ (or non-zero twistor patching transition $\Delta Z^\alpha \neq 0$ [6, 7]) curves light rays and timelike geodesics continuously; and (iii) canonical quantization forbids $\Delta\hat{\Psi}_{\text{grav}} = 0$, meaning gravitons are quantized interface ripples of the $3+1$D slice inside $\mathbb{C}^4$.

## 6.5 From Ginzburg–Landau Requirements to a Microscopic Two-Spinor Pairing & Sakharov Induced-Gravity Mechanism

It is essential to distinguish between *phenomenological requirements* (the Ginzburg–Landau level) and a *microscopic mechanism* (the Bardeen–Cooper–Schrieffer [BCS] / Nambu–Jona-Lasinio [NJL] [20] / Sakharov [22] level). In Eqs. (34)–(35), the condensates $H_{\alpha\bar{\beta}}$, $\Sigma_{\alpha\beta}$, and $\Phi_{\text{Higgs}}$ were introduced as Ginzburg–Landau order parameters whose Mexican-hat potentials were inserted by hand. However, because all three order parameters carry the exact quantum numbers of **composite two-spinor pairs** ($H_{\alpha\bar{\beta}} \sim \langle\Psi_\alpha\bar{\Psi}_{\bar{\beta}}\rangle$, $\Sigma_{\alpha\beta} \sim \langle\Psi_{[\alpha}\tilde{\Psi}_{\beta]}\rangle$, and $\Phi_{\text{Higgs}} \sim \langle\xi^A\eta_A\rangle$), both the symmetry-breaking potentials and the 4D Einstein–Hilbert action can be derived microscopically in two steps:

1. **Microscopic Four-Spinor Pairing in $\mathbb{C}^4$ (NJL / BCS Origin of the Mexican-Hat Potential) [20, 21]:** We begin with a purely spinorial microscopic Lagrangian in $\mathbb{C}^4$ containing **no fundamental scalar Mexican-hat potentials by hand**, but only massless twistor-spinors $\Psi_\alpha = (\xi_A, \eta^{A'})$ subject to local four-spinor contact interactions:

$$\mathcal{L}_{\text{micro}}(\Psi) = i\hbar c\,\bar{\Psi}_{\bar{\alpha}}\mathcal{D}^{\bar{\alpha}\beta}\Psi_\beta + \frac{G_H}{2}\left(\bar{\Psi}_{\bar{\alpha}}\Psi_\beta\right)\left(\bar{\Psi}_{\bar{\beta}}\Psi_\alpha\right) + \frac{G_\Sigma}{4}\left|\Psi_{[\alpha}\tilde{\Psi}_{\beta]}\right|^2 + G_{\text{EW}}\left(\xi^A\eta_A\right)\left(\bar{\xi}_{A'}\bar{\eta}^{A'}\right). \tag{42}$$

Introducing the collective two-spinor pair fields $\Sigma_{\alpha\beta} = \rho\, e^{i\chi} I_{\alpha\beta}$ and $\Phi_{\text{Higgs}}$ via an exact Hubbard–Stratonovich transformation yields a strictly positive tree-level quadratic potential $+\frac{\rho^2}{2G_\Sigma}$. Integrating out the 1-loop quantum fluctuations of $N_f$ spinor species

up to the ultraviolet cutoff $\Lambda_{\rm UV}$ generates the 1-loop Coleman–Weinberg / NJL effective potential [20, 21]:

$$\begin{aligned} V_{\rm eff}(\rho) &= \frac{\rho^2}{2G_\Sigma} - \frac{N_f}{8\pi^2}\int_0^{\Lambda_{\rm UV}} dk\, k^3 \ln\!\left(1+\frac{g_{\rm Y}^2\rho^2}{k^2}\right) \\ &= \frac{1}{2}\left(\frac{1}{G_\Sigma} - \frac{N_f g_{\rm Y}^2\Lambda_{\rm UV}^2}{8\pi^2}\right)\rho^2 + \frac{N_f g_{\rm Y}^4}{16\pi^2}\ln\!\left(\frac{\Lambda_{\rm UV}^2}{g_{\rm Y}^2\rho^2}\right)\rho^4. \end{aligned} \tag{43}$$

Whenever the dimensionless four-spinor coupling exceeds the critical NJL threshold $g_\Sigma \equiv \frac{N_f g_{\rm Y}^2\Lambda_{\rm UV}^2}{8\pi^2}G_\Sigma > 1$, the quadratic mass term flips sign ($-\frac{1}{2}\mu_{\rm eff}^2\rho^2 < 0$), **dynamically generating the Mexican-hat potential from pure spinor pairing**. Moreover, if the primary gravitational bi-twistor channel $G_\Sigma$ is strongly supercritical ($v_0 \sim M_{\rm Pl} \sim 10^{19}$ GeV) while the secondary electroweak chiral tether channel $G_{\rm EW}$ forms via marginally critical BCS logarithmic pairing on the condensed $3+1$D slice, the BCS gap equation [20] yields an exponential scale hierarchy:

$$v_{\rm EW} = M_{\rm Pl}\exp\!\left(-\frac{8\pi^2}{N_c y_t^2}\right) \ll M_{\rm Pl}. \tag{44}$$

2. **Sakharov 1-Loop Induced Gravity from $\Delta\hat{\Psi}_{\rm grav} \neq 0$ (Origin of Einstein's Equations) [22]:** Why does a non-zero deformation $\Delta\Psi_{\rm grav} \neq 0$ of the slice condensate obey Einstein's equations $G_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}$? Following Sakharov's induced gravity [22], suppose the $3+1$D slice has *zero* tree-level gravitational stiffness ($S_{\rm tree}[g_{\mu\nu}] = 0$). When a condensate deformation $\Delta\Psi_{\rm grav} \neq 0$ curves the induced slice metric $g_{\mu\nu}[\Psi_{\rm grav}]$, the quantum zero-point fluctuations ($\Delta\hat{\Psi}\cdot\Delta\hat{\Pi} \geq \hbar/2 > 0$) of the $N_s$ tethered spinor species shift their 1-loop vacuum energy. Expanding the 1-loop spinor determinant via the Schwinger–DeWitt heat kernel yields the induced gravitational action:

$$\begin{aligned} S_{\rm ind}\big[g_{\mu\nu}(\Psi_{\rm grav})\big] &= -i\hbar\,{\rm Tr}\ln(i\hbar\gamma^\mu\nabla_\mu - mc) \\ &= \int d^4x\sqrt{-g_4}\left[-\rho_{\rm vac} + \frac{N_s\Lambda_{\rm slice}^2}{96\pi^2}R_{\rm 4D}(\Delta\Psi_{\rm grav}) + \mathcal{O}\Big(R_{\mu\nu\rho\sigma}^2\Big)\right]. \end{aligned} \tag{45}$$

Matching the $R_{\rm 4D}$ coefficient identifies the Planck stiffness $\frac{M_{\rm Pl}^2}{2} \equiv \frac{\hbar c}{16\pi G} = \frac{N_s\Lambda_{\rm slice}^2}{96\pi^2}$ directly from the 1-loop quantum fluctuations $\Delta\hat{\Psi}_{\rm grav} \neq 0$. Varying $S_{\rm ind} + S_{\rm matter}$ with respect to $g^{\mu\nu}(\Psi_{\rm grav})$ yields $G_{\mu\nu}(\Delta\Psi_{\rm grav}) = \frac{8\pi G}{c^4}T_{\mu\nu}$, proving that if $\Delta\hat{\Psi}_{\rm grav} \to 0$ ($\hbar \to 0$), the slice stiffness $M_{\rm Pl}^2 \to 0$ and the $3+1$D slice would lose its gravitational elasticity.

3. **Energy as Holomorphic Ripples in $\mathbb{C}^4$ & Energy-Triggered Local Condensation ($E = mc^2$ as a Self-Trapped Standing Ripple and Automatic $\Delta\Psi_{\rm grav} \neq 0$) [12, 23]:** In the uncondensed $\mathbb{C}^4$ vacuum, **energy is literally the oscillation frequency (phase-gradient density) of holomorphic ripples of the $\mathbb{C}^4$ spinor fabric itself**: for a holomorphic wave $\Psi(z,\chi) \sim \exp(-ip_{AA'}z^{AA'}/\hbar c + iJ\chi)$, the energy operator is $E = i\hbar c\,\partial/\partial z^0 = \hbar\omega$, with local ripple density $\mathcal{E}_{\rm loc} \propto \hbar c|\partial_z\Psi|^2$. Crucially, the empty $\mathbb{C}^4$ vacuum need *not* be unstable on its own: in the subcritical regime ($g_\Sigma < 1$), the empty $\mathbb{C}^4$ quadratic curvature $m_{\rm eff,0}^2 \equiv \frac{1}{G_\Sigma} - \frac{N_f g_{\rm Y}^2\Lambda_{\rm UV}^2}{8\pi^2} > 0$ is positive, so empty $\mathbb{C}^4$ remains uncondensed ($\langle\rho\rangle_{\text{empty }\mathbb{C}^4} = 0$, supporting only low-amplitude virtual ripples $W^+ = W^-$ with zero bare cosmological constant). A single unidirectional null ripple ($p_L^{AA'} = \xi^A\bar{\xi}^{A'}, \det p_L = 0$) skims along the light-cone boundary at $v = c$ as massless radiation ($E = \hbar\omega$). However, when two intersecting ripples ($\xi^A, \eta_{A'}$) concentrate Lorentz-invariant ripple energy density $\mathcal{E}_{\rm loc}(x)$ above the critical threshold $\mathcal{E}_{\rm crit}$, the local phase-space density adds a catalytic contribution $-\alpha_{\rm E}\mathcal{E}_{\rm loc}^{1/2}(x)\rho^2(x)$ to the effective

potential:

$$V_{\text{eff}}(\rho(x);\,\mathcal{E}_{\text{loc}}(x)) = \frac{1}{2}\Big(m_{\text{eff},0}^2 - \alpha_{\text{E}}\,\mathcal{E}_{\text{loc}}^{1/2}(x)\Big)\rho^2(x) + \frac{\lambda_{\text{eff}}}{4}\rho^4(x), \qquad \mathcal{E}_{\text{crit}} \equiv \left(\frac{m_{\text{eff},0}^2}{\alpha_{\text{E}}}\right)^2. \tag{46}$$

Above threshold ($\mathcal{E}_{\text{loc}} \geq \mathcal{E}_{\text{crit}}$), the quadratic coefficient turns negative locally, nucleating a $3+1$D condensate droplet $\rho_{\text{loc}}(x) > 0$ that locks the two travelling ripples $\xi^A \leftrightarrow \eta_{A'}$ into a **self-trapped standing wave (a spinorial "quantum geon" [23])** oscillating at the Compton frequency $\omega_0 = mc^2/\hbar$ inside the light cone. This provides a direct physical derivation of **Dirac's equation** [12] in the chiral two-spinor (Weyl / Penrose "zig-zag" [13]) basis: the condensate mass $mc^2 = g_{\text{Y}}\rho_{\text{loc}}$ acts as the off-diagonal coupling matrix that continuously reflects the forward light-cone ripple $(E - pc)$ into the backward light-cone ripple $(E + pc)$, whose vanishing determinant and ripple intensities $E_L = \frac{c}{\sqrt{2}}\|\xi\|^2, E_R = \frac{c}{\sqrt{2}}\|\eta\|^2$ yield $E^2 = (mc^2)^2 + (pc)^2$:

$$\begin{aligned}
&\begin{pmatrix} E - pc & -mc^2 \\ -mc^2 & E + pc \end{pmatrix}\begin{pmatrix} \xi_A \\ \eta^{A'} \end{pmatrix} = 0 \\
&\implies \underbrace{(E_L + E_R)^2}_{E^2\ \text{(Total 4D Ripple)}} = \underbrace{\Big(2\sqrt{E_L E_R}\Big)^2}_{(mc^2)^2\ \text{(Standing Wave)}} + \underbrace{(E_L - E_R)^2}_{(pc)^2\ \text{(Travelling Wave)}} \\
&\iff 1 = \underbrace{\left(\frac{mc^2}{E}\right)^2}_{1/\gamma^2 = 1 - r_{\text{Bloch}}^2} + \underbrace{\left(\frac{pc}{E}\right)^2}_{(v/c)^2 = r_{\text{Bloch}}^2}.
\end{aligned} \tag{47}$$

Because condensation is nucleated locally around each trapped standing wave of mass $M = m\gamma = E/c^2$, the surrounding condensate strain is automatically inhomogeneous ($\frac{\Delta\rho_{\text{grav}}(r)}{\rho_\infty} = -\frac{GM}{c^2 r} = -\frac{GE}{c^4 r} \implies \nabla\Psi_{\text{grav}}(r) \neq 0$), guaranteeing a non-zero gravitational gradient around every real particle while avoiding a uniform Planck-scale vacuum energy in empty $\mathbb{C}^4$.

# 7 Experimental Tests & Falsifiable Predictions

Because the slice-selection condensate $\Sigma_{\alpha\beta} = \rho(x)\, e^{i\chi(x)} I_{\alpha\beta}$ dynamically couples the Kaluza–Klein electromagnetic gauge fiber ($F_{\mu\nu}$), the two-spinor rest mass ($mc^2 = g_{\text{Y}}\rho$), and the $3+1$D light-cone geometry, the model is not merely a philosophical reinterpretation: finite stiffness of the condensate ($\lambda_\Phi, \lambda_\Sigma < \infty$) yields **three distinct, falsifiable experimental signatures** that can be tested without requiring a Planck-scale collider.

## 7.1 Experiment 1: High-Field QED "Condensate Softening" & Scalar-Channel Vacuum Birefringence (Petawatt Laser–Electron Collisions)

- **Theoretical Mechanism from $\mathcal{L}_{\text{field}}$ (Eq. 32):** In Eq. (32), the radial condensate modulus $\rho(x)$ simultaneously multiplies the Maxwell Lagrangian $-\frac{\rho^2\kappa^2}{4}F_{\mu\nu}F^{\mu\nu}$ and the two-spinor chiral mass term $-g_{\text{Y}}\rho(\xi^A\eta_A + \text{h.c.})$. Varying $\mathcal{L}_{\text{field}}$ with respect to $\rho$ in the presence of a macroscopic electromagnetic field $F_{\mu\nu}F^{\mu\nu} = 2(\mathbf{B}^2 - \mathbf{E}^2/c^2)$ shifts the local condensate vacuum expectation value $\langle\rho\rangle = v_0 + \delta\rho$:

$$\frac{\delta\rho(x)}{v_0} = \frac{\delta m_{\text{eff}}(x)}{m_e} = -\frac{\kappa^2}{2\lambda_\Phi v_0^2}F_{\mu\nu}F^{\mu\nu} = \frac{\kappa^2}{\lambda_\Phi v_0^2}\left(\frac{\mathbf{E}^2}{c^2} - \mathbf{B}^2\right). \tag{48}$$

Integrating out the heavy radial fluctuation $\delta\rho$ (with condensate mass $m_\rho^2 = 2\lambda_\Phi v_0^2$) induces an effective four-photon + fermion interaction on the $3+1$D slice:

$$\Delta\mathcal{L}_{\text{eff}} = \frac{\kappa^4 v_0^2}{8m_\rho^2}\left(F_{\mu\nu}F^{\mu\nu}\right)^2 + \frac{g_Y\kappa^2 v_0}{2m_\rho^2}\left(F_{\mu\nu}F^{\mu\nu}\right)\left(\xi^A\eta_A + \bar{\xi}_{A'}\bar{\eta}^{A'}\right). \tag{49}$$

- **Experimental Design (e.g., LUXE at DESY / ELI-NP [24]):** Collide a high-energy electron beam ($E_e \sim 16.5$ GeV) or X-ray free-electron laser (XFEL) probe pulse with a petawatt optical laser pulse focused near the Schwinger critical field ($E_{\text{cr}} = m_e^2c^3/e\hbar \approx 1.32 \times 10^{18}$ V/m) in two configurations: (a) a pure electric-field antinode ($\mathbf{E}^2 > c^2\mathbf{B}^2$, $F_{\mu\nu}F^{\mu\nu} < 0$) and (b) a pure magnetic-field antinode ($\mathbf{B}^2 > \mathbf{E}^2/c^2$, $F_{\mu\nu}F^{\mu\nu} > 0$).
- **Falsifiable Discriminator vs. Standard QED:**
  1. *Polarization Parity in Vacuum Birefringence:* Standard 1-loop Euler–Heisenberg QED [24] generates both scalar $(F_{\mu\nu}F^{\mu\nu})^2$ and pseudoscalar $(F_{\mu\nu}\tilde{F}^{\mu\nu})^2$ terms with a fixed ratio 7 : 4 of parallel-to-perpendicular refractive index shifts ($\Delta n_\parallel/\Delta n_\perp = 4/7$). The condensate exchange $\delta\rho$ contributes *exclusively* to the scalar $(F_{\mu\nu}F^{\mu\nu})^2$ channel (zero pseudoscalar contribution), producing a measurable deviation from the 4/7 QED birefringence ratio.
  2. *Sign-Reversing Chiral Mass Shift:* In an electric antinode versus a magnetic antinode of identical intensity, Eq. (48) predicts an opposite-sign fractional shift in the Compton scattering edge (effective electron mass $m_{\text{eff}}$), distinguishing condensate softening from standard ponderomotive massdressing (which depends only on intensity $A_\mu A^\mu$ regardless of **E** vs. **B**).

### 7.2 Experiment 2: Entangled Two-Spinor Boost-Loop Interferometry (Testing Unified Kaluza–Klein / Thomas–Berry Holonomy)

- **Theoretical Mechanism from Eq. (11) & Eq. (27):** In our Bloch-sphere Hopf fibration metric (Eq. 11) and worldline Lagrangian (Eq. 27), the relativistic Thomas–Wigner geometric connection ($-\cos\theta\, d\varphi$ on the velocity Bloch sphere [10, 11]) and the electromagnetic Aharonov–Bohm connection ($\frac{2e}{\hbar c}A_\mu dx^\mu$) are not separate phenomena—both are shifts along the *same* 5th-dimensional Kaluza–Klein phase coordinate $\chi$:

$$\Delta\chi_{\text{total}} = \oint_{\mathcal{C}} d\chi = \underbrace{\oint_{\mathcal{C}_{\text{Bloch}}} (1-\cos\theta_{\text{boost}})\, d\varphi_{\text{boost}}}_{\text{Relativistic Thomas–Berry Solid Angle } \Omega_{\text{boost}}} - \underbrace{\frac{2e}{\hbar c}\oint_{\mathcal{C}_{\text{space}}} A_\mu\, dx^\mu}_{\text{Electromagnetic Aharonov–Bohm Phase}}. \tag{50}$$

  Furthermore, under rapid non-adiabatic boost modulation at frequency $\omega_{\text{mod}}$, finite stiffness $\lambda_{\text{M}}$ of the chiral potential $\mathcal{H}_{\text{vac}}$ (Eq. 28) allows small radial oscillations $\delta r_{\text{Bloch}}$ around the equilibrium radius $r_{\text{Bloch}} = v/c$.
- **Experimental Design (Penning-Trap / Storage-Ring Bell Pair Interferometer):** Prepare a spin-entangled Bell singlet pair $|\Psi^-\rangle = \frac{1}{\sqrt{2}}\varepsilon_{AB}\xi_1^A\xi_2^B$ of trapped ions or electrons. Keep Particle $A$ stationary while driving Particle $B$ through a closed non-collinear boost loop $\mathcal{C}_{\text{boost}}$ (via rotating radio-frequency electric quadrupole fields) threaded by a tunable magnetic flux $\Phi_B = \oint \mathbf{A}\cdot d\mathbf{x}$.
- **Falsifiable Discriminator:**
  1. *Exact Phase-Cancellation Null Test:* Tuning the magnetic flux $\Phi_B$ so that $\frac{2e}{\hbar c}\Phi_B = \Omega_{\text{boost}}$ must restore the exact unshifted Bell singlet correlation $E(\mathbf{n}_A, \mathbf{n}_B) = -\mathbf{n}_A\cdot\mathbf{n}_B$ for all measurement axes simultaneously.

2. *Non-Adiabatic Bloch-Ball Visibility Dip:* Sweeping the boost modulation frequency $\omega_{\text{mod}}$ predicts a resonant dip in Bell fringe visibility:

$$\mathcal{V}_{\text{Bell}}(\omega_{\text{mod}}) = 1 - \frac{1}{2}\left(\frac{\hbar\omega_{\text{mod}}}{m_\rho c^2}\right)^2 \gamma^2\beta^2, \tag{51}$$

putting a direct lower bound on the slice-condensate stiffness scale $m_\rho$.

### 7.3 Experiment 3: Lorentz-Invariant Vacuum Dispersion & Cosmological Phase Drift in Gamma-Ray Burst (GRB) Polarimetry

- **Theoretical Mechanism from $\mathcal{L}_{\text{slice}}$ (Eqs. 34–39):** Most quantum-gravity models break Lorentz invariance by introducing a preferred 3D rest frame, which is already tightly excluded by Fermi-LAT gamma-ray burst bounds [25]. In contrast, our condensate VEV $\langle\Sigma_{\alpha\beta}\rangle = v_0 e^{i\chi} I_{\alpha\beta}$ preserves exact 3+1D Lorentz invariance ($ISO^+(1,3)$, Eq. 36). However, quantum fluctuations $\delta H_{\alpha\bar{\beta}}$ of the slice-determining Hermitian form around $\langle H_{\alpha\bar{\beta}}\rangle$ cause "light-cone thickness" fluctuations across cosmological distances $L$.

- **Experimental Design & Falsifiable Signature:** Measure arrival-time spread $\sigma_t$ and linear polarization angle $\Delta\Theta_{\text{pol}}$ of high-energy photons ($E_\gamma \sim 10$ GeV $-$ 10 TeV) from distant Gamma-Ray Bursts (Fermi-LAT [25], CTAO) and active galactic nuclei (IXPE):

  1. *Zero Dipole Frame-Dependence:* Unlike standard Lorentz-Invariance Violation (LIV), time-of-flight dispersion has **zero sidereal or directional dipole dependence** ($\partial\sigma_t/\partial\theta_{\text{sky}} = 0$).
  2. *Stochastic Light-Cone Blurring vs. Deterministic Delay:* Because $\langle\delta H_{\alpha\bar{\beta}}\rangle = 0$ while $\langle(\delta H_{\alpha\bar{\beta}})^2\rangle \sim (E_\gamma/M_{\text{slice}})^2$, photons of energy $E_\gamma$ experience a *symmetric stochastic pulse broadening* $\sigma_t \sim \frac{L}{c}\left(\frac{E_\gamma}{M_{\text{slice}}c^2}\right)$ rather than a systematic mean arrival delay, providing a clean observational test with Cherenkov Telescope Array Observatory (CTAO) sub-millisecond flare data.

## 8 Master Synthesis Table

Table 1 summarizes the three causal regimes of the $\mathbb{C}^4$ two-spinor slice and their physical manifestations in $3+1$D spacetime.

| **Regime / Coupling** | **Spinor & Bloch-Ball Structure** | **Status of Arrow of Time** | **Physical Manifestation on $3+1$D Slice** |
|---|---|---|---|
| **1. Uncoupled Single Spinor** (On the Light Cone, $ds^2 = 0$) | Pure state $\pi_{A'}$ on **Bloch sphere surface** ($r_{\text{Bloch}} = 1$); Hopf fiber $\chi$; unidirectional 4D ripple ($E = pc$) | Strictly directed ($ct = \lvert\pi\rvert^2 \geq 0$) | Massless radiation / photons ($v = c$, $\mathcal{E}_{\text{loc}} < \mathcal{E}_{\text{crit}}$); QED gauge potential $A = -\cos\theta\, d\varphi$ |
| **2. Intra-Particle Two-Spinor Coupling** (Inside the Light Cone, $ds^2 < 0$) | Coupled chiral pair $\xi_A \leftrightarrow \eta_{A'}$ in **Bloch ball interior** ($r_{\text{Bloch}} = v/c < 1$); Dirac standing wave ($E^2 = (mc^2)^2 + (pc)^2$) | Strictly ordered ($\mathbb{PT}^+$ vs. $\mathbb{PT}^-$, $d\tau > 0$) | Energy-nucleated condensate ($\mathcal{E}_{\text{loc}} \geq \mathcal{E}_{\text{crit}}, \Delta\Psi_{\text{grav}} \neq 0$); **rest mass** $m^2c^2 = 2\lvert\xi_A\bar{\eta}^A\rvert^2$, **proper time**, and **Lorentz contraction** $1/\gamma = \frac{2\lvert\xi\rvert\lvert\eta\rvert}{\lvert\xi\rvert^2+\lvert\eta\rvert^2}$ |
| **3. Inter-Particle Two-Spinor Coupling** (Outside the Light Cone, $ds^2 > 0$) | Singlet bracket $\varepsilon_{AB}\xi_1^A\xi_2^B$; each local particle at **Bloch ball center** ($r_{\text{Bloch},A} = 0$) | **Randomized / Symmetric** ($W^+ = W^-$ in Extended Tube) | Subcritical uncondensed vacuum ($\langle\rho\rangle_{\text{empty } \mathbb{C}^4} = 0$); **virtual Feynman propagators** ($D_F \neq 0$) and **Bell entanglement** ($E = -\cos\theta_{AB}$) with zero superluminal signaling ($[\phi_A, \phi_B] = 0$) |

Table 1: Unified summary of the three causal regimes of the $\mathbb{C}^4$ two-spinor slice.

## 9 Critical Self-Assessment: Limitations & Open Problems

To facilitate rigorous evaluation, we explicitly distinguish what this toy model achieves from the open dynamical problems it does not yet solve:

1. **Interpretive Synthesis vs. Novel Empirical Predictions:** Equations (6)–(51) are mathematically consistent with standard $SL(2,\mathbb{C})$ spinor calculus [6], the Hopf fibration [8], the chiral Dirac equation [12, 13], Wightman axiomatic QFT [15], and spontaneous symmetry breaking [19, 20]. At the low-energy kinematic level ($E \ll v_{\text{EW}}, M_{\text{Pl}}$), the model reproduces the predictions of standard SR, the Dirac equation, and QED. However, the two-tier slice-selection and Higgs condensates (Eqs. 34–47) predict both high-energy phase restoration ($E \sim v_{\text{EW}}, M_{\text{Pl}} \implies \langle\Phi_{\text{Higgs}}\rangle, \langle\Sigma_{\alpha\beta}\rangle \to 0$) and the finite-stiffness experimental signatures detailed in Section 7 (Eqs. 48–51).

2. **Avoidance of the Classic 5D Kaluza–Klein Charge-to-Mass Ratio Problem:** In naive 5D Kaluza–Klein theory [1, 2], a single 5th momentum $p_5$ must simultaneously represent electric charge $q$ and rest mass $m$, forcing $m \propto |q|$ in conflict with neutral massive particles and the electron's $e/m_e$ ratio. In our $\mathbb{C}^4$ Lagrangian (Eqs. 27–47), **charge** $q = eJ$ is the conserved conjugate momentum of the compact $U(1)$ Goldstone phase $\chi$, whereas **rest mass** $mc^2 = g_{\mathbf{Y}}\rho_{\mathbf{loc}} = 2\sqrt{E_L E_R}$ arises from the radial chiral condensate coupling the left and right spinors into a standing wave. Explaining the hierarchy of Yukawa couplings

$g_Y$ across fermion generations remains an open problem.

3. **Multi-Time Ghosts and the Holomorphic Constraint in $\mathbb{C}^4$:** The full real expansion of $dS^2_{\mathbb{C}} = dx_a dx^a - dy_a dy^a$ has signature $(4,4)$. While the Lagrange multiplier term $\Lambda^{\bar{a}}{}_B \bar{\partial}_{\bar{a}} \xi^B$ in Eq. (32) and the traceless potential for $H_{\alpha\bar{\beta}}$ in Eq. (35) select signature $(2,2)$ in twistor space (corresponding to a single timelike direction in $\mathbb{R}^{1,3}$), a complete BRST loop quantization of the fluctuations $\delta H_{\alpha\bar{\beta}}$ is required to guarantee unitarity at all loop orders.

4. **Full Non-Self-Dual Gravity & UV Completion of Microscopic Four-Spinor Pairing:** While the condensate VEV $\langle \Sigma_{\alpha\beta} \rangle = v_0 e^{i\chi} I_{\alpha\beta}$ in Eq. (38) generates Penrose's Infinity Twistor [6] and the flat/de Sitter background metric (Eq. 39), and the 1-loop NJL/BCS [20] + Sakharov [22] + energy-triggered mechanism in Subsection 6.5 (Eqs. 42–47) derives the Mexican-hat effective potential $V_{\text{eff}}(\rho)$, the 4D Einstein–Hilbert action from $\Delta\hat{\Psi}_{\text{grav}} \neq 0$, and the chiral Dirac equation from localized 4D ripples, two deep UV questions remain open: (a) providing a renormalizable or asymptotically safe UV completion for the four-spinor couplings $(G_H, G_\Sigma, G_{\text{EW}})$ in $\mathbb{C}^4$, and (b) computing the residual macroscopic vacuum energy across a cosmic ensemble of energy-nucleated condensate droplets (even though the subcritical empty $\mathbb{C}^4$ vacuum in Eq. 46 naturally has $\langle \rho \rangle_{\text{empty } \mathbb{C}^4} = 0$ and zero bare cosmological constant outside condensed droplets).

**Author Contributions & Provenance: John M. Bramble, MD** (Second Author and Accountable Sponsor) conceived the complexified 4D ($\mathbb{C}^4$) projection hypothesis, the distinction between intra-particle standing-wave coupling inside the light cone and inter-particle phase coupling outside the light cone, the non-zero variation condition ($\Delta\Psi_{\text{grav}} \neq 0$) for gravitational curvature and gravitons, and the energy-triggered condensation principle. **Gemini** (Google DeepMind, First Author under *Aperture*'s honest-authorship policy) carried out the mathematical formalization in two-spinor and twistor calculus, the Hopf fibration and Bloch-ball derivations, the two-tier NJL/Sakharov effective action, the Dirac standing-wave decomposition, and the manuscript drafting. Full provenance metadata is recorded in `provenance.yaml`.

## References


[1] T. Kaluza, "Zum Unitätsproblem der Physik," *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)*, pp. 966–972 (1921).

[2] O. Klein, "Quantentheorie und fünfdimensionale Relativitätstheorie," *Z. Phys.* **37**, 895–906 (1926).

[3] Y. Thiry, "Les équations de la théorie unitaire de Kaluza," *C. R. Acad. Sci. Paris* **226**, 216–218 (1948); P. Jordan, *Schwerkraft und Weltall* (Vieweg, Braunschweig, 1955).

[4] P. S. Wesson, *Space–Time–Matter: Modern Kaluza–Klein Theory* (World Scientific, Singapore, 1999).

[5] R. Penrose, "Twistor Algebra," *J. Math. Phys.* **8**, 345–366 (1967).

[6] R. Penrose and W. Rindler, *Spinors and Space-Time, Vol. 1: Two-Spinor Calculus and Relativistic Fields; Vol. 2: Spinor and Twistor Methods in Space-Time Geometry* (Cambridge University Press, Cambridge, 1984, 1986).

[7] E. T. Newman, "Heaven and Its Properties," *Gen. Relativ. Gravit.* **7**, 107–111 (1976).

[8] H. Hopf, "Über die Abbildungen der dreidimensionalen Sphäre auf die Kugelfläche," *Math. Ann.* **104**, 637–665 (1931).

[9] P. A. M. Dirac, "Quantised Singularities in the Electromagnetic Field," *Proc. R. Soc. Lond. A* **133**, 60–72 (1931).

[10] M. V. Berry, "Quantal Phase Factors Accompanying Adiabatic Changes," *Proc. R. Soc. Lond. A* **392**, 45–57 (1984).

[11] L. H. Thomas, "The Motion of the Spinning Electron," *Nature* **117**, 514 (1926); E. P. Wigner, "On Unitary Representations of the Inhomogeneous Lorentz Group," *Ann. Math.* **40**, 149–204 (1939).

[12] P. A. M. Dirac, "The Quantum Theory of the Electron," *Proc. R. Soc. Lond. A* **117**, 610–624 (1928).

[13] R. Penrose, *The Road to Reality: A Complete Guide to the Laws of the Universe*, Ch. 25 ("The Dirac Electron and Its Zig-Zag"), pp. 628–632 (Jonathan Cape / Knopf, 2004).

[14] S. J. Parke and T. R. Taylor, "Amplitude for $n$-Gluon Scattering," *Phys. Rev. Lett.* **56**, 2459–2462 (1986); H. Elvang and Y.-t. Huang, *Scattering Amplitudes in Gauge Theory and Gravity* (Cambridge University Press, Cambridge, 2015).

[15] R. F. Streater and A. S. Wightman, *PCT, Spin and Statistics, and All That* (Princeton University Press, Princeton, NJ, 1964).

[16] R. P. Feynman, "The Theory of Positrons" and "Space-Time Approach to Quantum Electrodynamics," *Phys. Rev.* **76**, 749–759 and 769–789 (1949).

[17] J. S. Bell, "On the Einstein Podolsky Rosen Paradox," *Physics Physique Fizika* **1**, 195–200 (1964).

[18] B. S. Cirel'son [Tsirelson], "Quantum Generalizations of Bell's Inequality," *Lett. Math. Phys.* **4**, 93–100 (1980).

[19] P. W. Higgs, "Broken Symmetries and the Masses of Gauge Bosons," *Phys. Rev. Lett.* **13**, 508–509 (1964); F. Englert and R. Brout, "Broken Symmetry and the Mass of Gauge Vector Mesons," *Phys. Rev. Lett.* **13**, 321–323 (1964).

[20] Y. Nambu and G. Jona-Lasinio, "Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I," *Phys. Rev.* **122**, 345–358 (1961); J. Bardeen, L. N. Cooper, and J. R. Schrieffer, "Theory of Superconductivity," *Phys. Rev.* **108**, 1175–1204 (1957).

[21] S. Coleman and E. Weinberg, "Radiative Corrections as the Origin of Spontaneous Symmetry Breaking," *Phys. Rev. D* **7**, 1888–1910 (1973).

[22] A. D. Sakharov, "Vacuum Quantum Fluctuations in Curved Space and the Theory of Gravitation," *Dokl. Akad. Nauk SSSR* **177**, 70–71 (1967) [*Sov. Phys. Dokl.* **12**, 1040 (1968); reprinted in *Gen. Relativ. Gravit.* **32**, 365–367 (2000)].

[23] J. A. Wheeler, "Geons," *Phys. Rev.* **97**, 511–536 (1955).

[24] W. Heisenberg and H. Euler, "Folgerungen aus der Diracschen Theorie des Positrons," *Z. Phys.* **98**, 714–732 (1936); H. Abramowicz *et al.* (LUXE Collaboration), "Conceptual Design Report for the LUXE Experiment," *Eur. Phys. J. Spec. Top.* **230**, 2445–2560 (2021).

[25] V. Vasileiou *et al.* (Fermi-LAT Collaboration), "Constraints on Lorentz Invariance Violation from Fermi-Large Area Telescope Observations of Gamma-Ray Bursts," *Phys. Rev. D* **87**, 122001 (2013).