

Linear Programming: Solving Linear Systems

Daniel Kane

Department of Computer Science and Engineering
University of California, San Diego

Advanced Algorithms and Complexity
Data Structures and Algorithms

Learning Objectives

- Solve a system of linear equations.
- Say something about what the set of solution to a system of linear equations looks like.

Last Time

Linear programming: Dealing with systems of linear inequalities.

Linear Algebra

Today, we will deal with the simpler case, of systems of linear **equalities**.

Linear Algebra

Today, we will deal with the simpler case, of systems of linear **equalities**.

For example:

$$\begin{aligned}x + y &= 5 \\ 2x + 4y &= 12.\end{aligned}$$

Method of Substitution

- Use first equation to solve for one variable in terms of the others.
- Substitute into other equations.
- Solve recursively.
- Substitute back in to first equation to get initial variable.

Example

$$x + y = 5$$

$$2x + 4y = 12.$$

Example

$$\begin{aligned}x + y &= 5 \\ 2x + 4y &= 12.\end{aligned}$$

First equation implies

$$x = 5 - y.$$

Example

$$\begin{aligned}x + y &= 5 \\ 2x + 4y &= 12.\end{aligned}$$

First equation implies

$$x = 5 - y.$$

Substituting into second:

$$12 = 2x + 4y = 2(5 - y) + 4y = 10 + 2y.$$

Example

$$\begin{aligned}x + y &= 5 \\ 2x + 4y &= 12.\end{aligned}$$

First equation implies

$$x = 5 - y.$$

Substituting into second:

$$12 = 2x + 4y = 2(5 - y) + 4y = 10 + 2y.$$

So $y = 1, x = 5 - 1 = 4$.

Problem

What is the value of x in the solution to the following linear system?

$$\begin{aligned}x + 2y &= 6 \\ 3x - y &= -3.\end{aligned}$$

Solution

From the first equation, we get

$$x = 6 - 2y.$$

Solution

From the first equation, we get

$$x = 6 - 2y.$$

Substituting into the second,

$$-3 = 3(6 - 2y) - y = 18 - 7y.$$

Solution

From the first equation, we get

$$x = 6 - 2y.$$

Substituting into the second,

$$-3 = 3(6 - 2y) - y = 18 - 7y.$$

Solving gives, $y = 3$, so $x = 6 - 2 \cdot 3 = 0$.

Another Example

Consider the following system of equations:

$$x + y + z = 5$$

$$2x + y - z = 1.$$

Another Example

Consider the following system of equations:

$$x + y + z = 5$$

$$2x + y - z = 1.$$

Solve by substitution.

Solution

From first equation:

$$x = 5 - y - z.$$

Solution

From first equation:

$$x = 5 - y - z.$$

Substitute into second.

$$2(5 - y - z) + y - z = 1,$$

or

$$y = 9 + 3z.$$

Cannot Solve for z !

No equations left.

Cannot Solve for z !

No equations left. However, for any z have solution

$$y = 9 + 3z$$

$$x = 5 - y - z = -4 - 4z.$$

Cannot Solve for z !

No equations left. However, for any z have solution

$$y = 9 + 3z$$

$$x = 5 - y - z = -4 - 4z.$$

Have entire family of solutions. z is a free variable.

Degrees of Freedom

- Your solution set will be a subspace.
- Dimension = number of free variables.
- Each equation gives one variable in terms of others.
- Generally, dimension equals
$$\text{num. variables} - \text{num. equations}.$$

Summary

- Can solve systems using method of substitution.
- Each equation reduces degrees of freedom by one.

Next Time

Systematize this to simplify notation and make into an algorithm.