



MELANOTAN II RESEARCH HUB • RESEARCH USE ONLY

DEEPER TAN. HIGHER DRIVE.

FOR MEN & WOMEN. Get a deeper tan, protect your skin from UV damage, and boost your libido.

Get Melanotan II 10 mg at BioLinx

Read the guide first

Also shown to help with: chronic fatigue • angina • concussion / tbi • heart failure • macular degeneration

Daily until 50 mg total (13 days) • any time – fatigue is a common side effect • Difficulty: beginner

WHAT'S INSIDE

- 1** Shopping list — how many vials to buy GET p.2
- 2** Quick summary of your lab items GET p.3
- 3** Forming your peptide — “reconstitution” PREP p.4
- 4** How to inject DOSE p.5
- 5** Melanotan II dosing schedule DOSE p.6
- 6** What to expect + side effects AFTER p.7
- 7** Recap checklist AFTER p.8
- +** Appendix: ramp-up schedule + sourcing guide BONUS p.9–10

RESEARCH SHOWED THAT...

17 / 20 men

responded
in the erectile-dysfunction
study.

Int J Impotence Res, 2000

Sexual desire that was absent
came back in 68% of men.

Wessells et al., Int J

Impotence Res, 2000



STEP 1 OF 4 · WHAT TO BUY

SHOPPING LIST

Easy-to-follow shopping list to safely practice Melanotan II. Match your commitment level — don't skip anything.



PEPTIDE · ONE ORDER

MELANOTAN II 10 MG

Get a deeper tan, protect your skin from UV damage, and boost your libido. Common starting dose: **4 mg**, daily until 50 mg total (13 days), any time — fatigue is a common side effect.

Get Melanotan II 10 mg at BioLinX

biolinxlabs.com/products/melanotan-ii-10-mg

HOW MANY VIALS TO BUY · BEST VALUE BY GOAL

GOAL	VIAL	QTY	WHERE
1 CYCLE	10 mg vial	5×	<u>See it</u> biolinxlabs.com/products/melanotan-ii-10-mg
2 CYCLES RECOMMENDED	10 mg vial	10×	<u>See it</u> biolinxlabs.com/products/melanotan-ii-10-mg
4 CYCLES	10 mg vial	20×	<u>See it</u> biolinxlabs.com/products/melanotan-ii-10-mg

MATERIALS ≈ \$41 ON TOP — FULL LIST ON THE NEXT PAGE



PRODUCTS ARRIVED?

QUICK SUMMARY OF YOUR LAB ITEMS:

TWO VIALS IN THE BOX • MELANOTAN II 10 MG + BACTERIOSTATIC WATER 10 ML

VIAL 1	Melanotan II 10 mg. Arrives as a powder. Don't touch it until you've read page 4.
VIAL 2	Bacteriostatic water — 10 mL. This is what turns the powder into a liquid. Sterility and pH matter — don't substitute.

GROCERY LIST • ≈ \$41

ITEM	SPEC	QTY	WHERE
INSULIN SYRINGES BUY	1cc U100, 30G, 5/16"	50–100 ct	Any pharmacy
ALCOHOL PREP PADS BUY	Any brand, 70% isopropyl	50–100 ct	Any pharmacy
SHARPS CONTAINER BUY	1 quart, for needle disposal	1×	Any pharmacy
BAC WATER BUY	Bacteriostatic water, 10 mL — sterility and pH matter, don't substitute	1 vial	<u>See it</u> biolinxlabs.com/products/bacteriostatic-water-10-ml

Any brand works for syringes, pads and sharps — but not BAC water. Use 2 fresh syringes and 2 fresh pads every session: one to mix, one to inject.

HOW TO STORE IT


POWDER	Freezer — lasts years
RECONSTITUTED	Fridge — about 28 days. Write the mix date on the vial.
WATER	Room temperature — check the expiration date



STEP 2 OF 4 · RECONSTITUTION – FROM POWDER TO LIQUID

LET'S “FORM” YOUR PEPTIDE!

“Form” means “reconstitute”. Your product arrives as a powder — you need to liquify it before “you begin your research.” 100 units = one full syringe. **10 mg vial** → **add 100 units (1 mL)**. Every dose in this guide assumes exactly that ratio, so measure it — don't eyeball it.



YOUR SIMPLIFIED UNITS

DOSE	FILL TO
1 mg	10 units
2 mg	20 units
3 mg	30 units
4 mg	40 units
5 mg	50 units

Change the vial, water or dose →
[interactive calculator](#)

1

Remove the plastic caps & wipe both rubber vial tops

Fresh alcohol pad on the MT-II vial and the bac water vial.

2

Draw 100 units of bac water

One full pull on a 100-unit insulin syringe (1 mL total).

3

Inject it slowly down the side of the 10 mg vial

Let it run down the glass. Don't blast the powder.

4

Swirl gently. Never shake. Place it in the fridge!

Slow and gentle wins this one — shaking wastes peptide. Discard that syringe into the sharps container. Let it rest until completely clear (36–46 °F); use within ~28 days; never freeze it.



STEP 3 OF 4 · INJECT

WHERE TO INJECT

Time to inject · clean, pinch, poke, done. Daily until 50 mg total (13 days), any time — fatigue is a common side effect.

1

Take MT-II out of the refrigerator and extract your dose

Your first dose is **4 mg = 40 units** on a 100-unit syringe. Wipe the vial top with a fresh alcohol pad, let it air-dry a few seconds, then draw your units.

2

Pinch a fold of skin on the stomach + use an alcohol wipe

Two inches away from the belly button. Wipe with a fresh alcohol pad, let it air-dry, and don't touch that spot again before injecting. Rotate sites.

3

Needle in at the base of the fold

Push the plunger slowly and steadily. Hold for a few seconds, then withdraw.

4

That's it — you just did your first shot!

Drop the needle straight into the sharps bin. Vial goes back in the fridge.



STOMACH · PINCH A FOLD · NEEDLE AT THE BASE

WATCH FIRST


Mixing and injecting from start to finish, on video: professorpeptides.com. Use 2 fresh syringes and 2 fresh pads every session — one to mix, one to inject.



STEP 4 OF 4 · DOSING & TIMING SCHEDULE

MELANOTAN II SCHEDULE (PROTOCOL)

“Schedule” means “protocol”. Everything below assumes the 10 mg vial was reconstituted with exactly 100 units (1 mL), read off a standard 100-unit insulin syringe.



FIRST DOSE

How much, how often & when?

WHEN

Any time — fatigue is a common side effect. Daily until 50 mg total (13 days).

HOW MUCH

4 mg to start.

Dose runs in on/off blocks with breaks between — this is **cycling**, not titration. Full schedule on page 9.

UNITS

Extract **40 units** on a 100-unit insulin syringe.

40 units = 4 mg. Exact variables: 100-unit syringe · 10 mg vial reconstituted with 100 units · 0.1 mg per unit.

HOW OFTEN

Daily until 50 mg total (13 days). Hold each step for its full block before moving up.

CYCLE

Not a monthly cycle: run daily to 50 mg, then 6 weeks off. Repeat 2–4× per year..

Don't rush it — start at the lowest amount, hold each step for its full block, and only step up if the current dose is comfortable. Most people never need the top doses.



STEP 4 OF 4 · DOCUMENT

WHAT TO EXPECT + SIDE EFFECTS

Mostly mild, mostly early, mostly right after a dose increase. Know the three tiers before the first shot, not after.

NORMAL / EXPECTED

First few days

- Nausea and facial flushing within minutes of the shot, easing within 1–2 hours — the most common effects
- Fatigue and yawning, especially the first few days
- Darkening of existing moles and freckles as the tan develops

COMMON, WORTH NOTING

Manage it, don't push the dose

- Vomiting or loss of appetite — take the dose before bed and lower it
- Spontaneous erections or increased libido (a melanocortin effect)
- New freckles or darkening in patches — rotate sun exposure, use SPF

STOP & CALL A DOCTOR

Don't push through it

- A new mole, or a mole that grows, changes shape or colour, itches or bleeds — get it checked (melanoma has been reported in users)
- An erection lasting more than 4 hours — medical emergency
- A sustained rise in blood pressure, chest pain, or a pounding heartbeat
- Signs of an allergic reaction — swelling, rash, difficulty breathing

NOTE: for this one it is **HIGHLY RECOMMENDED** to start at the lowest amount first and work up over time, as needed. Melanotan II is unapproved everywhere and health regulators warn against it. Not for anyone with a history of melanoma or many atypical moles, high blood pressure, or who is pregnant — talk to a doctor before use and photograph your moles before you start.



FIRST-TIME TL;DR

RECAP: CHECKLIST

Everything from pages 2–7, in the order it happens.

BEFORE

- ☐ Vial reconstituted with exactly 100 units of bac water, resting in the fridge, mix date written on it
 - ☐ Supplies laid out: new syringe, alcohol pad, sharps container
 - ☐ Dose set at **4 mg = 40 units** — not higher, no matter what a forum says
-

DURING

- ☐ Inject daily until 50 mg total (13 days), any time — fatigue is a common side effect — stomach, rotate the spot
 - ☐ Note the date and dose somewhere — a phone note is fine
-

AFTER

- ☐ Expect possible redness, mild irritation or a headache — it settles
 - ☐ Mild nausea can happen after a dose increase — sit tight, it typically passes
 - ☐ If anything feels more than mild — dizziness, real nausea, allergic reaction — stop and get medical advice
-

NEXT BLOCK


- ☐ Hold each step for its full block before stepping up — never skip a step (not a monthly cycle: run daily to 50 mg, then 6 weeks off. repeat 2–4× per year.)
 - ☐ Only step up if the current dose is tolerated; if side effects are rough, stay another block
 - ☐ Fresh syringe, fresh pad, needle in the sharps bin — every single time
-

UNIT CHEAT-SHEET · 100-UNIT SYRINGE · 10 MG VIAL RECONSTITUTED WITH 100 UNITS

HOW TO CYCLE (RAMP-UP & CYCLE SCHEDULE)

Assumes the 10 mg vial was mixed with exactly 100 units, so **1 unit = 0.1 mg**.

Units shown are per shot.



WHEN	DOSE	SYRINGE · PER SHOT	NOTES
Days 1–12	4 mg per day	40 units	START HERE
Day 13	2 mg	20 units	
Then	Off — 6 weeks	60 units	REST
Repeat	2–4× per year	20 units	

Rule of thumb: don't move up a step until you know how the current dose feels. Only step up if the previous block was tolerated. Most people never need to go past the middle of the ramp.



THE MT-II SOURCING GUIDE

PROFESSOR PEPS: SOURCING GUIDE (APPENDIX)

How to verify any supplier in 90 seconds. A legitimate source clears all three checks below in under two minutes — no back-and-forth required.



1

Ask for the COA

A batch-specific Certificate of Analysis — not a generic PDF reused across batches — showing purity $\geq 98\%$ via HPLC/MS.

2

Match the lot number on the vial

It must match the lot number on the COA. If they don't match, walk away.

3

Check the track record

Searchable reviews or forum mentions older than roughly six months. Brand-new storefronts with no history are the highest-risk category.

DON'T WORRY — WE'VE ALREADY DONE THE VETTING

BioLinx Labs clears the list: third-party HPLC testing, a 99%+ purity guarantee and a COA included with every order. biolinxlabs.com/best-sellers

biolinxlabs.com/best-sellers

Educational and research purposes only — not medical advice. Professor Peptides does not sell peptides. Products referenced are sold for laboratory research use only and are not for human consumption. Want to learn more? Join the Professor Peptides community — 30 days free.

STAY CURIOUS, PROFESSOR PEPTIDES

Melanotan II Research Hub · companion to the email series