



NAD+ RESEARCH HUB · RESEARCH USE ONLY

# MORE ENERGY. SHARPER MIND.

FOR MEN & WOMEN. Get more energy, think sharper, and slow down aging.

Get NAD+ 500 mg at BioLinx

Read the guide first

Also shown to help with: chronic fatigue · aging skin · brain fog  
2x per week · in the morning · Difficulty: beginner

## WHAT'S INSIDE

- |   |                                             |       |        |
|---|---------------------------------------------|-------|--------|
| 1 | Shopping list — how many vials to buy       | GET   | p.2    |
| 2 | Quick summary of your lab items             | GET   | p.3    |
| 3 | Forming your peptide — “reconstitution”     | PREP  | p.4    |
| 4 | How to inject                               | DOSE  | p.5    |
| 5 | NAD+ dosing schedule                        | DOSE  | p.6    |
| 6 | What to expect + side effects               | AFTER | p.7    |
| 7 | Recap checklist                             | AFTER | p.8    |
| + | Appendix: ramp-up schedule + sourcing guide | BONUS | p.9-10 |

## RESEARCH SHOWED THAT...

**Every** human

study  
raised age-declined NAD+  
levels.

Aging Res Rev, 2026

NAD+ levels that declined with  
age were successfully raised in  
every human study that  
measured them.

Gallagher & Emmanuel,

Aging Res Rev, 2026



STEP 1 OF 4 · WHAT TO BUY

# SHOPPING LIST

Easy-to-follow shopping list to safely practice NAD+. Match your commitment level — don't skip anything.



PEPTIDE · ONE ORDER

## NAD+ 500 MG

Get more energy, think sharper, and slow down aging. Common starting dose: **25 mg**, 2× per week, in the morning.

**Get NAD+ 500 mg at BioLinx**

[biolinxlabs.com/products/nad-500-mg](https://biolinxlabs.com/products/nad-500-mg)

HOW MANY VIALS TO BUY · BEST VALUE BY GOAL

| GOAL                               | VIAL        | QTY | WHERE                                                                                                                 |
|------------------------------------|-------------|-----|-----------------------------------------------------------------------------------------------------------------------|
| <b>1 MONTH</b>                     | 500 mg vial | 1×  | <b><u>See it</u></b><br><a href="https://biolinxlabs.com/products/nad-500-mg">biolinxlabs.com/products/nad-500-mg</a> |
| <b>3 MONTHS</b> <b>RECOMMENDED</b> | 500 mg vial | 2×  | <b><u>See it</u></b><br><a href="https://biolinxlabs.com/products/nad-500-mg">biolinxlabs.com/products/nad-500-mg</a> |
| <b>FULL CYCLE</b>                  | 500 mg vial | 4×  | <b><u>See it</u></b><br><a href="https://biolinxlabs.com/products/nad-500-mg">biolinxlabs.com/products/nad-500-mg</a> |

MATERIALS ≈ \$41 ON TOP — FULL LIST ON THE NEXT PAGE



## PRODUCTS ARRIVED?

# QUICK SUMMARY OF YOUR LAB ITEMS:

TWO VIALS IN THE BOX • NAD+ 500 MG + BACTERIOSTATIC WATER 10 ML

VIAL 1      **NAD+ 500 mg.** Arrives as a powder. Don't touch it until you've read page 4.

VIAL 2      **Bacteriostatic water — 10 mL.** This is what turns the powder into a liquid.  
Sterility and pH matter — don't substitute.

## GROCERY LIST • ≈ \$41

| ITEM                                | SPEC                                                                             | QTY       | WHERE                                                                                                                                          |
|-------------------------------------|----------------------------------------------------------------------------------|-----------|------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>INSULIN SYRINGES</b> <b>BUY</b>  | 1cc U100, 30G, 5/16"                                                             | 50–100 ct | Any pharmacy                                                                                                                                   |
| <b>ALCOHOL PREP PADS</b> <b>BUY</b> | Any brand, 70% isopropyl                                                         | 50–100 ct | Any pharmacy                                                                                                                                   |
| <b>SHARPS CONTAINER</b> <b>BUY</b>  | 1 quart, for needle disposal                                                     | 1×        | Any pharmacy                                                                                                                                   |
| <b>BAC WATER</b> <b>BUY</b>         | Bacteriostatic water,<br>10 mL — sterility and<br>pH matter, don't<br>substitute | 1 vial    | <b>See it</b><br><a href="https://biolinxlabs.com/products/bacteriostatic-water-10-ml">biolinxlabs.com/products/bacteriostatic-water-10-ml</a> |

Any brand works for syringes, pads and sharps — but not BAC water. Use 2 fresh syringes and 2 fresh pads every session: one to mix, one to inject.

## HOW TO STORE IT


|               |                                                         |
|---------------|---------------------------------------------------------|
| POWDER        | Freezer — lasts years                                   |
| RECONSTITUTED | Fridge — about 28 days. Write the mix date on the vial. |
| WATER         | Room temperature — check the expiration date            |



STEP 2 OF 4 · RECONSTITUTION – FROM POWDER TO LIQUID

# LET'S “FORM” YOUR PEPTIDE!

“Form” means “reconstitute”. Your product arrives as a powder — you need to liquify it before “you begin your research.” 100 units = one full syringe. **500 mg vial** → **add 300 units (3 mL)**. Every dose in this guide assumes exactly that ratio, so measure it — don't eyeball it.



## YOUR SIMPLIFIED UNITS

DOSE      FILL TO

**25 mg**    15 units

**50 mg**    30 units

**75 mg**    45 units

**100 mg**   60 units

**150 mg**   90 units

**200 mg**   120 units – 100 + 20

Change the vial, water or dose → [interactive calculator](#)

1

### Remove the plastic caps & wipe both rubber vial tops

Fresh alcohol pad on the NAD+ vial and the bac water vial.

2

### Draw 300 units of bac water

That's 3 full pulls on a 100-unit insulin syringe (3 mL total).

3

### Inject it slowly down the side of the 500 mg vial

Let it run down the glass. Don't blast the powder.

4

### Swirl gently. Never shake. Place it in the fridge!

Slow and gentle wins this one — shaking wastes peptide. Discard that syringe into the sharps container. Let it rest until completely clear (36–46 °F); use within ~28 days; never freeze it.



STEP 3 OF 4 · INJECT

# WHERE TO INJECT

Time to inject · clean, pinch, poke, done. 2× per week, in the morning.

1

## Take NAD+ out of the refrigerator and extract your dose

Your first dose is **25 mg = 15 units** on a 100-unit syringe. Wipe the vial top with a fresh alcohol pad, let it air-dry a few seconds, then draw your units.

2

## Pinch a fold of skin on the stomach + use an alcohol wipe

Two inches away from the belly button. Wipe with a fresh alcohol pad, let it air-dry, and don't touch that spot again before injecting. Rotate sites.

3

## Needle in at the base of the fold

Push the plunger slowly and steadily. Hold for a few seconds, then withdraw.

4

## That's it — you just did your first shot!

Drop the needle straight into the sharps bin. Vial goes back in the fridge.



STOMACH · PINCH A FOLD · NEEDLE AT THE BASE

### WATCH FIRST


Mixing and injecting from start to finish, on video: [professorpeptides.com](https://professorpeptides.com). Use 2 fresh syringes and 2 fresh pads every session — one to mix, one to inject.



STEP 4 OF 4 · DOSING & TIMING SCHEDULE

# NAD+ SCHEDULE (PROTOCOL)

“Schedule” means “protocol”. Everything below assumes the 500 mg vial was reconstituted with exactly 300 units (3 mL), read off a standard 100-unit insulin syringe.



## FIRST DOSE

How much, how often & when?

WHEN

**In the morning. 2× per week.**

HOW MUCH

**25 mg** to start.

Dose runs in on/off blocks with breaks between — this is **cycling**, not titration. Full schedule on page 9.

UNITS

Extract **15 units** on a 100-unit insulin syringe.

15 units = 25 mg. Exact variables: 100-unit syringe · 500 mg vial reconstituted with 300 units · 1.67 mg per unit.

HOW OFTEN

**2× per week.** Hold each step for its full block before moving up.

**CYCLE**

**1 month on, 2 months off.**

**Don't rush it — start at the lowest amount, hold each step for its full block, and only step up if the current dose is comfortable. Most people never need the top doses.**



STEP 4 OF 4 · DOCUMENT

# WHAT TO EXPECT + SIDE EFFECTS

Mostly mild, mostly early, mostly right after a dose increase. Know the three tiers before the first shot, not after.

## NORMAL / EXPECTED

First few days

- Flushing, warmth or a red face for 30–60 minutes after the shot (NAD+ widens blood vessels)
- Mild nausea, headache or fatigue that fades within a few hours
- Soreness at the injection site

## COMMON, WORTH NOTING

Manage it, don't push the dose

- Stomach cramping or a "heavy chest" feeling — more likely above 100 mg per shot; inject slowly and split the dose
- A faster heartbeat or light-headedness from the drop in blood pressure — sit down, hydrate
- Anxiety or restlessness during the first hour

## STOP & CALL A DOCTOR

Don't push through it

- Chest tightness or pressure that doesn't ease within the hour, or chest pain
- Fainting, or a racing heart that won't settle
- Signs of an allergic reaction — swelling, rash, difficulty breathing
- Vomiting or cramping that doesn't stop

**NOTE:** for this one it is **HIGHLY RECOMMENDED** to start at the lowest amount first and work up over time, as needed. NAD+ is not for anyone with a heart condition, low blood pressure, or who is pregnant — talk to a doctor before use. Never push a dose faster than your body tolerates; the side effects are dose- and speed-dependent.



---

FIRST-TIME TL;DR

# RECAP: CHECKLIST

Everything from pages 2–7, in the order it happens.

---

## BEFORE

- ☐ Vial reconstituted with exactly 300 units of bac water, resting in the fridge, mix date written on it
  - ☐ Supplies laid out: new syringe, alcohol pad, sharps container
  - ☐ Dose set at **25 mg = 15 units** — not higher, no matter what a forum says
- 

## DURING

- ☐ Inject 2× per week, in the morning — stomach, rotate the spot
  - ☐ Note the date and dose somewhere — a phone note is fine
- 

## AFTER

- ☐ Expect possible redness, mild irritation or a headache — it settles
  - ☐ Mild nausea can happen after a dose increase — sit tight, it typically passes
  - ☐ If anything feels more than mild — dizziness, real nausea, allergic reaction — stop and get medical advice
- 

## NEXT BLOCK

- ☐ Hold each step for its full block before stepping up — never skip a step (1 month on, 2 months off)
  - ☐ Only step up if the current dose is tolerated; if side effects are rough, stay another block
  - ☐ Fresh syringe, fresh pad, needle in the sharps bin — every single time
-






UNIT CHEAT-SHEET · 100-UNIT SYRINGE · 500 MG VIAL RECONSTITUTED WITH 300 UNITS

# HOW TO CYCLE (RAMP-UP & CYCLE SCHEDULE)

Assumes the 500 mg vial was mixed with exactly 300 units, so **1 unit = 1.67 mg**.

Units shown are per shot.



| WHEN                | DOSE                       | SYRINGE · PER SHOT          | NOTES                      |
|---------------------|----------------------------|-----------------------------|----------------------------|
| Month 1 · weeks 1-2 | <b>25 mg, 2× per week</b>  | <b>15 units</b>             | <a href="#">START HERE</a> |
| Month 1 · weeks 3-4 | <b>50 mg, 2× per week</b>  | <b>30 units</b>             |                            |
| Months 2-3          | <b>Off</b>                 | —                           | <a href="#">REST</a>       |
| Month 4 · weeks 1-2 | <b>100 mg, 2× per week</b> | <b>60 units</b>             |                            |
| Month 4 · weeks 3-4 | <b>200 mg, 3× per week</b> | <b>120 units — 100 + 20</b> | <a href="#">MAX DOSE</a>   |

**Rule of thumb:** don't move up a step until you know how the current dose feels. Only step up if the previous block was tolerated. Most people never need to go past the middle of the ramp.



## THE NAD+ SOURCING GUIDE

# PROFESSOR PEPS: SOURCING GUIDE (APPENDIX)

How to verify any supplier in 90 seconds. A legitimate source clears all three checks below in under two minutes — no back-and-forth required.



1

### Ask for the COA

A batch-specific Certificate of Analysis — not a generic PDF reused across batches — showing purity  $\geq 98\%$  via HPLC/MS.

2

### Match the lot number on the vial

It must match the lot number on the COA. If they don't match, walk away.

3

### Check the track record

Searchable reviews or forum mentions older than roughly six months. Brand-new storefronts with no history are the highest-risk category.

DON'T WORRY — WE'VE ALREADY DONE THE VETTING

BioLinx Labs clears the list: third-party HPLC testing, a 99%+ purity guarantee and a COA included with every order. [biolinxlabs.com/best-sellers](https://biolinxlabs.com/best-sellers)

[biolinxlabs.com/best-sellers](https://biolinxlabs.com/best-sellers)

Educational and research purposes only — not medical advice. Professor Peptides does not sell peptides. Products referenced are sold for laboratory research use only and are not for human consumption. Want to learn more? Join the Professor Peptides community — 30 days free.

**STAY CURIOUS,  
PROFESSOR PEPTIDES**

NAD+ Research Hub · companion to the email series