



SEMEX RESEARCH HUB · RESEARCH USE ONLY

THINK SHARPER. FOCUS LONGER.

FOR MEN & WOMEN. Think sharper, focus better, and recover from stress and neurological injury.

Get Semax 30 mg at BioLinx

Read the guide first

Also shown to help with: anxiety · depression · brain fog · adhd · stress resilience
Daily – 5 days on, 2 days off · before activities that need focus · Difficulty: beginner

WHAT'S INSIDE

- | | | | |
|---|---|-------|--------|
| 1 | Shopping list — how many vials to buy | GET | p.2 |
| 2 | Quick summary of your lab items | GET | p.3 |
| 3 | Forming your peptide — “reconstitution” | PREP | p.4 |
| 4 | How to inject | DOSE | p.5 |
| 5 | Semex dosing schedule | DOSE | p.6 |
| 6 | What to expect + side effects | AFTER | p.7 |
| 7 | Recap checklist | AFTER | p.8 |
| + | Appendix: ramp-up schedule + sourcing guide | BONUS | p.9–10 |

RESEARCH SHOWED THAT...

3× memory protein
in the brain after a single dose.

Brain Res, 2006

Rats given Semax learned faster and were less afraid during the tests.

Dolotov et al., Brain Res, 2006



STEP 1 OF 4 · WHAT TO BUY

SHOPPING LIST

Easy-to-follow shopping list to safely practice Semax. Match your commitment level — don't skip anything.



PEPTIDE · ONE ORDER

SEMAX 30 MG

Think sharper, focus better, and recover from stress and neurological injury. Common starting dose: **0.2 mg**, daily — 5 days on, 2 days off, before activities that need focus.

Get Semax 30 mg at BioLinx

biolinxlabs.com/products/semex-30-mg

HOW MANY VIALS TO BUY · BEST VALUE BY GOAL

GOAL	VIAL	QTY	WHERE
1 MONTH	30 mg vial	1×	<u>See it</u> biolinxlabs.com/products/semex-30-mg
FULL CYCLE RECOMMENDED	30 mg vial	2×	<u>See it</u> biolinxlabs.com/products/semex-30-mg

MATERIALS ≈ \$41 ON TOP — FULL LIST ON THE NEXT PAGE



PRODUCTS ARRIVED?

QUICK SUMMARY OF YOUR LAB ITEMS:

TWO VIALS IN THE BOX • SEMAX 30 MG + BACTERIOSTATIC WATER 10 ML

VIAL 1 **Semax 30 mg.** Arrives as a powder. Don't touch it until you've read page 4.

VIAL 2 **Bacteriostatic water — 10 mL.** This is what turns the powder into a liquid.
Sterility and pH matter — don't substitute.

GROCERY LIST • ≈ \$41

ITEM	SPEC	QTY	WHERE
INSULIN SYRINGES BUY	1cc U100, 30G, 5/16"	50–100 ct	Any pharmacy
ALCOHOL PREP PADS BUY	Any brand, 70% isopropyl	50–100 ct	Any pharmacy
SHARPS CONTAINER BUY	1 quart, for needle disposal	1×	Any pharmacy
BAC WATER BUY	Bacteriostatic water, 10 mL — sterility and pH matter, don't substitute	1 vial	See it biolinxlabs.com/products/bacteriostatic-water-10-ml

Any brand works for syringes, pads and sharps — but not BAC water. Use 2 fresh syringes and 2 fresh pads every session: one to mix, one to inject.

HOW TO STORE IT


POWDER	Freezer — lasts years
RECONSTITUTED	Fridge — about 28 days. Write the mix date on the vial.
WATER	Room temperature — check the expiration date



STEP 2 OF 4 · RECONSTITUTION – FROM POWDER TO LIQUID

LET'S “FORM” YOUR PEPTIDE!

“Form” means “reconstitute”. Your product arrives as a powder — you need to liquify it before “you begin your research.” 100 units = one full syringe. **30 mg vial** → **add 300 units (3 mL)**. Every dose in this guide assumes exactly that ratio, so measure it — don't eyeball it.



1 Remove the plastic caps & wipe both rubber vial tops

Fresh alcohol pad on the Semax vial and the bac water vial.

2 Draw 300 units of bac water

That's 3 full pulls on a 100-unit insulin syringe (3 mL total).

3 Inject it slowly down the side of the 30 mg vial

Let it run down the glass. Don't blast the powder.

4 Swirl gently. Never shake. Place it in the fridge!

Slow and gentle wins this one — shaking wastes peptide. Discard that syringe into the sharps container. Let it rest until completely clear (36–46 °F); use within ~28 days; never freeze it.

YOUR SIMPLIFIED UNITS

DOSE	FILL TO
------	---------

0.2 mg	2 units
--------	---------

0.4 mg	4 units
--------	---------

0.5 mg	5 units
--------	---------

0.6 mg	6 units
--------	---------

0.8 mg	8 units
--------	---------

Change the vial, water or dose →
[interactive calculator](#)

STEP 3 OF 4 · INJECT

WHERE TO INJECT

Time to inject · clean, pinch, poke, done. Daily — 5 days on, 2 days off, before activities that need focus.

1 Take Semax out of the refrigerator and extract your dose

Your first dose is **0.2 mg = 2 units** on a 100-unit syringe. Wipe the vial top with a fresh alcohol pad, let it air-dry a few seconds, then draw your units.

2 Pinch a fold of skin on the stomach + use an alcohol wipe

Two inches away from the belly button. Wipe with a fresh alcohol pad, let it air-dry, and don't touch that spot again before injecting. Rotate sites.

3 Needle in at the base of the fold

Push the plunger slowly and steadily. Hold for a few seconds, then withdraw.

4 That's it — you just did your first shot!

Drop the needle straight into the sharps bin. Vial goes back in the fridge.



STOMACH · PINCH A FOLD · NEEDLE AT THE BASE

WATCH FIRST


Mixing and injecting from start to finish, on video: professorpeptides.com. Use 2 fresh syringes and 2 fresh pads every session — one to mix, one to inject.



STEP 4 OF 4 · DOSING & TIMING SCHEDULE

SEMAX SCHEDULE (PROTOCOL)

“Schedule” means “protocol”. Everything below assumes the 30 mg vial was reconstituted with exactly 300 units (3 mL), read off a standard 100-unit insulin syringe.



FIRST DOSE

How much, how often & when?

WHEN	Before activities that need focus. Daily — 5 days on, 2 days off.
HOW MUCH	0.2 mg to start. Dose runs in on/off blocks with breaks between — this is cycling , not titration. Full schedule on page 9.
UNITS	Extract 2 units on a 100-unit insulin syringe. 2 units = 0.2 mg. Exact variables: 100-unit syringe · 30 mg vial reconstituted with 300 units · 0.1 mg per unit.
HOW OFTEN	Daily — 5 days on, 2 days off. Hold each step for its full block before moving up.
CYCLE	1 month on, 1 month off.

Don't rush it — start at the lowest amount, hold each step for its full block, and only step up if the current dose is comfortable. Most people never need the top doses.



STEP 4 OF 4 · DOCUMENT

WHAT TO EXPECT + SIDE EFFECTS

Mostly mild, mostly early, mostly right after a dose increase. Know the three tiers before the first shot, not after.

NORMAL / EXPECTED

First few days

- Mild headache or a "wired" feeling for a few hours after dosing
- Mild fatigue after the first doses as your body adjusts
- Slight redness at the injection site

COMMON, WORTH NOTING

Manage it, don't push the dose

- Feeling jittery or anxious — more common at higher doses; drop back a step
- Trouble falling asleep if dosed late in the day — keep it to mornings
- Mild nausea or dizziness

STOP & CALL A DOCTOR

Don't push through it

- Anxiety or agitation that doesn't settle
- Persistent insomnia, or a racing heart
- Signs of an allergic reaction — swelling, rash, difficulty breathing

NOTE: for this one it is **HIGHLY RECOMMENDED** to start at the lowest amount first and work up over time, as needed. Human data comes from short Russian clinical courses; long-term safety is unknown. Not for anyone with an anxiety disorder that isn't controlled, or who is pregnant or breastfeeding — talk to a doctor before use.



FIRST-TIME TL;DR

RECAP: CHECKLIST

Everything from pages 2–7, in the order it happens.

BEFORE

- ☐ Vial reconstituted with exactly 300 units of bac water, resting in the fridge, mix date written on it
- ☐ Supplies laid out: new syringe, alcohol pad, sharps container
- ☐ Dose set at **0.2 mg = 2 units** — not higher, no matter what a forum says

DURING

- ☐ Inject daily — 5 days on, 2 days off, before activities that need focus — stomach, rotate the spot
- ☐ Note the date and dose somewhere — a phone note is fine

AFTER

- ☐ Expect possible redness, mild irritation or a headache — it settles
- ☐ Mild nausea can happen after a dose increase — sit tight, it typically passes
- ☐ If anything feels more than mild — dizziness, real nausea, allergic reaction — stop and get medical advice

NEXT BLOCK

- ☐ Hold each step for its full block before stepping up — never skip a step (1 month on, 1 month off)
 - ☐ Only step up if the current dose is tolerated; if side effects are rough, stay another block
 - ☐ Fresh syringe, fresh pad, needle in the sharps bin — every single time
-




UNIT CHEAT-SHEET · 100-UNIT SYRINGE · 30 MG VIAL RECONSTITUTED WITH 300 UNITS

HOW TO CYCLE (RAMP-UP & CYCLE SCHEDULE)

Assumes the 30 mg vial was mixed with exactly 300 units, so **1 unit = 0.1 mg**.

Units shown are per shot.

START · 2 UNITS = 0.2 MG



WHEN	DOSE	SYRINGE · PER SHOT	NOTES
Month 1	0.2 mg/day · 1 mg/week	2 units	START HERE
Month 2	Off	—	REST
Month 3	0.4 mg/day · 2 mg/week	4 units	
Month 4	Off	—	REST
Month 5	0.5 mg/day · 2.5 mg/week	5 units	
Month 6	Off	—	REST
Month 7	0.6 mg/day · 3 mg/week	6 units	
Month 8	Off	—	REST
Month 9	0.8 mg/day · 4 mg/week	8 units	MAX DOSE

Rule of thumb: don't move up a step until you know how the current dose feels. Only step up if the previous block was tolerated. Most people never need to go past the middle of the ramp.



THE SEMAX SOURCING GUIDE

PROFESSOR PEPS: SOURCING GUIDE (APPENDIX)

How to verify any supplier in 90 seconds. A legitimate source clears all three checks below in under two minutes — no back-and-forth required.



1

Ask for the COA

A batch-specific Certificate of Analysis — not a generic PDF reused across batches — showing purity $\geq 98\%$ via HPLC/MS.

2

Match the lot number on the vial

It must match the lot number on the COA. If they don't match, walk away.

3

Check the track record

Searchable reviews or forum mentions older than roughly six months. Brand-new storefronts with no history are the highest-risk category.

DON'T WORRY — WE'VE ALREADY DONE THE VETTING

BioLinx Labs clears the list: third-party HPLC testing, a 99%+ purity guarantee and a COA included with every order. biolinxlabs.com/best-sellers

biolinxlabs.com/best-sellers

Educational and research purposes only — not medical advice. Professor Peptides does not sell peptides. Products referenced are sold for laboratory research use only and are not for human consumption. Want to learn more? Join the Professor Peptides community — 30 days free.

STAY CURIOUS, PROFESSOR PEPTIDES

Semax Research Hub · companion to the email series