



SS-31 RESEARCH HUB · RESEARCH USE ONLY

POWER THE CELLS. PROTECT THE HEART.

FOR MEN & WOMEN. Boost mitochondrial energy, slow down aging, and protect your heart and brain.

Get SS-31 vial at BioLinx

Read the guide first

Also shown to help with: chronic fatigue · heart failure · macular degeneration · muscle loss
Daily for 2 weeks · morning · Difficulty: beginner

WHAT'S INSIDE

- | | | | |
|---|---|-------|--------|
| 1 | Shopping list — how many vials to buy | GET | p.2 |
| 2 | Quick summary of your lab items | GET | p.3 |
| 3 | Forming your peptide — “reconstitution” | PREP | p.4 |
| 4 | How to inject | DOSE | p.5 |
| 5 | SS-31 dosing schedule | DOSE | p.6 |
| 6 | What to expect + side effects | AFTER | p.7 |
| 7 | Recap checklist | AFTER | p.8 |
| + | Appendix: ramp-up schedule + sourcing guide | BONUS | p.9–10 |

RESEARCH SHOWED THAT...

8 wks muscle

preserved
age-related muscle loss
halted.

Free Radic Biol Med, 2019

People who couldn't exercise
without exhaustion started
going farther at every dose.

Karaa et al., Neurology,


2018



STEP 1 OF 4 · WHAT TO BUY

SHOPPING LIST

Easy-to-follow shopping list to safely practice SS-31. Match your commitment level — don't skip anything.



VIAL 1 · PEPTIDE

PEPTIDE · ONE ORDER

SS-31 VIAL

Boost mitochondrial energy, slow down aging, and protect your heart and brain. Common starting dose: **0.1 mg**, daily for 2 weeks, morning.

Get SS-31 vial at BioLinx

biolinxlabs.com/best-sellers

HOW MANY VIALS TO BUY · BEST VALUE BY GOAL

GOAL

VIAL

QTY

WHERE

FULL CYCLE

RECOMMENDED

10 mg vial

1×

Best sellers

not listed yet ·
biolinxlabs.com/best-sellers

MATERIALS ≈ \$41 ON TOP — FULL LIST ON THE NEXT PAGE



PRODUCTS ARRIVED?

QUICK SUMMARY OF YOUR LAB ITEMS:

TWO VIALS IN THE BOX • SS-31 VIAL + BACTERIOSTATIC WATER 10 ML

VIAL 1 **SS-31 vial.** Arrives as a powder. Don't touch it until you've read page 4.

VIAL 2 **Bacteriostatic water — 10 mL.** This is what turns the powder into a liquid. Sterility and pH matter — don't substitute.

GROCERY LIST • ≈ \$41

ITEM	SPEC	QTY	WHERE
INSULIN SYRINGES BUY	1cc U100, 30G, 5/16"	50–100 ct	Any pharmacy
ALCOHOL PREP PADS BUY	Any brand, 70% isopropyl	50–100 ct	Any pharmacy
SHARPS CONTAINER BUY	1 quart, for needle disposal	1×	Any pharmacy
BAC WATER BUY	Bacteriostatic water, 10 mL — sterility and pH matter, don't substitute	1 vial	<u>See it</u> biolinxlabs.com/products/bacteriostatic-water-10-ml

Any brand works for syringes, pads and sharps — but not BAC water. Use 2 fresh syringes and 2 fresh pads every session: one to mix, one to inject.

HOW TO STORE IT


POWDER	Freezer — lasts years
RECONSTITUTED	Fridge — about 28 days. Write the mix date on the vial.
WATER	Room temperature — check the expiration date



STEP 2 OF 4 · RECONSTITUTION – FROM POWDER TO LIQUID

LET'S “FORM” YOUR PEPTIDE!

“Form” means “reconstitute”. Your product arrives as a powder — you need to liquify it before “you begin your research.” 100 units = one full syringe. **10 mg vial** → **add 300 units (3 mL)**. Every dose in this guide assumes exactly that ratio, so measure it — don't eyeball it.



1 Remove the plastic caps & wipe both rubber vial tops

Fresh alcohol pad on the SS-31 vial and the bac water vial.

2 Draw 300 units of bac water

That's 3 full pulls on a 100-unit insulin syringe (3 mL total).

3 Inject it slowly down the side of the 10 mg vial

Let it run down the glass. Don't blast the powder.

4 Swirl gently. Never shake. Place it in the fridge!

Slow and gentle wins this one — shaking wastes peptide. Discard that syringe into the sharps container. Let it rest until completely clear (36–46 °F); use within ~28 days; never freeze it.

YOUR SIMPLIFIED UNITS

DOSE	FILL TO
0.05 mg	1.5 units
0.075 mg	2.25 units
0.1 mg	3 units
0.125 mg	3.75 units
0.15 mg	4.5 units
0.2 mg	6 units



STEP 3 OF 4 · INJECT

WHERE TO INJECT

Time to inject · clean, pinch, poke, done. Daily for 2 weeks, morning.

1

Take SS-31 out of the refrigerator and extract your dose

Your first dose is **0.1 mg = 3 units** on a 100-unit syringe. Wipe the vial top with a fresh alcohol pad, let it air-dry a few seconds, then draw your units.

2

Pinch a fold of skin on the stomach + use an alcohol wipe

Two inches away from the belly button. Wipe with a fresh alcohol pad, let it air-dry, and don't touch that spot again before injecting. Rotate sites.

3

Needle in at the base of the fold

Push the plunger slowly and steadily. Hold for a few seconds, then withdraw.

4

That's it — you just did your first shot!

Drop the needle straight into the sharps bin. Vial goes back in the fridge.



STOMACH · PINCH A FOLD · NEEDLE AT THE BASE

WATCH FIRST


Mixing and injecting from start to finish, on video: professorpeptides.com. Use 2 fresh syringes and 2 fresh pads every session — one to mix, one to inject.



STEP 4 OF 4 · DOSING & TIMING SCHEDULE

SS-31 SCHEDULE (PROTOCOL)

“Schedule” means “protocol”. Everything below assumes the 10 mg vial was reconstituted with exactly 300 units (3 mL), read off a standard 100-unit insulin syringe.



FIRST DOSE

How much, how often & when?

WHEN

Morning. Daily for 2 weeks.

HOW MUCH

0.1 mg to start.

Dose runs in on/off blocks with breaks between — this is **cycling**, not titration. Full schedule on page 9.

UNITS

Extract **3 units** on a 100-unit insulin syringe.

3 units = 0.1 mg. Exact variables: 100-unit syringe · 10 mg vial reconstituted with 300 units · 0.03 mg per unit.

HOW OFTEN

Daily for 2 weeks. Hold each step for its full block before moving up.

CYCLE

Do not run longer than 2 weeks per cycle · 12 weeks off between cycles.

Do not run longer than 2 weeks per cycle. Don't rush it — start at the lowest amount, hold each step for its full block, and only step up if the current dose is comfortable. Most people never need the top doses.



STEP 4 OF 4 · DOCUMENT

WHAT TO EXPECT + SIDE EFFECTS

Mostly mild, mostly early, mostly right after a dose increase. Know the three tiers before the first shot, not after.

NORMAL / EXPECTED

First few days

- Redness, swelling, itching or bruising at the injection site — reported by most subjects in the elamipretide trials, usually gone within 4 hours
- Mild headache
- Mild fatigue in the first days

COMMON, WORTH NOTING

Manage it, don't push the dose

- Injection-site hives or hardness that lasts days — rotate sites, keep cycles to 2 weeks
- Dizziness that passes within the hour
- Injection-site reactions that linger up to 2 weeks after a cycle

STOP & CALL A DOCTOR

Don't push through it

- Signs of an allergic reaction beyond the injection site — swelling, rash, difficulty breathing
- Dizziness that doesn't pass, or fainting
- Any injection site that becomes hot, very painful or oozes

NOTE: for this one it is **HIGHLY RECOMMENDED** to start at the lowest amount first and work up over time, as needed. Over 500 people have received elamipretide in trials at doses far above these; the main issue is injection-site reactions. Keep each cycle to 2 weeks. Talk to a doctor before use if you are pregnant or have a heart or kidney condition.



FIRST-TIME TL;DR

RECAP: CHECKLIST

Everything from pages 2–7, in the order it happens.

BEFORE

- ☐ Vial reconstituted with exactly 300 units of bac water, resting in the fridge, mix date written on it
- ☐ Supplies laid out: new syringe, alcohol pad, sharps container
- ☐ Dose set at **0.1 mg = 3 units** — not higher, no matter what a forum says

DURING

- ☐ Inject daily for 2 weeks, morning — stomach, rotate the spot
- ☐ Note the date and dose somewhere — a phone note is fine

AFTER

- ☐ Expect possible redness, mild irritation or a headache — it settles
- ☐ Mild nausea can happen after a dose increase — sit tight, it typically passes
- ☐ If anything feels more than mild — dizziness, real nausea, allergic reaction — stop and get medical advice

NEXT BLOCK


- ☐ Hold each step for its full block before stepping up — never skip a step (do not run longer than 2 weeks per cycle · 12 weeks off between cycles)
 - ☐ Only step up if the current dose is tolerated; if side effects are rough, stay another block
 - ☐ Fresh syringe, fresh pad, needle in the sharps bin — every single time
-

UNIT CHEAT-SHEET · 100-UNIT SYRINGE · 10 MG VIAL RECONSTITUTED WITH 300 UNITS

HOW TO CYCLE (RAMP-UP & CYCLE SCHEDULE)

Assumes the 10 mg vial was mixed with exactly 300 units, so **1 unit = 0.03 mg**.

Units shown are per shot.



WHEN	DOSE	SYRINGE · PER SHOT	NOTES
Cycle 1 · weeks 1-2	0.1 mg per day	3 units	START HERE
Off	12 weeks	—	REST
Cycle 2 · weeks 1-2	0.15 mg per day	4.5 units	
Off	12 weeks	—	REST
Cycle 3 · weeks 1-2	0.2 mg per day	6 units	MAX DOSE
Off	12 weeks	—	REST

Rule of thumb: don't move up a step until you know how the current dose feels. Only step up if the previous block was tolerated — Do not run longer than 2 weeks per cycle. Most people never need to go past the middle of the ramp.



THE SS-31 SOURCING GUIDE

PROFESSOR PEPS: SOURCING GUIDE (APPENDIX)

How to verify any supplier in 90 seconds. A legitimate source clears all three checks below in under two minutes — no back-and-forth required.



1

Ask for the COA

A batch-specific Certificate of Analysis — not a generic PDF reused across batches — showing purity $\geq 98\%$ via HPLC/MS.

2

Match the lot number on the vial

It must match the lot number on the COA. If they don't match, walk away.

3

Check the track record

Searchable reviews or forum mentions older than roughly six months. Brand-new storefronts with no history are the highest-risk category.

DON'T WORRY — WE'VE ALREADY DONE THE VETTING

BioLinx Labs clears the list: third-party HPLC testing, a 99%+ purity guarantee and a COA included with every order. biolinxlabs.com/best-sellers

biolinxlabs.com/best-sellers

Educational and research purposes only — not medical advice. Professor Peptides does not sell peptides. Products referenced are sold for laboratory research use only and are not for human consumption. Want to learn more? Join the Professor Peptides community — 30 days free.

**STAY CURIOUS,
PROFESSOR PEPTIDES**

SS-31 Research Hub · companion to the email series