



BPC-157 + TB-500 BLEND RESEARCH HUB · RESEARCH USE ONLY

HEAL INJURIES. REBUILD TISSUE.

FOR MEN & WOMEN. Heal injuries faster, reduce inflammation, and repair tendons, ligaments, and muscles.

[Get BPC-157 / TB-500 5/5 mg at BioLinx](#)[Read the guide first](#)

Also shown to help with: joint pain · tendinitis · lower back strain · osteoarthritis · plantar fasciitis

Once daily · time of day doesn't matter · Difficulty: beginner

WHAT'S INSIDE

- 1** Shopping list — how many vials to buy GET p.2
- 2** Quick summary of your lab items GET p.3
- 3** Forming your peptide — “reconstitution” PREP p.4
- 4** How to inject DOSE p.5
- 5** BPC-157 + TB-500 Blend dosing schedule DOSE p.6
- 6** What to expect + side effects AFTER p.7
- 7** Recap checklist AFTER p.8
- +** Appendix: ramp-up schedule + sourcing guide BONUS p.9–10

RESEARCH SHOWED THAT...

42% faster

healing
wounds treated with thymosin
beta-4, by day 4.

J Invest Dermatol, 1999

Heart cells that stopped
functioning after coronary
ligation survived longer when
treated with thymosin beta-4.

Bock-Marquette et al.,
Nature, 2004



STEP 1 OF 4 · WHAT TO BUY

SHOPPING LIST

Easy-to-follow shopping list to safely practice BPC-157 + TB-500 Blend. Match your commitment level — don't skip anything.



PEPTIDE · ONE ORDER

BPC-157 / TB-500 5/5 MG

Heal injuries faster, reduce inflammation, and repair tendons, ligaments, and muscles. Common starting dose: **0.25 mg of each**, once daily, time of day doesn't matter.

Get BPC-157 / TB-500 5/5 mg at BioLinX

biolinxlabs.com/products/bpc-157-tb-500-blend-10-mg

HOW MANY VIALS TO BUY · BEST VALUE BY GOAL

GOAL	VIAL	QTY	WHERE
1 MONTH	5/5 mg vial	2×	<u>See it</u> biolinxlabs.com/products/bpc-157-tb-500-blend-10-mg
3 MONTHS	10/10 mg vial ×2 + 5/5 mg vial ×1	3 vials	<u>See it</u> biolinxlabs.com/products/bpc-157-tb-500-blend-20-mg
FULL CYCLE	10/10 mg vial	7×	<u>See it</u> biolinxlabs.com/products/bpc-157-tb-500-blend-20-mg

MATERIALS ≈ \$41 ON TOP — FULL LIST ON THE NEXT PAGE



PRODUCTS ARRIVED?

QUICK SUMMARY OF YOUR LAB ITEMS:

TWO VIALS IN THE BOX • BPC-157 / TB-500 5/5 MG + BACTERIOSTATIC WATER
10 ML

VIAL 1 **BPC-157 / TB-500 5/5 mg.** Arrives as a powder. Don't touch it until you've read page 4.

VIAL 2 **Bacteriostatic water — 10 mL.** This is what turns the powder into a liquid. Sterility and pH matter — don't substitute.

GROCERY LIST • ≈ \$41

ITEM	SPEC	QTY	WHERE
INSULIN SYRINGES BUY	1cc U100, 30G, 5/16"	50–100 ct	Any pharmacy
ALCOHOL PREP PADS BUY	Any brand, 70% isopropyl	50–100 ct	Any pharmacy
SHARPS CONTAINER BUY	1 quart, for needle disposal	1×	Any pharmacy
BAC WATER BUY	Bacteriostatic water, 10 mL — sterility and pH matter, don't substitute	1 vial	<u>See it</u> biolinxlabs.com/products/bacteriostatic-water-10-ml

Any brand works for syringes, pads and sharps — but not BAC water. Use 2 fresh syringes and 2 fresh pads every session: one to mix, one to inject.

HOW TO STORE IT


POWDER	Freezer — lasts years
RECONSTITUTED	Fridge — about 28 days. Write the mix date on the vial.
WATER	Room temperature — check the expiration date



STEP 2 OF 4 · RECONSTITUTION – FROM POWDER TO LIQUID

LET'S “FORM” YOUR PEPTIDE!

“Form” means “reconstitute”. Your product arrives as a powder — you need to liquify it before “you begin your research.” 100 units = one full syringe. **5/5 mg vial → add 100 units (1 mL) · 10/10 mg vial → add 200 units (2 mL)**. Every dose in this guide assumes exactly that ratio, so measure it — don't eyeball it.



YOUR SIMPLIFIED UNITS

DOSE	FILL TO
0.25 mg of each	5 units
0.5 mg of each	10 units
0.75 mg of each	15 units
1 mg of each	20 units

1

Remove the plastic caps & wipe both rubber vial tops

Fresh alcohol pad on the BPC + TB vial and the bac water vial.

2

Draw 100 units of bac water

One full pull on a 100-unit insulin syringe (1 mL total).

3

Inject it slowly down the side of the 5/5 mg vial

Let it run down the glass. Don't blast the powder.

4

Swirl gently. Never shake. Place it in the fridge!

Slow and gentle wins this one — shaking wastes peptide. Discard that syringe into the sharps container. Let it rest until completely clear (36–46 °F); use within ~28 days; never freeze it.



STEP 3 OF 4 · INJECT

WHERE TO INJECT

Time to inject · clean, pinch, poke, done. Once daily, time of day doesn't matter.

- 1 Take BPC + TB out of the refrigerator and extract your dose**
Your first dose is **0.25 mg of each = 5 units** on a 100-unit syringe. Wipe the vial top with a fresh alcohol pad, let it air-dry a few seconds, then draw your units.
- 2 Pinch a fold of skin on the stomach + use an alcohol wipe**
Two inches away from the belly button. Wipe with a fresh alcohol pad, let it air-dry, and don't touch that spot again before injecting. Rotate sites.
- 3 Needle in at the base of the fold**
Push the plunger slowly and steadily. Hold for a few seconds, then withdraw.
- 4 That's it — you just did your first shot!**
Drop the needle straight into the sharps bin. Vial goes back in the fridge.



STOMACH · PINCH A FOLD · NEEDLE AT THE BASE

WATCH FIRST


Mixing and injecting from start to finish, on video: professorpeptides.com. Use 2 fresh syringes and 2 fresh pads every session — one to mix, one to inject.



STEP 4 OF 4 · DOSING & TIMING SCHEDULE

BPC-157 + TB-500 BLEND SCHEDULE (PROTOCOL)

“Schedule” means “protocol”. Everything below assumes the 5/5 mg vial was reconstituted with exactly 100 units (1 mL), read off a standard 100-unit insulin syringe.



FIRST DOSE

How much, how often & when?

WHEN

Time of day doesn't matter. Once daily.

HOW MUCH

0.25 mg of each to start.

Dose runs in on/off blocks with breaks between — this is **cycling**, not titration. Full schedule on page 9.

UNITS

Extract **5 units** on a 100-unit insulin syringe.

5 units = 0.25 mg of each. Exact variables: 100-unit syringe · 5/5 mg vial reconstituted with 100 units · 0.05 mg per unit.

HOW OFTEN

Once daily. Hold each step for its full block before moving up.

CYCLE

1 month on, 1 month off.

Don't rush it — start at the lowest amount, hold each step for its full block, and only step up if the current dose is comfortable. Most people never need the top doses.



STEP 4 OF 4 · DOCUMENT

WHAT TO EXPECT + SIDE EFFECTS

Mostly mild, mostly early, mostly right after a dose increase. Know the three tiers before the first shot, not after.

NORMAL / EXPECTED

First few days

- Redness, mild swelling or a sting at the injection site — the most common effect of both peptides
- Mild headache 1–4 hours after a shot (TB-500 widens blood vessels)
- Slight fatigue during the first week

COMMON, WORTH NOTING

Manage it, don't push the dose

- Mild nausea when the daily dose goes above the starting range
- Brief dizziness or light-headedness early on
- Hot or cold flushes that pass within the hour

STOP & CALL A DOCTOR

Don't push through it

- Dizziness that doesn't pass, or fainting
- Persistent nausea or vomiting
- Signs of an allergic reaction — swelling, rash, difficulty breathing
- Injection site that becomes hot, hard, very painful or oozes

NOTE: for this one it is **HIGHLY RECOMMENDED** to start at the lowest amount first and work up over time, as needed. Neither peptide has long-term human safety data. Not for anyone with an active cancer or tumour history, or who is pregnant or breastfeeding — talk to a doctor before use.



FIRST-TIME TL;DR

RECAP: CHECKLIST

Everything from pages 2–7, in the order it happens.

BEFORE

- ☐ Vial reconstituted with exactly 100 units of bac water, resting in the fridge, mix date written on it
 - ☐ Supplies laid out: new syringe, alcohol pad, sharps container
 - ☐ Dose set at **0.25 mg of each = 5 units** — not higher, no matter what a forum says
-

DURING

- ☐ Inject once daily, time of day doesn't matter — stomach, rotate the spot
 - ☐ Note the date and dose somewhere — a phone note is fine
-

AFTER

- ☐ Expect possible redness, mild irritation or a headache — it settles
 - ☐ Mild nausea can happen after a dose increase — sit tight, it typically passes
 - ☐ If anything feels more than mild — dizziness, real nausea, allergic reaction — stop and get medical advice
-

NEXT BLOCK


- ☐ Hold each step for its full block before stepping up — never skip a step (1 month on, 1 month off)
 - ☐ Only step up if the current dose is tolerated; if side effects are rough, stay another block
 - ☐ Fresh syringe, fresh pad, needle in the sharps bin — every single time
-

UNIT CHEAT-SHEET · 100-UNIT SYRINGE · 5/5 MG VIAL RECONSTITUTED WITH 100 UNITS

HOW TO CYCLE (RAMP-UP & CYCLE SCHEDULE)

Assumes the 5/5 mg vial was mixed with exactly 100 units, so **1 unit = 0.05 mg**.

Units shown are per shot.



WHEN	DOSE	SYRINGE · PER SHOT	NOTES
Month 1	0.25 mg of each/day · 1.75 mg/week	5 units	START HERE
Month 2	Off	—	REST
Month 3	0.5 mg of each/day · 3.5 mg/week	10 units	
Month 4	Off	—	REST
Month 5	0.75 mg of each/day · 5.25 mg/week	15 units	
Month 6	Off	—	REST
Month 7	1 mg of each/day · 7 mg/week	20 units	MAX DOSE
Month 8	Off	—	REST

Rule of thumb: don't move up a step until you know how the current dose feels. Only step up if the previous block was tolerated. Most people never need to go past the middle of the ramp.



THE BPC + TB SOURCING GUIDE

PROFESSOR PEPS: SOURCING GUIDE (APPENDIX)

How to verify any supplier in 90 seconds. A legitimate source clears all three checks below in under two minutes — no back-and-forth required.



1

Ask for the COA

A batch-specific Certificate of Analysis — not a generic PDF reused across batches — showing purity $\geq 98\%$ via HPLC/MS.

2

Match the lot number on the vial

It must match the lot number on the COA. If they don't match, walk away.

3

Check the track record

Searchable reviews or forum mentions older than roughly six months. Brand-new storefronts with no history are the highest-risk category.

DON'T WORRY — WE'VE ALREADY DONE THE VETTING

BioLinx Labs clears the list: third-party HPLC testing, a 99%+ purity guarantee and a COA included with every order. biolinxlabs.com/best-sellers

biolinxlabs.com/best-sellers

Educational and research purposes only — not medical advice. Professor Peptides does not sell peptides. Products referenced are sold for laboratory research use only and are not for human consumption. Want to learn more? Join the Professor Peptides community — 30 days free.

BPC-157 + TB-500 Blend Research Hub · companion to the email series

**STAY CURIOUS,
PROFESSOR PEPTIDES**