

# STRENGTH *of* BALANCE

## Strong RA therapy with a proven safety profile<sup>1-5,\*</sup>

JYSELECA is indicated for the treatment of moderate to severe active rheumatoid arthritis in adult patients who have responded inadequately to, or who are intolerant to one or more disease-modifying antirheumatic drugs (DMARDs). JYSELECA may be used as monotherapy or in combination with methotrexate (MTX).<sup>1</sup>

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the [national reporting systems](#) and to [DrugSafety.Global@glpg.com](mailto:DrugSafety.Global@glpg.com).

EXPLORE HOW JYSELECA MAY HELP YOUR PATIENTS



RA, rheumatoid arthritis.  
\* The most frequently reported adverse reactions were nausea (3.5%), upper respiratory tract infection (3.3%), urinary tract infection (1.7%), and dizziness (1.2%).<sup>1</sup>  
**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.  
Materials have been prepared for an international audience and are intended for healthcare professionals, excluding those from France. JYSELECA is approved but not reimbursed in Denmark.  
Data are correct as of 01/06/2022.





# Galapagos at EULAR 2022

We look forward to seeing you at the following events

## INDUSTRY CONSULT-THE-EXPERT SESSION

JUNE  
01  
WED

15:45 – 16:15 CEST

### Patient-centred care: cutting through the jargon

**Dr Lars Erik Kristensen** The Parker Institute, Copenhagen University Hospital, Copenhagen, Denmark

#### Learning objectives

- Review the **unmet needs** experienced by patients with RA and the importance of a comprehensive approach to care
- Explore the role of **JAK inhibitors** in addressing unmet needs for patients with RA
- Consider how to facilitate a **comprehensive approach to patient care** and discuss any potential barriers to clinical implementation
- Discuss how to integrate concrete steps towards **shared decision-making** in RA

## INDUSTRY SYMPOSIUM

JUNE  
02  
THURS

17:30 – 18:45 CEST

### Evolving patient care: can JAK inhibitors meet patient and physician expectations for RA treatment?

**Prof. Roberto Caporali** University of Milan, Milan, Italy  
**Prof. Bruno Fautrel** Pitié-Salpêtrière Hospital, Paris, France  
**Dr James Galloway** King's College London, London, UK

#### Learning objectives

- Explore **efficacy and safety goals** for patients and physicians in the treatment of RA
- Explore the **efficacy and safety profiles of JAK inhibitors**, including data from the filgotinib clinical trial programme
- Review the importance of a **comprehensive management approach** for patients with RA
- Discuss how to optimally **align patient and physician treatment goals** to achieve favourable outcomes in RA



RA, rheumatoid arthritis.  
\* The most frequently reported adverse events.  
**Reference: 1.** JYSELECA (filgotinib) clinical trial programme.  
Materials have been prepared for the purpose of this presentation.  
Data are correct as of 01/06/2022.



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI







# Evaluated across a comprehensive clinical trial programme

A solid foundation of clinical evidence in RA<sup>1-5</sup>

3

Phase 3 trials<sup>1</sup>  
FINCH 1-3

**FINCH 1-3<sup>1</sup>**  
Three Phase 3, randomised, double-blind, placebo-controlled studies in patients with moderate to severe active RA.

**DARWIN 1 and 2<sup>2,3</sup>**  
Two Phase 2b, randomised, double-blind, placebo-controlled, dose-ranging studies in patients with moderate to severe active RA.

2

Phase 2b trials<sup>2,3</sup>  
DARWIN 1 and 2

**DARWIN 3<sup>4</sup>**  
Ongoing open-label extension trial of DARWIN 1 and 2 studies.

**FINCH 4<sup>5</sup>**  
Ongoing double-blind extension trial of FINCH 1-3.

2

Long-term extension trials<sup>4,5</sup>  
DARWIN 3 and FINCH 4

OVER  
**8000**  
patient-years of exposure<sup>6</sup>

RA, rheumatoid arthritis.

**References:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. **2.** Westhovens R, Taylor PC, Alten R, et al. *Ann Rheum Dis.* 2017;76(6):998-1008. **3.** Kavanaugh A, Kremer J, Ponce L, et al. *Ann Rheum Dis.* 2017;76(6):1009-1019. **4.** Kavanaugh A, Westhovens R, Winthrop K, et al. *J Rheumatol.* 2021;48(8):1230-1238. **5.** Long term extension study to assess the safety and efficacy of filgotinib in adults with rheumatoid arthritis (FINCH 4). Clinical trials identifier: NCT03025308. Updated January 14, 2021. Accessed January 20, 2022. <https://www.clinicaltrials.gov/ct2/show/NCT03025308>. **6.** Winthrop K, Tanaka Y, Takeuchi T, et al. *Arthritis Rheumatol.* 2021;73(suppl 10).

RA, rheumatoid arthritis.  
\* The most frequently reported adverse events were headache, nasopharyngitis, upper respiratory tract infection, and back pain.  
**Reference:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.  
Materials have been prepared for an international audience and are intended for healthcare professionals, excluding those from France. JYSELECA is approved but not reimbursed in Denmark.  
Data are correct as of 01/06/2022.





- Clinical Trial Programme
- Study Design
- Primary Endpoint

## Study design

Three Phase 3, randomised, double-blind, placebo-controlled trials in adult patients with moderate to severe active RA.<sup>1</sup>



THE FINCH 1  
STUDY

Patients with inadequate response to MTX (**MTX-IR**) randomised to<sup>1</sup>:

- JYSELECA 200 mg + MTX
- JYSELECA 100 mg + MTX
- Adalimumab + MTX
- Placebo + MTX



THE FINCH 2  
STUDY

Patients with inadequate response to bDMARDs (**Biologic-IR**) randomised to<sup>1</sup>:

- JYSELECA 200 mg + csDMARD
- JYSELECA 100 mg + csDMARD
- Placebo + csDMARD



THE FINCH 3  
STUDY

Patients naïve to MTX therapy (**MTX-Naïve\***) randomised to<sup>1</sup>:

- JYSELECA 200 mg + MTX
- JYSELECA 100 mg + MTX
- JYSELECA 200 mg monotherapy
- MTX alone

ACR, American College of Rheumatology; bDMARD, biologic disease-modifying antirheumatic drug; csDMARD, conventional synthetic disease-modifying antirheumatic drug; DMARD, disease-modifying antirheumatic drug; IR, intolerance or inadequate response; MTX, methotrexate; RA, rheumatoid arthritis.

\* JYSELECA is not indicated for use in DMARD-naïve patients.

**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.

RA, rheumatoid arthritis.  
\* The most frequently reported adverse events were headache, nasopharyngitis, upper respiratory tract infection, influenza, and diarrhoea.  
**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.  
Materials have been prepared for an international audience and are intended for healthcare professionals, excluding those from France. JYSELECA is approved but not reimbursed in Denmark.  
Data are correct as of 01/06/2022.








# JYSELECA met its primary endpoint of ACR20 across all Phase 3 trials<sup>1</sup>

Across all Phase 3 trials, JYSELECA demonstrated significant ACR20 response at Week 12 (FINCH 1 and FINCH 2) and Week 24 (FINCH 3) vs placebo + MTX, placebo + csDMARD, or MTX alone.<sup>1</sup>



<sup>a</sup> JYSELECA is not indicated for use in DMARD-naïve patients.  
\*\*\*  $P \leq .001$  vs placebo + MTX, placebo + csDMARD, or MTX alone.

ACR, American College of Rheumatology; bDMARD, biologic disease-modifying antirheumatic drug; csDMARD, conventional synthetic disease-modifying antirheumatic drug; DMARD, disease-modifying antirheumatic drug; IR, intolerance or inadequate response; MTX, methotrexate; RA, rheumatoid arthritis.

**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.

RA, rheumatoid arthritis.  
\* The most frequently reported adverse events were  
**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.  
Materials have been prepared for the purpose of this presentation.  
Data are correct as of 01/06/2022.





PRODUCT INFORMATION FOR HEALTHCARE PROFESSIONALS

▼ This medicinal product is subject to additional monitoring. Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting systems and to DrugSafety.Global@glpg.com.

**Abbreviated Summary of Product Characteristics: Jyseleca® (filgotinib)**, 100 mg or 200 mg film-coated tablets.

**Indications:** Treatment of moderate to severe rheumatoid arthritis in adults who have had an inadequate response to, or who are intolerant to, one or more disease-modifying anti-rheumatic drugs (DMARDs). Can be used as monotherapy or in combination with methotrexate.

**Dosage:** Treatment should be initiated by a physician experienced in the treatment of rheumatoid arthritis. The tablets should be swallowed whole. For guidance on laboratory monitoring and dose initiation or discontinuation, see the Summary of Product Characteristics. Treatment should be discontinued if a patient develops a serious infection until the infection is under control. Adults: 200 mg once daily. Elderly ≥75 years: Recommended starting dose: 100 mg once daily. Moderate or severe renal impairment (CrCl 15 to <60 mL/min): 100 mg once daily. End-stage renal disease (CrCl <15 mL/min): Filgotinib has not been studied and is therefore not recommended. Severe hepatic impairment (Child-Pugh C): Filgotinib has not been studied and is therefore not recommended. Children <18 years: The safety and efficacy of filgotinib have not yet been established. No data available.

**Contraindications:** Hypersensitivity to the active substance or to any of the excipients, active tuberculosis (TB), active serious infections, pregnancy.

**Warnings and precautions:** Immunosuppressants: Combination with other potent immunosuppressants such as ciclosporin, tacrolimus, biological DMARDs or other Janus kinase (JAK) inhibitors is not recommended. Infections, including severe infections (most commonly pneumonia), have been reported. Patients should be closely monitored for the development of signs and symptoms of infection during and after treatment with filgotinib. Elderly ≥75 years: Caution should be exercised due to higher incidence of serious infections. Tuberculosis: Patients should be screened for TB before initiating filgotinib. Filgotinib must not be used during active TB. In latent TB, standard antimycobacterial therapy should be given before filgotinib administration. Viral reactivation (including herpesvirus reactivation) has been reported in clinical trials. If the patient develops shingles, treatment should be temporarily withheld until the episode is resolved. Malignancy: Malignancies have been observed in clinical trials with filgotinib. Risks and benefits should be considered before initiating treatment for known malignancy other than well-treated non-melanoma skin cancer (NMSC), or when considering continuing treatment in patients who develop a malignancy. Patients at increased risk of skin cancer should have regular skin examinations. Haematological abnormalities: Treatment should not be initiated or should be discontinued temporarily in patients with absolute neutrophil count (ANC) <1 x 10<sup>9</sup> cells/L, absolute lymphocyte count (ALC) <0.5 x 10<sup>9</sup> cells/L or haemoglobin <8 g/dL observed during routine treatment. Vaccinations: Use of live vaccines during/immediately before treatment with filgotinib is not recommended. Cardiovascular risk: Risk factors (eg, hypertension, hyperlipidaemia) should be treated as part of normal treatment. Venous thromboembolism: JAK inhibitors should be used with caution in risk factors for DVT/PE or in connection with surgery and prolonged immobilisation. In case of clinical signs of DVT/PE, treatment should be discontinued and the patient evaluated immediately, followed by appropriate treatment.

**Interactions:** P-gp or BCRP inhibition: Caution should be exercised when administering substrates with a narrow therapeutic index concomitantly with filgotinib.

**Side effects:** Common: Urinary tract infection, upper respiratory tract infection, dizziness, nausea.

**Overdose:** In case of overdose, the patient should be monitored for signs and symptoms of side effects. Treatment: General supportive measures, including monitoring of vital signs as well as observation of the patient's clinical status. It is not known if filgotinib can be removed by dialysis.

**Pregnancy:** Filgotinib is contraindicated during pregnancy. Women of childbearing potential must use effective contraception during treatment and for at least 1 week after stopping treatment.

**Breastfeeding:** Jyseleca must not be used during breastfeeding.

**Fertility:** The potential risk of reduced fertility/infertility should be discussed with male patients before initiating treatment.

**Driving/operating machinery:** Filgotinib has no or negligible influence on the ability to drive and operate machinery. However, patients should be informed that dizziness has been reported during treatment with Jyseleca.

**Pack sizes:** 30 and 3 × 30 tablets.

**Price:** See current medicine prices at [www.medicinpriser.dk](http://www.medicinpriser.dk).

**Grant status:** Not eligible.

**Delivery:** NBS (may only be dispensed to hospitals or on the prescription of specific specialists).

**Marketing authorisation number:** EU/1/20/1480/003, EU/1/20/1480/004

**Marketing authorisation holder:** Galapagos NV, Belgium.

**The product information is abbreviated in relation to the product summary approved by the European Medicines Agency (EMA) dated 16.12.2021.** A complete product summary can be requested from the marketing authorisation holder or the Danish representative:

**Galapagos, tel. +800 7878 1345** or viewed on EMA's website.

**PLEASE READ THE SUMMARY OF PRODUCT CHARACTERISTICS BEFORE PRESCRIBING, IN PARTICULAR WITH REGARD TO SIDE EFFECTS, WARNINGS, AND CONTRAINDICATIONS.**

RA, rheumatoid arthritis.

\* The most frequently reported adverse reactions were nausea (3.5%), upper respiratory tract infection (3.3%), urinary tract infection (1.7%), and dizziness (1.2%).<sup>1</sup>

**Reference: 1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022.

Materials have been prepared for an international audience and are intended for healthcare professionals, excluding those from France. JYSELECA is approved but not reimbursed in Denmark.

Data are correct as of 01/06/2022.



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI



**Jyseleca®**  
filgotinib  
JAK1-preferential inhibitor



# Strong RA therapy across all key clinical outcomes<sup>1,2</sup>

With JYSELECA 200 mg + MTX at Week 24<sup>1</sup>:

58%

of patients achieved **ACR50**  
vs 52% with adalimumab + MTX

36%<sup>§</sup>

of patients achieved **ACR70**  
vs 30% with adalimumab + MTX

88%

of patients had **no**  
**radiographic progression**  
vs 86% with adalimumab + MTX

48%<sup>§§§</sup>

of patients achieved  
**remission** (DAS28-CRP <2.6)  
vs 36% with adalimumab + MTX

## JYSELECA achieved nominally significant improvements vs adalimumab across key clinical outcomes<sup>1,\*</sup>

ACR, American College of Rheumatology; CRP, C-reactive protein; DAS28, Disease Activity Score 28 joints; IR, inadequate response; MTX, methotrexate; RA, rheumatoid arthritis.

§  $P \leq .05$ , §§§  $P \leq .001$  vs adalimumab + MTX (nominal  $P$  values).

\* Patients in the study were MTX-IR. The FINCH 1 trial was not powered to detect superiority of JYSELECA vs adalimumab for ACR responses, radiographic progression, or remission.

**References:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. **2.** Combe B, Kivitz A, Tanaka Y, et al. *Ann Rheum Dis.* 2021;80(7):848-858.



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI

 **Jyseleca**<sup>®</sup>  
filgotinib  
JAK1-preferential inhibitor

# JAK1-preferential inhibitor with a proven safety profile<sup>1-5,\*</sup>

- › Generally well tolerated<sup>1</sup>
- › Low rates of JAKi-associated AEs, including serious infections, herpes zoster, and VTE, similar to adalimumab<sup>2,4,\*</sup>
- › Consistent safety profile up to 6.8 years<sup>5</sup>

Throughout 1 year of treatment with JYSELECA<sup>6</sup>:

3.0/100 PY

Serious infections EAIR  
vs 3.4/100 PY adalimumab + MTX

1.4/100 PY

Herpes zoster EAIR  
vs 0.7/100 PY adalimumab + MTX

0.2/100 PY

VTE EAIR  
vs 0.3/100 PY adalimumab + MTX

AE, adverse event; EAIR, exposure-adjusted incidence rate; JAK, Janus kinase; JAKi, Janus kinase inhibitor; MTX, methotrexate; PY, patient-years; VTE, venous thromboembolism.  
\* The most frequently reported adverse reactions were nausea (3.5%), upper respiratory tract infection (3.3%), urinary tract infection (1.7%), and dizziness (1.2%).<sup>1</sup>

**References:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. **2.** Combe B, Kivitz A, Tanaka Y, et al. *Ann Rheum Dis.* 2021;80(7):848-858. **3.** Committee for Medicinal Products for Human Use (CHMP). *Jyseleca Assessment Report.* European Medicines Agency; 2020:1-170. **4.** Genovese MC, Winthrop K, Tanaka Y, et al. *Ann Rheum Dis.* 2020;79:324-325. **5.** Winthrop K, Tanaka Y, Takeuchi T, et al. *Arthritis Rheumatol.* 2021;73(suppl 10). **6.** Data on file. Galapagos NV; 2019.



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI





# A convenient dosing regimen<sup>1,2</sup>

## Once-daily dosing<sup>1</sup>

- Recommended dose: 200 mg daily\* with MTX or as monotherapy
- 100 mg available for patients with moderate or severe renal impairment (CrCl 15 to <60 mL/min)<sup>†</sup> and as a starting dose for patients ≥75 years

## Low risk of drug-drug interactions<sup>1</sup>

- JYSELECA is the **only JAK inhibitor** that is not metabolised through CYP450<sup>1,3-5</sup>

## Easy-to-open bottle cap<sup>6</sup>



Combining JYSELECA with other potent immunosuppressants such as ciclosporin, tacrolimus, biologics, or other JAK inhibitors is not recommended, as a risk of additive immunosuppression cannot be excluded.<sup>1</sup>

\* No dose adjustment required in patients with mild renal impairment or mild or moderate hepatic impairment (Child-Pugh A or B).

† JYSELECA has not been studied in patients with end-stage renal disease (CrCl <15 mL/min) or in patients with severe hepatic impairment (Child-Pugh C) and is therefore not recommended for use in these patients.

**References:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. **2.** Taylor PC, Betteridge N, Brown TM, et al. *Patient Prefer Adherence*. 2020;14:119-131. **3.** RINVOQ (upadacitinib) Summary of Product Characteristics. AbbVie Inc.; 2021. **4.** XELJANZ (tofacitinib) Summary of Product Characteristics. Pfizer Inc.; 2021. **5.** OLUMIANT (baricitinib) Summary of Product Characteristics. Eli Lilly and Company; 2021. **6.** Data on file. Galapagos NV; 2019.



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI

 **Jyseleca**<sup>®</sup>  
filgotinib  
JAK1-preferential inhibitor



# STRENGTH *of* BALANCE

- Second-generation, JAK1-preferential inhibitor<sup>1</sup>
- Strong RA therapy<sup>1,2</sup>
- Proven safety profile<sup>1-5,\*</sup>

JAK, Janus kinase; RA, rheumatoid arthritis.

\* The most frequently reported adverse reactions were nausea (3.5%), upper respiratory tract infection (3.3%), urinary tract infection (1.7%), and dizziness (1.2%).<sup>1</sup>

**References:** **1.** JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. **2.** Combe B, Kivitz A, Tanaka Y, et al. *Ann Rheum Dis.* 2021;80(7):848-858. **3.** Committee for Medicinal Products for Human Use (CHMP). *Jyseleca Assessment Report*. European Medicines Agency; 2020:1-170. **4.** Genovese MC, Winthrop K, Tanaka Y, et al. *Ann Rheum Dis.* 2020;79:324-325. **5.** Winthrop K, Tanaka Y, Takeuchi T, et al. *Arthritis Rheumatol.* 2021;73(suppl 10).



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI

 **Jyseleca**<sup>®</sup>  
filgotinib  
JAK1-preferential inhibitor







## Resources

REAL-WORLD EXPERIENCE

CLINICAL DATA

MOA

COMMITMENT TO INFLAMMATION



Talking JYSELECA  
Series Trailer



Discovery  
Featuring René Galien, PhD  
Translational Research Lead,  
Galapagos France



Clinical Trials  
Featuring Dr David Walker  
Rheumatologist and Clinical  
Trial Investigator, UK



Bio-IR Patients  
Featuring Prof. Dr X. Baraliakos  
Rheumatologist, Germany



MTX-IR Patients  
Featuring Prof. Dr Torsten Witte  
Rheumatologist, Germany



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered  
trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI









## Resources

REAL-WORLD EXPERIENCE

CLINICAL DATA

MOA

COMMITMENT TO INFLAMMATION



FINCH 1: MTX-IR



FINCH 2: Bio-IR



FINCH 3: Monotherapy



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI






Resources

REAL-WORLD EXPERIENCE


CLINICAL DATA

MOA

COMMITMENT TO INFLAMMATION



JYSELECA: Mechanism of Action



Striking a Balance in JAK Inhibition  
Featuring Dr Paqui Gonzalez Traves  
Research Scientist, United States

What is the role of JAK1 in RA?

Balancing inhibition of proinflammatory cytokine signalling via JAK1 and limiting impact on JAK2- and JAK3-related physiological functions<sup>2,\*</sup>





Image is illustrative and based on in vitro findings; clinical relevance is unknown. There are currently no head-to-head trials between JAK inhibitors.

\* The role of JAK1 is not limited to proinflammatory cytokine signalling. These cytokines signal via JAK pairs, though they may depend predominantly on one JAK more than another for signalling. For example, IL-6 and IFN-γ both signal through JAK1/JAK2, but IL-6 may predominantly signal through JAK1, whereas IFN-γ is more dependent on JAK2.<sup>2</sup>

JYSELECA is a JAK1-preferential inhibitor<sup>1</sup>


JYSELECA has a 5-fold potency for JAK1 vs JAK2, JAK3, and TYK2 in biochemical assays<sup>1</sup>



The relevance of inhibition of specific JAK isoforms to therapeutic effectiveness or safety is not currently known.

Differential cytokine impact of JAK inhibitors

JYSELECA's inhibition of JAK1-dependent cytokines IFN-α and IL-6 is comparable to other JAK inhibitors, but with less inhibition of JAK2- and JAK3-dependent cytokines<sup>2</sup>



Adapted from Traves et al.<sup>2</sup>  
Data are based on in vitro whole-blood assays; clinical relevance is unknown. There are currently no head-to-head trials between JAK inhibitors.

<sup>a</sup> IC<sub>50</sub> indicates how much of a specific pharmacologic agent is required to inhibit a given biological activity by half.

<sup>b</sup> Data are normalised against IFN-α.

**References:** 1. JYSELECA (filgotinib) Summary of Product Characteristics. Galapagos NV; 2022. 2. Traves PG, Murray B, Campigotto F, Galien R, Meng A, Di Paolo JA. *Ann Rheum Dis*. 2021;80(7):865-875. 3. Malemud CJ. *Int J Mol Sci*. 2017;18(3):1-9. 4. Tan S, Xu J, Lai A, et al. *Mol Med Rep*. 2019;19(3):2057-2064. 5. Clark JD, Flanagan ME, Telliez JB. *J Med Chem*. 2014;57(12):5023-5038. 6. Schwartz DM, Kanno Y, Villarino A, Ward M, Gadina M, O'Shea JJ. *Nat Rev Drug Discov*. 2017;16(12):843-862. 7. Virtanen AT, Haikarainen T, Raivola J, Silvennoinen O. *BioDrugs*. 2019;33(1):15-32.





## Resources

REAL-WORLD EXPERIENCE

CLINICAL DATA

MOA

COMMITMENT TO INFLAMMATION

## Commitment to Inflammation



HOME



GALAPAGOS  
EVENT SCHEDULE



RESOURCES

JYSELECA, GALAPAGOS, and the GALAPAGOS logo are registered trademarks of Galapagos NV. © 2022 Galapagos NV. All rights reserved.  
GL-RA-JY-202204-00006 04/22

STUDY DETAILS

PI

