

AQUABC v0.3 — Ecological Model Reference Manual

ESTAS-AQUABC Development Team

September 2026

Contents

1	Introduction	3
1.1	Model Architecture	3
1.2	Source Code Organisation	3
2	Pelagic Model	4
2.1	State Variables	4
2.2	Phytoplankton Submodels	5
2.2.1	Temperature Limitation (CTMI)	5
2.2.2	Light Limitation	6
2.2.3	Nutrient Limitation (Synthesizing Unit)	6
2.2.4	Growth, Respiration, Mortality	6
2.2.5	Hypoxia Stress: Three-Regime System	7
2.3	Zooplankton Dynamics	7
2.4	Organic Matter Cycling	8
2.4.1	POM Dissolution	8
2.4.2	DOM Mineralisation	8
2.5	Dissolved Oxygen	8
2.6	Nutrient Cycling	8
2.7	Redox Chemistry	8
3	Sediment Diagenesis Model	9
3.1	Overview	9
3.2	Sediment State Variables (per layer)	9
3.3	Transport Processes	9
3.3.1	Advection	9
3.3.2	Diffusion	9
3.3.3	Particle Mixing (Bioturbation)	9
3.3.4	Burial	10
3.3.5	Combined Transport	10
3.4	Erosion and Deposition	10
3.5	Sediment–Water Fluxes	10
3.6	Bioturbation and Bioirrigation	10
3.6.1	Biodiffusion Coefficient	10
3.6.2	Bioturbation Parameters	11
3.6.3	Bioirrigation	11

3.6.4	Lower Boundary Condition	11
3.7	Sediment pH Correction	11
4	CO₂ System	13
4.0.1	Equilibrium Constant Selection	13
5	Macroalgae Model	14
5.1	Droop Quota Nutrient Limitation	14
5.2	Growth	14
6	Allelopathy Module	15
7	Optional Mechanisms	16
7.1	Zooplankton Food Limitation (ZOO_FOOD_MODEL)	16
7.2	Cyanobacterial Surface Positioning (CYANO_POS_MODEL, CYANO_POS_GUILDS)	16
7.3	Nostocales Akinete Staging (NOST_STAGE_MODEL)	16
7.4	Nostocales Fixation Switch (NOST_FIX_SWITCH, K_FIX_NOST, R_FIX_NOST)	17
7.5	Density-Dependent Phytoplankton Closure (PHYTO_CLOSURE_MODEL)	17
7.6	CYN Nitrogen Quota (CYN_VARIABLE_N) — VARN build only	17
7.7	Day-Length Weighting (LIGHT_DAYLENGTH_OPTION)	17
8	Model Constants	18
8.1	Pelagic Model Constants (324 parameters)	18
8.2	Sediment Model Constants (171 parameters)	19
9	Numerical Methods	20
9.1	Safety Functions	20
9.2	Time Integration	20
9.3	Mass Balance Safeguards	20
10	References	21

1 Introduction

AQUABC v0.3 is a comprehensive aquatic biogeochemical model that simulates coupled pelagic–sediment dynamics in lakes, reservoirs, estuaries, and coastal waters. The model is designed to be embedded in hydrodynamic transport frameworks (such as ESTAS or SHYFEM) that provide advection, dispersion, and external forcing.

1.1 Model Architecture

AQUABC consists of five coupled sub-models:

1. **Pelagic model:** 32 state variables covering phytoplankton (4 groups), zooplankton, organic matter, nutrients, dissolved oxygen, carbonate chemistry, and redox-sensitive metals.
2. **Sediment diagenesis model:** 24 state variables per sediment layer for early diagenesis with full redox sequencing and bioturbation.
3. **Macroalgae model:** 6 state variables using Droop quota kinetics.
4. **Allelopathy module:** 4 secondary metabolite pools for inter-species chemical inhibition.
5. **CO₂SYS:** Full carbonate chemistry solver (CDIAC implementation).

1.2 Source Code Organisation

Component	File
Pelagic kinetics	aquabc_II_pelagic_model.f90
Phytoplankton libraries	aquabc_II_pelagic_lib_*.f90
Sediment model	aquabc_II_sediment_model_1_fast.f90
Bioturbation module	aquabc_II_sediment_bioturbation.f90
Sediment model constants	aquabc_II_sediment_model_constants.f90
CO ₂ SYS	aquabc_II_co2sys.f90
Macroalgae	mod_MACROALGAE.f90
Allelopathy	mod_ALLELOPATHY.f90
Precision definitions	precision_kinds.f90
Parameter dictionary	STRING_UTILS.f90 (para_aqua module)

2 Pelagic Model

2.1 State Variables

The pelagic model tracks 32 state variables in each computational box. An additional 4 allelopathic metabolite variables (indices 33–36) are appended when the allelopathy module is active, giving 36 total transported variables.

Note: The indices below correspond to the actual code assignments in `aquabc_II_pelagic_svindex.f90`. The paper (Ertürk et al., 2023) uses a different numbering (1–21) reflecting a simplified configuration.

Index	Code Name	Unit	Description
1	NH4_N	mg N/L	Ammonium nitrogen
2	NO3_N	mg N/L	Nitrate nitrogen
3	PO4_P	mg P/L	Orthophosphate phosphorus
4	DOXY	mg O ₂ /L	Dissolved oxygen
5	DIA_C	mg C/L	Diatom carbon
6	ZOO_C	mg C/L	Zooplankton carbon
7	ZOO_N	mg N/L	Zooplankton nitrogen
8	ZOO_P	mg P/L	Zooplankton phosphorus
9	DET_PART_ORG_C	mg C/L	Detrital particulate organic carbon
10	DET_PART_ORG_N	mg N/L	Detrital particulate organic nitrogen
11	DET_PART_ORG_P	mg P/L	Detrital particulate organic phosphorus
12	DISS_ORG_C	mg C/L	Dissolved organic carbon
13	DISS_ORG_N	mg N/L	Dissolved organic nitrogen
14	DISS_ORG_P	mg P/L	Dissolved organic phosphorus
15	CYN_C	mg C/L	Non-fixing cyanobacteria carbon
16	OPA_C	mg C/L	Other planktonic algae carbon
17	DISS_Si	mg Si/L	Dissolved silica
18	PART_Si	mg Si/L	Particulate (biogenic) silica
19	FIX_CYN_C	mg C/L	N-fixing cyanobacteria carbon
20	INORG_C	mg C/L	Dissolved inorganic carbon (DIC)
21	TOT_ALK	meq/L	Alkalinity
22	FE_II	mg Fe/L	Dissolved ferrous iron
23	FE_III	mg Fe/L	Particulate ferric iron
24	MN_II	mg Mn/L	Dissolved manganese(II)
25	MN_IV	mg Mn/L	Particulate manganese(IV) oxide
26	CA	mg Ca/L	Calcium
27	MG	mg Mg/L	Magnesium
28	S_PLUS_6	mg S/L	Dissolved sulphate
29	S_MINUS_2	mg S/L	Dissolved sulphide
30	CH4_C	mg C/L	Dissolved methane

Index	Code Name	Unit	Description
31	NOST_VEG_HET_C	mg C/L	Nostocales vegetative cells + heterocysts
32	NOST_AKI_C	mg C/L	Nostocales akinetes
33	SEC_METAB_1	mg/L	Allelopathic secondary metabolite 1
34	SEC_METAB_2	mg/L	Allelopathic secondary metabolite 2
35	SEC_METAB_3	mg/L	Allelopathic secondary metabolite 3
36	SEC_METAB_4	mg/L	Allelopathic secondary metabolite 4

Legacy naming note: Variables 31–32 use two naming conventions in the code: the legacy `FIX_CYN_HET_C_INDEX` / `FIX_CYN_AK_C_INDEX` and the preferred `NOST_VEG_HET_C_INDEX` / `NOST_AKI_C_INDEX`. Both refer to the same state variables. The `NOST_*` convention is recommended.

2.2 Phytoplankton Submodels

Four phytoplankton groups share a common kinetic framework:

2.2.1 Temperature Limitation (CTMI)

The Cardinal Temperature Model with Inflection (Rosso et al., 1993):

$$\mu(T) = \begin{cases} \mu_{\max} \cdot \frac{(T - T_{\max})(T - T_{\min})^2}{(T_{\text{opt}} - T_{\min})[(T_{\text{opt}} - T_{\min})(T - T_{\text{opt}}) - (T_{\text{opt}} - T_{\max})(T_{\text{opt}} + T_{\min} - 2T)]} & T_{\min} < T < T_{\max} \\ 0 & \text{otherwise} \end{cases}$$

2.2.1.1 CTMI Parameter Name Mapping The parameter names in the constant file (`WCONST_04.txt`) have counter-intuitive mappings to the CTMI cardinal temperatures. The following table clarifies the relationship (see `aquabc_II_pelagic_auxillary.f90`):

WCONST Parameter Name	CTMI Role	Description
OPT_TEMP_LR (mapped to <code>Lower_TEMP</code>)	$T_{\min}$	Minimum cardinal temperature (growth = 0 below)
OPT_TEMP_UR (mapped to <code>Upper_TEMP</code>)	T_{opt}	Optimal temperature (growth peaks)
KAPPA_OVER_OPT_TEMP	$T_{\max}$	Maximum cardinal temperature (growth = 0 above)
KAPPA_UNDER_OPT_TEMP	<i>unused</i>	Not used in the CTMI formulation

2.2.2 Light Limitation

Steele/Smith formulation with optional Platt-style photoinhibition:

$$f_I = \frac{I}{I_s} \exp\left(1 - \frac{I}{I_s}\right) \cdot \exp(-\beta \cdot I)$$

where I is the depth-averaged PAR (Photosynthetically Active Radiation), I_s is the light saturation parameter, and β controls photoinhibition intensity.

Adaptive saturation. I_s is not a free constant: it is derived as $I_s = G_{max} \theta_{C:Chl} e / (0.083 \phi_{max} X_{KC})$. Because I_s scales with the maximum growth rate, G_{max} **cancels in the light-limited regime** — raising a group's growth constant cannot increase production where light is the binding constraint. This is a frequent source of confusion when calibrating: an apparently insensitive growth constant may simply be operating below light saturation.

Ice attenuation. Incident PAR is reduced before it enters the light limitation as an areal blend of open and ice-covered water,

$$I_{eff} = I \cdot [(1 - f) + f T] = I \cdot [1 - f(1 - T)]$$

where f is the ice areal fraction (driving function 10) and T is ICE_LIGHT_TRANS (model constant 324). With either $f = 0$ everywhere or $T = 1$ the multiplier is exactly unity and results are unchanged. Note that the ice fraction also damps surface aeration, ammonia volatilisation and atmospheric CO₂ exchange; before the light coupling was added, those three gas-exchange terms were its **only** consumers.

2.2.3 Nutrient Limitation (Synthesizing Unit)

The Synthesizing Unit colimitation model (Saito et al., 2008) is used for N–P colimitation in the **non-fixing** growth pathway of all phytoplankton groups:

$$f_N = \frac{1}{\frac{1}{f_{N-lim}} + \frac{1}{f_{P-lim}} + \frac{1}{f_{Si-lim}} - \frac{1}{f_{N-lim} \cdot f_{P-lim}} - \frac{1}{f_{N-lim} \cdot f_{Si-lim}} - \frac{1}{f_{P-lim} \cdot f_{Si-lim}} + \frac{1}{f_{N-lim} \cdot f_{P-lim} \cdot f_{Si-lim}}}$$

where each individual limitation follows Monod kinetics: $f_{N-lim} = [DIN] / ([DIN] + K_{HS,N})$.

Important: The Synthesizing Unit replaces Liebig's minimum **only** for N–P nutrient colimitation in the non-fixing pathway. Liebig's minimum is still used in: (1) the **fixing fraction** of N-fixing cyanobacteria, where the nitrogen term is a DIN-inhibition switch rather than colimitation; (2) combining nutrient limitation with oxygen limitation; (3) diatom Si colimitation cascaded with the N–P SU result. There is no user-configurable switch between SU and Liebig — the choice is hardcoded per pathway.

2.2.4 Growth, Respiration, Mortality

$$\frac{d[\text{PHYTO}]}{dt} = (\mu - R - D) \cdot [\text{PHYTO}]$$

where μ is the temperature-, light-, and nutrient-limited growth rate, R is Arrhenius-corrected respiration $R = k_{R,20} \cdot \theta^{T-20}$, and D is mortality (enhanced under hypoxia).

2.2.5 Hypoxia Stress: Three-Regime System

All phytoplankton groups implement a three-regime dissolved oxygen stress system that modulates mortality, respiration, and growth. Given a group-specific DO threshold `DO_STR_HYPOX`:

Regime 1 — Normal ($[O_2] > DO_STR_HYPOX$):

$$FAC_HYPOX = 1.0$$

No enhancement of mortality. Normal growth and respiration.

Regime 2 — Hypoxia ($0.1 \cdot DO_STR_HYPOX < [O_2] \leq DO_STR_HYPOX$):

$$FAC_HYPOX = \theta_{\text{hypox}}^{\epsilon_{\text{hypox}} \cdot (DO_STR_HYPOX - [O_2])}$$

Mortality is exponentially enhanced as DO drops below the threshold.

Regime 3 — Crash ($[O_2]/DO_STR_HYPOX \leq 0.1$):

$$FAC_HYPOX = \min\left(\frac{\Delta t}{0.5 \cdot K_D}, \frac{0.9}{K_D \cdot \Delta t}\right)$$

$$R_{\text{growth}} = 0, \quad R_{\text{respiration}} = 0$$

This crash regime is a critical **numerical safeguard** that prevents mass-balance overshoot under near-anoxia. Growth and respiration are zeroed, and mortality is capped to ensure no more than 90% of biomass can be removed in a single time step.

The parameters `THETA_HYPOX` and `EXPON_HYPOX` are defined per phytoplankton group in the `WCONST` constants file.

2.3 Zooplankton Dynamics

Zooplankton feeding uses the Active Switching Model (Gentleman et al., 2003):

$$I_i = I_{\text{max}} \cdot \frac{\rho_i \cdot [F_i] / \sum_j (\rho_j \cdot [F_j])}{K_F + \sum_j [F_j]} \cdot \sum_j [F_j]$$

where I_i is ingestion rate of food source i , ρ_i are preference weights, $[F_i]$ are food concentrations, and K_F is the half-saturation constant.

The dynamic preferences are computed using a switching power exponent:

$$\rho_i^{\text{dyn}} = \rho_i \cdot \left(\frac{\rho_i \cdot [F_i]}{\sum_j (\rho_j \cdot [F_j])} \right)^{n-1}$$

where $n = 1.5$ is the **switching power** (hardcoded in `aquabc_II_pelagic_lib_ZOOPLANKTON.f90`, line 192). Values: $n = 1.0$ gives linear (no switching), $n = 2.0$ gives strong switching. The default $n = 1.5$ provides moderate prey-switching behaviour.

2.4 Organic Matter Cycling

2.4.1 POM Dissolution

Particulate organic matter (C, N, P, Si) dissolves into dissolved forms:

$$R_{\text{diss}} = k_{\text{diss},20} \cdot \theta^{T-20} \cdot \frac{[\text{POM}]}{[\text{POM}] + K_{\text{HS}}} \cdot [\text{POM}]$$

2.4.2 DOM Mineralisation

DOC mineralisation follows a six-pathway redox sequence, each with its own rate constant, temperature correction, and half-saturation kinetics:

1. Aerobic respiration (O_2)
2. Denitrification (NO_3^-)
3. Manganese reduction (Mn(IV))
4. Iron reduction (Fe(III))
5. Sulphate reduction (S(VI))
6. Methanogenesis (DOC as terminal acceptor)

Each pathway is regulated by:

- Substrate half-saturation: $f = [\text{DOC}] / ([\text{DOC}] + K_{\text{HS,DOC}})$
- Electron acceptor availability: $f = [\text{EA}] / ([\text{EA}] + K_{\text{HS,EA}})$
- Inhibition by higher-priority acceptors (reversed Monod)
- pH correction (Gaussian)

2.5 Dissolved Oxygen

$$\frac{d[\text{O}_2]}{dt} = R_{\text{reaeration}} + R_{\text{photosynthesis}} - R_{\text{respiration}} - R_{\text{nitrification}} - R_{\text{oxidation}}$$

Reaeration follows the O'Connor–Dobbins or Wanninkhof formulation depending on wind speed input.

2.6 Nutrient Cycling

- **Nitrification:** $\text{NH}_4^+ \rightarrow \text{NO}_3^-$ (Monod kinetics in O_2 and NH_4 , pH-corrected)
- **Denitrification:** $\text{NO}_3^- \rightarrow \text{N}_2$ (only under low- O_2 conditions)
- **Phosphate speciation:** pH-dependent sorption with Fe(III)

2.7 Redox Chemistry

Full thermodynamic hierarchy: $\text{O}_2 > \text{NO}_3^- > \text{Mn(IV)} > \text{Fe(III)} > \text{SO}_4^{2-} > \text{methanogenesis}$.

Metal speciation (Fe, Mn) includes oxidation–reduction kinetics and equilibrium solubility calculations.

3 Sediment Diagenesis Model

3.1 Overview

The sediment model tracks 24 state variables per layer in a multi-layer vertical discretisation. It shares the same biogeochemical reaction framework as the pelagic model (POM dissolution, DOM mineralisation, nitrification, denitrification, full redox sequence) applied to sediment porewater and solid phases.

File: `aquabc_II_sediment_model_1_fast.f90`

3.2 Sediment State Variables (per layer)

The 24 sediment state variables mirror the pelagic model with sediment-specific organic matter pools, dissolved nutrients, and redox species. Each variable exists in dissolved (porewater) and/or particulate (solid) phase.

3.3 Transport Processes

Four transport mechanisms operate on each sediment layer:

3.3.1 Advection

Porewater flow between layers:

$$R_{\text{adv,in}} = \frac{C_{\text{entering}}}{\max(\phi, 10^{-20})} \cdot |v_{\text{adv}}|$$

$$R_{\text{adv,out}} = \frac{C_{\text{current}} \cdot f_{\text{sol}}}{\max(\phi, 10^{-20})} \cdot |v_{\text{adv}}|$$

where ϕ is porosity and f_{sol} is the dissolved fraction.

3.3.2 Diffusion

Concentration gradient-driven with tortuosity correction:

$$R_{\text{diff}} = \frac{D_{\text{eff}}}{1 + 3(1 - \phi)} \cdot \frac{\Delta C}{\max(\ell, 10^{-20})}$$

where $\ell = 0.5 \cdot (h_{i-1} + h_i)$ is the mixing length between adjacent layers.

3.3.3 Particle Mixing (Bioturbation)

For solid-phase species:

$$R_{\text{pmix}} = \frac{D_b \cdot \Delta C_{\text{solid}}}{\max(\ell, 10^{-20})}$$

where D_b is the biodiffusion coefficient (see §3.5).

3.3.4 Burial

Permanent removal to deep sediment:

$$R_{\text{burial}} = \frac{C \cdot w_{\text{burial}}}{\max(h, 10^{-20})}$$

where w_{burial} is the burial rate (m/day) and h is the layer thickness.

3.3.5 Combined Transport

$$\left. \frac{dC}{dt} \right|_{\text{transport}} = R_{\text{diff}} \cdot \phi + R_{\text{burial}} + R_{\text{pmix}} + R_{\text{adv,in}} - R_{\text{adv,out}}$$

3.4 Erosion and Deposition

When the sediment transport model is active (`isedi` > 0), layer concentrations are adjusted for sediment–water interface movement:

- **Deposition** ($H_{\text{erodep}} \leq 0$): Upper layer material mixed into lower layers
- **Erosion** ($H_{\text{erodep}} > 0$): Lower layer material exposed at the surface

In ESTAS standalone mode, `isedi` = 0 (prescribed settling used instead).

3.5 Sediment–Water Fluxes

24 dissolved fluxes are exchanged at the sediment–water interface based on concentration gradients between porewater and overlying water. Positive flux = release from sediment to water column.

Flux Index	Species	Direction
1	NH ₄ -N	Typically upward (nutrient release)
2	NO ₃ -N	Variable (nitrification/denitrification)
3	PO ₄ -P	Typically upward
4	DOXY	Typically downward (sediment O ₂ demand)
9	DOC	Typically upward
15	ALK	Typically upward

3.6 Bioturbation and Bioirrigation

File: `aquabc_II_sediment_bioturbation.f90`

When enabled (`switch_partmixing` = 1, default), particle mixing coefficients are dynamically computed based on local conditions rather than using a uniform constant.

3.6.1 Biodiffusion Coefficient

$$D_b(z, t) = D_{b0} \cdot \exp\left(-\frac{z}{z_{\text{mix}}}\right) \cdot \frac{[\text{O}_2]}{[\text{O}_2] + K_{\text{HS},\text{O}_2}} \cdot \left[1 + A \cos\left(\frac{2\pi(d - d_{\text{peak}})}{365}\right)\right]$$

The three multiplicative factors represent:

1. **Depth attenuation:** Exponential decay reflecting the decrease of macrofauna density with depth in sediments (Boudreau, 1997).
2. **Oxygen dependence:** Monod kinetics reflecting that benthic fauna require oxygen; under anoxia, bioturbation ceases.
3. **Seasonal modulation:** Sinusoidal annual cycle with summer peak, reflecting temperature-driven activity patterns (Soetaert et al., 1996).

3.6.2 Bioturbation Parameters

Parameter	Symbol	Default	Unit
Surface biodiffusion	D_{b0}	from input	m^2/day
Mixing depth	z_{mix}	0.05	m
O ₂ half-saturation	K_{HS,O_2}	2.0	mg/L
Seasonal amplitude	A	0.3	—
Peak day	d_{peak}	200	day
Irrigation enhancement	α_0	3.0	—
Irrigation depth	z_{irr}	0.04	m

3.6.3 Bioirrigation

Burrow-dwelling organisms pump overlying water through the sediment, enhancing porewater solute exchange beyond molecular diffusion:

$$\alpha(z, t) = 1 + \alpha_0 \cdot \exp\left(-\frac{z}{z_{\text{irr}}}\right) \cdot \frac{[\text{O}_2]}{[\text{O}_2] + K_{\text{HS},\text{O}_2}} \cdot f_{\text{season}}(t)$$

The irrigation factor α multiplies the molecular diffusion coefficient for dissolved-phase species:

$$D_{\text{eff,enhanced}} = \alpha \cdot D_{\text{eff,molecular}}$$

Bioirrigation is only applied to dissolved and mixed-phase species (not purely particulate). It is controlled by `switch_bioirrigation` (default = 1, enabled).

3.6.4 Lower Boundary Condition

A zero-flux (Neumann) boundary condition at the bottom of the deepest layer:

$$\left. \frac{\partial C}{\partial z} \right|_{z_{\text{bottom}}} = 0 \quad \Rightarrow \quad R_{\text{pmix}}(N) = -R_{\text{pmix,in}}(N)$$

This prevents artificial mass loss through the lower boundary.

3.7 Sediment pH Correction

Gaussian formulation (different from pelagic exponential):

$$f_{\text{pH}} = \exp\left(-\frac{(\text{pH} - \text{pH}_{\text{opt}})^2}{2\sigma^2}\right)$$

where $\text{pH}_{\text{opt}} = (\text{pH}_{\text{min}} + \text{pH}_{\text{max}})/2$ and $\sigma = (\text{pH}_{\text{max}} - \text{pH}_{\text{min}})/4$.

4 CO₂ System

The full carbonate chemistry system follows the CDIAC CO2SYS implementation, computing equilibrium speciation of dissolved inorganic carbon:



Given any two of (pH, pCO₂, DIC, ALK, CO₃²⁻), the system solves for all others using iterative Newton–Raphson methods with temperature- and salinity-dependent equilibrium constants.

4.0.1 Equilibrium Constant Selection

The code uses `K1K2CONSTANTS = 4`, which selects the **Mehrbach (1973) constants refit by Dickson & Millero (1987)**. This is the standard choice for estuarine and coastal applications. The selection is hardcoded in `aquabc_II_pelagic_model.f90` (line 390).

5 Macroalgae Model

5.1 Droop Quota Nutrient Limitation

$$Q_N = \frac{[\text{MAC-N}]}{[\text{MAC-C}]}, \quad \Phi_N = \max\left(0, 1 - \frac{Q_{0,N}}{Q_N}\right)$$

5.2 Growth

$$R_{\text{growth}} = k_{G,20} \cdot \theta^{T-20} \cdot \Phi_N \cdot \Phi_P \cdot \Phi_L \cdot \Phi_S \cdot [\text{MAC-C}]$$

where $\Phi_S = 1 - [\text{MAC-C}]/C_{\text{max}}$ is the logistic space limitation.

6 Allelopathy Module

Four secondary metabolite pools (state variable indices **33–36**, appended after the 32 standard pelagic variables) model inter-species chemical inhibition. Each phytoplankton group produces a specific metabolite that inhibits growth of competing groups via Monod-type dose-response:

$$f_{\text{inhib}} = 1 - \frac{[\text{SEC-METAB}]}{[\text{SEC-METAB}] + K_{\text{HS,inhib}}}$$

7 Optional Mechanisms

The mechanisms below are opt-in. Each defaults to a setting that is **byte-identical** to the model without it, so a configuration that enables none of them reproduces the legacy formulation exactly. All are selected in `PELAGIC_MODEL_OPTIONS.txt` (see the ESTAS manual for the file's positional format and the ordering constraint).

7.1 Zooplankton Food Limitation (`ZOO_FOOD_MODEL`)

The legacy formulation sums preference-weighted per-prey Monod terms. Because each term is diluted by its preference, the sum saturates well below unity even at high total food, which can cap ingestion below the mortality rate at *any* food concentration — a ceiling no coefficient can lift. Setting `ZOO_FOOD_MODEL = 1` replaces it with a saturating response to total preferred food,

$$f_{food} = \frac{F_{tot}}{K_{food} + F_{tot}}, \quad F_{tot} = \sum_j p_j C_j$$

with $K_{food} = \text{KHS_FOOD_TOT_ZOO}$.

Warning — The quadratic closure is not optional with this setting. Removing the legacy ceiling also removes the implicit stabilisation it provided; with a linear closure the zooplankton state diverges (Steele-type runaway). `ZOO_FOOD_MODEL = 1` therefore also applies a density-dependent closure referenced to `ZOO_CLOSURE_REF`, at which biomass the specific loss equals the calibrated rate. The ceiling and the stability were the same artifact, and the two changes must be tested as a pair.

7.2 Cyanobacterial Surface Positioning (`CYANO_POS_MODEL`, `CYANO_POS_GUILDS`)

Depth-averaged light limitation cannot represent a buoyant guild that escapes the average by accumulating at the surface. Option 1 applies a sub-daily calm-fraction surface blend; option 2 adds a persistence state — a per-box surface-accumulated fraction S that builds during calm spells and is dispersed by storms:

$$\frac{dS}{dt} = K_{up} F_{calm} (1 - S) - K_{disp} F_{storm} S$$

with formation and dispersal wind thresholds (`W_CRIT_POS_MIN`, `W_DISP_POS`) giving hysteresis, and self-shading applied to the concentrated fraction over depth `H_SURF_POS`. `CYANO_POS_GUILDS` restricts the mechanism to a subset of guilds; note that the ratchet keeps a separate S per guild but its rate constants are global, so all selected guilds surface identically.

7.3 Nostocales Akinete Staging (`NOST_STAGE_MODEL`)

Replaces the legacy akinete gates with an explicit benthic resting-stage bank: akinetes settle at `V_SETTLE_AKI`, germinate from the bed at `KR_GERM_BED` subject to a temperature guard (`T_GERM_AKI_STAGE`), are lost to burial at `K_MORT_BED_AKI`, and form under a declining-radiation latch (`I_FORM_AKI`). The bed bank is conservative and is reported in `NOST_STAGING.out`.

7.4 Nostocales Fixation Switch (NOST_FIX_SWITCH, K_FIX_NOST, R_FIX_NOST)

By default the fixation share of Nostocales growth is the fixed constant `FRAC_NOST_GROWTH`, independent of available nitrogen — so the guild grows predominantly on DIN and competes directly with non-fixers rather than occupying a diazotroph niche. Setting `NOST_FIX_SWITCH = 1` makes the share an inverse Monod in dissolved nitrogen, mirroring the switch the facultative fixer already uses (heterocyst induction):

$$\phi_{fix} = \frac{K_{fix}}{K_{fix} + \text{DIN} + \text{DON}}$$

`R_FIX_NOST` scales the productivity of the fixing channel only; values below 1 impose an energetic cost on N_2 fixation. Because it multiplies the fixing channel, it acts where nitrogen is scarce and has essentially no effect where nitrogen is replete.

7.5 Density-Dependent Phytoplankton Closure (PHYTO_CLOSURE_MODEL)

Scales the cyanobacterial specific death rate by C/C_{ref} (`PHYTO_CLOSURE_REF`), making total loss quadratic in biomass — the same closure the zooplankton use. It stands in for losses that scale with bloom density and are otherwise absent (colony aggregation and sinking, cyanophage lysis, scum senescence). This changes the *stability* of the equilibrium, not merely its position: stable coexistence of competing guilds requires each to limit itself more than it limits the other, which a linear loss cannot supply.

7.6 CYN Nitrogen Quota (CYN_VARIABLE_N) — VARN build only

Replaces Monod nitrogen limitation for non-fixing cyanobacteria with a Droop quota: an internal nitrogen store between `CYN_N_QMIN` and `CYN_N_QMAX`, filled by uptake at `CYN_N_VMAX` with half saturation `CYN_N_KHS_UPT`. **Requires the VARN build** (`nstate = 33`, adding the quota state variable); enabling it against a standard build halts the run with an explicit **error stop** rather than proceeding with a mis-indexed state array.

7.7 Day-Length Weighting (LIGHT_DAYLENGTH_OPTION)

I_A is a daily integral, so a photoperiod correction can enter two ways. Form A ($f_I \rightarrow F_{day} f(I_A)$) weights by the photoperiod without concentrating the dose into it, and therefore discards $(1 - F_{day})$ of each day's light; it is retained **only** to reproduce a published comparison and is not intended for use. Form B ($f_I \rightarrow F_{day} f(I_A/F_{day})$) concentrates the daily dose into the photoperiod before weighting, following WASP/EUTRO, and is the correct construction. Note that while light-limited the P–I curve is near-linear, so F_{day} largely cancels in Form B — it departs from the default only through curvature.

Note: Evaluating either form on the instantaneous rate is misleading. In a dynamical model the quantity that matters is the *integrated* effect on biomass, which can differ from the rate multiplier by a large factor once growth and loss re-equilibrate.

8 Model Constants

8.1 Pelagic Model Constants (324 parameters)

The pelagic model uses 324 parameters: 318 stored in the `MODEL_CONSTANTS(1:318)` array, 5 additional photoinhibition parameters (`BETA_*`) accessed via the named-parameter dictionary (`para_get_value`), and the under-ice PAR transmittance. Organised in 21 categories:

1. **General** (1–4): Temperature ranges, O₂ tolerances
2. **Diatoms** (5–31): Growth, respiration, mortality, nutrient kinetics, Si:C ratio
3. **Non-fixing cyanobacteria** (29–50): Buoyancy regulation, light adaptation
4. **N-fixing cyanobacteria** (51–74): Nitrogen fixation rates, heterocyst parameters
5. **Other planktonic algae** (75–96): Growth and loss parameters
6. **Zooplankton** (97–133): Feeding preferences, active switching parameters
7. **POM dissolution** (134–146): Rate constants per organic matter type
8. **DOM mineralisation** (147–152): Aerobic pathway parameters
9. **Nitrification** (152–157): Rate and half-saturation constants
10. **Redox chemistry** (158–209): All six electron acceptor pathways
11. **Methane** (210–234): CH₄ oxidation and production
12. **Settling** (235–250): Settling velocities per state variable
13. **pH correction** (251–266): Optimal pH ranges per process
14. **Dissolved metals** (267–276): Fe/Mn oxidation and dissolution rates
15. **Nostocales** (276–298): Heterocyst-forming cyanobacteria parameters
16. **Nostocales extended** (299–306): Akinete temperature/nitrogen thresholds
17. **POM dissolution saturation** (307–309): POM dissolution saturation kinetics
18. **DOM availability fractions** (310–315): DON/DOP availability fractions per process
19. **Phytoplankton mineralisation caps** (316–318): Max phytoplankton-mediated mineralisation rates
20. **Photoinhibition BETA** (319–323): Five β photoinhibition constants for each phytoplankton group, accessed via `para_get_value` using names `BETA_*`
21. **Under-ice light** (324): `ICE_LIGHT_TRANS` — PAR transmittance through ice cover (0–1). Default **1.00** (no attenuation, byte-identical to a build without the mechanism). Both readers guard on its presence, so constants files that predate it load unchanged; values outside [0, 1] are clamped with a warning

Warning — Constants reach the model through two parallel readers — the AQUABC name-based `para_get_value` path and the ESTAS index-based `INIT_PELAGIC_MODEL_CONSTANTS` path. A constant registered in only one of them is present in the file, readable by name, and **silently inert in the executing path**. Both the `BETA` block (categories 20) and `ICE_LIGHT_TRANS` (21) must appear in both. The diagnostic for this failure is a deliberate perturbation returning bit-identical output: a parameter whose variation changes nothing is not insensitive, it is disconnected.

Note on paper vs. code: Ert00ferk et al. (2023) report 183 calibrated constants — this refers to the subset actively calibrated for the Curonian Lagoon case study, not the full model constant set.

All constants are read from an external file and stored in the `para_aqua` name-value dictionary for runtime access.

8.2 Sediment Model Constants (171 parameters)

Sediment-specific rate constants mirror the pelagic model with separate values optimised for sediment conditions (lower temperatures, higher organic matter concentrations, different pH ranges). Read from `BOTTOM_SEDIMENT_MODEL_INPUT.txt` via the `WCONST` constants file.

9 Numerical Methods

9.1 Safety Functions

All division operations use `max(divisor, 10^{-20})` floors to prevent NaN/Inf. Temperature-dependent exponentials use `safe_exp()` with overflow protection. pH is clamped to $[4, 11]$.

9.2 Time Integration

The host transport framework (ESTAS or SHYFEM) provides the numerical time integration. AQUABC returns instantaneous rate derivatives which are integrated by the host using Euler or second-order Runge–Kutta methods.

9.3 Mass Balance Safeguards

- Maximum 50% loss per timestep for zooplankton and phytoplankton
- Concentration floors at 0 (non-negative constraint)
- Derivative clamping to prevent numerical overshoot

10 References

- Boudreau, B.P. (1997). *Diagenetic Models and Their Implementation*. Springer.
- Gentleman, W., Leising, A., Frost, B., Strom, S., and Murray, J. (2003). Functional responses for zooplankton feeding on multiple resources. *J. Plankton Res.* 25:1215–1234.
- Middelburg, J.J., Soetaert, K., and Herman, P.M.J. (1997). Empirical relationships for use in global diagenetic models. *Deep-Sea Res. I* 44:327–344.
- Morgan, B. and Lahav, O. (2007). The effect of pH on the kinetics of spontaneous Fe(II) oxidation. *Chemosphere*.
- Platt, T., Gallegos, C.L., and Harrison, W.G. (1980). Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton. *J. Mar. Res.* 38:687–701.
- Rosso, L., Lobry, J.R., and Flandrois, J.P. (1993). An unexpected correlation between cardinal temperatures of microbial growth highlighted by a new model. *J. Theor. Biol.* 162:447–463.
- Saito, M.A., Goepfert, T.J., and Ritt, J.T. (2008). Some thoughts on the concept of colimitation. *Limnol. Oceanogr.* 53:276–290.
- Soetaert, K., Herman, P.M.J., and Middelburg, J.J. (1996). A model of early diagenetic processes from the shelf to abyssal depths. *Geochim. Cosmochim. Acta* 60:1019–1040.
- Steele, J.H. (1962). Environmental control of photosynthesis in the sea. *Limnol. Oceanogr.*
- Stumm, W. and Morgan, J.J. (1996). *Aquatic Chemistry*. Wiley.
- CDIAC CO2SYS program documentation.
- EPA-829-R-14-007. AQUATOX Technical Documentation.