

# One Integer, Three Generations: $\sim 38$ Standard Model Quantities from $a_1 = 5$

Razvan-Constantin Anghelina  
contact@webdesignmedia.ro

[Interactive visualization](#) · [Reproducible code \(GitHub\)](#)

*Version 4.5 – March 2026*

## Abstract

We show that  $\sim 38$  quantities associated with Standard Model structure—the gauge group, three coupling constants, three generations, nine fermion mass ratios, six mixing angles, two CP phases, the Higgs-to- $W$  mass ratio, the electron-to-Planck mass ratio, a discrete PPN analogue, and more—are traced in this framework to a single integer  $a_1 = 5$ , the unique positive solution of  $a_1! = 4 a_1(a_1 + 1)$ . All dimensionless relations are internal to the framework, with zero adjustable parameters; in Planck units, the electron mass is fixed by the hierarchy identity  $m_e/m_P = \alpha^{4\phi^2}$  (0.24%), so only the conversion from Planck units to laboratory units remains external.

From  $a_1$  we obtain  $\phi = (1 + \sqrt{5})/2$ , the 600-cell (120 vertices), and  $E_8$  via the icosian construction. We prove  $N_{\text{gen}} = 3$  from Galois invariance on  $\mathbb{Z}[\phi]$ , derive the gauge group  $SU(3) \times SU(2) \times U(1)$  from two independent routes—the  $A_5$  decomposition of the vertex figure, and the kernel of the wave operator on the edge space ( $\ker(\square_1) = \rho_0 \oplus 2\rho_5$ , the  $E_8$  branching node)—and obtain all three couplings algebraically:  $\alpha^{-1} = 137.036$  (0.0001%),  $\alpha_s = 1/(2\phi^3) = 0.1180$  (0.11%),  $\sin^2 \theta_W = 6/26$  (0.19%). Nine fermion masses follow from  $m_f = m_e \phi^{5a+6b+\delta_1+\delta_2}$  with zero-parameter norm-log, Galois-conjugate, and Verlinde radiative corrections ( $c_{\text{eff}} = \sqrt{2/a_1}(1 - b_1\alpha^2)$ , one-loop from the  $S$ -matrix prefactor, two-loop vertex from the fusion matrix Frobenius norm  $\|N_{1/2}\|_F^2 = b_1$ ), achieving  $\chi^2 = 2.2/8$  dof ( $p = 0.97$ ). CKM and PMNS mixing angles, CP phases ( $\delta_{\text{CKM}} = \arctan \sqrt{5}$ ,  $\delta_{\text{PMNS}} = 3\arctan \sqrt{5}$ ), and  $\theta_{\text{QCD}} = 0$  are derived within the stated assumptions. The bare CKM exponents  $\{3, 7, 12\}$  satisfy three algebraic identities whose intersection singles out  $a_1 = 5$  (thirteenth characterisation); the Cayley–CKM vacuum offset  $\delta = 1$  provides a fourteenth. The Higgs mass ratio  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$  (0.008%) and the hierarchy  $m_e/m_P = \alpha^{4\phi^2}$  (0.24%) connect electroweak and gravitational scales through a single identity  $z_{\text{Planck}} = 1/\alpha_s + 2$ . The spectral action on the 600-cell simplicial complex reproduces the coefficient pattern of the bosonic sector, with all discrete coefficients determined by  $a_1$ .

The electroweak exponent is not chosen by hand: an RG-bootstrap cost function over integer pairs  $(a_1, n)$  has a unique isolated minimum at  $(5, 25)$  on the scanned discrete domain, and remains there under broad reweightings and after adding a cosmological term. Interpreting  $\Psi(a_1, n) \propto \exp(-S_{\text{eff}})$  yields an exponentially dominant saddle at  $(5, 25)$ , with the partition function numerically saturated by that saddle to machine precision, suggesting an effective vacuum-selection principle.

Galois conjugation  $\phi \rightarrow -1/\phi$  generates a dark sector with complex  $\alpha'$  and negative  $\alpha'_s$ : nine stable, electromagnetically dark particles interacting only gravitationally. The same Galois automorphism acts as  $q \rightarrow q^2$  on  $SU(2)$  Chern–Simons theory at level  $k = a_1 - 2 = 3$ , whose Fibonacci anyons have quantum dimension  $d = \phi$  and whose Verlinde fusion ring is isomorphic to  $\mathbb{Z}[\phi]$ . The topological entanglement entropy difference between physical and dark sectors is exactly  $\ln \phi$ . The cosmological constant  $\Lambda_P = \alpha^{57-\alpha_s}$  matches observation to  $0.13\sigma$ .

*Falsifiable predictions testable by 2030:*  $\sin^2 \theta_{13} = 1/45$  (JUNO);  $\sum m_\nu = 0.058$  eV (DESI/Euclid);  $\delta_{\text{PMNS}} = 197.7^\circ$  (DUNE/Hyper-K);  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$  (FCC-ee). Any single miss falsifies the entire framework.

## Contents

|          |                                                            |           |
|----------|------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                        | <b>5</b>  |
| <b>2</b> | <b>The Unique Integer</b>                                  | <b>7</b>  |
| 2.1      | Self-Reference and the Golden Ratio . . . . .              | 7         |
| 2.2      | Vacuum Selection: the Bootstrap Theorem . . . . .          | 7         |
| 2.3      | Geometric Characterisation . . . . .                       | 8         |
| 2.4      | What $a_1 = 5$ Determines . . . . .                        | 9         |
| <b>3</b> | <b>Geometric Framework</b>                                 | <b>9</b>  |
| 3.1      | The 600-Cell Polytope . . . . .                            | 9         |
| 3.2      | Spectral Structure . . . . .                               | 10        |
| 3.3      | The $E_8$ Connection . . . . .                             | 10        |
| <b>4</b> | <b>Gauge Group from <math>D_4</math></b>                   | <b>11</b> |
| 4.1      | From Discrete Symmetry to Continuous Gauge Group . . . . . | 14        |
| 4.2      | Edge Gauge Spectrum: Second Derivation . . . . .           | 15        |
| <b>5</b> | <b>Coupling Constants from <math>a_1</math></b>            | <b>16</b> |
| 5.1      | The Fine Structure Constant . . . . .                      | 16        |
| 5.2      | The Strong Coupling Constant . . . . .                     | 21        |
| 5.3      | The Weinberg Angle . . . . .                               | 22        |
| 5.4      | Summary of Coupling Constants . . . . .                    | 23        |
| <b>6</b> | <b>Why Exactly Three Generations</b>                       | <b>23</b> |
| <b>7</b> | <b>The Complete Fermion Mass Spectrum</b>                  | <b>25</b> |
| 7.1      | The Mass Formula . . . . .                                 | 25        |
| 7.2      | The Fibonacci Mass Formula . . . . .                       | 25        |
| 7.3      | Quark Sector: Offsets from Leptons . . . . .               | 25        |
| 7.4      | The Complete Mass Table . . . . .                          | 27        |
| 7.5      | Uniqueness of the Quantum Number Assignment . . . . .      | 28        |
| 7.6      | Norm-Log Mass Corrections . . . . .                        | 30        |
| 7.7      | The Galois Cascade . . . . .                               | 36        |
| 7.8      | Galois Norm Sum Rules . . . . .                            | 37        |
| 7.9      | Radiative Corrections from the Verlinde Theory . . . . .   | 37        |

|           |                                                                          |           |
|-----------|--------------------------------------------------------------------------|-----------|
| <b>8</b>  | <b>The Electroweak Sector</b>                                            | <b>39</b> |
| 8.1       | The $Z$ Boson Mass . . . . .                                             | 39        |
| 8.2       | The $W$ Boson and Higgs Mass . . . . .                                   | 40        |
| 8.3       | The Planck Mass . . . . .                                                | 42        |
| 8.4       | The Spectral Weight and the Electromagnetic–Gravitational Link . . . . . | 43        |
| <b>9</b>  | <b>The Gravitational Field Equation</b>                                  | <b>44</b> |
| 9.1       | Hodge Decomposition on the 600-Cell . . . . .                            | 44        |
| 9.2       | Gauge Structure and Physical Degrees of Freedom . . . . .                | 44        |
| 9.3       | Pair Creation and the Wronskian Selection Rule . . . . .                 | 45        |
| 9.4       | The Linearized Einstein Equation . . . . .                               | 45        |
| 9.5       | Discrete Scalar Response . . . . .                                       | 46        |
| 9.6       | The Galois Kernel Theorem . . . . .                                      | 47        |
| 9.7       | The Spectral Action and the Bosonic Lagrangian . . . . .                 | 51        |
| <b>10</b> | <b>Neutrino Masses</b>                                                   | <b>54</b> |
| 10.1      | Algebraic structure of the exponent . . . . .                            | 54        |
| 10.2      | The complete neutrino spectrum . . . . .                                 | 55        |
| <b>11</b> | <b>Mixing Angles</b>                                                     | <b>57</b> |
| 11.1      | CKM Matrix (Quarks) . . . . .                                            | 57        |
| 11.2      | CKM Concordance: Thirteenth Characterisation . . . . .                   | 58        |
| 11.3      | Cayley–Concordance Vacuum Offset: Fourteenth Characterisation . . . . .  | 58        |
| 11.4      | CKM Exponents from $E_8$ Decomposition Overlap . . . . .                 | 59        |
| 11.5      | PMNS Matrix (Neutrinos) . . . . .                                        | 60        |
| <b>12</b> | <b>CP Violation from Galois Theory</b>                                   | <b>62</b> |
| 12.1      | The Galois Angle . . . . .                                               | 62        |
| 12.2      | PMNS CP Phase . . . . .                                                  | 63        |
| 12.3      | The Strong CP Problem: $\theta_{\text{QCD}} = 0$ . . . . .               | 64        |
| 12.4      | The Jarlskog Invariant . . . . .                                         | 64        |
| <b>13</b> | <b>Extensions: Dark Sector, TQFT, and Cosmology</b>                      | <b>64</b> |
| 13.1      | The Galois Dark Sector . . . . .                                         | 65        |
| 13.2      | Topological Entanglement . . . . .                                       | 65        |
| 13.3      | The Cosmological Constant . . . . .                                      | 65        |
| 13.4      | Cosmological Consistency . . . . .                                       | 66        |
| <b>14</b> | <b>Testable Predictions</b>                                              | <b>66</b> |
| <b>15</b> | <b>Discussion and Conclusions</b>                                        | <b>67</b> |
| 15.1      | Summary of Predictions . . . . .                                         | 67        |
| 15.2      | Limitations . . . . .                                                    | 68        |
| 15.3      | Statistical Robustness . . . . .                                         | 70        |
| 15.4      | Uniqueness Among Regular Polytopes . . . . .                             | 70        |
| 15.5      | Variational Selection . . . . .                                          | 70        |
| 15.6      | Vacuum Selection by Effective Action . . . . .                           | 70        |
| 15.7      | Conclusions . . . . .                                                    | 72        |
| 15.8      | Related Work . . . . .                                                   | 74        |



# 1 Introduction

The Standard Model of particle physics is the most precisely tested physical theory in history. Yet it contains approximately 19 free parameters—coupling constants, mixing angles, and mass ratios—whose values must be determined experimentally. Why does the fine structure constant equal  $1/137.036$  rather than some other value? Why are there exactly three generations of fermions? Why does the Higgs boson weigh 125 GeV? These questions remain unanswered.

We propose a radical answer: **every** dimensionless parameter of the Standard Model is determined by a single self-referential equation,

$$x^2 = x + 1. \tag{1}$$

Its positive root is the golden ratio  $\phi = (1 + \sqrt{5})/2$ , the simplest number that equals one plus its own inverse. Interpreted as a fusion rule— $\tau \otimes \tau = \mathbf{1} \oplus \tau$ —it describes a particle that, upon self-interaction, produces the vacuum plus a copy of itself. Realized as a topological quantum field theory ( $SU(2)$  Chern–Simons), this self-referential structure selects a *unique* integer  $a_1 = 5$  through a bootstrap self-consistency condition: the quantum dimension of the theory must equal the algebraic parameter it defines.

From  $a_1 = 5$  we derive  $\phi$ , the 600-cell polytope (120 vertices), the  $E_8$  root system (240 roots), the gauge group  $SU(3) \times SU(2) \times U(1)$ , two gauge coupling constants algebraically and a third structurally, the number of fermion generations ( $N_{\text{gen}} = 3$ ), the masses of all nine charged fermions, and the mixing angles of both CKM and PMNS matrices. The mass spectrum is a *fractal* consequence of (1): every fermion mass is  $m_e \cdot \phi^n$ , and every power of  $\phi$  decomposes as  $\phi^n = F_n \phi + F_{n-1}$  (Fibonacci), reproducing the same self-referential pattern at every scale.

The logical flow is shown in Figure 1: from the single root  $a_1 = 5$ , three algebraic branches ( $\phi$ ,  $b_1$ ,  $N$ ) seed the geometric, spectral, and Galois structures that determine all physical parameters.




Figure 1: Derivation tree from  $a_1 = 5$ . Green: algebraic consequences. Orange: geometric/structural objects. Red: derived physics. Violet: observables. Every arrow represents a deduction with zero free parameters (except  $m_e$ , the dimensional anchor).

**Related work.** Several programs share the philosophy that Standard Model structure should follow from mathematical constraints rather than free parameters. Lisi [12] proposed embedding all gauge fields and fermions in  $E_8$ ; Furey [13] derived gauge quantum numbers from the algebra  $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$ ; and Chamseddine, Connes, and Marcolli [14] showed that the spectral action on a noncommutative geometry reproduces the full Standard Model Lagrangian coupled to gravity. The present work differs in that the input is a single integer  $a_1 = 5$  rather than an algebra or a finite geometry chosen to match observations; the 600-cell and  $E_8$  arise as *consequences* of this integer. We use the Chamseddine–Connes spectral action as the dynamical principle (Section 9), but the finite geometry is determined by  $a_1$ , not selected by hand.

**Guiding principle.** The framework rests on a single principle: *self-referential consistency*. The equation  $x^2 = x + 1$ —the requirement that the whole equal its part plus unity—generates the golden ratio  $\phi$ , whose Fibonacci fusion rule  $\tau \otimes \tau = \mathbf{1} \oplus \tau$  defines the simplest non-trivial topological quantum field theory. Demanding that the algebraic parameter of this theory equal its own quantum dimension selects  $a_1 = 5$  uniquely (Theorem 1). From there, the 600-cell,  $E_8$ , and the Standard Model follow without further input. Fermion masses are powers of  $\phi$  because each mass scale is a level of the same self-similar structure; the mass formula  $m_f = m_e \phi^n$  is the fractal decomposition  $\phi^n = F_n \phi + F_{n-1}$  made physical. Dark matter is not an addition: it is the second root of the same equa-

tion,  $\phi' = -1/\phi$ , the Galois conjugate that shares the algebraic structure but lacks a real electromagnetic coupling. The sole dimensional input  $m_e$  anchors the fractal to a physical scale.

The paper follows a single logical chain:  $a_1 = 5$  (Section 2)  $\rightarrow$  600-cell geometry (Section 3)  $\rightarrow$  gauge group (Section 4)  $\rightarrow$  coupling constants (Section 5)  $\rightarrow$   $N_{\text{gen}} = 3$  (Section 6)  $\rightarrow$  fermion masses (Section 7)  $\rightarrow$  electroweak bosons (Section 8)  $\rightarrow$  gravity and spectral action (Section 9)  $\rightarrow$  neutrinos (Section 10)  $\rightarrow$  mixing angles (Section 11)  $\rightarrow$  CP violation (Section 12)  $\rightarrow$  testable predictions (Section 14). Extensions to the dark sector, TQFT entanglement, the cosmological constant, and inflation are summarized in Section 13 with full details in the Supplementary Material.

At the end of this chain, the framework admits a second layer of selection: the same data that determine the electroweak and cosmological sectors define an effective cost functional with a sharply isolated minimum at the observed vacuum  $(a_1, n) = (5, 25)$  on the scanned domain. We treat this as an effective vacuum-selection principle, not yet as a microscopic Euclidean path integral derivation.

## 2 The Unique Integer

### 2.1 Self-Reference and the Golden Ratio

The framework begins with the observation that the simplest self-referential algebraic equation,

$$x^2 = x + 1,$$

already encodes the structure of a topological quantum field theory. Its positive root  $\phi = (1 + \sqrt{5})/2$  satisfies  $\phi = 1 + 1/\phi$ : the whole equals one plus its own inverse. Reinterpreted as a fusion rule for anyonic excitations,

$$\tau \otimes \tau = \mathbf{1} \oplus \tau, \tag{2}$$

this equation defines the Fibonacci anyon category: two copies of the fundamental excitation combine to produce either the vacuum ( $\mathbf{1}$ ) or a new copy of themselves ( $\tau$ ). The quantum dimension is  $d_\tau = \phi$ , and every iterated fusion decomposes via  $\phi^n = F_n \phi + F_{n-1}$  (Fibonacci numbers), giving the theory a *fractal* self-similar structure at every level.

The fusion matrix

$$N_\tau = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

has eigenvalues  $\phi$  and  $\phi' = -1/\phi$ : the physical sector and the Galois-conjugate (dark) sector are the two eigenvalues of self-observation.

### 2.2 Vacuum Selection: the Bootstrap Theorem

The Fibonacci fusion rule (2) selects a *unique* integer. Each positive integer  $a_1 \geq 3$  determines both an algebraic parameter  $\phi(a_1) = (1 + \sqrt{a_1})/2$  and an  $SU(2)$  Chern–Simons theory at level  $k = a_1 - 2$ , whose  $j = 1$  quantum dimension is

$$d_1(a_1) = \frac{\sin(3\pi/a_1)}{\sin(\pi/a_1)} = 1 + 2 \cos\left(\frac{2\pi}{a_1}\right). \tag{3}$$

Demanding that the algebraic parameter equal the quantum dimension of the theory it generates—a *bootstrap* self-consistency condition—gives

$$\boxed{d_1 = \phi(a_1) \iff \cos\left(\frac{2\pi}{a_1}\right) = \frac{\sqrt{a_1} - 1}{4}} \quad (4)$$

**Theorem 1** (TQFT Bootstrap Vacuum Selection). *The only positive integer  $a_1$  satisfying (4) is  $a_1 = 5$ .*

*Proof.* Write  $f(a_1) = \cos(2\pi/a_1) - (\sqrt{a_1} - 1)/4$ .

*Upper bound.* For  $a_1 \geq 26$ ,  $(\sqrt{a_1} - 1)/4 > 1 \geq |\cos(\cdot)|$ , so  $f < 0$ .

*Finite check.* For  $a_1 = 1, \dots, 25$ , exhaustive evaluation shows  $f(a_1) = 0$  only at  $a_1 = 5$ .

*Exactness.* For  $a_1 = 5$ :  $\cos(2\pi/5)$  is a root of  $4x^2 + 2x - 1 = 0$ , whose positive root is  $(\sqrt{5} - 1)/4$ . Hence  $f(5) = 0$  exactly.  $\square$

**Corollary 1** (Double embedding).  $d_{1/2}(a_1) = 2\cos(\pi/a_1) = \phi(a_1)$  also holds if and only if  $a_1 = 5$ . Thus both the  $j = \frac{1}{2}$  and  $j = 1$  quantum dimensions equal  $\phi$ , giving the spectrum  $\{d_j\} = \{1, \phi, \phi, 1\}$ .

*Proof.*  $2\cos(\pi/a_1) = \phi(a_1)$  iff  $\cos(\pi/5) = (1 + \sqrt{5})/4$ , the positive root of  $4x^2 - 2x - 1 = 0$ . The equality  $d_{1/2} = d_1$  reduces to  $x^2 - x - 1 = 0$  (the golden ratio equation), which requires  $a_1 = 5$  uniquely.  $\square$

**Physical interpretation.** The parameter  $a_1$  simultaneously defines three structures:

- (i) the number field  $\mathbb{Q}(\sqrt{a_1})$  with ring of integers  $\mathbb{Z}[\phi]$ ;
- (ii) a topological quantum field theory via  $SU(2)$  Chern–Simons at level  $k = a_1 - 2$ ;
- (iii) the quantum dimension  $d_1$  of the  $j = 1$  representation.

Self-consistency requires the *number* from (i) to equal the *quantum dimension* from (iii). This is a fixed-point condition  $\phi(a_1) = d_1(a_1)$  with a unique solution  $a_1 = 5$ . The theory selects itself: the algebraic parameter *is* the quantum dimension. The Chern–Simons theory and its physical consequences (Fibonacci anyons, Verlinde ring, topological entanglement) are developed in Section 13.2.

## 2.3 Geometric Characterisation

The integer  $a_1 = 5$  selected by the bootstrap admits a Diophantine characterisation:

$$\boxed{a_1! = 4a_1(a_1 + 1)} \quad (5)$$

**Unique solution:**  $5! = 120 = 4 \times 5 \times 6$ . The left side is the order of the binary icosahedral group  $2I$ , which permutes the 120 vertices of the 600-cell. The right side is  $4 \times (\text{diameter}) \times (\text{distance classes})$  of the 600-cell Cayley graph. This is a self-consistency condition between the symmetry group and the geometry it acts on: the group order equals a product of its own graph invariants. For  $a_1 \geq 6$  the factorial outgrows  $4a_1^2$ , so no further solutions exist.



## 2.4 What $a_1 = 5$ Determines

Define  $b_1 \equiv a_1 + 1 = 6$ . The cascade of derived quantities is:

| Quantity                        | Formula                                                         | Value              | Physical role                |
|---------------------------------|-----------------------------------------------------------------|--------------------|------------------------------|
| $\phi$                          | $(1 + \sqrt{a_1})/2$                                            | $(1 + \sqrt{5})/2$ | Golden ratio, scaling factor |
| $\phi'$                         | $(1 - \sqrt{a_1})/2$                                            | $(1 - \sqrt{5})/2$ | Galois conjugate             |
| $\text{disc}(\mathbb{Z}[\phi])$ | $a_1$                                                           | 5                  | Ring discriminant            |
| $N$                             | $a_1!$                                                          | 120                | 600-cell vertices = $ 2I $   |
| $h(E_8)$                        | $a_1 \cdot b_1$                                                 | 30                 | $E_8$ Coxeter number         |
| $N_{\text{eig}}$                | sol. of $\binom{n}{2} = b_1^2$                                  | 9                  | Distinct eigenvalues         |
| $\text{rank}(E_8)$              | $N_{\text{eig}} - 1$                                            | 8                  | $E_8$ rank                   |
| $\dim(E_8)$                     | $(N_{\text{eig}} - 1)(h + 1)$                                   | 248                | $E_8$ dimension              |
| $\sin^2 \theta_W$               | $b_1/(a_1^2 + 1)$                                               | 6/26               | Weinberg angle               |
| $\alpha$                        | sol. of $2\pi x^2 - 4a_1\phi^4 x + 1 = 0$                       | 1/137.036          | Fine structure constant      |
| $\alpha_s$                      | $1/z$ : $\text{Tr} = 8$ , $ N  = 4$                             | $1/(2\phi^3)$      | Strong coupling              |
| $N_{\text{gen}}$                | from Fibonacci on $\mathbb{Z}[\phi]$                            | 3                  | Fermion generations          |
| $m_e/m_P$                       | $\alpha^{z_P}$ , $z_P = 12\phi^2/3$ (spectral weight of $R_3$ ) | $\alpha^{4\phi^2}$ | Planck hierarchy             |

Table 1: The cascade from  $a_1 = 5$ . Every entry is determined, with zero free parameters.

A remarkable secondary Diophantine reinforces the uniqueness:

$$(a_1 - 2)! = a_1 + 1 \quad \implies \quad 3! = 6 \quad \checkmark \quad (6)$$

which also has  $a_1 = 5$  as its unique solution (re-derived independently from the Galois norm cubic identity in Section 7.8). The number 5 is simultaneously the graph diameter, the discriminant of  $\mathbb{Z}[\phi]$ , and  $|2I/2T|$  (cosets of the binary tetrahedral in the binary icosahedral group).

A complete notation table is provided in Supplementary Material S1.

## 3 Geometric Framework

### 3.1 The 600-Cell Polytope

The 600-cell  $\{3, 3, 5\}$  is a regular 4-dimensional polytope:

|                      |                         |
|----------------------|-------------------------|
| Vertices             | $120 = a_1!$            |
| Edges                | $720 = 120 \times 12/2$ |
| Faces (triangles)    | 1200                    |
| Cells (tetrahedra)   | 600                     |
| Symmetry group $H_4$ | Order 14,400            |
| Graph diameter       | $5 = a_1$               |
| Vertex degree        | $12 = 2 b_1$            |

**Universal optimality.** By the Cohn–Kumar theorem [1, 3], the 120 vertices of the 600-cell form the *unique* universally optimal spherical code on  $S^3$ : they minimize the energy for *every* completely monotonic potential simultaneously. This is not an assumption

but a theorem—any attractive pairwise interaction on  $S^3$  produces this configuration. The discreteness is forced by energy minimization, just as the hexagonal lattice (2D) and FCC structure (3D) are forced by close-packing. The 600-cell is not chosen; it is the unique answer to “what is the optimal discrete geometry on  $S^3$ ?”

### 3.2 Spectral Structure

The 600-cell graph is distance-transitive with an association scheme of  $N_{\text{eig}} = 9$  classes. Its adjacency matrix has exactly 9 distinct eigenvalues, all in  $\mathbb{Z}[\phi]$ :

| Index | Eigenvalue $\lambda_i$ | Multiplicity | Structure       |
|-------|------------------------|--------------|-----------------|
| 0     | 12                     | $1 = 1^2$    | Trivial         |
| 1     | $6\phi$                | $4 = 2^2$    | $\phi$ -sector  |
| 2     | $4\phi$                | $9 = 3^2$    | $\phi$ -sector  |
| 3     | 3                      | $16 = 4^2$   | Rational        |
| 4     | 0                      | $25 = 5^2$   | Kernel          |
| 5     | -2                     | $36 = 6^2$   | Rational        |
| 6     | $4 - 4\phi$            | $9 = 3^2$    | $\phi'$ -sector |
| 7     | -3                     | $16 = 4^2$   | Rational        |
| 8     | $6 - 6\phi$            | $4 = 2^2$    | $\phi'$ -sector |

Table 2: 600-cell adjacency spectrum. All eigenvalues are of the form  $a + b\phi$  with  $a, b \in \mathbb{Z}$ , and all multiplicities are perfect squares  $n^2$ . The intersection numbers are  $a_1 = 5$ ,  $b_1 = 6$ .

**Key structural facts:** (i) Every multiplicity is a perfect square—the irreducible representation dimensions of  $2I$  squared. (ii) The  $\phi$ -dependent eigenvalues (those with  $b \neq 0$ ) have total dimension  $4 + 9 + 9 + 4 = 26$ , split symmetrically between  $\phi$ - and  $\phi'$ -sectors ( $13 + 13$ ). (iii) The intersection numbers  $a_1 = 5$  and  $b_1 = 6$  determine the mass formula (Section 7).

### 3.3 The $E_8$ Connection

The  $E_8$  root system is constructed from the 600-cell via the icosian lattice [7, 8]. Each vertex  $v = (v_0, v_1, v_2, v_3)$  with  $v_i = a_i + b_i\phi$  maps to an 8-tuple  $(a_0, b_0, \dots, a_3, b_3) \in \mathbb{R}^8$ . The *Galois conjugation*  $\sigma : \phi \rightarrow \phi' = (1 - \sqrt{5})/2$  is the unique non-trivial automorphism of the number field  $\mathbb{Q}(\sqrt{5})$ ; it sends  $\sqrt{5} \rightarrow -\sqrt{5}$  while fixing all rational numbers. Physically,  $\sigma$  will play the role of a discrete symmetry that distinguishes the visible and dark sectors (Section 13.1). Applying  $\sigma$  to the vertex coordinates defines a second copy  $T = \phi' \cdot S$ . Then:

$$\boxed{E_8 \text{ roots} = S \cup T, \quad |S| = |T| = 120, \quad S \cap T = \emptyset, \quad |S \cup T| = 240} \quad (7)$$

The inner product structure matches  $E_8$  exactly (240/240 roots verified by Procrustes alignment). The 24 Galois-fixed vertices ( $b_i = 0$  for all  $i$ ) form the  $D_4$  root system, providing a second derivation of the gauge/fermion split:

$$120 = \underbrace{24}_{\text{Galois-fixed} = D_4 = \text{gauge}} + \underbrace{96}_{\text{Galois-moving} = \text{fermions}} \quad (8)$$

The Coxeter number  $h(E_8) = 30 = a_1 \cdot b_1$ , the rank  $= 8 = N_{\text{eig}} - 1$ , and  $\dim(E_8) = 248 = 8 \times 31 = (N_{\text{eig}} - 1)(h + 1)$ .

## 4 Gauge Group from $D_4$

The 24 non-fermionic vertices (Types A and B: cross-polytope + tesseract) form the  $D_4$  root system, which is the 24-cell. The 96 remaining vertices decompose as

$$96 = 16 \times 3 \times 2 = (\text{fermions per generation}) \times (\text{generations}) \times (\text{chiralities}) \quad (9)$$

**Representation-theoretic derivation.** The 12 nearest neighbors of any vertex form an *icosahedron* (the vertex figure of the 600-cell), whose rotation group is  $A_5 \cong 2I/\mathbb{Z}_2$ . The permutation representation of  $A_5$  on the 12 icosahedron vertices has character  $\chi(e) = 12$ ,  $\chi(\text{double transposition}) = 0$ ,  $\chi(3\text{-cycle}) = 0$ ,  $\chi(5\text{-cycle}) = 2$ , which decomposes as

$$12 = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5} \quad (10)$$

where  $\mathbf{3}$  and  $\mathbf{3}'$  are the two inequivalent 3-dimensional irreps of  $A_5$ , distinguished by the Galois automorphism  $\phi \leftrightarrow \phi'$  (their characters on 5-cycles are  $\phi$  and  $\phi'$  respectively).

Now embed  $A_5 \subset SO(3) \subset SU(3)$  via the  $\mathbf{3}$ -irrep. The adjoint of  $SU(3)$  restricts to  $A_5$  as

$$\text{ad}(SU(3))|_{A_5} = \mathbf{3} \otimes \mathbf{3} - \mathbf{1} = \mathbf{3} \oplus \mathbf{5} \quad (\dim = 8) \quad (11)$$

since  $\mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{5}$  (verified via character inner products). The permutation decomposition (10) therefore becomes

$$12 = \underbrace{\mathbf{1}}_{U(1)} + \underbrace{\mathbf{3}'}_{=\text{ad}(SU(2))} + \underbrace{\mathbf{3} \oplus \mathbf{5}}_{=\text{ad}(SU(3))} = 1 + 3 + 8 \quad (12)$$

This is the *only* valid grouping. To see why, enumerate all partitions of  $\{\mathbf{1}, \mathbf{3}, \mathbf{3}', \mathbf{5}\}$  into sub-sums and check which produce adjoint dimensions of simple Lie groups ( $\text{ad}(SU(n)) = n^2 - 1$ , so the allowed adjoint dimensions  $\leq 12$  are 3, 8):

| Grouping                                                        | Dimensions  | Adjoint?                                |
|-----------------------------------------------------------------|-------------|-----------------------------------------|
| $\{\mathbf{1}\} + \{\mathbf{3}'\} + \{\mathbf{3}, \mathbf{5}\}$ | $1 + 3 + 8$ | <b>Yes:</b> $U(1) + SU(2) + SU(3)$      |
| $\{\mathbf{1}\} + \{\mathbf{3}\} + \{\mathbf{3}', \mathbf{5}\}$ | $1 + 3 + 8$ | <b>Yes:</b> Galois conjugate of above   |
| $\{\mathbf{1}\} + \{\mathbf{5}\} + \{\mathbf{3}, \mathbf{3}'\}$ | $1 + 5 + 6$ | No: $5 \neq n^2 - 1$ ; $6 \neq n^2 - 1$ |
| $\{\mathbf{1}\} + \{\mathbf{3}, \mathbf{3}', \mathbf{5}\}$      | $1 + 11$    | No: $11 \neq n^2 - 1$                   |
| $\{\mathbf{1}, \mathbf{3}\} + \{\mathbf{3}', \mathbf{5}\}$      | $4 + 8$     | No: $4 \neq n^2 - 1$                    |
| $\{\mathbf{1}, \mathbf{3}'\} + \{\mathbf{3}, \mathbf{5}\}$      | $4 + 8$     | No: $4 \neq n^2 - 1$                    |
| $\{\mathbf{1}, \mathbf{5}\} + \{\mathbf{3}, \mathbf{3}'\}$      | $6 + 6$     | No: $6 \neq n^2 - 1$                    |
| $\{\mathbf{1}, \mathbf{3}, \mathbf{3}'\} + \{\mathbf{5}\}$      | $7 + 5$     | No: neither is $n^2 - 1$                |

Only the first two rows produce valid adjoint decompositions, and they are related by the Galois automorphism  $\phi \leftrightarrow \phi'$  (which swaps  $\mathbf{3} \leftrightarrow \mathbf{3}'$ ). The physical choice  $\phi > 0$  breaks this discrete symmetry, selecting a unique assignment.

The connection to the McKay correspondence is immediate: the first three nodes of  $\widetilde{E}_8$  carry marks 1, 2, 3 (irrep dimensions of  $2I$ ), and the corresponding  $SU(d)$  adjoints have dimensions  $1, 2^2 - 1, 3^2 - 1 = 1, 3, 8$ . The identity  $1 + 3 + 8 = 12 = \text{vertex degree}$  is thus not a coincidence but a consequence of the fact that the permutation representation on the vertex figure decomposes into exactly the adjoint representations selected by the McKay marks. It is essential to distinguish this *gauge dimension* decomposition  $12 = 1 + 3 + 8$  from the  $A_5$  *irrep* decomposition  $12 = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5}$  (Eq. 15): the bridge is that

$\text{ad}(\mathfrak{su}(3))|_{A_5} = \mathbf{3}' \oplus \mathbf{5}$  has dimension  $3 + 5 = 8$  (proven in Theorem 3). The  $E_8$  Dynkin diagram branches at the  $D_4$  node into three legs with dimensions Leg A =  $11 = a_1 + b_1$ , Leg B =  $6 = b_1$ , and Leg C =  $9 = N_{\text{eig}}$ , with triality broken by the Kac labels  $\{3, 4, 5\}$ .

**Computational verification (edge splitting).** Each fermionic vertex (type C, 96 total) has 12 neighbors decomposing by vertex type as  $1A + 2B + 9C$ , confirmed by direct geometry. The SM assignment follows:  $U(1)$ : 1 AC edge (perfect matching, topological confinement with gap = 2);  $SU(2)$ : 2 BC edges (charged weak) + 1  $CC^*$  edge (neutral current criterion: shared A neighbor, no shared B);  $SU(3)$ : 8 regular CC edges (color sector, 288 pure plaquettes reproducing QCD self-interaction). Thus

$$12 = \underbrace{1}_{U(1)} + \underbrace{2+1}_{SU(2)} + \underbrace{8}_{SU(3)} = \dim(\text{SM gauge group}) \quad (13)$$

verified for all 96 fermion vertices.

The 96 fermionic vertices admit a natural  $48 + 48$  split under the Galois involution  $\sigma : \phi \rightarrow \phi'$ , with the 24 gauge vertices being Galois-fixed (their coordinates lie in  $\mathbb{Q}$ ). This provides a candidate chirality grading algebraically derived from the framework. We summarize below what is established and what remains open.

**What is derived.** (i) The  $48 + 48$  Galois split exists and is *exact*: each fermionic vertex  $v$  has a unique Galois partner  $\sigma(v) \neq v$ , giving  $96/2 = 48$  pairs. (ii) The counting  $48 = 16 \times 3$  matches the SM content: 16 Weyl fermions per generation times 3 generations. (iii) All six anomaly cancellation conditions— $\text{Tr}(Y^3) = 0$ ,  $\text{Tr}(Y) = 0$ ,  $[SU(3)]^2 U(1) = 0$ ,  $[SU(2)]^2 U(1) = 0$ ,  $[SU(3)]^3 = 0$ , and Witten’s global  $SU(2)$  anomaly (4 doublets, even)—are satisfied for the standard hypercharge assignments with 16 left-handed Weyl fermions per generation (verified by exact arithmetic). The hypercharge values are *uniquely* determined by anomaly cancellation plus charge quantization ( $Q = T_3 + Y$ ), with only one free parameter (overall scale  $y_1 = 1/6$ ). The 16th fermion—the right-handed neutrino  $\nu_R$  as a gauge singlet  $(1, 1, 0)$ —is required for anomaly cancellation and enables the seesaw mechanism.

**Why chirality is hard.** The 600-cell lives on  $S^3$ , which is odd-dimensional. Two no-go results apply: (i) The simplicial Dirac operator  $D = d + d^*$  has index zero on  $S^3$  (the Euler characteristic vanishes:  $\chi(S^3) = 0$ ). No choice of combinatorial Dirac operator can produce a nonzero index on a triangulated  $S^3$ . (ii) In the continuum, the chirality operator  $\gamma_5 = \gamma^0 \gamma^1 \gamma^2 \gamma^3$  requires four spacetime dimensions. On the 3-sphere, there is no intrinsic  $\gamma_5$ ; the Atiyah–Singer index theorem gives  $\text{ind}(\not{D}) = 0$  for any Dirac operator on  $S^3$ . These are *topological* obstructions, not limitations of our framework.

**Possible resolution.** The Hopf fibration  $S^3 \rightarrow S^2$  with  $S^1$  fibers provides a natural lift from 3 to 4 dimensions: the total space  $S^3$  is the principal  $U(1)$  bundle over  $S^2$ , and the extra  $S^1$  direction could serve as the “time-like” dimension needed for  $\gamma_5$ . The Hopf quadrupole  $T_3 = a^2 + b^2 - c^2 - d^2$  on  $S^3$  produces six distinct values  $\pm\phi/2, \pm 1/2, \pm 1/(2\phi)$  on the 96 fermionic vertices, with 48 positive and 48 negative—the correct counting for chirality.

**Status.** Three ingredients for chirality on the *manifold* are in place: (i) the Galois grading provides an exact  $48 + 48$  split with the correct fermion count  $48 = 16 \times N_{\text{gen}}$ ; (ii) anomaly cancellation is verified for all six conditions with the standard hypercharge assignments, and the 16th fermion  $\nu_R$  is uniquely required; (iii) the spectral structure (the Hopf quadrupole  $T_3$ , with 16 vertices per level and 3 levels per sign) reproduces the generation structure  $48 = 3 \times 16$ .

A fourth ingredient now comes from the *internal* space. The McKay graph of  $2I$

is a tree (9 nodes, 8 edges), hence bipartite. The unique bipartition assigns chirality  $\gamma_F = +1$  (WHITE) to the five integer-spin irreps  $\{\rho_1(1), \rho_4(3), \rho_5(3), \rho_6(4), \rho_8(5)\}$  with total dimension 16, and  $\gamma_F = -1$  (BLACK) to the four half-integer-spin irreps  $\{\rho_2(2), \rho_3(2), \rho_7(4), \rho_9(6)\}$  with total dimension 14. Crucially:

$$\dim(\text{WHITE}) = 16 = (a_1 - 1)^2 = \text{fermions per SM generation.} \quad (14)$$

The operator  $\gamma_F$  satisfies  $\gamma_F^2 = 1$  and  $\{\gamma_F, D_F\} = 0$  exactly (a consequence of the tree/bipartite structure: every McKay edge connects WHITE to BLACK). This identification  $\gamma_F = (-1)^{2j}$ , where  $j$  is the  $SU(2)$  spin of the  $2I$  irrep, holds for all 9 irreps. The representation ring  $\text{Rep}(2I)$  respects the  $\mathbb{Z}/2$  grading:  $\text{WHITE} \otimes \text{WHITE} = \text{WHITE}$ ,  $\text{BLACK} \otimes \text{BLACK} = \text{WHITE}$ ,  $\text{WHITE} \otimes \text{BLACK} = \text{BLACK}$  (verified for all  $9 \times 9 = 81$  tensor products using  $A_5$  decomposition).

Taking  $T_3(\text{WHITE}) = -\frac{1}{2}$  and  $T_3(\text{BLACK}) = +\frac{1}{2}$  reproduces the SM left-handed weak isospin  $T_{3,L}$  for 8 out of 9 fermions; the single failure ( $s$  quark) is structural, arising from the  $\mu$ - $s$  degeneracy at the  $w = 0$  McKay edge. The  $SU(2)$  Casimir sum  $\sum C_2 = 26 = a_1^2 + 1$  (Eq. 41) connects  $\gamma_F$  to the Weinberg angle.

This internal chirality  $\gamma_F$ , combined with the product structure, yields the Standard Model chiral gauge coupling.

**Theorem 2** (Chiral gauge coupling). *In the product spectral triple  $(\mathcal{A}, \mathcal{H}_M \otimes \mathcal{H}_F, D, \gamma)$ , the Standard Model chirality pattern— $SU(2)$  chiral,  $SU(3)$  and  $U(1)$  vector-like—is a consequence of the McKay graph structure.*

*Proof.* The gauge group factors correspond to McKay irreps near the trivial node (Theorem 3):  $U(1) \leftrightarrow \rho_1$  (dim 1, spin 0, WHITE),  $SU(2) \leftrightarrow \rho_2$  (dim 2, spin  $\frac{1}{2}$ , BLACK),  $SU(3) \leftrightarrow \rho_4$  (dim 3, spin 1, WHITE). For each irrep  $\sigma$ , the tensor product operator  $T_\sigma : \mathcal{H}_F \rightarrow \mathcal{H}_F$  maps  $V_{\rho_k} \rightarrow V_{\rho_k \otimes \sigma}$ . Since  $\gamma_F$  respects the  $\mathbb{Z}/2$  tensor-product grading (verified for all 81 products):

- $\gamma_F(\sigma) = +1$  (WHITE): every constituent of  $\rho_k \otimes \sigma$  has  $\gamma_F = \gamma_F(\rho_k)$ , so  $[T_\sigma, \gamma_F] = 0$  (vector-like coupling).
- $\gamma_F(\sigma) = -1$  (BLACK): every constituent of  $\rho_k \otimes \sigma$  has  $\gamma_F = -\gamma_F(\rho_k)$ , so  $\{T_\sigma, \gamma_F\} = 0$  (chiral coupling).

The McKay adjacency  $D_F$  encodes  $T_{\rho_2}$  (the standard representation, spin  $\frac{1}{2}$ , BLACK):  $\{D_F, \gamma_F\} = 0$  gives chiral  $SU(2)$  coupling. The  $SU(3)$  direction  $T_{\rho_4} \sim D_F^2$  and the  $U(1)$  direction  $T_{\rho_1} = \text{Id}$  both commute with  $\gamma_F$ , giving vector-like coupling. Robustness: both dimension-2 irreps ( $\rho_2, \rho_3$ ) are BLACK and both dimension-3 irreps ( $\rho_4, \rho_5$ ) are WHITE, so the Galois choice  $\rho_2 \leftrightarrow \rho_3, \rho_4 \leftrightarrow \rho_5$  does not affect the result.  $\square$

**Corollary 2.** *In the physical Hilbert space  $\mathcal{H}_+ = \{\psi : \gamma\psi = \psi\}$ , where  $\gamma = \gamma_{\text{form}} \otimes \gamma_F$ :*

$$\mathcal{H}_+ = \underbrace{(\mathcal{H}_{\text{even}} \otimes \mathcal{H}_W)}_{\text{right-handed, } SU(2) \text{ singlet}} \oplus \underbrace{(\mathcal{H}_{\text{odd}} \otimes \mathcal{H}_B)}_{\text{left-handed, } SU(2) \text{ doublet}}.$$

*Left-handed fermions occupy the BLACK sector ( $SU(2)$  doublets) and right-handed fermions the WHITE sector ( $SU(2)$  singlets), with  $\dim(\mathcal{H}_W) = 16$  fermions per generation.*

*Remark.* The obstacle to chirality on  $S^3$  alone is topological ( $\text{ind}(D) = 0$ ). The theorem shows that the chiral mechanism resides entirely in the *finite* space  $F$ —specifically in the bipartite structure of the McKay graph—and is therefore independent of the manifold dimension. The identification  $\gamma_{\text{form}} = (-1)^p$  is a theorem about the simplicial Dirac operator  $d + d^*$ , not an assumption about the spinor  $\gamma_5$ ; the continuum limit (whether  $\gamma_{\text{form}}$  reduces to  $\gamma_5$  in 4D) remains open (Section 14).

*Remark (Lorentzian signature).* The McKay graph structure permits Lorentzian signature through the KO-dimension mechanism. On the edge space ( $\dim 120 = |2I|$ ), the charge conjugation  $J$  satisfies  $J^2 = \nu(\rho_i) \nu(\rho_j)$  on each edge  $(\rho_i, \rho_j)$ , where  $\nu$  is the Frobenius–Schur indicator. Since all edges connect nodes of opposite dimension parity (bipartite graph),  $J^2$  splits into a dominant sector ( $J^2 = -1$ , six edges,  $\dim 88$ ) and a subdominant sector ( $J^2 = +1$ , two edges,  $\dim 32$ ). Both sectors yield  $\varepsilon' = +1$  (from graph symmetry) and  $\varepsilon'' = -1$  (from chirality flip under  $J$ ), giving KO-dimension 2 or 6 respectively. In either case, the Poincaré duality condition  $\text{KO}(M \times F) \equiv 0 \pmod{8}$  forces the spacetime manifold  $M$  to have Lorentzian signature  $(1, 3)$ , making the result robust under sector selection. This parallels the algebra choice  $\mathcal{A}_F = \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C})$  in standard NCG, where the signature also follows from KO-dimension matching rather than being postulated independently.

## 4.1 From Discrete Symmetry to Continuous Gauge Group

The decomposition (12) shows that the  $A_5$ -module structure of the vertex figure *selects* the Standard Model gauge group from the classification of compact Lie groups. We now prove this rigorously, addressing the natural objection that a finite group cannot generate a continuous gauge symmetry.

**Theorem 3** (Gauge group identification). *Let  $G$  be a compact Lie group with  $\dim(G) = 12$ , and let  $A_5 \subset G$  be a subgroup such that*

$$\text{ad}(G)|_{A_5} = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5} \quad (15)$$

*as  $A_5$ -modules. Then  $G \cong U(1) \times SU(2) \times SU(3)$ .*

*Proof.* The trivial summand  $\mathbf{1}$  in (15) corresponds to an abelian factor:  $\mathfrak{g} = \mathfrak{u}(1) \oplus \mathfrak{h}$  where  $\mathfrak{h}$  is semisimple with  $\dim(\mathfrak{h}) = 11$  and  $\text{ad}(\mathfrak{h})|_{A_5} = \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5}$ . Write  $\mathfrak{h} = \mathfrak{h}_1 \oplus \cdots \oplus \mathfrak{h}_k$  as a direct sum of simple compact Lie algebras. The dimensions of the simple compact Lie algebras with  $\dim \leq 11$  are:

$$\mathfrak{su}(2) : 3, \quad \mathfrak{su}(3) : 8, \quad \mathfrak{sp}(2) : 10 \quad (16)$$

(and  $\mathfrak{g}_2 = 14$ ,  $\mathfrak{su}(4) = 15$ , etc.). The *only* partition of 11 into such dimensions is  $11 = 3+8$ , giving  $\mathfrak{h} = \mathfrak{su}(2) \oplus \mathfrak{su}(3)$  and hence  $G = U(1) \times SU(2) \times SU(3)$ . (The partition  $3+3+5$  fails because no simple Lie algebra has dimension 5;  $10+1$  fails because  $\mathfrak{sp}(2)$  is simple, not  $\mathfrak{sp}(2) \oplus \mathfrak{u}(1)$  with  $\dim 11$ .)

For compatibility with (15), note that  $A_5 \subset SO(3) = SU(2)/\mathbb{Z}_2$  via the icosahedral rotation group, giving  $\text{ad}(\mathfrak{su}(2))|_{A_5} = \mathbf{3}$  (or  $\mathbf{3}'$ ). Meanwhile, the tensor product  $\mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{5}$  (verified by character inner products) gives  $\text{ad}(\mathfrak{su}(3))|_{A_5} = \mathbf{3} \otimes \mathbf{3}^* - \mathbf{1} = \mathbf{3} \oplus \mathbf{5}$  (or  $\mathbf{3}' \oplus \mathbf{5}$ ), using the Frobenius–Schur indicator  $\nu(\mathbf{3}) = +1$  (real/orthogonal type, so  $\mathbf{3}^* \cong \mathbf{3}$ ). The two valid assignments are exchanged by the Galois automorphism  $\sigma : \sqrt{5} \rightarrow -\sqrt{5}$ , and  $\phi > 0$  selects a unique one.  $\square$

The bridge from discrete to continuous symmetry operates through three complementary mechanisms: (1) the representation-theoretic decomposition (10) *identifies* the gauge group uniquely (Theorem 3); (2) Burnside’s theorem shows the discrete symmetries *algebraically span* the full gauge Lie algebra (see Supplementary Material S2); (3) the spectral action (Section 9.7) provides the Yang–Mills *dynamics* with continuous gauge invariance. The hypothesis  $\dim(G) = 12$  is not external input: the vertex degree  $d = 2(a_1 + 1) = 12$  determines it.

*Remark* (Distler–Garibaldi). The impossibility theorem of [17] does *not* apply here: the gauge group is identified via the  $A_5$  decomposition, and fermions arise from  $2I$  irreps—not from  $E_8$  representations. The  $E_8$  structure enters exclusively through the McKay correspondence (see Supplementary Material S2 for details).

## 4.2 Edge Gauge Spectrum: Second Derivation

The vertex-figure argument of Section 4 identifies the gauge group from the local geometry around each vertex. We now give an independent, *global* derivation from the spectrum of the wave operator  $\square$  lifted to the edge space.

**The line-graph operator.** Let  $L(G)$  be the line graph of the 600-cell: its 720 nodes are the edges of the original graph, connected when they share a vertex. The Hopf fibration classifies these 720 edges as 120 *fiber edges* (both endpoints in the same Hopf fiber) and 600 *cross edges*. In  $L(G)$ , the fiber adjacency  $A_{\text{fiber}}$  connects consecutive fiber edges within each  $C_{10}$  cycle. Define

$$\square_1 = L_{\text{cross}} - a_1 L_{\text{fiber}} \quad (17)$$

on  $L(G)$ , in exact analogy with  $\square_0 = L_{\text{cross}} - a_1 L_{\text{fiber}}$  on the vertex graph. The matrix  $\square_1$  is  $720 \times 720$  with *integer* entries.

**Theorem 4** (Edge gauge spectrum). *The kernel of  $\square_1$  has dimension  $13 = (a_1^2 + 1)/2$  and decomposes under  $2I$  as*

$$\ker(\square_1) = \rho_0 \oplus 2\rho_5 \quad (18)$$

where  $\rho_0$  is the trivial representation ( $\dim 1$ ) and  $\rho_5$  is the 6-dimensional irrep corresponding to the branching node of the affine  $E_8$  Dynkin diagram.

*Proof.* The matrix  $\square_1$  has integer entries (verified), so  $\ker(\square_1) = 720 - \text{rank}(\square_1) = 720 - 707 = 13$  is exact. For the irrep decomposition, the character is computed on each of the 9 conjugacy classes of  $2I$ :  $\chi_{\ker}(1) = 13$ ,  $\chi_{\ker}(-1) = -11$ . Applying the decomposition formula  $n_\rho = \frac{1}{|2I|} \sum_g \chi_\rho(g) \chi_{\ker}(g)$  yields  $n_{\rho_0} = 1$  and  $n_{\rho_5} = 2$ , with all other multiplicities zero. The dimension check  $1 + 2 \times 6 = 13$  confirms the decomposition.  $\square$

The 12 gauge modes (the  $2\rho_5$  subspace) have a striking geometric character:

**Proposition 1** (Fiber localisation and Hodge eigenvalues). *The 12-dimensional gauge subspace  $2\rho_5 \subset \ker(\square_1)$  satisfies:*

- (i) **Fiber localisation:** *the 12 modes have support exclusively on the 120 fiber edges (cross-edge weight  $< 10^{-29}$ );*
- (ii) **Coexact eigenvalue:**  $C|_{2\rho_5} = a_1 \mathbf{I}_{12}$ , *where  $C = d_1^\top d_1$  is the coexact Hodge Laplacian;*

(iii) **Exact eigenvalue:**  $B|_{2\rho_5} = \frac{16}{a_1} \mathbf{I}_{12}$ , where  $B = d_0 d_0^\top$  is the exact Hodge Laplacian;

(iv) **Trace identity:**  $\text{Tr}(\square_1^2) / \text{Tr}(\square_0^2) = 9 a_1 = 45$ .

*Proof.* Each claim is verified by direct computation on the 720-dimensional edge space (see `reproducible/verify_edge_gauge_spectrum.py`). For (i), the  $\rho_0$ -projector is built from the group average; the complementary 12-dimensional subspace is then tested against the fiber projector  $P_{\text{fiber}}$ , giving fiber fraction 1.000... to machine precision. For (ii)–(iii), the Hodge operators  $B = d_0 d_0^\top$  and  $C = d_1^\top d_1$  are restricted to the gauge subspace and diagonalised; all 12 eigenvalues equal 5 and 16/5 respectively. For (iv),  $\text{Tr}(\square_1^2) = 324,000$  and  $\text{Tr}(\square_0^2) = 7,200$ .  $\square$

The 12 Hopf fibers project (via  $2I \rightarrow A_5$ ) to 12 points forming a 5-regular graph on  $S^2$  with Laplacian eigenvalues  $\{0, 5-\sqrt{5}, 6, 5+\sqrt{5}\}$  and multiplicities  $\{1, 3, 5, 3\}$ .

**Corollary 3** (Gauge algebra from edge spectrum). *The  $A_5$  permutation representation on the 12 Hopf fibers decomposes as*

$$12 = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5} \quad (19)$$

*reproducing the vertex-figure decomposition (10) and hence the gauge algebra identification  $1 + 3 + 8 = \mathfrak{u}(1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(3)$  of Theorem 3.*

*Proof.* The central element  $-1 \in 2I$  acts trivially on fibers (verified for all 12), so the fiber permutation factors through  $A_5 = 2I/\{\pm 1\}$ . The character is computed from fixed-fiber counts and decomposed using the irrep character table; the multiplicities are  $n_{\rho_0} = n_{\rho_2} = n_{\rho_4} = n_{\rho_6} = 1$ , giving dimensions  $1 + 3 + 5 + 3 = 12$ . The grouping  $\mathbf{3}' \oplus \mathbf{5} = \text{ad}(\mathfrak{su}(3))|_{A_5}$  follows from Theorem 3.  $\square$

*Remark* (Scope). The edge spectrum reproduces the gauge *dimensions*  $1 + 3 + 8$  through the  $A_5$  fiber decomposition, independently confirming Theorem 3. The Lie bracket itself does not emerge at the discrete level:  $\Lambda^2(\rho_5)$  does not contain  $\rho_5$  as a  $2I$ -submodule, and the simplicial wedge product vanishes on fiber-localised modes (every triangle has at most one fiber edge). This is consistent with standard lattice gauge theory, where the continuous gauge algebra appears only in the continuum limit; the spectral action (Section 9.7) supplies the Yang–Mills dynamics.

*Remark* (The  $\square$  hierarchy). Extending  $\square$  to the full simplicial complex via the face adjacency graph (1200 nodes, degree 12, fiber degree 4) yields a hierarchy of integer operators:

| Space                | dim  | $\ker(\square_p)$                          | Physical content                        |
|----------------------|------|--------------------------------------------|-----------------------------------------|
| Vertices ( $p = 0$ ) | 120  | $9 = \rho_0 \oplus 2\rho_1 \oplus 2\rho_8$ | $c^2 = a_1$ , photon, bootstrap         |
| Edges ( $p = 1$ )    | 720  | $13 = \rho_0 \oplus 2\rho_5$               | 12 gauge bosons ( $E_8$ branching node) |
| Faces ( $p = 2$ )    | 1200 | $1 = \rho_0$                               | vacuum singlet                          |

The vertex kernel sits at the *endpoints* of the  $E_8$  Dynkin diagram (the Galois pair  $\rho_1, \rho_8$ ); the edge kernel sits at the *branching node* ( $\rho_5$ ); the face kernel reduces to the trivial representation alone. All three operators share the pattern  $\ker \ni \rho_0$  (a universal vacuum mode). The gauge content  $1 + 3 + 8 = 12$  emerges exclusively at  $p = 1$ , while  $c^2 = a_1$  is a vertex-level identity. The trace ratios  $\text{Tr}(\square_1^2) / \text{Tr}(\square_0^2) = 9 a_1 = 45$  and  $\text{Tr}(\square_2^2) / \text{Tr}(\square_0^2) = 34$  are exact integers.



## 5 Coupling Constants from $a_1$

### 5.1 The Fine Structure Constant

Both non-trivial coefficients of the  $\alpha$  equation arise from the *icosahedral Laplacian*. The vertex figure of the 600-cell is an icosahedron (12 vertices, degree  $a_1 = 5$ ). Its graph Laplacian  $L = a_1 I - A_{\text{ico}}$  decomposes under  $A_5$  into eigenvalues

$$L(\mathbf{1}) = 0, \quad L(\mathbf{3}) = a_1 - \sqrt{a_1}, \quad L(\mathbf{5}) = b_1, \quad L(\mathbf{3}') = a_1 + \sqrt{a_1} \quad (20)$$

These follow from standard spectral graph theory on Cayley graphs: for each irrep  $\rho$  of  $A_5$ , the Laplacian eigenvalue is  $L(\rho) = d - \chi_\rho(S)/\dim(\rho)$ , where  $\chi_\rho(S)$  is the character summed over the generating set (nearest-neighbor rotations) and  $d = a_1$  is the vertex degree. The two 3-dimensional irreps  $\mathbf{3}$  and  $\mathbf{3}'$  have characters  $\phi$  and  $\phi' = -1/\phi$  on the generating class, giving  $L(\mathbf{3}) = 5 - \sqrt{5}$  and  $L(\mathbf{3}') = 5 + \sqrt{5}$ . These are related by the Galois automorphism  $\phi \leftrightarrow \phi'$  (Section 8 derives additional consequences of this spectrum).

**Tree-level coupling.** The product of the Galois-conjugate eigenvalues is

$$L(\mathbf{3}) \cdot L(\mathbf{3}') = (a_1 - \sqrt{a_1})(a_1 + \sqrt{a_1}) = a_1^2 - a_1 = a_1(a_1 - 1) = 4a_1 \quad (21)$$

where the last step uses  $a_1 - 1 = 4$  (specific to  $a_1 = 5$ , the same condition underlying the Higgs mass ratio). The eigenvalue ratio is  $L(\mathbf{3}')/L(\mathbf{3}) = (a_1 + \sqrt{a_1})/(a_1 - \sqrt{a_1}) = \phi^2$  (holding if and only if  $a_1 = 5$ ; see Section 8 for the Higgs mass consequence).

*Geometric identification.* The Cayley graph of  $2I$  (the 600-cell 1-skeleton) has eigenvalues  $\lambda_k = \chi_{\text{std}}(k) = \phi$  for the  $\rho_1$  representation. The spectral gap diffusion factor  $e^{-t\lambda_1}$  at unit lattice time gives  $\phi^{-1}$  per step, so the  $a_1 = 5$  nearest-neighbor interaction on the icosahedral vertex figure contributes  $\phi^{-a_1+1} = \phi^{-4}$ . The tree-level coupling combines the Galois product with this spectral factor:

$$\frac{1}{\alpha_0} = L(\mathbf{3}) L(\mathbf{3}') \cdot \phi^4 = 4a_1 \phi^4 \quad (22)$$

The identification of  $\phi^4$  with the spectral diffusion factor remains **structural**: the eigenvalue product  $4a_1$  and the scaling  $\phi^4$  are both individually determined by the icosahedral Laplacian, but the interpretation of the full combination  $L(\mathbf{3})L(\mathbf{3}') \cdot \phi^4$  as the inverse coupling strength still uses the Kaluza–Klein step discussed below. What is no longer merely algebraic is the denominator  $b_1$ : it has an exact Yang–Mills interpretation on the simplicial complex (Proposition 2).

**Product identity.** Including the singlet eigenvalue  $L(\mathbf{5}) = b_1$ :

$$L(\mathbf{3}) \cdot L(\mathbf{5}) \cdot L(\mathbf{3}') = b_1(a_1^2 - a_1) = b_1 \cdot a_1(a_1 - 1) = 6 \cdot 5 \cdot 4 = 120 = N \quad (23)$$

This is equivalent to  $a_1(a_1^2 - 1) = a_1!$ , which holds *only* for  $a_1 = 5$  (and trivially  $a_1 = 1$ ). Here the equality  $L(\mathbf{3})L(\mathbf{5})L(\mathbf{3}') = 120$  is a direct algebraic identity for the *distinct* nonzero icosahedral Laplacian eigenvalues; it is *not* an application of the Kirchhoff matrix-tree theorem, which uses the full Laplacian spectrum with multiplicities. This identity allows rewriting the linear coefficient as  $4a_1 = N/b_1$ , so the tree-level coupling becomes  $1/\alpha_0 = (N/b_1) \phi^4$ .

**Proposition 2** (Discrete Yang–Mills stiffness of the Hopf twist). *Let  $d_0(\theta)$  be the simplicial  $U(1)$  connection obtained by assigning the phase  $e^{i\theta/10}$  to each oriented fiber edge of a Hopf decagon and 1 to each cross-fiber edge, and let  $F(\theta) = d_1 d_0(\theta)$ . Then:*

- (i)  $\|F(\theta)\|^2 = \text{Tr}(F(\theta)^* F(\theta))$  is gauge-invariant;
- (ii) only the  $T_1 = 600$  triangles with exactly one fiber edge contribute;
- (iii) the Yang–Mills action has the exact closed form

$$S_{\text{YM}}(\theta) \equiv \|F(\theta)\|^2 = 4T_1 \sin^2\left(\frac{\theta}{2|\text{fiber}|}\right) = 2400 \sin^2\left(\frac{\theta}{20}\right), \quad (24)$$

with  $|\text{fiber}| = 10 = 2a_1$ ;

- (iv) its quadratic stiffness is

$$S_{\text{YM}}(\theta) = K_{\text{YM}} \theta^2 + O(\theta^4), \quad K_{\text{YM}} = \frac{T_1}{|\text{fiber}|^2} = \frac{600}{100} = 6 = b_1. \quad (25)$$

*Interpretation.* Proposition 2 upgrades the denominator in the tree-level coefficient from a bare algebraic symbol to an exact discrete gauge-theoretic quantity:  $b_1 = K_{\text{YM}}$ . The count  $T_1 = 600$  and therefore the stiffness  $K_{\text{YM}} = 6$  are the same for all Hopf fibrations of the 600-cell tested in the discrete construction, so this is a fibration-stable invariant rather than an artifact of a particular ordering. Therefore

$$\frac{1}{\alpha_0} = \frac{N}{K_{\text{YM}}} \phi^4 = \frac{N}{b_1} \phi^4. \quad (26)$$

The residual structural step is the identification of the spectral factor  $\phi^4$  with the tree-level diffusion weight; once that is made, the denominator is fixed by the Yang–Mills action of the simplicial Hopf twist.

**Fully spectral form of the tree-level coefficient.** The same factor  $\phi^4$  can be written in terms of the Hopf-fiber spectral gap. For the decagonal fiber graph,

$$\lambda_1(L_{\text{fiber}}) = 2 - \phi = \frac{1}{\phi^2}, \quad (27)$$

so

$$\phi^4 = \frac{1}{\lambda_1(L_{\text{fiber}})^2}. \quad (28)$$

Hence the tree-level coefficient admits the purely discrete spectral expression

$$\frac{1}{\alpha_0} = \frac{N}{K_{\text{YM}} \lambda_1(L_{\text{fiber}})^2} = \frac{120}{6(2 - \phi)^2} = 4a_1 \phi^4. \quad (29)$$

All three ingredients are therefore internal to the 600-cell with Hopf fibration:  $N = |2I|$  is combinatorial,  $K_{\text{YM}} = b_1$  is gauge-theoretic, and  $\lambda_1(L_{\text{fiber}})$  is spectral.

**Spectral action origin.** The tree-level coefficient also admits a direct spectral action characterization. Project  $\square$  to the gauge sector (fiber zero mode) via  $\square_{\text{gauge}} = P_0 \square P_0$ , where  $P_0$  averages over each Hopf fiber. The resulting  $12 \times 12$  operator on the icosahedral base has eigenvalues  $\{0^{(\times 1)}, 10 - 2\sqrt{5}^{(\times 3)}, 12^{(\times 5)}, 10 + 2\sqrt{5}^{(\times 3)}\}$ —*exactly* the Laplacian eigenvalues  $L(\rho_2), L(\rho_4), L(\rho_6)$  of the 600-cell restricted to the base. The Frobenius norm satisfies  $\text{Tr}(\square_{\text{gauge}}^2) = 1440 = \deg \times N = 12 \times 120$ , giving

$$\frac{1}{\alpha_0} = \frac{\text{Tr}(\square_{\text{gauge}}^2)}{n_{\text{base}} b_1 \lambda_1(L_{\text{fiber}})^2} = \frac{1440}{12 \cdot 6 \cdot \phi^{-4}} = \frac{N}{b_1} \phi^4 = 4a_1 \phi^4. \quad (30)$$

The gauge-sector spectral action on  $\square$  thus produces the tree-level coupling directly, with the ratio  $\text{Tr}(\square_{\text{full}}^2)/\text{Tr}(\square_{\text{gauge}}^2) = a_1$  confirming that the gauge sector is exactly  $1/a_1$  of the full spectrum.

**Holonomy coefficient.** The sum of the same eigenvalues equals

$$L(\mathbf{3}) + L(\mathbf{3}') = 2a_1 = 10 \quad (31)$$

which is precisely the number of edges in a Hopf fiber (the decagonal geodesic on the 600-cell). The angle  $\pi/a_1$  per edge is *derived* from the spectral gap of the fiber graph Laplacian: the  $C_{2a_1}$  cycle has smallest nonzero eigenvalue  $\lambda_1 = 2(1 - \cos(\pi/a_1)) = 1/\phi^2$ , and inverting the dispersion relation gives

$$\theta_{\text{edge}} = \arccos(1 - \frac{\lambda_1}{2}) = \arccos(\frac{\phi}{2}) = \frac{\pi}{a_1} \quad (32)$$

where the identity  $\cos(\pi/a_1) = \phi/2$  is *specific to*  $a_1 = 5$  (the golden ratio angle; for no other integer  $a_1$  does the spectral gap of  $C_{2a_1}$  equal  $1/\phi^2$ ). The total  $U(1)$  holonomy around the fiber is  $(2a_1) \cdot (\pi/a_1) = 2\pi$ . The holonomy correction to the coupling is the product of the fiber edge sum, the per-edge angle, and the coupling itself:

$$\delta\left(\frac{1}{\alpha}\right) = -[L(\mathbf{3}) + L(\mathbf{3}')] \frac{\pi}{a_1} \cdot \alpha = -2\pi \alpha \quad (33)$$

**Algebraic self-consistency.** The equation  $1/\alpha = 1/\alpha_0 - 2\pi\alpha$  is an algebraic relation whose three coefficients each have a geometric origin (tree-level product, Hopf holonomy, and normalization). It is *not* a perturbative QFT one-loop calculation—no Feynman diagrams or renormalization are involved. The holonomy  $2\pi$  is an exact geometric fact (Eq. 32); its entrance into the coupling equation is motivated by the Kaluza–Klein identification of  $\alpha$  with a  $U(1)$  gauge coupling on the Hopf fiber (see below). Substituting (22) yields:

$$\boxed{2\pi\alpha^2 - 4a_1\phi^4\alpha + 1 = 0} \quad (34)$$

The constant term equals 1 by the normalization  $\alpha \cdot (1/\alpha) = 1$ , which fixes the equation's scale. By Vieta's formulas, the product of the two roots is  $\alpha\alpha' = 1/(2\pi)$ —a universal topological constant independent of  $a_1$ .

Using the product identity (23), the  $\alpha$  equation takes the manifestly geometric form:

$$\text{Vol}(S^1)\alpha^2 - \frac{N}{b_1}\phi^4\alpha + 1 = 0 \quad (35)$$

where  $\text{Vol}(S^1) = 2\pi$  is the Hopf fiber holonomy. Each coefficient has a distinct geometric origin: the quadratic term is *topological* (first Chern class  $c_1 = 1$  of the Hopf bundle), the linear term is *spectral* (Cayley eigenvalue product), and the constant is *normalization*.

**Kaluza–Klein interpretation.** The Vieta product  $\alpha\alpha' = 1/(2\pi) = 1/\text{Vol}(S^1)$  is precisely the standard Kaluza–Klein result for a  $U(1)$  gauge coupling on a fiber of volume  $\text{Vol}(S^1)$ : the gauge coupling squared equals  $1/\text{Vol}(\text{fiber})$ . This identification is structural—it follows from the Hopf fibration  $S^3 \rightarrow S^2$  with  $c_1 = 1$ —and is tightly constrained: an exhaustive scan over algebraic alternatives (replacing  $2\pi$  by elements of  $\mathbb{Z}[\phi]$  with coefficients  $\leq 200$ ) shows that no purely algebraic coefficient reproduces  $1/\alpha$  to experimental precision; the transcendental factor  $2\pi$  is the unique value consistent with the data at the  $10^{-4}\%$  level, confirming its topological necessity. The two roots  $\alpha$  and  $\alpha'$

correspond to the two orientations of the  $U(1)$  fiber; their product  $1/(2\pi)$  is a topological invariant (depending only on  $c_1 = 1$ ), while their sum  $B/(2\pi)$  encodes the spectral data.

**Fiber–Cayley duality.** Both non-trivial coefficients are natural operations on the *same* Galois-conjugate pair of Laplacian eigenvalues: the linear coefficient is their *product*  $L(\mathbf{3})L(\mathbf{3}') = a_1^2 - a_1 = 4a_1$  (spectral/metric), while the quadratic coefficient involves their *sum*  $L(\mathbf{3}) + L(\mathbf{3}') = 2a_1 = 10$  (topological/holonomy). The sum counts Hopf fiber edges; the product gives the tree-level coupling strength. The Hopf fibration decomposes the 720 edges of the 600-cell into 120 *fiber* edges (12 fibers  $\times$  10 edges) and 600 *cross-fiber* edges, with ratio  $600/120 = a_1$ . Each vertex has 2 fiber neighbors and  $2a_1 = 10$  cross-fiber neighbors, summing to  $2 + 2a_1 = 12 = \dim(G)$ .

**Derivation status.** The three coefficients of (34) decompose as follows: the quadratic term  $2\pi$  is **topological** (the holonomy  $2\pi$  is an exact consequence of  $c_1 = 1$  for the Hopf bundle; the Vieta constraint  $\alpha\alpha' = 1/(2\pi c_1)$  is the unique normalization compatible with charge quantization on a fiber with first Chern class 1; its role as the coupling coefficient follows from the Kaluza–Klein identification, which is the sole structural postulate—not a first-principles lattice derivation, but tightly constrained by the topological necessity of  $\pi$ ); the linear term  $4a_1\phi^4$  is **spectral**: Eq. (30) derives it as  $\text{Tr}(\Box_{\text{gauge}}^2)/[n_{\text{base}} b_1 \lambda_1^{-2}]$ , where every factor is computable from the wave operator  $\Box$  and the Hopf fiber projection—the same spectral data that produce  $c^2 = a_1$ , the graviton propagator, and the Seeley–DeWitt coefficients; the constant term 1 is **normalization** (Vieta). The equation is a geometric self-consistency relation on the 600-cell lattice; the structural parallel with one-loop QED vacuum polarization ( $1/\alpha = 1/\alpha_0 - \delta$ ) is suggestive but the origin is topological (fiber holonomy), not perturbative. The only transcendental input is  $\pi$ , which reflects the topology of  $S^3$  and places  $\alpha$  in the number field  $\mathbb{Z}[\phi, \pi]$ . An exhaustive search over simple formulas built from the discrete invariants of  $2I$  (subgroup indices, irrep dimensions, character sums, Laplacian traces) confirms that all *algebraic* coefficients of the  $\alpha$  equation are already present in the discrete data, while the factor  $2\pi$  constitutes an irreducible transcendental/topological input.

$$\alpha(0) = \frac{4a_1\phi^4 - \sqrt{16a_1^2\phi^8 - 8\pi}}{4\pi} = \frac{1}{137.036\dots} \quad (36)$$

This is  $\alpha$  at zero momentum transfer (Thomson limit). The running of  $\alpha$  with energy is *partially* derived: one-loop QED vacuum polarization with framework-derived fermion masses and beta coefficients ( $b_1, b_2, b_3$  all determined by  $N_{\text{gen}} = 3$  and the gauge group) gives a self-consistent ratio  $\alpha(m_Z)/\alpha(0) = 1.084$  (experimental: 1.071, error 1.2%). The residual 1.2% reflects the standard limitation of one-loop QED (missing hadronic vacuum polarization and higher-order corrections), not a framework-specific gap. Where sub-percent precision is needed (Section 8), we use the experimental ratio.

|                  | Predicted | CODATA 2022         |
|------------------|-----------|---------------------|
| $\alpha^{-1}(0)$ | 137.036   | 137.035 999 084(21) |
| <b>Error</b>     |           | <b>0.0001%</b>      |

**Coherence, not just numerics.** A potential objection is that  $B = 4a_1\phi^4 \approx 137.08$  was *chosen* because it approximates  $1/\alpha$ . Two facts argue against this. First, an exhaustive scan of all “natural” combinations  $c a_1^p b_1^q \phi^r$  (with  $c \leq 30$ ,  $p, q \leq 2$ ,  $r \in [-2, 7]$ ), supplemented by  $N$ ,  $h$ ,  $\text{rank}(E_8)$ ,  $\dim(E_8)$ , and  $\pi$ , finds **exactly 8 expressions** in [136, 138],

of which only **3 are within 0.1% of  $1/\alpha$** —all three being rearrangements of the same quantity:  $4a_1\phi^4 = 20\phi^4 = (N/b_1)\phi^4$ .

Second, and more importantly, an isolated numerical coincidence would not propagate coherently through independent sectors of the framework. The *same* spectral ingredients that enter the  $\alpha$  equation reappear elsewhere with different physical roles:

- The eigenvalue ratio  $L(\mathbf{3}')/L(\mathbf{3}) = \phi^2$  determines the Higgs mass (Section 8:  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$ );
- The Vieta product  $\alpha\alpha' = 1/(2\pi)$  is the Kaluza–Klein coupling on the Hopf fiber;
- The eigenvalue product  $L(\mathbf{3})L(\mathbf{3}') = 4a_1$  fixes  $\sin^2\theta_W = b_1/(a_1^2 + 1)$  through the singlet eigenvalue  $L(\mathbf{5}) = b_1$ ;
- The hierarchy exponent  $z_{\text{Planck}} = 4\phi^2$  has trace 12 (vertex degree) and norm 16 (multiplicity of  $\lambda_3$ ).

A fit would have no reason to be coherent with the Higgs mass, the Weinberg angle, or the gravitational hierarchy. The  $\alpha$  equation is not derived from a single variational principle—this remains an open problem—but its three coefficients are individually determined: the quadratic  $2\pi$  by the topology of the Hopf bundle ( $c_1 = 1$ ), the linear  $4a_1\phi^4$  by the spectral geometry of the fiber ( $N/b_1 = 20$ ,  $\lambda_1^{-2} = \phi^4$ ), and the constant 1 by Vieta normalization. The sole residual gap is the Kaluza–Klein identification step (coupling = fiber geometry), which is structural but not yet derived from the spectral action.

## 5.2 The Strong Coupling Constant

We derive  $\alpha_s$  from the requirement that  $1/\alpha_s$  be a “natural” element of the number ring  $\mathbb{Z}[\phi]$ —the same ring in which all 600-cell spectral quantities live. The three conditions below are not arbitrary: (i) fixes the algebraic form, (ii) connects the trace to the  $E_8$  rank via the McKay correspondence (the trace of a  $\mathbb{Z}[\phi]$  element counts its “integer shadow”), and (iii) encodes asymptotic freedom.

**Factor structure.** Every element of  $\mathbb{Z}[\phi]$  with positive norm is a product of a rational integer  $c$  and a unit  $\phi^k$ . We require  $1/\alpha_s$  to have this form, with the trace fixed by the gauge sector:

- (i)  $z = c\phi^k$ ,  $c \in \mathbb{Z}_{>0}$ ,  $k \geq 0$ ,
- (ii)  $\text{Tr}(z) = c\text{Tr}(\phi^k) = \text{rank}(E_8) = 8$ ,
- (iii)  $z' < 0$  (asymptotic freedom: Galois conjugate has opposite sign).

Condition (ii) forces  $c = 8/\text{Tr}(\phi^k)$ , requiring  $\text{Tr}(\phi^k) \mid 8$ . Among the first powers of  $\phi$ :

| $k$      | $\phi^k$                      | Tr       | Tr $\mid 8$ ? | $c$      | $z' < 0$ ?    |                                       |
|----------|-------------------------------|----------|---------------|----------|---------------|---------------------------------------|
| 0        | 1                             | 2        | yes           | 4        | no (rational) | eliminated by (iii)                   |
| 1        | $\phi$                        | 1        | yes           | 8        | yes           | $\alpha_s = 0.077$ (too small)        |
| 2        | $\phi + 1$                    | 3        | no            | —        | —             |                                       |
| <b>3</b> | <b><math>2\phi + 1</math></b> | <b>4</b> | <b>yes</b>    | <b>2</b> | <b>yes</b>    | <b><math>\alpha_s = 0.1180</math></b> |
| 4        | $3\phi + 2$                   | 7        | no            | —        | —             |                                       |

Only  $k = 0, 1, 3$  satisfy (i)–(ii); condition (iii) eliminates  $k = 0$ ; among  $k = 1$  and  $k = 3$ , the latter has the *minimal* integer prefactor ( $c = 2$  vs.  $c = 8$ ), selecting it uniquely. Note that  $\text{Tr}(\phi^3) = 4$  equals the spacetime dimension (the 600-cell lives in  $\mathbb{R}^4$ ), and  $k = 3 = N_{\text{gen}}$ . The norm follows as a consequence:

$$|N(z)| = c^2 \cdot |N(\phi^3)| = 2^2 \times 1 = 4 = \frac{N}{h} = \frac{120}{30} \quad (37)$$

reproducing the number of Coxeter orbits of  $H_4$ .

**Proposition 3** (Algebraic uniqueness). *The element  $z = 2\phi^3 = 2 + 4\phi$  is the unique positive element of  $\mathbb{Z}[\phi]$  with  $\text{Tr}(z) = 8$  and  $N(z) = -4$ .*

*Proof.* The conditions give  $z^2 - 8z - 4 = 0$ , with discriminant  $80 = 16 \times 5$ . Since  $\sqrt{5} = 2\phi - 1 \in \mathbb{Z}[\phi]$ , the positive root is  $z = 4 + 2\sqrt{5} = 2 + 4\phi$ . The other root  $6 - 4\phi < 0$ . (The case  $N = +4$  gives  $z = 4 \pm 2\sqrt{3}$ , but  $\sqrt{3} \notin \mathbb{Z}[\phi]$ .)  $\square$

**Physical interpretation.** The trace condition ties  $\alpha_s$  to the gauge sector ( $\text{Tr} = \dim(SU(3)) = \text{rank}(E_8)$ ). The factor  $c = 2 = \text{rank}(E_8)/\text{Tr}(\phi^3)$  is the ratio of the gauge rank to the spacetime dimension. The negative norm  $N = -4$  means  $z' < 0$ : the Galois conjugate sector has opposite sign, reflecting asymptotic freedom of  $SU(3)$ .

**Hierarchy identity.** The  $\alpha_s$  exponent  $z = 2\phi^3$  and the Planck exponent  $z_{\text{Planck}} = 4\phi^2$  (derived in Section 8.4) are related by an exact algebraic identity:

$$\boxed{z_{\text{Planck}} = \frac{1}{\alpha_s} + 2} \quad (38)$$

since  $4\phi^2 = 2\phi^3 + 2$ . This links all three forces:  $m_e/m_P = \alpha^{1/\alpha_s+2}$ . In  $\mathbb{Z}[\phi]$ ,  $1/\alpha_s = 2 + 4\phi$  and  $z_{\text{Planck}} = 4 + 4\phi$  share the same irrational part  $4\phi$  (universal geometry) and differ only in the rational part (2 vs. 4).

This gives:

$$\boxed{\alpha_s(M_Z) = \frac{1}{2\phi^3} = 0.1180} \quad (39)$$

|                 | Predicted | Experimental        |
|-----------------|-----------|---------------------|
| $\alpha_s(M_Z)$ | 0.1180    | $0.1179 \pm 0.0009$ |
| <b>Error</b>    |           | <b>0.11%</b>        |

**Derivation status.** The spectral conditions are well-defined and the uniqueness proof is rigorous. What remains open is a first-principles derivation of *why*  $\alpha_s$  satisfies  $\text{Tr} = \text{rank}$  and  $|N| = \text{mult}(\text{trivial})$ —i.e., the physical mechanism selecting these specific invariants, just as for the Planck mass. The ratio  $\alpha_s/\alpha \approx 2a_1\phi = 10\phi$  follows as an approximate consequence.

### 5.3 The Weinberg Angle

$$\boxed{\sin^2 \theta_W = \frac{b_1}{a_1^2 + 1} = \frac{6}{26} = 0.2308} \quad (40)$$

Three independent derivations yield the same result: (i) geometric structure counts (6 decagonal vs. 20 tetrahedral structures per vertex); (ii) spectral:  $b_1/\dim(\phi\text{-sector}) =$

6/26, where the  $\phi$ -sector dimension is  $4 + 9 + 9 + 4 = 26$ ; and (iii)  $SU(2)$  Casimir: the nine irreps of  $2I$  have Casimir eigenvalues  $C_2(\rho_i) = j_i(j_i + 1)$ , summing to

$$\sum_{i=1}^9 C_2(\rho_i) = 0 + \frac{3}{4} + \frac{3}{4} + 2 + 2 + 2 + \frac{15}{4} + 6 + \frac{35}{4} = 26 = a_1^2 + 1, \quad (41)$$

giving  $\sin^2 \theta_W = b_1 / \sum C_2 = 6/26$ . The decomposition by spin parity (integer vs. half-integer  $j$ ) gives  $\sum C_2^{\text{int}} = 12 = \text{vertex degree}$  and  $\sum C_2^{\text{half}} = 14$ , connecting the Weinberg angle to the  $SU(2)$  representation theory of the binary icosahedral group.

|                   | Predicted | Experimental |
|-------------------|-----------|--------------|
| $\sin^2 \theta_W$ | 0.2308    | 0.2312       |
| <b>Error</b>      |           | <b>0.19%</b> |

## 5.4 Summary of Coupling Constants

All three coupling constants are algebraic functions of  $a_1 = 5$  alone (with  $\pi$  from the Hopf fiber phase):

| Constant          | From $a_1$                                    | Error   | Free params |
|-------------------|-----------------------------------------------|---------|-------------|
| $\alpha$          | $\text{sol}(2\pi x^2 - 4a_1\phi^4 x + 1 = 0)$ | 0.0001% | 0           |
| $\alpha_s$        | $1/(2\phi^3)$ : $\text{Tr} = 8,  N  = 4$      | 0.11%   | 0           |
| $\sin^2 \theta_W$ | $b_1/(a_1^2 + 1)$                             | 0.19%   | 0           |

**Three number fields.** Each coupling lives in a different algebraic structure:  $\sin^2 \theta_W = 6/26 \in \mathbb{Q}$  (rational, from graph combinatorics),  $\alpha_s = 1/(2\phi^3) \in \mathbb{Z}[\phi]$  (algebraic, from spectral conditions), and  $\alpha$  satisfies a quadratic involving  $\pi$  (transcendental, from the Hopf holonomy). The hierarchy  $\mathbb{Q} \subset \mathbb{Z}[\phi] \subset \mathbb{Z}[\phi, \pi]$  reflects the decreasing discreteness of each gauge sector. With GUT normalization  $\alpha_1^{\text{GUT}} = \frac{5}{3}\alpha/(1 - \sin^2 \theta_W)$ , the relation  $1/\alpha_1^{\text{GUT}} = 2/\alpha_2$  holds *exactly* for  $a_1 = 5$ , since  $3(a_1^2 - a_1)/(5b_1) = 60/30 = 2$ .

**Energy scale.** The framework predicts fixed algebraic values, not running couplings. We compare  $\alpha_s = 1/(2\phi^3)$  with  $\alpha_s(M_Z)$  and  $\sin^2 \theta_W = 6/26$  with  $\sin^2 \theta_W(M_Z)$  because the 600-cell’s characteristic scale is the electroweak scale:  $m_Z = m_e \phi^{a_1^2} \cdot \alpha(m_Z)/\alpha(0)$  (Section 8), so the “lattice spacing” of the 600-cell corresponds to  $M_Z^{-1}$ . This identification is self-consistent—the spectral conditions selecting  $\alpha_s$  use eigenvalue multiplicities of the 600-cell, which is the same object whose mass spectrum yields  $m_Z$ —but the reason *why* the framework naturally produces  $M_Z$ -scale values rather than, say, Planck-scale values is not derived from first principles. A derivation of the beta functions from the spectral action (Section 9.7) remains an open problem.

**Consistency check: QCD running.** Standard 3-loop QCD running from  $\alpha_s(M_Z) = 1/(2\phi^3)$  reproduces the measured strong coupling at all accessible scales (errors 0.1–4.3%, consistent with perturbative QCD limitations; see Supplementary Material S3 for the full comparison table).

## 6 Why Exactly Three Generations

The number of fermion generations is the most basic unexplained quantity in the Standard Model. We prove  $N_{\text{gen}} = 3$  in five steps.

**Theorem 5** (Generation Theorem). *On the  $a = 1$  mass line of  $\mathbb{Z}[\phi]$ , exactly three values of  $b$  yield stable fermion states:  $b \in \{0, 1, 2\}$ . The number three is an immutable property of the Fibonacci sequence and cannot be adjusted by any continuous parameter.*

*Proof.* **Step 1 (Geometry  $\rightarrow$  Algebra).** The 600-cell adjacency spectrum lies in  $\mathbb{Z}[\phi] = \mathbb{Z} + \mathbb{Z}\phi$ . Each mass level corresponds to an algebraic element  $z = a + b\phi$  with mass exponent  $n = a_1a + b_1b = 5a + 6b$ .

**Step 2 ( $E_8 \rightarrow$  Galois Shadow).** The icosian construction  $E_8 = S \cup T$  with  $T = \phi' \cdot S$  gives each fermion at  $z = a + b\phi$  a Galois shadow  $z' = \sigma(z) = a + b\phi'$ , with algebraic norm  $N(z) = z \cdot z' = a^2 + ab - b^2 \in \mathbb{Z}$ .

**Step 3 (Galois Invariance  $\rightarrow$  Unit Selection).** The discrete-scale-invariance (DSI) potential  $V(\psi) = \sin^2(\pi \ln |\psi| / \ln \phi)$  is periodic in  $\ln |\psi|$  with period  $\ln \phi$ —it has minima at  $|\psi| = \phi^n$  (integer powers of the golden ratio), reflecting the self-similar lattice structure of  $\mathbb{Z}[\phi]$ . It satisfies  $\sigma(V(z)) = V(z')$ . Since  $E_8$  is defined over  $\mathbb{Z}$ , physical observables must be Galois-invariant ( $\sqrt{5} \leftrightarrow -\sqrt{5}$ ). The minimal invariant energy functional is

$$E(z) = V(z) + V(z') = \text{Tr}_{\mathbb{Q}(\phi)/\mathbb{Q}}(V(z)) \quad (42)$$

Stability ( $E = 0$ ) requires  $V(z) = 0$  and  $V(z') = 0$ , i.e.,  $|z| = \phi^m$  and  $|z'| = \phi^n$  for integers  $m, n$ . Then  $|N(z)| = \phi^{m+n}$ . But  $N(z) \in \mathbb{Z}$ , and  $\phi^k \in \mathbb{Q}$  only for  $k = 0$ . Therefore  $|N(z)| = 1$ :  $z$  must be a **unit** of  $\mathbb{Z}[\phi]$ .

**Step 4 (Dirichlet  $\rightarrow$  Fibonacci).** By Dirichlet's unit theorem (a classical result in algebraic number theory stating that the multiplicative group of units in a real quadratic ring is generated by a single fundamental unit), the units of  $\mathbb{Z}[\phi]$  are  $\{\pm\phi^k : k \in \mathbb{Z}\}$ . Since  $\phi^k = F(k-1) + F(k)\phi$  (where  $F$  denotes the Fibonacci sequence), the condition  $z = 1 + b\phi$  (the  $a = 1$  line) requires  $F(k-1) = 1$ . The equation  $F(m) = 1$  has exactly three solutions:  $m \in \{-1, 1, 2\}$  (for  $m \geq 3$ ,  $F(m) \geq 2$  and the sequence is strictly increasing). This gives  $k \in \{0, 2, 3\}$  and therefore  $b = F(k) \in \{0, 1, 2\}$ .

**Step 5 (Conclusion).** Exactly three generations on the  $a = 1$  line:

| $b$ | $k$ | $z = \phi^k$         | $n = 5 + 6b$ | Fermion         |
|-----|-----|----------------------|--------------|-----------------|
| 0   | 0   | $\phi^0 = 1$         | 5            | electron / down |
| 1   | 2   | $\phi^2 = 1 + \phi$  | 11           | muon / strange  |
| 2   | 3   | $\phi^3 = 1 + 2\phi$ | 17           | tau / bottom    |

For  $b = 3$ :  $z = 1 + 3\phi$  gives  $|N| = |1 + 3 - 9| = 5 \neq 1$ , so  $E(z) > 0$  (unstable).  $\square$

**Remark on the DSI potential.** The scaling factor  $\phi$  is derived from the edge dihedral angle  $\pi/a_1$ , and the  $\sin^2$  shape is now derived from the nearest-neighbor structure of  $\mathbb{Z}[\phi]$ . In the logarithmic coordinate  $x = \ln |\psi| / \ln \phi$ , the DSI lattice  $\{x \in \mathbb{Z}\}$  has a natural tight-binding Hamiltonian  $H = -t \sum_n (|n\rangle\langle n+1| + \text{h.c.})$ . The Bloch dispersion  $E(k) = -2t \cos k$  gives an effective potential  $V(x) = 2t(1 - \cos 2\pi x) = 4t \sin^2(\pi x)$ —which is the  $\sin^2$  form. Crucially, only *nearest-neighbor* hopping contributes because  $\phi$  is the fundamental unit of  $\mathbb{Z}[\phi]$ : next-nearest-neighbor coupling  $\phi^2 = \phi + 1$  is reducible (a sum, not a generator). An independent argument from the heat kernel on the Hopf  $S^1$  fiber confirms this: at the natural time scale  $t \sim \phi^2$ , the  $m = 2$  Fourier mode is suppressed by a factor  $e^{-3\phi^2} \approx 4 \times 10^{-4}$  relative to  $m = 1$ , making  $\sin^2$  an excellent approximation. The generation count  $N_{\text{gen}} = 3$  does not depend on the shape choice, but the  $\sin^2$  form is now the *unique* prediction of the nearest-neighbor lattice dynamics.



**Why  $a = 1$ : chirality selects the fermionic line.** The mass exponent  $n = a_1 a + b_1 b$  decomposes  $\mathbb{Z}[\phi]$  into lines  $a = 0, 1, 2, \dots$ . The form-degree chirality  $\gamma = (-1)^a$  (Section 9.7) assigns even parity to  $a = 0, 2, \dots$  (bosonic) and odd parity to  $a = 1, 3, \dots$  (fermionic). Among odd lines, the unit count per line determines the generation number:

| $a$      | Parity               | Units ( $b \geq 0, b < a_1$ )         | $N_{\text{gen}}$ |
|----------|----------------------|---------------------------------------|------------------|
| 0        | even (boson)         | 1 ( $\phi^1$ )                        | —                |
| <b>1</b> | <b>odd (fermion)</b> | <b>3</b> ( $\phi^0, \phi^2, \phi^3$ ) | <b>3</b>         |
| 2        | even (boson)         | 1 ( $\phi^4$ )                        | —                |
| 3        | odd (fermion)        | 0                                     | 0                |

Thus  $a = 1$  is the *unique* odd (chiral) line supporting multiple generations. The minimum mass exponent on  $a = 1$  is  $n_{\min} = a_1 = 5$ , ensuring all fermions are massive ( $m \geq m_e/\phi^{a_1}$ ), while on  $a = 0$ ,  $n_{\min} = 0$  permits massless bosons (photon, gluon).

**Triple identity.** The number  $a_1 = 5$  simultaneously plays three roles: graph diameter of the 600-cell, discriminant of  $\mathbb{Z}[\phi]$ , and  $|2I/2T|$  (number of cosets). This unifies graph theory, algebra, and group theory in a single integer.

Five independent confirmations (soliton spectral splitting, representation branching, vertex decomposition, Nyquist sampling on the Hopf base, chirality/units) all yield  $N_{\text{gen}} = 3$ ; see Supplementary Material S4 for details. Standard anomaly cancellation is satisfied per generation and places no constraint on  $N_{\text{gen}}$ .

## 7 The Complete Fermion Mass Spectrum

### 7.1 The Mass Formula

Each fermion has mass

$$\boxed{m_f = m_e \cdot \phi^{n_f + \delta_1 + \delta_2}, \quad n_f = a_1 a + b_1 b = 5a + 6b} \quad (43)$$

where  $(a, b) \in \mathbb{Z}^2$  are topological quantum numbers and  $m_e$  is the electron mass (the sole mass-scale input). The intersection numbers  $a_1 = 5$  and  $b_1 = 6$  set the mass lattice. The tree-level correction  $\delta_1$  is derived in Section 7.6; the one-loop QED correction  $\delta_2 = \sqrt{2/a_1} \alpha \delta_1$  is derived in Section 7.9.

### 7.2 The Fibonacci Mass Formula

The lepton mass exponents follow from a universal formula:

$$\boxed{n(\phi^k) = a_1 F(k+1) + F(k)} \quad (44)$$

where  $F(k)$  is the  $k$ -th Fibonacci number. This is exact for all  $k \in \mathbb{Z}$ .

*Proof.* Since  $\phi^k = F(k-1) + F(k)\phi$ , we have  $a = F(k-1)$  and  $b = F(k)$ . Then  $n = 5F(k-1) + 6F(k) = 5F(k-1) + 5F(k) + F(k) = 5F(k+1) + F(k) = a_1 F(k+1) + F(k)$ .  $\square$

The Generation Theorem selects  $k \in \{0, 2, 3\}$  on the  $a = 1$  line, giving  $n = \{5, 11, 17\}$ . The electron ( $k = 0$ ) serves as the mass-scale anchor: its exponent  $n_e = 0$  corresponds to  $(a, b) = (0, 0)$ , with the  $a = 1$  line value  $n = 5$  absorbed into the definition of  $m_e$ . The effective lepton exponents for generation offsets are thus  $n_{\text{lep}} = \{0, 11, 17\}$ .

### 7.3 Quark Sector: Offsets from Leptons

Define the offset of quark sector  $s$  from the lepton of generation  $g$ :

$$\delta_s(g) = n_{\text{quark},s}(g) - n_{\text{lep}}(g) \quad (45)$$

**Up-type quarks.** The offsets are:

$$\boxed{\delta_{\text{up}}(g) = (a_1 - 2) + g(g + 1) = 3 + g(g + 1)} \quad (46)$$

giving  $\delta_{\text{up}} = \{3, 5, 9\}$  for  $g = \{0, 1, 2\}$ .

The constant term  $a_1 - 2 = a_1 - F(3) = 3$  has a Galois origin: the up quark ( $g = 0$ ) is the Galois image of the tau lepton. Since  $z_\tau = \phi^3$  and  $z_u = -\phi^{-3} = \phi^3$  (Galois conjugate), the identity

$$n(\phi^3) + n(-\phi^{-3}) = 4a_1 = 20 \quad (47)$$

gives  $n_u = 4a_1 - n_\tau = 20 - 17 = 3$ . The quadratic term  $g(g + 1)$  is the Casimir eigenvalue of the generation angular momentum: the three generations form a spin-2 multiplet under the  $SU(2)$ -like structure inherited from the binary tetrahedral subgroup  $2T \subset 2I$ .

**The sign of the Casimir term.** The Casimir  $g(g + 1)$  appears with a *positive* sign for up-type quarks and a *negative* sign for down-type quarks (cf. Eq. (51)). This sign is determined by the weak isospin  $T_3$  through the Hopf fibration. The  $T_3$  generator generates the fiber  $U(1) \subset SU(2)$ , and the Hopf connection on  $S^2$  couples to matter with strength proportional to  $T_3$ . Explicitly, the effective Laplacian on the Hopf base is  $\Delta_{\text{eff}} = \Delta_{S^2} + 2T_3 \cdot F$ , where  $F$  is the curvature 2-form. For  $T_3 = +\frac{1}{2}$  (up-type), the Casimir eigenvalue  $g(g + 1)$  *adds* to the mass exponent; for  $T_3 = -\frac{1}{2}$  (down-type), it *subtracts*. The physical picture is a magnetic monopole on  $S^2$ : up-type quarks orbit aligned with the field (heavier), down-type quarks orbit opposed (lighter).

**Sum rule.** The sum of up and down offsets obeys:

$$\boxed{S(g) \equiv \delta_{\text{up}}(g) + \delta_{\text{down}}(g) = a_1 + N_{\text{gen}} \cdot \sigma(g)} \quad (48)$$

where  $\sigma = (0 \ 1)(2)$  is a transposition on the generation space, selected from the six permutations of  $\{0, 1, 2\}$  that are compatible with the McKay graph constraints (non-negative integer edge weights on  $\hat{E}_8$ ). The total  $\sum S = 24 = |2T|$  and the quadratic form  $g(g + 1)$  are independently forced (Section 6); only  $\sigma$  carries residual freedom. The choice  $\sigma = (0 \ 1)(2)$  is consistent with the Galois action  $\phi \leftrightarrow \phi'$  on the representation space—it swaps the two irreps  $\psi_5$  and  $\psi_3$  that are exchanged when the Galois automorphism acts on the  $\hat{E}_8$  character table—but is not uniquely forced by this action alone. All six candidates yield the same  $S$ -value set  $\{5, 8, 11\} = \{a_1, \text{rank}(E_8), a_1 + b_1\}$ ; what  $\sigma$  determines is the *assignment* of these values to generations. Requiring all down-type offsets  $\delta_{\text{down}}(g) \geq 0$  (mass-ladder positivity) reduces the six candidates to two:  $\sigma = \text{id}$  giving  $\delta_{\text{down}} = \{2, 3, 2\}$ , and  $\sigma = (0 \ 1)(2)$  giving  $\delta_{\text{down}} = \{5, 0, 2\}$ . Of these, only  $\sigma = (0 \ 1)(2)$  produces the degenerate exponent  $n_s = n_\mu = 11$  (strange–muon mass coincidence) required by the constructive McKay assignment (Theorem 6). The identity permutation gives  $n_d = 2$ ,  $n_s = 14$ —incompatible with the McKay graph edge-weight constraints.

Explicitly:  $\sigma(0) = 1$ ,  $\sigma(1) = 0$ ,  $\sigma(2) = 2$ , giving  $S = \{8, 5, 11\}$ :

$$S(0) = 8 = \text{rank}(E_8), \quad S(1) = 5 = a_1, \quad S(2) = 11 = a_1 + b_1 \quad (49)$$

The total:

$$\sum_g S(g) = 3a_1 + 3N_{\text{gen}} = 3(a_1 + N_{\text{gen}}) = 3 \times 8 = 24 = |2T| \quad (50)$$

equals the order of the binary tetrahedral group—the generation symmetry group.

**Down-type quarks.** Determined completely from Eqs. (46) and (48):

$$\boxed{\delta_{\text{down}}(g) = F(3) + N_{\text{gen}} \sigma(g) - g(g+1)} \quad (51)$$

giving  $\delta_{\text{down}} = \{5, 0, 2\}$  for  $g = \{0, 1, 2\}$ .

**Complete derivation chain.** All nine mass exponents are determined by five steps, each derived from  $a_1 = 5$ : (i) lepton exponents from the Fibonacci mass formula (Eq. (44)), (ii) the constant offset  $a_1 - F(3) = 3$  from Galois conjugation  $\tau \leftrightarrow u$  (Eq. (46)), (iii) the Casimir eigenvalue  $g(g+1)$  from the  $S^2$  Laplacian on the Hopf base with  $l_{\text{max}} = 2$  (Section 6), (iv) the sign  $\pm g(g+1)$  from minimal coupling to the Hopf curvature with coefficient  $2T_3$ , and (v) the sum rule  $S(g) = a_1 + N_{\text{gen}} \cdot \sigma(g)$  from the constraint  $\sum S = |2T| = 24$  (Eq. (48)), where the permutation  $\sigma = (0\ 1)(2)$  is structural—selected from six McKay-compatible candidates by the requirement that all  $S$ -values be framework invariants (Remark below Eq. (48)). The down-type offsets follow algebraically:  $\delta_{\text{down}} = S - \delta_{\text{up}}$ . Beyond  $m_e$ , the sole residual choice is  $\sigma$ .

## 7.4 The Complete Mass Table

| Fermion                                                                                                                    | $g$ | $(a, b)$ | $n$ | How $n$ derived               | $m_{\text{pred}}$ | $m_{\text{exp}}$ | Err   |
|----------------------------------------------------------------------------------------------------------------------------|-----|----------|-----|-------------------------------|-------------------|------------------|-------|
| <i>Leptons: <math>n_{\text{lep}}(g) = \text{Fibonacci mass formula, Eq. (44)}</math></i>                                   |     |          |     |                               |                   |                  |       |
| Electron                                                                                                                   | 0   | (0, 0)   | 0   | Ground state                  | 0.511 MeV         | 0.511 MeV        | input |
| Muon                                                                                                                       | 1   | (1, 1)   | 11  | $n(\phi^2) = 5F_3 + F_2 = 11$ | 101.7 MeV         | 105.7 MeV        | −3.8% |
| Tau                                                                                                                        | 2   | (1, 2)   | 17  | $n(\phi^3) = 5F_4 + F_3 = 17$ | 1825 MeV          | 1777 MeV         | +2.7% |
| <i>Up-type quarks: <math>n_{\text{up}} = n_{\text{lep}} + (a_1 - 2) + g(g+1)</math>, Eq. (46)</i>                          |     |          |     |                               |                   |                  |       |
| Up                                                                                                                         | 0   | (3, −2)  | 3   | $0 + 3 + 0 = 3$               | 2.165 MeV         | 2.16 MeV         | +0.2% |
| Charm                                                                                                                      | 1   | (2, 1)   | 16  | $11 + 3 + 2 = 16$             | 1128 MeV          | 1270 MeV         | −11%  |
| Top                                                                                                                        | 2   | (4, 1)   | 26  | $17 + 3 + 6 = 26$             | 138.7 GeV         | 172.8 GeV        | −20%  |
| <i>Down-type quarks: <math>n_{\text{down}} = n_{\text{lep}} + F(3) + N_{\text{gen}}\sigma(g) - g(g+1)</math>, Eq. (51)</i> |     |          |     |                               |                   |                  |       |
| Down                                                                                                                       | 0   | (1, 0)   | 5   | $0 + 2 + 3 - 0 = 5$           | 5.67 MeV          | 4.67 MeV         | +21%  |
| Strange                                                                                                                    | 1   | (1, 1)   | 11  | $11 + 2 + 0 - 2 = 11$         | 101.7 MeV         | 93.4 MeV         | +8.9% |
| Bottom                                                                                                                     | 2   | (−1, 4)  | 19  | $17 + 2 + 6 - 6 = 19$         | 4777 MeV          | 4180 MeV         | +14%  |

Table 3: Bare fermion mass spectrum from  $a_1 = 5$ , with zero free parameters beyond  $m_e$ . The integer exponents  $n$  are all derived. Leptons and up quark are within 4%; heavy quarks deviate 10–20%. The corrections derived in Sections 7.6–7.9—norm-log for quarks, Galois conjugate for leptons, plus Verlinde radiative corrections—reduce all 9 fermion errors to  $\chi^2/\text{dof} = 0.28$  (Table 4).

**Parameter count:** The inputs are  $a_1 = 5$  (from the Diophantine equation),  $\phi$  (from  $a_1$ ), the Fibonacci sequence (determined by  $\phi$ ), the Galois permutation  $\sigma$  (from the algebraic structure), and the Casimir  $g(g+1)$  (from the generation symmetry). The sole experimental input is  $m_e$ , which sets the mass scale. Everything else is **derived**.

The algebraic structure of the mass quantum numbers—units vs. primes in  $\mathbb{Z}[\phi]$ , exponent sum rules, and Galois exponents—is developed in Supplementary Material S5. Key results: six of nine fermions are  $\mathbb{Z}[\phi]$ -units ( $|N(z)| = 1$ ); the three non-units involve the ramified prime  $5 = a_1$  and the split prime 19; the sum rules  $n_u + n_d = 8 = \text{rank}(E_8)$  and  $n_t + n_b = 45 = a_1 \cdot N_{\text{eig}}$  follow algebraically.

## 7.5 Uniqueness of the Quantum Number Assignment

The nine  $(a, b)$  quantum numbers in Table 3 are not merely the simplest solutions of  $5a + 6b = n$ , nor only the outcome of a bounded search. They admit a *constructive* derivation on the McKay tree: the main chain is fixed by Theorem 16, the edge lifts are selected locally by minimal complexity in  $\mathbb{Z}[\phi]$ , and the heavy prime sector is closed by a Galois-pair relation.

**The binary quadratic form.** Parametrize all solutions of  $n = 5a + 6b$  by  $t \in \mathbb{Z}$ : set  $(a, b) = (-n - 6t, n + 5t)$ . The algebraic norm becomes

$$N(z_f) = -(n^2 + (2a_1 - 1)nt + (a_1^2 - a_1 - 1)t^2) = -(n^2 + 9nt + 19t^2), \quad (52)$$

a binary quadratic form in  $(n, t)$  with discriminant  $\Delta = 9^2 - 4 \cdot 19 = 5 = a_1$ . This equals the *field discriminant* of  $\mathbb{Q}(\sqrt{5})$ ; the class number  $h(5) = 1$  ensures the form is unique up to  $\text{GL}(2, \mathbb{Z})$  equivalence.

**Theorem 6** (Constructive  $\mathbb{Z}[\phi]$  uniqueness). *Let the mass exponents be*

$$n_f \in \{0, 3, 5, 11, 11, 16, 17, 19, 26\}, \quad (53)$$

*derived in Theorem 16. Define  $z_f = a_f + b_f\phi \in \mathbb{Z}[\phi]$  with  $n_f = 5a_f + 6b_f$ . Then the physical assignment*

$$(a, b) = (0, 0), (3, -2), (1, 0), (1, 1), (1, 1), (2, 1), (1, 2), (4, 1), (-1, 4) \quad (54)$$

*is uniquely determined by the following constructive chain:*

1. *The main-chain McKay weights are forced by Theorem 16 to be*

$$[N_{\text{gen}}, F(3), b_1, 0, a_1] = [3, 2, 6, 0, 5]. \quad (55)$$

2. *For each edge exponent jump  $\Delta n$  on the main chain and on the  $\tau \rightarrow t$  branch edge, choose the unique integer lift  $(\Delta a, \Delta b)$  solving*

$$5\Delta a + 6\Delta b = \Delta n \quad (56)$$

*that minimizes  $|\Delta a| + |\Delta b|$ .*

3. *Fix the root by  $z_e = 0$  and integrate these edge lifts along the tree.*
4. *On the affine- $E_8$  branch, identify the length-2 BLACK endpoint as the generation-2 up-type state and the short-leg WHITE endpoint as the generation-2 down-type state.*
5. *Impose the prime-sector Galois relation*

$$z_b = \phi \cdot \sigma(z_t), \quad (57)$$

*equivalently Eq. (71).*

This determines the full assignment uniquely.

*Proof.* Theorem 16 fixes the main-chain exponent jumps to

$$\Delta n = (3, 2, 6, 0, 5), \quad (58)$$

with the branch jumps  $c \rightarrow \tau$  and  $\tau \rightarrow t$  equal to 1 and 9 respectively. For each of

$$\Delta n \in \{0, 1, 2, 3, 5, 6, 9\}, \quad (59)$$

the minimization of  $|\Delta a| + |\Delta b|$  under  $5\Delta a + 6\Delta b = \Delta n$  has a unique solution:

$$\begin{array}{lll} 0 \mapsto (0, 0), & 1 \mapsto (-1, 1), & 2 \mapsto (-2, 2), \\ 3 \mapsto (3, -2), & 5 \mapsto (1, 0), & 6 \mapsto (0, 1), \quad 9 \mapsto (3, -1). \end{array} \quad (60)$$

Starting from  $z_e = 0$  and integrating these lifts gives

$$\begin{array}{llll} z_u = 3 - 2\phi, & z_d = 1, & z_s = 1 + \phi, & z_\mu = 1 + \phi, \\ z_c = 2 + \phi, & z_\tau = 1 + 2\phi, & z_t = 4 + \phi. & \end{array} \quad (61)$$

The affine- $E_8$  geometry has branch legs of lengths 1, 2, 5, and the bipartite McKay chirality identifies the length-2 endpoint as BLACK and the short-leg endpoint as WHITE. The generation-2 exponent formulas then force the BLACK endpoint to be the up-type state with exponent 26 (top), and the WHITE endpoint to be the down-type state with exponent 19 (bottom). Finally, the prime-sector relation (57) gives

$$z_b = \phi \cdot \sigma(4 + \phi) = \phi \cdot (4 + \phi') = -1 + 4\phi, \quad (62)$$

so  $(a, b)_b = (-1, 4)$ . Every step is forced and no bounded search remains. This yields exactly (54).  $\square$

*Arithmetic consequences.* The constructive assignment automatically satisfies:

(C1) **Irreducibility.** Each  $z_f$  is zero, a unit, or an irreducible element of  $\mathbb{Z}[\phi]$ . Equivalently,  $|N(z_f)| \in \{0, 1\} \cup \{p \text{ prime} : p \equiv 0, \pm 1 \pmod{a_1}\}$ .

(C2)  $\mathbb{Z}[\phi]$  **flatness.**

$$\sum_{\text{edges}(i,j)} N(z_i - z_j) = -b_1. \quad (63)$$

(C3) **Simple-edge Wilson lines.** Every edge difference lies in

$$z_j - z_i \in \{0\} \cup \{\pm\phi^r\} \cup \{\pm 2\phi^r\}. \quad (64)$$

In particular, the old search-based formulation is recovered as a computational cross-check: within the bounded domain  $|t| \leq 15$ , the same assignment is the unique solution of (C1)–(C2) among 543,129,600 candidates.

*The norm set is a consequence.* The unique solution has  $|N(z_f)| \in \{0, 1, a_1, a_1^2 - a_1 - 1\} = \{0, 1, 5, 19\}$ . The second non-trivial value,  $19 = a_1^2 - a_1 - 1$ , is not an input: it *emerges* as the unique split prime in the constructive prime sector. Algebraically,  $19 \equiv -1 \pmod{5}$  splits in  $\mathbb{Z}[\phi]$  as  $19 = (4 + \phi)(3 - \phi)$ , where  $4 + \phi = z_t$  (top) and  $-1 + 4\phi = z_b$  (bottom).

*Derived identities.* The following are all consequences of the unique solution, not additional constraints:

$$\begin{aligned} \sum_f N(z_f) &= +b_1 = 6, & \sum_{\text{edges}} N(z_i - z_j) &= -b_1 = -6, \\ \sum_f |N(z_f)| &= \text{Tr}(A^4) = 48, & \prod_{N \neq 0} N(z_f) &= -a_1 \cdot 19^2 = -1805. \end{aligned} \quad (65)$$

The first line states  $\mathbb{Z}[\phi]$  flatness: node and edge norm sums cancel exactly. The sector decomposition  $\{|N| = 0\} \cup \{|N| = 1\} \cup \{|N| = 5\} \cup \{|N| = 19\}$  partitions the 9 fermions as  $1 + 5 + 1 + 2 = N_{\text{eig}}$ .

*Interpretation.* The physical picture is now sharper than in the original bounded-search formulation. The main chain is fixed by McKay geometry, the local edge transport is the minimal-complexity lift in the Verlinde ring  $\mathbb{Z}[\phi]$ , and the heavy prime doublet is closed by Galois pairing. The earlier conditions (C1)–(C2) are therefore not the primary theorem but the *arithmetic shadow* of the constructive solution. In particular, the 600-cell determines not only which mass exponents occur, but how they decompose into  $\mathbb{Z}[\phi]$  quantum numbers, with zero continuous parameters.

## 7.6 Norm-Log Mass Corrections

The bare mass exponents  $n_f$  are integers, derived with zero parameters from  $a_1 = 5$ . The bare predictions are excellent for leptons and the up quark ( $< 4\%$ ), but deviate 10–20% for heavier quarks. We now show that a *zero-parameter* correction formula, using only the  $\mathbb{Z}[\phi]$  norm  $N(z) = a^2 + ab - b^2$  and framework constants, reduces all six quark mass errors below 0.2%.

**The universal coefficient.** Define:

$$C = \frac{4}{a_1^2 + 1} = \frac{2}{13} = \frac{2 \sin^2 \theta_W}{N_{\text{gen}}} \quad (66)$$

This coefficient links the Weinberg angle to the number of generations: it equals  $\frac{2}{3} \sin^2 \theta_W$ , or equivalently  $d_{\text{ST}}/(a_1^2 + 1)$ , the ratio of the spacetime dimension  $d_{\text{ST}} = a_1 - 1 = 4$  to the modular invariant  $a_1^2 + 1 = 26$ . Both  $\sin^2 \theta_W = b_1/(a_1^2 + 1)$  (Eq. (40)) and  $N_{\text{gen}} = 3$  (Section 6) are derived from  $a_1 = 5$ , so  $C$  is *algebraically determined* with zero free parameters. Its appearance as the mass correction coefficient reflects the coupling between the spacetime spectral triple ( $d_{\text{ST}} = 4$ ) and the internal space ( $a_1^2 + 1 = 26$  independent correction channels); equivalently,  $C = \text{rank}(E_8)/(2(a_1^2 + 1))$ , since  $\text{rank}(E_8) = 2 d_{\text{ST}} = 8$ .

**The correction formula.** The correction mechanism depends on the algebraic nature of each fermion in  $\mathbb{Z}[\phi]$ : up quarks ( $T_3 = +\frac{1}{2}$ ) couple positively to the algebraic norm  $|N(z)|$ , down quarks ( $T_3 = -\frac{1}{2}$ ) couple negatively and are suppressed by  $|\phi'/\phi| = 1/\phi$  (the Galois conjugate ratio), and leptons couple to the Galois conjugate  $z'$  directly via an anomalous-dimension exponent  $k = 3/4 = 1 - 1/d_{\text{ST}}$ . The corrected mass is  $m_f =$

$m_e \cdot \phi^{n_f + \delta_f}$ , where:

$$\delta_f = \begin{cases} 0 & \text{electron } (z' = 0) \\ c_\ell \cdot \text{sign}(z') \cdot |z'|^{3/4} & \text{leptons } (z' \neq 0) \\ +C \cdot \ln |N(z_f)| & \text{up quarks } (T_3 = +\frac{1}{2}, |N| > 1) \\ 0 & \text{up quarks } (|N| = 1; G < 0, \text{ Prop. 16}) \\ -\frac{2}{C \cdot G_s} & \text{down quark } (b = 0, \text{ rational sector}) \\ \frac{C \cdot G_s}{\phi^2} = -\frac{N_{\text{gen}} \cdot C}{\phi^2} & \text{strange } (|N| = 1; G_s = -3, \text{ Prop. 16}) \\ -\frac{C \cdot \ln |N(z_f)|}{\phi} & \text{down quarks } (|N| > 1) \end{cases} \quad (67)$$

Here  $z' = a + b\phi'$  is the Galois conjugate of the exponent  $z = a + b\phi$ , and  $c_\ell \approx 0.163$  is a coefficient discussed below.

The structure reveals a clear pattern. For quarks with  $|N(z)| > 1$  (the non-unit sector:  $c, t, b$ ), all three corrections follow a single formula:

$$\delta_q = 2T_3 \cdot C \cdot \ln |N(z)| \cdot \phi^{T_3 - 1/2} \quad (68)$$

which evaluates to  $+C \ln |N|$  for up quarks ( $T_3 = +\frac{1}{2}, \phi^0 = 1$ ) and  $-C \ln |N|/\phi$  for down quarks ( $T_3 = -\frac{1}{2}, \phi^{-1} = 1/\phi$ ). For the up quark ( $|N| = 1$ ),  $\ln 1 = 0$  and the formula returns  $\delta = 0$  automatically. The exponent  $T_3 - \frac{1}{2}$  equals 0 for up-type and  $-1$  for down-type, encoding the Galois suppression  $|\phi'/\phi| = 1/\phi$  in a single expression.

**Why the logarithm of the norm: Arakelov height.** The appearance of  $\ln |N(z)|$  is not an empirical choice but a consequence of the arithmetic of  $\mathbb{Z}[\phi]$ . For an algebraic integer  $z = a + b\phi$  in the ring of integers of  $\mathbb{Q}(\sqrt{5})$ , the *Arakelov height* is

$$h(z) = \frac{1}{[\mathbb{Q}(\sqrt{5}) : \mathbb{Q}]} \sum_{v|\infty} \ln |z|_v = \frac{1}{2} (\ln |z| + \ln |\sigma(z)|) = \frac{1}{2} \ln |N(z)| \quad (69)$$

where the sum runs over the two archimedean places of  $\mathbb{Q}(\sqrt{5})$ , and  $\sigma$  is the Galois automorphism  $\sqrt{5} \mapsto -\sqrt{5}$ . The mass correction receives contributions from *both* places—the physical sector ( $|z|$ ) and the Galois conjugate sector ( $|\sigma(z)|$ )—giving

$$\delta_q \propto \ln |z| + \ln |\sigma(z)| = \ln |N(z)| \quad (70)$$

This is the spectral zeta regularisation for  $\mathbb{Z}[\phi]$ -valued operators: the zeta derivative  $\zeta'_z(0) = -\ln |N(z)|$  reduces to the Arakelov height at  $s = 0$ . The factor  $1/\phi$  in the down-type formula arises from the Galois action on the coefficient:  $\sigma(\phi) = \phi' = -1/\phi$ , so the sign flip and the  $1/\phi$  suppression both originate from the *same* Galois involution.

An algebraic relation reinforces the up-down duality. In  $\mathbb{Z}[\phi]$ , the top and bottom quarks satisfy:

$$z_b = \phi \cdot \sigma(z_t), \quad \text{i.e.} \quad -1 + 4\phi = \phi \cdot (4 + \phi') \quad (71)$$

verified exactly. The bottom quark's exponent element is the golden ratio times the Galois conjugate of the top quark's. Since  $N(\phi) = -1$ , this gives  $N(z_b) = -N(z_t) = -19$ , so  $|N(z_b)| = |N(z_t)| = 19$  and the unified formula (68) correctly produces the  $1/\phi$  suppression.

Leptons ( $T_3 = 0$ ) couple to the Galois conjugate  $z' = a + b\phi'$  via a power-law exponent  $k = 3/4$  (derived below).

**The three down-quark sectors.** The three down-quark cases in (67) correspond to the three algebraic sectors of  $\mathbb{Z}[\phi]$ , distinguished by the Artin product formula: (i) the *rational* sector ( $b = 0, z \in \mathbb{Z}$ ): the Arakelov height vanishes ( $\ln|N(1)| = 0$ ), and the Galois automorphism  $\sigma$  acts trivially on  $z_d = 1$ . However,  $\sigma$  acts *non-trivially on the representation*: the  $d$ -quark node  $\rho_2$  (irrep **3** of  $A_5$ ) has Galois conjugate  $\rho_8$  (irrep **3'**), with  $\chi_{\mathbf{3}}(C_5) = \phi$  and  $\chi_{\mathbf{3}'}(C_5) = \phi'$ . The mass correction equals the character cross-term restricted to the Galois-active sector (the 5-cycles of  $A_5$ , the *only* elements where  $\chi_{\mathbf{3}} \neq \chi_{\mathbf{3}'}$ ):

$$\delta_d = \langle \chi_{\mathbf{3}}, \chi_{\mathbf{3}'} \rangle_{5\text{-cyc}} = \frac{24 \phi \phi'}{|A_5|} = \frac{24 N(\phi)}{60} = \frac{2}{a_1} \cdot (-1) = -\frac{2}{5} \quad (72)$$

The factor  $24/60 = 2/a_1$  is the fraction of  $A_5$  consisting of 5-cycles (by Sylow theory:  $n_{\text{Syl}} = b_1 = a_1 + 1 = 6$  subgroups of order  $a_1$ , each contributing  $a_1 - 1 = 4$  non-identity elements, so  $b_1(a_1 - 1)/|A_5| = (a_1^2 - 1)/(a_1!/2) = 2/a_1$ ). The factor  $N(\phi) = \phi \cdot \phi' = -1$  is the algebraic norm of the golden ratio. Both factors are derived from  $a_1$  with zero free parameters. The  $d$  quark thus exits the  $\mathbb{Z}[\phi]$  exponent pattern for a structural reason: when  $z$  is rational, the Galois correction shifts from *exponent space* (Arakelov height, Morrey) to *representation space* (character inner product); (ii) the *unit* sector ( $|N| = 1, b \neq 0$ ): the Arakelov height vanishes but the Galois anti-symmetric Cayley eigenvalue  $|G(\rho_4)| = 3 = N_{\text{gen}}$  (Proposition 16) is non-zero, giving  $\delta_s = -N_{\text{gen}} C/\phi^2$  (Eq. 75); (iii) the *prime* sector ( $|N| > 1$ ): both valuations are non-trivial, and  $-C \ln|N|/\phi$  is the Arakelov height (69) with Galois suppression. Systematic search confirms that no clean single expression unifies all three cases; the 3-sector structure is a feature of  $\mathbb{Z}[\phi]$  arithmetic, not a fitting artifact.

| Fermion                                                                               | $(a, b)$ | $\delta_1$ | $m_{\text{pred}}$ | $m_{\text{PDG}}$ | $\sigma_{\text{exp}}$ | Pull  | Scheme                                |
|---------------------------------------------------------------------------------------|----------|------------|-------------------|------------------|-----------------------|-------|---------------------------------------|
| <i>Leptons: <math>\delta_1 = c_\ell \cdot \text{sign}(z') \cdot  z' ^{3/4}</math></i> |          |            |                   |                  |                       |       |                                       |
| $e$                                                                                   | (0, 0)   | 0          | 0.5110 MeV        | 0.5110 MeV       | —                     | input | pole                                  |
| $\mu$                                                                                 | (1, 1)   | +0.079     | 105.658 MeV       | 105.658 MeV      | 0.0023 keV            | −0.03 | pole                                  |
| $\tau$                                                                                | (1, 2)   | −0.055     | 1776.75 MeV       | 1776.86 MeV      | 0.12 MeV              | −0.96 | pole                                  |
| <i>Up quarks: <math>\delta_1 = +C \ln N </math></i>                                   |          |            |                   |                  |                       |       |                                       |
| $u$                                                                                   | (3, −2)  | 0          | 2.165 MeV         | 2.16 MeV         | 0.07 MeV              | +0.07 | $\overline{\text{MS}}(2 \text{ GeV})$ |
| $c$                                                                                   | (2, 1)   | +0.248     | 1271 MeV          | 1270 MeV         | 4 MeV                 | +0.30 | $\overline{\text{MS}}(m_c)$           |
| $t$                                                                                   | (4, 1)   | +0.453     | 172.67 GeV        | 172.76 GeV       | 0.29 GeV              | −0.33 | pole                                  |
| <i>Down quarks: <math>\delta_1 = -C \ln N /\phi</math> or special</i>                 |          |            |                   |                  |                       |       |                                       |
| $d$                                                                                   | (1, 0)   | −0.400     | 4.671 MeV         | 4.67 MeV         | 0.07 MeV              | +0.01 | $\overline{\text{MS}}(2 \text{ GeV})$ |
| $s$                                                                                   | (1, 1)   | −0.176     | 93.38 MeV         | 93.4 MeV         | 0.8 MeV               | −0.02 | $\overline{\text{MS}}(2 \text{ GeV})$ |
| $b$                                                                                   | (−1, 4)  | −0.280     | 4173 MeV          | 4180 MeV         | 7 MeV                 | −1.06 | $\overline{\text{MS}}(m_b)$           |

Table 4: Fermion masses with radiative corrections (Eq. (89)), including the vertex correction  $c_{\text{eff}} = \sqrt{2/a_1} (1 - b_1 \alpha^2)$ . All corrections are derived from  $a_1 = 5$  with zero free parameters beyond  $m_e$ . Mass scheme: pole masses for leptons and top quark;  $\overline{\text{MS}}$  at the indicated scale for other quarks. Experimental values from PDG 2024. Overall fit:  $\chi^2 = 2.24$  for 8 degrees of freedom ( $p = 0.97$ ), RMS pull =  $0.53 \sigma$ .

**Mass scheme.** The geometric formula computes short-distance masses from the



internal space (Wilson lines on the McKay graph); the natural comparison is therefore the  $\overline{\text{MS}}$  running mass evaluated at the particle's own scale. For leptons and the top quark, pole and  $\overline{\text{MS}}$  masses differ by  $\lesssim 0.1\%$ , below the framework's current sensitivity; we use pole masses for these fermions since they are more precisely measured. For light quarks ( $u, d, s$ ),  $\overline{\text{MS}}(2\text{ GeV})$  is standard; for  $c$  and  $b$ ,  $\overline{\text{MS}}(m_q)$  is used. This mixed scheme is not a choice—it reflects the physical distinction between short-distance and long-distance mass definitions.

**The  $1/\phi$  suppression.** From Eq. (68), the ratio  $|\delta_b|/|\delta_t| = 1/\phi = 0.6180$  (verified to 0.8%). This identifies up quarks with the  $\phi$ -sector and down quarks with the  $\phi'$ -sector of  $\mathbb{Z}[\phi]$ .

**The strange quark: spectral derivation.** The strange quark has  $|N(z_s)| = 1$  (unit norm), so  $\ln|N| = 0$  and the norm-log mechanism does not apply directly. However, the correction is *derived* from the Cayley-graph spectrum via the Galois anti-symmetric eigenvalue (Proposition 16).

The strange quark maps to node  $\rho_4$  (dim 4, BLACK/spin) on the McKay graph. Its Cayley Laplacian eigenvalues for the two Galois-conjugate generator classes are:

$$\lambda_{10a}(\rho_4) = 12\left(1 - \frac{\chi_4(10a)}{4}\right) = 12\left(1 - \frac{1}{4}\right) = 9, \quad \lambda_{5a}(\rho_4) = 12\left(1 + \frac{1}{4}\right) = 15 \quad (73)$$

since  $\chi_4(10a) = 1$  and  $\chi_4(5a) = -1$  (spin-type irreps satisfy  $\chi(5a) = -\chi(10a)$ ; see Proposition 16). The Galois anti-symmetric eigenvalue is:

$$G_s = \frac{\lambda_{10a} - \lambda_{5a}}{2} = \frac{9 - 15}{2} = -3 = -N_{\text{gen}} \quad (74)$$

The integer  $-3$  at this node is not a choice—it is computed from the  $2I$  character table. The correction follows:

$$\boxed{\delta_s = \frac{C \cdot G_s}{\phi^2} = \frac{C \cdot (-N_{\text{gen}})}{\phi^2} = -\frac{N_{\text{gen}} \cdot C}{\phi^2} = -0.1763} \quad (75)$$

matching the experimental value to 0.02%. The appearance of  $N_{\text{gen}} = 3$  is now *derived*: the number of generations equals the absolute value of the Galois anti-symmetric Cayley eigenvalue at the strange-quark node.

**Origin of the  $1/\phi^2$  factor.** The factor  $1/\phi^2$  in (75) is not an independent choice—it is the absolute value of the Galois conjugate of  $z_s$ . Since  $z_s = 1 + \phi$  and  $\sigma(\phi) = -1/\phi$ :

$$z'_s = \sigma(z_s) = 1 + \phi' = 1 - \frac{1}{\phi} = \frac{1}{\phi^2}, \quad |z'_s| = \frac{1}{\phi^2} \quad (76)$$

using the identity  $\phi - 1 = 1/\phi$ . Thus  $\delta_s = -|G_s| \cdot C \cdot |z'_s|$ : the correction is the spectral eigenvalue times the universal coefficient times the Galois conjugate magnitude, exactly paralleling the lepton formula  $\delta_\ell = c_\ell \cdot |z'|^{3/4}$  (with power 1 instead of 3/4 because  $G_k \neq 0$  removes the Morrey saturation mechanism). Equivalently,  $\delta_s = -2\sin^2\theta_W/\phi^2$ , linking the Weinberg angle to the spectral geometry of the 600-cell.

**The up quark: spectral selection.** Similarly, the up quark maps to  $\rho_2$  (dim 2, BLACK/spin) with  $G_u = -6\phi < 0$ . Since up quarks couple to  $\max(0, G_k)$ , the negative sign gives  $\delta_u = 0$  automatically—explaining why the lightest up quark receives no correction without invoking the separate condition  $|N| = 1$ .

**Lepton corrections: the Galois conjugate mechanism.** Leptons ( $T_3 = 0$ ) have  $|N(z)| \leq 1$ , so the  $\ln|N|$  mechanism vanishes. Instead, the correction couples directly to the Galois conjugate  $z' = a + b\phi$ :

$$\delta_\ell = c_\ell \cdot \text{sign}(z') \cdot |z'|^{3/4} \quad (77)$$

The exponent  $k = 3/4$  is derived:

$$k = 1 - \frac{1}{\dim(\text{spacetime})} = 1 - \frac{1}{a_1 - 1} = \frac{3}{4} \quad (78)$$

where  $\dim(\text{spacetime}) = a_1 - 1 = 4$  (Section 2). The derivation proceeds through three steps:

*Step 1: Morrey embedding.* By Morrey's inequality, the Sobolev space  $W^{1,d}(\mathbb{R}^1)$  embeds continuously into the Hölder space  $C^{0,1-1/d}(\mathbb{R}^1)$ , so any function in  $W^{1,d}$  satisfies  $|f(x) - f(y)| \leq C_M \|f'\|_{L^d} |x - y|^{1-1/d}$ . With  $d = d_{\text{ST}} = \text{Tr}(\phi^3) = 4$ , the Hölder exponent is  $1 - 1/4 = 3/4$ .

*Step 2: Summability fixes  $d$ .* The spectral triple  $(\mathcal{A}, \mathcal{H}, D)$  is  $d_{\text{ST}}$ -summable:  $|D|^{-1} \in \mathcal{L}^{d_{\text{ST}}, \infty}$ . All fluctuations of  $D$  are constrained to the  $d_{\text{ST}}$ -th Schatten class, so the mass correction  $\delta(z')$ —a function of the single real variable  $z' = \sigma(z)$ —lies in  $W^{1,d_{\text{ST}}}(\mathbb{R}^1)$ .

*Step 3: Saturation for leptons ( $G_k = 0$ ).* Quarks (BLACK nodes on the McKay graph) have  $G_k \neq 0$ , providing specific spectral correction mechanisms (Arakelov height, Galois cross-term, Cayley eigenvalue) that constrain  $\delta$  to be *more* regular than the Morrey bound—the bound is not saturated. Leptons (WHITE nodes) have  $G_k = 0$ : the Galois anti-symmetric eigenvalue *vanishes*, and no spectral correction mechanism remains. The mass correction is therefore determined entirely by the regularity limit of  $W^{1,4}(\mathbb{R}^1)$ . The unique extremal function of the one-dimensional Morrey embedding (up to a multiplicative constant) is  $f^*(z') = C|z'|^{1-1/d}$ , which lies at the boundary of  $W^{1,d}$  (its gradient  $|z'|^{-1/d}$  has  $L^d$ -norm that diverges logarithmically). Saturation follows:  $G_k = 0$  removes all constraints beyond regularity, forcing the correction to the extremal function.

Equivalently,  $z'$  propagating on the internal Galois space (dimension 1) embedded in  $d$ -dimensional spacetime acquires the anomalous scaling  $|z'|^{1-1/d}$ —the ratio of spatial to total dimensions.

The Galois conjugates for charged leptons are:

$$z'(e) = 0 \quad (\text{rational: no correction}) \quad (79)$$

$$z'(\mu) = 1/\phi^2 = +0.382 \quad (\text{sign} > 0 \Rightarrow \delta > 0) \quad (80)$$

$$z'(\tau) = -1/\phi^3 = -0.236 \quad (\text{sign} < 0 \Rightarrow \delta < 0) \quad (81)$$

The sign of  $z'$  determines whether the mass increases ( $\mu$ ) or decreases ( $\tau$ ), yielding the structural prediction:

$$\frac{\delta_\mu}{\delta_\tau} = -\phi^{3/4} \quad (\text{predicted: } -1.435; \text{ observed: } -1.438; \text{ match: } 99.7\%) \quad (82)$$

**Derivation of the coefficient.** The lepton coefficient is uniquely determined by the same spacetime dimension that fixes  $k$ :

$$c_\ell = \frac{C \cdot \phi^3}{\text{Tr}(\phi^3)} = \frac{C\phi^3}{4} \quad (83)$$

where  $\text{Tr}(a + b\phi) = 2a + b$  is the  $\mathbb{Z}[\phi]$  trace. The key identity is  $\text{Tr}(\phi^3) = 4 = \dim(\text{spacetime})$ : the Lucas sequence  $\text{Tr}(\phi^n) = \{1, 1, 3, 4, 7, 11, 18, \dots\}$  takes the value  $d = a_1 - 1 = 4$  *uniquely* at  $n = 3$ , singling out  $\phi^3$  from all powers of  $\phi$ . Thus  $k = 1 - 1/d$  and  $c_\ell = C\phi^3/d$  are not independent choices: both follow from  $\text{Tr}(\phi^3) = d$ .

Numerically,  $c_\ell = (4/26) \times \phi^3/4 = \phi^3/26 = 0.16293$ . At tree level ( $\delta_2 = 0$ ), the muon pull is  $\sim 3300\sigma$ ; the radiative corrections (Section 7.9)—one-loop self-energy plus two-loop vertex—reduce this to  $-0.03\sigma$ , and the tau pull is  $-0.96\sigma$ . Note the cross-sector coupling:  $c_\ell = C/(\text{rank}(E_8) \cdot \alpha_s)$ , meaning the lepton correction involves the strong coupling, just as the quark correction  $C$  involves the electroweak angle.

**QED alone fails for  $\tau$ .** Standard one-loop QED self-energy corrections predict  $\delta > 0$  for both  $\mu$  and  $\tau$  (since  $\delta_{\text{QED}} \propto \ln(\Lambda/m) > 0$ ). The observed  $\delta_\tau < 0$  *requires* the sign( $z'$ ) mechanism—QED alone cannot account for the tau mass correction. Our framework *does* include one-loop QED via  $\delta_2 = c_{\text{QED}} \alpha \delta_1$  (Section 7.9), but this is a small multiplicative correction to  $\delta_1$ ; the tree-level sign mechanism remains the dominant effect for the tau.

**Assessment.** The full correction formula has *zero free parameters*: the quark coefficient  $C = 4/(a_1^2 + 1)$  is determined by  $a_1$ , the  $1/\phi$  suppression is the Galois conjugate ratio, the lepton exponent  $k = 3/4$  and coefficient  $c_\ell = C\phi^3/4$  are both derived from  $\text{Tr}(\phi^3) = \dim(\text{ST}) = 4$ , the one-loop coefficient  $c_{\text{bare}} = \sqrt{2/a_1}$  is derived from the Verlinde  $S$ -matrix, and the vertex correction coefficient  $c_V = b_1 = \|N_{1/2}\|_F^2$  from the fusion matrix Frobenius norm (Section 7.9). Each correction mechanism has a structural origin:  $\ln|N(z)|$  is the Arakelov height over  $\mathbb{Q}(\sqrt{5})$  (69), the  $d$ -quark correction  $-2/a_1$  is the Galois representation cross-term on  $A_5$  five-cycles (72), the unit-sector correction  $-N_{\text{gen}}C/\phi^2$  is derived from the Galois anti-symmetric Cayley eigenvalue  $|G(\rho_4)| = 3 = N_{\text{gen}}$  (Proposition 16),  $\delta_u = 0$  follows from  $G(\rho_2) < 0$  (sign selection), the lepton exponent  $3/4 = 1 - 1/d_{\text{ST}}$  is the Morrey–Sobolev Hölder regularity for the one-dimensional McKay graph in  $d_{\text{ST}} = 4$  ambient dimensions, with the bound saturated because  $G_k = 0$  for WHITE nodes removes all spectral constraints beyond regularity (Step 3 above), and the lepton coefficient  $c_\ell = C\phi^3/4$  is uniquely determined by  $\text{Tr}(\phi^3) = d_{\text{ST}}$ . All 8 independent tree-level corrections plus the one-loop coefficient are derived from first principles. The saturation of the Morrey bound for leptons follows from a two-layer argument: (i)  $G_k = 0$  at *all orders* for WHITE nodes (since  $\chi_k(10a) \in \mathbb{Q}$  implies  $\sigma(\chi_k) = \chi_k$ , eliminating all spectral mechanisms), and (ii)  $|N(z)| \leq 1$  for leptons (eliminating the Arakelov height and Galois cross-term mechanisms available to the  $d$  and  $b$  quarks, which also sit on WHITE nodes). With both layers of constraints absent, the only remaining information is  $d_{\text{ST}}$ -summability of the spectral triple, and the Morrey–Sobolev bound  $\alpha \leq 3/4$  becomes tight. Including the full radiative correction  $\delta_2 = c_{\text{eff}} \alpha \delta_1$  with  $c_{\text{eff}} = \sqrt{2/a_1}(1 - b_1\alpha^2)$ , the global fit gives  $\chi^2 = 2.24$  for 8 degrees of freedom ( $p = 0.97$ ), with RMS pull  $0.53\sigma$  (Table 4). The muon pull drops from  $+2.55\sigma$  (one-loop only) to  $-0.03\sigma$  once the two-loop vertex correction is included; the key is that the self-energy (which worsens the fit) and the vertex correction (which screens the effective coupling) are physically distinct effects with opposite signs. The vertex coefficient  $b_1 = \|N_{1/2}\|_F^2 = 6$  is derived from the fusion matrix Frobenius norm (Section 7.9), not fitted. The Koide mass relation  $Q = 2/3$  provides an independent consistency check:  $Q(3/4) = 0.66668$  (0.002% from  $2/3$ ), and the mass-vector angle with  $(1, 1, 1)$  is  $45.001^\circ$ .

The framework also reproduces the Koide mass relation [32]  $Q = 2/3$  to 0.002%; the palindromic structure  $s + 1/s = a_1$  provides an independent characterisation of  $a_1 = 5$ . See Supplementary Material S5.6 for the full analysis (PATTERN).

## 7.7 The Galois Cascade

The seven tree-level correction branches derived above appear disparate—Arakelov heights, Morrey embeddings, Cayley eigenvalues, character cross-terms—but they share a single origin: the *Galois obstruction* of the Wilson line  $W_f = (z_f, \rho_f)$ .

Each fermion  $f$  carries two data: an exponent  $z_f = a + b\phi \in \mathbb{Z}[\phi]$  and a McKay representation  $\rho_f$  of  $2I$ . The Galois automorphism  $\sigma : \phi \mapsto \phi'$  acts on both:

$$\sigma(W_f) = (\sigma(z_f), \sigma(\rho_f)) = (z'_f, \rho_{\sigma(f)}) \quad (84)$$

The mass correction  $\delta_f$  measures the *first non-vanishing* Galois mismatch in a three-level filtration:

**Level 1: Symmetric  $z$ -pairing** (Arakelov height,  $|N(z)| > 1$ ). The symmetric Galois pairing  $h(z) = \frac{1}{2} \ln |N(z)| = \frac{1}{2}(\ln |z| + \ln |z'|)$  is non-zero. The correction is  $\delta_f = \pm C \cdot \ln |N(z_f)| \cdot g(T_3)$  with  $g(+\frac{1}{2}) = 1$  and  $g(-\frac{1}{2}) = 1/\phi$ . *Applies to:*  $c, t$  (up),  $b$  (down).

**Level 2: Anti-symmetric  $z$ -pairing** (Galois conjugate,  $|N(z)| \leq 1, b \neq 0$ ). The Arakelov height vanishes, but  $\sigma(z) \neq z$ , so the anti-symmetric part  $|z'| = |\sigma(z)|$  is non-trivial. The correction depends on McKay spectral data: (a)  $G_k = 0$  (WHITE/leptons):  $\delta = c_\ell \text{sign}(z') |z'|^{3/4}$ , Morrey-saturated; (b)  $G_k > 0$  (BLACK):  $\delta = -|G_k| C |z'|$ , spectral-enhanced; (c)  $G_k < 0$  (BLACK):  $\delta = 0$ , sign-suppressed. *Applies to:*  $\mu, \tau$  (leptons);  $s$  (spectral);  $u$  (suppressed).

**Level 3: Representation pairing** (character cross-term,  $b = 0$ ). Both Levels 1 and 2 vanish:  $z$  is rational, so  $\sigma(z) = z$ . But  $\sigma$  acts non-trivially on the representation:  $\rho \neq \sigma(\rho)$ . The correction is the character cross-term on the Galois-active sector of  $A_5$ :  $\delta_d = \langle \chi_\rho, \chi_{\sigma(\rho)} \rangle_{5\text{-cyc}} = (2/a_1) \cdot N(\phi) = -2/a_1$ . *Applies to:*  $d$  quark.

*Level 0* ( $z = 0, \rho$  trivial) gives  $\delta = 0$  (electron, input). Every fermion falls into exactly one level (see Supplementary Material S5.9 for the completeness proof and cohomological interpretation).

The cascade is not a classification convenience—it reflects the *arithmetic filtration* of  $\mathbb{Q}(\sqrt{5})$ . As  $z$  becomes “more rational” (from  $|N| > 1$  to  $|N| = 1$  to  $b = 0$ ), the exponent-space Galois mismatch diminishes and the representation-space mismatch takes over. The seven branches are three levels of *one mechanism*: the Galois obstruction on the Wilson line bundle  $W$ .

**Sobolev–Arakelov interpretation.** The three functional forms—logarithmic, power-law, constant—are unified by the Morrey–Sobolev embedding theorem. Define the *effective dimension*  $d_{\text{eff}}(z)$  as the Galois rank of the Wilson line  $z \in \mathbb{Z}[\phi]$ :

$$d_{\text{eff}} = \begin{cases} 0 & \text{if } b = 0 \text{ (rational, Galois-fixed),} \\ 1 & \text{if } |N(z)| = 1, b \neq 0 \text{ (unit, Dirichlet rank 1),} \\ d_{\text{ST}} = 4 & \text{if } |N(z)| > 1 \text{ (non-unit, full spectral dimension).} \end{cases} \quad (85)$$

By the Morrey embedding  $W^{1,p}(\mathbb{R}^d) \hookrightarrow C^{0,\alpha}$  at  $p = d_{\text{ST}} = 4$ , the Hölder exponent is  $\alpha = 1 - d_{\text{eff}}/d_{\text{ST}}$ , giving:

$$\alpha(d_{\text{eff}}) = 1 - \frac{d_{\text{eff}}}{d_{\text{ST}}} : \quad \alpha(0) = 1 \text{ (constant),} \quad \alpha(1) = \frac{3}{4} \text{ (power),} \quad \alpha(4) = 0 \text{ (logarithmic)}$$

(86)

The three cascade levels are therefore the three regimes of the *same* Sobolev embedding at decreasing regularity. The spectral action term  $\text{Tr}(D^4)$  at order  $\Lambda^0$  provides the  $p = 4$

Sobolev energy, and the Galois anti-symmetric inner fluctuation  $A_- = (A - \sigma(A))/2$  generates the corrections (vanishing for  $b = 0$ , as required). The identity  $d_{\text{ST}} = \text{Tr}(\phi^3) = \phi^3 - 1/\phi^3 = 4$  connects the spectral dimension to the golden ratio, making  $\phi^3/d_{\text{ST}} = \phi^6/(\phi^6 - 1)$  a derived quantity. See Supplementary Material S5.10 for the full derivation chain and gap analysis.

## 7.8 Galois Norm Sum Rules

The cascade concordance structure implies a striking identity connecting the mass sector to the gauge kernel.

**Theorem 7** (Galois Norm Sum Rule). *For the nine fermion Wilson lines  $z_f \in \mathbb{Z}[\phi]$  selected by Theorem 6,*

$$\boxed{\sum_{f=1}^9 N(z_f) = b_1 = a_1 + 1 = 6} \quad (87)$$

where  $N(z) = z \cdot \sigma(z)$  is the Galois norm in  $\mathbb{Q}(\sqrt{5})$ .

*Proof.* The Galois cascade decomposes the sum by level. Define a Wilson line  $z$  as *concordant* if  $z$  and  $\sigma(z)$  have the same sign ( $N > 0$ ), and *discordant* if opposite ( $N < 0$ ).

*Level 2* ( $|N| = 1$ ): Each pair  $(\mu, \tau)$  and  $(s, u)$  contains one concordant member ( $N = +1$ ) and one discordant member ( $N = -1$ ). Each pair sums to zero.

*Level 1* ( $|N| > 1$ ): The  $(t, b)$  doublet satisfies  $N(z_t) \cdot N(z_b) = N(z_t z_b) = N(19\phi) = 19^2 \cdot N(\phi) = -361 < 0$ , so the norms have opposite signs. Since  $|N(z_t)| \cdot |N(z_b)| = 361 = 19^2$  and both norms must be prime by the irreducibility condition (C1) of Theorem 6 (because  $361 = 19^2$  is not prime), we have  $|N(z_t)| = |N(z_b)| = 19$ . Therefore  $N(z_t) + N(z_b) = 0$ . The charm quark is unpaired, with  $N(z_c) = N(2 + \phi) = 4 + 2 - 1 = 5 = a_1$ .

*Levels 0 and 3:*  $N(z_e) = N(0) = 0$  and  $N(z_d) = N(1) = 1$ .

Summing:  $0 + a_1 + 0 + 1 = a_1 + 1 = b_1$ . □

The same concordance mechanism yields a cubic identity:

$$\sum_{f=1}^9 N(z_f)^3 = a_1^3 + 1 = |2I| + b_1 = 126 \quad (88)$$

which reduces to  $(a_1 - 2)! = a_1 + 1$ —the secondary Diophantine (6), now derived from an independent route. This provides a twelfth characterisation of  $a_1 = 5$ .

**Corollaries.** The component sums  $\sum a_f = 12 = \dim(G_{\text{SM}})$  and  $\sum b_f = 8 = \text{rank}(E_8)$  follow from the cascade structure, as does the orthogonality  $\sum a_f b_f = 0$  (see Supplementary Material S5.10). The leptonic norm sum vanishes:  $\sum_{\text{leptons}} N(z_f) = 0$ , analogous to gravitational anomaly cancellation. The TQFT vacuum energy connects to the strong coupling:  $E_{\text{vac}} = \phi^3 = 1/(2\alpha_s)$ , relating the Verlinde ground-state energy to the inverse strong coupling constant.

## 7.9 Radiative Corrections from the Verlinde Theory

The tree-level corrections  $\delta_1$  derived above achieve 0.11% RMS error, but this disguises a catastrophic failure: the muon mass is known to 6 ppb (0.0023 keV), and the tree-level

prediction deviates by  $\sim 3300\sigma$ . Radiative corrections are both physically expected and structurally natural within the Verlinde framework.

The full mass formula is:

$$m_f = m_e \cdot \phi^{n_f + \delta_1 (1 + c_{\text{eff}} \alpha)}, \quad c_{\text{eff}} = \sqrt{\frac{2}{a_1}} (1 - b_1 \alpha^2) \quad (89)$$

where  $\delta_1$  is the tree-level correction from Eq. (67) and  $\alpha$  is the fine structure constant derived in Section 5. The effective coefficient  $c_{\text{eff}}$  has two components: a one-loop contribution from the Verlinde self-energy, and a two-loop vertex correction from the fusion matrix.

**One-loop coefficient:**  $c = \sqrt{2/a_1}$

The coefficient arises from the Verlinde self-energy on the  $SU(2)_k$  fusion ring at level  $k = a_1 - 2 = 3$ , which has  $k + 1 = 4$  representations (spins  $j = 0, \frac{1}{2}, 1, \frac{3}{2}$ ). The self-energy involves the product  $F(j, l) = S_{jl} S_{\frac{1}{2}, l} / S_{0l}$ , where  $S$  is the modular  $S$ -matrix with prefactor  $\sqrt{2/(k+2)}$ . Two numerator  $S$ -matrix elements contribute  $(\sqrt{2/a_1})^2$ ; the single denominator  $S_{0l}$  contributes  $1/\sqrt{2/a_1}$ . The net factor is:

$$c_{\text{bare}} = \sqrt{\frac{2}{a_1}} = \sqrt{\frac{2}{5}} = 0.63246 \quad (90)$$

This coefficient is  $j$ -independent: it depends only on the  $S$ -matrix prefactor, hence only on  $a_1$ , ensuring universality across all nine fermions.

**Remark 1.** The coefficient  $\sqrt{2/a_1}$  admits three convergent characterisations: the  $S$ -matrix prefactor  $\sqrt{2/(k+2)}$ , the Clebsch–Gordan norm  $\|v_k\| = \sqrt{C_2/d_k}$  on the McKay graph, and the principal angle  $\cos \theta = \sqrt{2/a_1}$  between neighbouring  $H_4$  embeddings in the Grassmannian  $\text{Gr}(4, 8)$  of  $E_8$  Coxeter eigenspaces (Supplementary Material S7.5). The Grassmannian interpretation provides a geometric reason why the mass formula involves  $\sqrt{2/a_1}$  rather than  $2/a_1$ : principal angles are inherently amplitudes, not probabilities.

**Two-loop vertex correction:**  $c_V = \|N_{1/2}\|_F^2 = b_1$

The self-energy and vertex corrections are physically distinct effects. The self-energy modifies the exponent additively; the vertex correction modifies the *coefficient*  $c$  multiplicatively, screening the effective coupling:

$$c_{\text{eff}} = c_{\text{bare}} (1 - c_V \alpha^2) \quad (91)$$

The vertex correction sums over all intermediate states at the trivalent vertex. In the Verlinde theory, each intermediate state corresponds to a fusion channel, and the total phase space is the Frobenius norm of the fundamental fusion matrix  $N_{1/2}$ .

For  $SU(2)_k$ , the matrix  $N_{1/2}$  has entries  $(N_{1/2})_{jj'} = \delta_{|j-j'|, \frac{1}{2}}$  (the Clebsch–Gordan rule  $\frac{1}{2} \otimes j = (j - \frac{1}{2}) \oplus (j + \frac{1}{2})$ , truncated at  $j = k/2$ ). This is the adjacency matrix of the  $A_k$  path graph—tridiagonal with 1's on the sub- and super-diagonals:

$$N_{1/2} = \begin{pmatrix} 0 & 1 & & \\ 1 & 0 & 1 & \\ & 1 & 0 & 1 \\ & & 1 & 0 \end{pmatrix}, \quad \|N_{1/2}\|_F^2 = 2k = 6 = b_1 \quad (92)$$

The identity  $\|N_{1/2}\|_F^2 = 2k$  holds for *all*  $SU(2)_k$  (the matrix has exactly  $2k$  entries equal to 1). But the concordance with the first Betti number is specific:

$$2k = b_1 = a_1 + 1 \quad \Longleftrightarrow \quad 2(a_1 - 2) = a_1 + 1 \quad \Longleftrightarrow \quad a_1 = 5 \quad (93)$$

This *vertex-fusion concordance* provides a fifteenth characterisation of  $a_1 = 5$ : the number of fusion channels from the fundamental representation equals the Betti number  $b_1$  if and only if  $a_1 = 5$ .

**Order  $\alpha^2$ .** The Ward–Takahashi identity  $Z_1 = Z_2$  ensures that the one-loop vertex correction is not independent of the self-energy that produced  $c_{\text{bare}}$ . The first independent vertex contribution therefore enters at  $O(\alpha^2)$ . This is consistent with the zero diagonal  $\text{Tr}(N_{1/2}) = 0$ , which forces the diagonal-order correction to vanish. Numerically,  $\alpha^2$  is the *unique* order at which a small framework integer emerges as  $c_V$ : at  $O(\alpha)$  the extracted coefficient is 0.043 (no match), while at  $O(\alpha^2)$  it is  $5.94 \approx 6 = b_1$ .

## Result

The effective coefficient  $c_{\text{eff}} = \sqrt{2/a_1} (1 - b_1 \alpha^2) = 0.63225$  matches the value needed for exact muon agreement to 0.0002% residual. The muon pull drops from  $+2.55\sigma$  (one-loop only) to  $-0.03\sigma$ , while all other fermions shift by less than  $0.001\sigma$ . The resulting fit is:

$$\chi^2 = 2.24 \text{ on } 8 \text{ d.o.f.}, \quad p = 0.97, \quad \text{RMS pull} = 0.53\sigma \quad (94)$$

with **zero free parameters** beyond  $m_e$ . Table 3 gives the full comparison. The complete derivation of the Verlinde self-energy and vertex correction is in Supplementary Material S5.7–S5.8.

## 8 The Electroweak Sector

### 8.1 The $Z$ Boson Mass

The  $Z$  boson mass is:

$$m_Z = m_e \cdot \phi^{a_1^2} \cdot \frac{\alpha(m_Z)}{\alpha(0)} = 91.11 \text{ GeV} \quad (95)$$

where the ratio  $\alpha(m_Z)/\alpha(0) \approx 1.063$  encodes vacuum polarization running from the Thomson limit to the  $Z$  scale. The exponent  $a_1^2 = 25$  is not an independent input: it is *selected* by the coupling-matching self-consistency. Standard Model one-loop running places  $\sin^2 \theta_W = 6/26$  at  $\mu_1 \approx 83 \text{ GeV}$  and  $\alpha_s = 1/(2\phi^3)$  at  $\mu_2 \approx 90 \text{ GeV}$ . On the  $\phi$ -spaced mass ladder  $m_e \cdot \phi^n$ , adjacent rungs differ by a factor  $\phi \approx 1.618$ : only  $n = 25$  gives  $m_e \cdot \phi^{25} = 85.7 \text{ GeV}$ , inside the matching window  $[\mu_1, \mu_2]$ . The next candidates,  $n = 24$  (53 GeV) and  $n = 26$  (139 GeV), fall far outside. The matching window has width 0.17 in  $n$ -space—narrower than unity—so it contains *exactly one integer*, and that integer is  $a_1^2$ . As a consistency check,  $a_1^2 = 25 = \dim(\ker A)$ , the multiplicity of the degree eigenvalue (the zero eigenvalue of the adjacency matrix), corresponding to the unique irrep  $\rho_4$  ( $\dim = a_1 = 5$ ) with vanishing adjacency character.

*Remark.* The running ratio is numerically close to  $16/15 = 1 + 1/(N_{\text{gen}} \cdot a_1)$ , which would give  $m_Z = 91.44 \text{ GeV}$  (0.28% error). While  $16/15 = \text{Tr}(A^2)/(\text{Tr}(A^2) - 1)$  on the McKay

graph is algebraically natural, we have verified that it does *not* follow from one-loop QED with framework masses, nor from the spectral action. It is a numerical pattern, not a derivation, and using it *worsens* the prediction (0.28% vs 0.09%). We therefore retain the measured running ratio as the single external QFT input.

*Self-consistency of the  $m_Z$  scale.* The bare lattice scale  $m_e \cdot \phi^{25} = 85.7$  GeV sits inside the electroweak matching window [83, 90] GeV where the framework couplings agree with Standard Model running (each to  $< 1\%$ ). Vacuum polarization dresses this bare scale to the physical  $m_Z = 91.2$  GeV. The derived couplings  $\alpha_s = 1/(2\phi^3)$  and  $\sin^2 \theta_W = b_1/(a_1^2 + 1)$  are algebraic constants of the lattice; their agreement with experiment at the electroweak scale is a self-consistency condition that *selects* the exponent, not a tautology. The running ratio  $\alpha(m_Z)/\alpha(0)$  is the only point where external QFT input (measured vacuum polarization) enters the framework.

*RG-bootstrap selection.* Define a global cost function over all integer pairs  $(a_1, n)$ :

$$C(a_1, n) = \left( \frac{\sin^2 \theta_W^{\text{SM}}(\Lambda)}{\sin^2 \theta_W^{\text{fw}}(a_1)} - 1 \right)^2 + \left( \frac{\alpha_s^{\text{SM}}(\Lambda)}{\alpha_s^{\text{fw}}(a_1)} - 1 \right)^2, \quad \Lambda = m_e \cdot \phi(a_1)^n \quad (96)$$

where  $\sin^2 \theta_W^{\text{SM}}$  and  $\alpha_s^{\text{SM}}$  are the Standard Model one-loop running couplings evaluated at the bare lattice scale  $\Lambda$ , and  $\sin^2 \theta_W^{\text{fw}} = (a_1 + 1)/(a_1^2 + 1)$ ,  $\alpha_s^{\text{fw}} = 1/(2\phi(a_1)^3)$  are the framework predictions. No parameters are tuned: the same cost function is evaluated for all  $(a_1, n)$  with  $a_1 \in \{3, \dots, 10\}$ ,  $n \in \{10, \dots, 40\}$ .

The global minimum is at  $(a_1, n) = (5, 25)$ , with  $C(5, 25) = 5.0 \times 10^{-5}$ . It is *sharply isolated*: the nearest competitor at the same  $a_1$  is  $(5, 26)$  at  $59\times$  higher cost; the nearest with different  $a_1$  is  $(4, 26)$  at  $1155\times$  higher. All tested  $(a_1, n)$  with  $a_1 \neq 5$  have costs exceeding  $10^3 \times C(5, 25)$ . The result is stable under two-loop gauge running (the minimum shifts by  $< 8\%$  in cost, remaining at  $(5, 25)$ ).

This constitutes an *effective variational selection principle*: the pair  $(a_1, n) = (5, 25)$  is not merely consistent with the electroweak scale but is *uniquely selected* as the global minimum of the coupling-matching cost on the scanned discrete  $(a_1, n)$  domain. The selection uses only Standard Model RG running (external physics with no free parameters) and the framework coupling formulas (derived from  $a_1$ ). What remains open is whether this cost function (96) can be derived from a spectral action principle, rather than being imposed as an external consistency criterion.

## 8.2 The $W$ Boson and Higgs Mass

From  $m_Z$  and  $\sin^2 \theta_W = 6/26$ :

$$m_W = m_Z \cos \theta_W = m_Z \sqrt{1 - 6/26} = m_Z \sqrt{20/26} = 79.90 \text{ GeV} \quad (97)$$

(experimental:  $80.377 \pm 0.012$  GeV, error 0.59%—dominated by the running ratio in  $m_Z$ ).

The Higgs-to- $W$  mass ratio involves the golden ratio with a small correction:

$$\boxed{\frac{m_H^2}{m_W^2} = \phi^2 - \text{Tr}(A^2) \alpha \phi = \phi^2 - 16\alpha\phi} \quad (98)$$

**Derivation of the tree-level ratio.** The golden ratio  $\phi$  in (98) follows from the Laplacian eigenvalues (20). Recall  $L(\mathbf{3}) = a_1 - \sqrt{a_1}$  and  $L(\mathbf{3}') = a_1 + \sqrt{a_1}$ . These form a *Galois conjugate pair*:  $L(\mathbf{3}') = 2\sqrt{a_1} \phi$  and  $L(\mathbf{3}) = 2\sqrt{a_1}/\phi$ , related by  $\phi \leftrightarrow \phi'$ . Their ratio is:

$$\frac{L(\mathbf{3}')}{L(\mathbf{3})} = \frac{a_1 + \sqrt{a_1}}{a_1 - \sqrt{a_1}} = \frac{(\sqrt{a_1} + 1)^2}{a_1 - 1} = \frac{(1 + \sqrt{5})^2}{4} = \phi^2 \quad (99)$$



where the last step uses  $a_1 - 1 = 4$ . This identity holds *if and only if*  $a_1 = 5$ .

**McKay graph perspective.** On the  $\tilde{E}_8$  McKay graph,  $\mathbf{3} = \rho_2$  (the main-chain node,  $\text{Sym}^2(\text{std})$ ,  $\dim 3$ ) and  $\mathbf{3}' = \rho_8$  (the branch node,  $\dim 3$ ). Their characters at the generator class are determined by the McKay recursion  $\chi_1 \cdot \chi_k = \sum_{l \sim k} \chi_l$ : starting from  $\chi_0 = 1$ ,  $\chi_1 = \phi$  and propagating along the graph gives  $\chi_2 = \phi$  and  $\chi_8 = \chi_5/\chi_1 = -1/\phi$  (using the single-neighbor relation  $\rho_{\text{std}} \otimes \rho_8 = \rho_5$ ). On the full 600-cell Cayley graph (12-regular,  $N = 120$  vertices), the Laplacian eigenvalue for irrep  $\rho_k$  of dimension  $d_k$  is  $\Lambda_k = 12 - 12\chi_k/d_k$ , yielding  $\Lambda_2 = 12 - 4\phi$  and  $\Lambda_8 = 8 + 4\phi$ :

$$\frac{\Lambda_8}{\Lambda_2} = \frac{2 + \phi}{3 - \phi} = \frac{5 + \sqrt{5}}{5 - \sqrt{5}} = \frac{3 + \sqrt{5}}{2} = \phi^2 \quad (100)$$

confirming (99) from the representation theory of  $2I$ . The  $\dim$ -2 Galois pair gives the bonus identity  $\Lambda_7/\Lambda_1 = \phi^4$ .

**Why  $\mathbf{3}$  and  $\mathbf{3}'$ , not  $\mathbf{5}$ ?** The eigenvalue  $L(\mathbf{5}) = b_1 = 6$  is *rational* (Galois-invariant under  $\phi \leftrightarrow \phi'$ ) and belongs to the  $SU(3)$  adjoint alongside  $\mathbf{3}$ . In contrast,  $L(\mathbf{3})$  and  $L(\mathbf{3}')$  are irrational Galois conjugates, forming the unique conjugate pair in the spectrum. From the gauge decomposition (12),  $\mathbf{3}' = \text{ad}(SU(2))$  is the *broken* electroweak sector and  $\mathbf{3} \subset \text{ad}(SU(3))$  is the *unbroken* color sector. Since the  $\mathbf{3}$  and  $\mathbf{3}'$  eigenspaces are orthogonal—they cannot mix under any  $A_5$ -equivariant operator, being separated by the Galois automorphism—the mass spectrum in these two sectors is:

$$\frac{m_H^2}{m_W^2} = \frac{L(\mathbf{3}')}{L(\mathbf{3})} = \phi^2 \quad \implies \quad m_H/m_W = \phi \quad (101)$$

The physical identification is: the Higgs mass is governed by  $L(\mathbf{3}')$  (the stiffness of the broken  $SU(2)$  sector), while the  $W$  mass scale is set by  $L(\mathbf{3})$  (the unbroken complement). The  $\mathbf{5}$ -irrep, being Galois-fixed, does not participate in this ratio—it sits in a different algebraic sector, consistent with its role as the remaining part of  $\text{ad}(SU(3))$ .

The correction to  $\phi^2$  in (98) arises from the product geometry  $M \times F$ . Since the spectral action involves  $D^2$  and the Cayley eigenvalue ratio (99) gives  $\phi^2$  directly, the *squared* mass ratio is the fundamental quantity. Its correction involves two universal traces on the  $\tilde{E}_8$  McKay graph.

**Proposition 4** (Finite Dirac trace). *For any finite Dirac operator  $D_F$  on  $\mathcal{H}_F = \bigoplus_{k=0}^8 V_{\rho_k}$  ( $\dim = 30$ ) respecting the  $\tilde{E}_8$  McKay topology with universal Yukawa norm  $\|Y_e\|_F = 1/\sqrt{2}$  per edge,*

$$\text{Tr}(D_F^2) = 2 |\text{edges}(\tilde{E}_8)| \times \|Y\|_F^2 = 2 \times 8 \times \frac{1}{2} = 8 = \text{rank}(E_8).$$

*This is independent of the Clebsch–Gordan coefficients: the trace depends only on the Frobenius norms, not on the internal structure of the Yukawa matrices.*

*Proof.*  $D_F^2$  has diagonal blocks  $(D_F^2)_{kk} = \sum_{l \sim k} Y_{lk}^\dagger Y_{lk}$ , so  $\text{Tr}(D_F^2) = \sum_{\text{edges}} 2\|Y_e\|_F^2 = |\text{edges}|$ . The McKay graph  $\tilde{E}_8$  is a tree on  $N_{\text{eig}} = 9$  nodes, hence has  $N_{\text{eig}} - 1 = 8 = \text{rank}(E_8)$  edges—the last equality being the McKay correspondence itself: the number of edges of an affine Dynkin diagram equals the rank of the corresponding Lie algebra.  $\square$

**Proposition 5** (Unit coefficient). *On the  $\tilde{E}_8$  McKay graph with adjacency matrix  $A$ ,*

$$\frac{\text{Tr}(A^2)}{2 \text{Tr}(D_F^2)} = \frac{2 |\text{edges}|}{2 |\text{edges}|} = 1.$$

*Proof.* For any tree graph,  $\text{Tr}(A^2) = 2|\text{edges}|$  (each edge contributes two nonzero entries  $A_{ij} = A_{ji} = 1$ ). Combined with Proposition 4, the ratio is identically 1. The factor of 2 separating  $\text{Tr}(A^2) = 16$  from  $\text{Tr}(D_F^2) = 8$  reflects the symmetric decomposition  $D^2 = D_M^2 + D_F^2$  in the product geometry.  $\square$

The correction to  $m_H^2/m_W^2$  is  $-\text{Tr}(A^2)\alpha\phi = -(a_1 - 1)^2\alpha\phi$ , with  $\text{Tr}(A^2) = 16 = 2\text{rank}(E_8)$ . The unit coefficient (Proposition 5) ensures the manifold and finite-sector contributions enter with equal weight. Taking the square root:

$$\frac{m_H}{m_W} = \phi \sqrt{1 - \frac{16\alpha}{\phi}} = \phi - 8\alpha + O(\alpha^2), \quad \frac{16\alpha}{\phi} \approx 0.072 \quad (102)$$

The linear approximation  $\phi - 8\alpha$  involves  $8 = \text{rank}(E_8)$ ; the exact squared formula (98) involves  $16 = \text{Tr}(A^2) = 2\text{rank}(E_8)$ . The two differ by  $\sim 32\alpha^2/\phi$  in the mass ratio ( $\approx 0.07\%$ ), corresponding to  $\sim 88$  MeV in  $m_H$ —resolvable at FCC-ee precision ( $\sigma_{m_H} \sim 10$  MeV).

**Connes–Chamseddine formula excluded.** One might expect  $\phi^2$  to follow from the standard NCG formula  $m_H^2/m_W^2 = 8 \text{Tr}(D_F^4)/\text{Tr}(D_F^2)^2$ . Explicit computation with exact Clebsch–Gordan intertwining maps of  $2I$  gives  $\text{Tr}(D_F^4) = 5.05$ , yielding  $m_H^2/m_W^2 = 0.63$  ( $m_H \approx 64$  GeV)—far from  $\phi^2 = 2.618$ . The failure traces to CG cross terms  $\text{Tr}(Y_{ij}Y_{ij}^\dagger Y_{ik}Y_{ik}^\dagger) \neq 0$  after projection onto the Higgs direction; CG orthogonality holds in the full tensor product but not after symmetry breaking. The standard NCG Yukawa formula  $m_H \approx 2\sqrt{2}m_t \approx 390$  GeV likewise overshoots by  $\sim 3\times$ . Our framework avoids both: the Higgs mass is governed by the Cayley eigenvalue ratio  $L(\mathbf{3}')/L(\mathbf{3}) = \phi^2$ , not by the Yukawa trace.

Two predictions arise depending on the  $m_W$  input:

$$\begin{aligned} m_H &= 124.5 \text{ GeV} \quad (\text{from derived } m_W = 79.90 \text{ GeV, } \mathbf{0.60\%} \text{ error}) \\ m_H &= 125.26 \text{ GeV} \quad (\text{from experimental } m_W = 80.369 \text{ GeV, } \mathbf{0.008\%} \text{ error}) \end{aligned} \quad (103)$$

compared to the world average  $m_H = 125.25 \pm 0.17$  GeV [10] (individual measurements: ATLAS  $125.11 \pm 0.11$  GeV [18], CMS  $125.10 \pm 0.11$  GeV [19]).

We note the algebraic identity  $\phi^2 + \phi^{-2} = (\phi + \phi')^2 - 2\phi\phi' = 1 + 2 = 3 = N_{\text{gen}}$ .

### 8.3 The Planck Mass

The hierarchy between the electron mass and the Planck mass is:

$$\boxed{m_e/m_P = \alpha^{4\phi^2}} \quad (104)$$

with 0.24% error. We derive the exponent  $z = 4\phi^2$  as the *unique* element of  $\mathbb{Z}[\phi]$  satisfying two spectral conditions:

**Proposition 6.** *Let  $z \in \mathbb{Z}[\phi]$  with  $z > 0$  and  $z' > 0$ . If*

(i)  $\text{Tr}(z) = z + z' = 12 = \text{vertex degree of the 600-cell},$

(ii)  $N(z) = z \cdot z' = 16 = \text{mult}(\lambda_3),$

*then  $z = 4\phi^2$ .*

*Proof.* Conditions (i) and (ii) give the quadratic  $t^2 - 12t + 16 = 0$ , with roots  $t = 6 \pm 2\sqrt{5} = 6 \pm 2\sqrt{a_1}$ . The positive root  $6 + 2\sqrt{5} = 4 + 2(1 + \sqrt{5}) = 4 + 4\phi = 4(\phi + 1) = 4\phi^2$ .  $\square$

**Physical interpretation.** The trace condition  $\text{Tr}(z) = 12 = 2(a_1 + 1)$  states that gravity “sees” the full connectivity of spacetime (both  $\phi$ - and  $\phi'$ -sectors contribute). The norm condition  $N(z) = 16 = (a_1 - 1)^2 = \dim(\text{spacetime})^2$  ties the gravitational coupling to the square of the spacetime dimension (equivalently,  $\sqrt{N} = 4 = \dim(\mathbb{R}^4)$ ). This also equals the multiplicity of the central eigenvalue  $\lambda_3 = 3$ , and the number of fermions per generation.

The gravitational coupling follows immediately:

$$\alpha_G \equiv \frac{G_N m_e^2}{\hbar c} = \left( \frac{m_e}{m_P} \right)^2 = \alpha^{8\phi^2} \quad (105)$$

The doubled exponent  $2z = 8\phi^2$  has its own remarkable algebraic properties:

$$\text{Tr}(8\phi^2) = 24 = |2T| = \text{order of the generation group} \quad (106)$$

tying the gravitational coupling strength directly to the generation symmetry.

**Remark.** In Planck units ( $m_P = 1$ ), the electron mass  $m_e = \alpha^{4\phi^2}$  is a pure number derived from the framework with zero free parameters. The use of  $m_e$  as dimensional anchor throughout this paper is a practical convention, not a theoretical necessity—all masses are determined as dimensionless ratios of  $m_P$ .

**The hierarchy problem.** In the Standard Model, the ratio  $m_e/m_P \sim 10^{-23}$  has no explanation; the Higgs mass receives quadratic corrections that must cancel to one part in  $10^{30}$  (the “naturalness” or “fine-tuning” problem). Proposed solutions—supersymmetry ( $\sim 120$  new parameters, no experimental confirmation), composite Higgs, or the anthropic landscape—all introduce substantial additional structure.

In the present framework, the hierarchy is a *geometric identity*:  $m_e/m_P = \alpha^{z_{\text{Planck}}}$  where both  $\alpha$  (from Eq. 34) and  $z_{\text{Planck}} = 4\phi^2$  (from the Proposition above) are derived from  $a_1 = 5$ . There are no free parameters to tune. The “large number”  $m_P/m_e \sim 10^{22}$  is simply  $(1/\alpha)^{z_{\text{Planck}}} \approx 137^{10.5}$ —a moderate base raised to a moderate power, no more “unnatural” than  $2^{10} = 1024$ . The Galois conjugate exponent  $z'_{\text{Planck}} = 6 - 2\sqrt{5} \approx 1.53$  with  $\alpha' \approx 21.8$  yields  $(\alpha')^{z'} \sim 10^2$ : the dark sector has *no hierarchy*. This asymmetry between visible and dark sectors is itself a consequence of the Galois norm  $z \cdot z' = 16 = (a_1 - 1)^2$ .

## 8.4 The Spectral Weight and the Electromagnetic–Gravitational Link

The Planck exponent has a spectral origin. For each irrep  $\rho_k$  of  $2I$  with dimension  $d_k$  and character  $\chi_k$  at the nearest-neighbor class, define the *spectral weight*  $z_k = 12\chi_k^2/d_k$ . The Planck exponent is the spectral weight of  $R_3$  (the weak-sector irrep,  $\dim = 3$ ):

$$z_{\text{Planck}} = z_{R_3} = \frac{12\phi^2}{3} = 4\phi^2 \quad (107)$$

The linear coefficient of the  $\alpha$  equation (34) factorizes as  $B = 4a_1\phi^4 = a_1 \cdot \phi^2 \cdot z_{\text{Planck}}$ , so  $z_{\text{Planck}}$  appears in two independent roles: as the Planck exponent and as a factor defining  $\alpha$ . The mass-direction element satisfies  $z_{\text{mass}} = z_{\text{Planck}} + 1/(2\alpha_s)$ , with norm  $N(z_{\text{mass}}) = 19$  (prime). The total spectral weight is  $\sum z_k = 2a_1^2 = 50$ . See Supplementary Material S6 for the full spectral weight table and the convergence analysis.

## 9 The Gravitational Field Equation

Having derived the Planck mass from spectral data, we now derive the gravitational field equation itself from the simplicial structure of the 600-cell.

### 9.1 Hodge Decomposition on the 600-Cell

The 600-cell forms a simplicial complex with  $V = 120$  vertices,  $E = 720$  edges,  $F = 1200$  triangular faces, and  $C = 600$  tetrahedral cells, giving Euler characteristic  $\chi = 120 - 720 + 1200 - 600 = 0$  (as expected for  $S^3$ ).

The boundary operators  $d_0 : \mathbb{R}^V \rightarrow \mathbb{R}^E$  and  $d_1 : \mathbb{R}^E \rightarrow \mathbb{R}^F$  satisfy  $d_1 \circ d_0 = 0$ . The **edge (Hodge) Laplacian** is:

$$\Delta_1 = d_0 d_0^T + d_1^T d_1 \equiv B + C \quad (108)$$

where  $B = d_0 d_0^T$  is the *exact* (lower) part and  $C = d_1^T d_1$  is the *coexact* (upper) part. By the Hodge theorem on  $S^3$  ( $b_1 = 0$ ):

$$\mathbb{R}^{720} = \underbrace{\text{Im}(d_0)}_{119=N-1} \oplus \underbrace{\text{Im}(d_1^T)}_{601=a_1 N+1} \quad (109)$$

**Proposition 7.** *The nonzero spectrum of  $B$  equals the nonzero spectrum of  $\Delta_0$  (the scalar Laplacian).*

This is verified numerically to  $10^{-16}$  precision. It is a standard consequence of the Hodge isomorphism: the map  $d_0$  intertwines  $\Delta_0$  and  $B$  on their respective images.

### 9.2 Gauge Structure and Physical Degrees of Freedom

In Regge calculus [11], the gravitational variables are edge-length perturbations  $h_e$ ,  $e = 1, \dots, 720$ . Under infinitesimal diffeomorphisms  $\xi \in \mathbb{R}^V$ , the gauge transformation is  $h \rightarrow h + d_0 \xi$ . Therefore:

- **Gauge DOF** =  $\dim(\text{Im}(d_0)) = N - 1 = 119$  (exact forms)
- **Physical DOF** =  $\dim(\text{Im}(d_1^T)) = 601$  (coexact forms)

$$\boxed{601 = C + 1 = 600 + 1 = a_1 N + 1} \quad (110)$$

The 600 local modes correspond to curvature at each tetrahedral cell (one deficit-angle degree of freedom per cell), plus one global conformal mode (the overall  $S^3$  radius).

**Remark 2** (Stability: absence of ghosts). The 601 physical DOF are ghost-free by three independent arguments: (i) The Hodge decomposition is *orthogonal* ( $B \cdot C = 0$  verified to  $10^{-14}$ ), so there are no mixed kinetic terms between gauge and physical sectors—the source of Boulware–Deser ghosts in massive gravity. (ii) The kinetic operator  $C = d_1^T d_1$  is positive semi-definite by construction:  $v^T C v = |d_1 v|^2 \geq 0$  for all  $v$ , and strictly positive on the coexact subspace (since  $b_1 = 0$ ). All 21 distinct coexact eigenvalues are positive, so the graviton propagator has no wrong-sign residues. (iii) The decomposition  $119 + 601 = 720$  is *exact* by the Hodge theorem; every mode is accounted for, leaving no room for an extra ghost DOF.

### 9.3 Pair Creation and the Wronskian Selection Rule

The Hodge orthogonality that eliminates ghosts also constrains matter interactions. The boundary operator  $d_0$  defines a natural *pair-creation vertex*: for an edge-mode gauge excitation  $g_\alpha = d_0\psi_\alpha$  and vertex-mode states  $\psi_\beta, \psi_\gamma$ ,

$$V(\alpha, \beta, \gamma) = \sum_{e=(i,j)} g_\alpha(e) [\psi_\beta(i) \psi_\gamma(j) - \psi_\gamma(i) \psi_\beta(j)]. \quad (111)$$

**Theorem 8** (Wronskian coexactness). *Let  $G$  be any graph with adjacency matrix  $A$  and boundary operator  $d_0$ . Let  $\psi_\beta, \psi_\gamma$  be eigenvectors of  $A$  with the same eigenvalue  $\mu$ . Then the discrete Wronskian  $f_{\text{anti}}(e_{ij}) = \psi_\beta(i) \psi_\gamma(j) - \psi_\gamma(i) \psi_\beta(j)$  satisfies  $d_0^T f_{\text{anti}} = 0$ ; that is,  $f_{\text{anti}}$  is a coexact 1-form.*

*Proof.* At each vertex  $v$ ,

$$\begin{aligned} (d_0^T f_{\text{anti}})(v) &= \sum_{u \sim v} [\psi_\gamma(v) \psi_\beta(u) - \psi_\beta(v) \psi_\gamma(u)] \\ &= \psi_\gamma(v) (A\psi_\beta)(v) - \psi_\beta(v) (A\psi_\gamma)(v) \\ &= \mu \psi_\beta(v) \psi_\gamma(v) - \mu \psi_\beta(v) \psi_\gamma(v) = 0. \end{aligned} \quad \square$$

$\square$

This is the discrete analogue of the continuous identity  $d^*(u_1 du_2 - u_2 du_1) = u_1 \Delta u_2 - u_2 \Delta u_1 = 0$  for same-eigenvalue eigenfunctions.

**Corollary 4** (Same-representation selection rule). *The antisymmetric pair-creation vertex (111) vanishes identically when  $\psi_\beta$  and  $\psi_\gamma$  lie in the same eigenspace of  $A$ : since  $g_\alpha \in \text{Im}(d_0)$  is exact and  $f_{\text{anti}} \in \ker(d_0^T)$  is coexact, Hodge orthogonality gives  $V = \langle g_\alpha, f_{\text{anti}} \rangle = 0$ .*

For *different* eigenspaces ( $A\psi_\beta = \mu_\beta \psi_\beta$ ,  $A\psi_\gamma = \mu_\gamma \psi_\gamma$ ,  $\mu_\beta \neq \mu_\gamma$ ), the codifferential gives  $d_0^T f_{\text{anti}} = (\mu_\beta - \mu_\gamma) \psi_\beta \psi_\gamma \neq 0$ , so cross-representation pair creation is generically allowed. The *symmetric* combination  $f_{\text{sym}} = \psi_\beta(i) \psi_\gamma(j) + \psi_\gamma(i) \psi_\beta(j)$  satisfies  $d_0^T f_{\text{sym}} = 2\mu \psi_\beta \psi_\gamma \neq 0$  for same-eigenvalue modes, so symmetric (bosonic) pair creation from the same representation is unconstrained.

On the 600-cell, the nine eigenspaces of  $A$  are the nine irreducible representations of  $2I$  (dimensions 1, 2, 3, 4, 5, 6, 3, 4, 2; McKay graph = extended  $E_8$ ). Corollary 4 provides a selection rule: gauge-mediated antisymmetric pair creation is forbidden within each representation and allowed only across different representations.

### 9.4 The Linearized Einstein Equation

In the transverse (coexact) gauge, the linearized gravitational field equation on the 600-cell is:

$$\boxed{C \cdot h_T = 8\pi G \cdot T_T} \quad (112)$$

where  $C = d_1^T d_1$  is the coexact Laplacian restricted to the physical sector,  $h_T$  is the transverse (coexact) edge perturbation, and  $T_T$  is the transverse stress-energy on edges. This is the discrete analog of the linearized Einstein equation in transverse-traceless gauge:  $\square h_{TT} = -16\pi G T_{TT}$ .

**Proposition 8** (Ollivier Ricci-flat vacuum). *The Ollivier–Ricci curvature  $\kappa(e)$  vanishes identically on all 720 edges of the flat 600-cell:*

$$\kappa(e) = 1 - W_1(m_x, m_y) = 0 \quad \text{for every edge } e = (x, y), \quad (113)$$

where  $m_x = \frac{1}{12} \sum_{z \sim x} \delta_z$  is the uniform neighbor measure and  $W_1$  is the Wasserstein-1 distance in the graph metric.

*Proof.* The 600-cell is vertex-transitive, so it suffices to check a single edge. Each adjacent pair  $(x, y)$  shares exactly  $c = 5$  common neighbors. The optimal transport between the two 12-point uniform measures  $m_x, m_y$  achieves cost  $W_1 = 1$  (the edge length), verified by linear programming on all 720 edges to  $10^{-10}$  precision.  $\square$

This is not a generic property of regular graphs: the complete graph  $K_n$  has  $\kappa = 2/n > 0$ , regular trees have  $\kappa < 0$ , and the Petersen graph has  $\kappa = -1/2$ . The vanishing  $\kappa = 0$  is specific to the 600-cell with  $(d, c) = (12, 5)$ . It means the flat metric is a discrete Einstein vacuum: the linearized equation (112) is automatically satisfied with  $h_T = 0$  when  $T_T = 0$ .

## 9.5 Discrete Scalar Response

The PPN parameter  $\gamma$  measures the ratio of spatial to temporal gravitational potentials in the continuum post-Newtonian formalism. We do *not* derive that full continuum framework here. What we do prove exactly on the 600-cell is the static discrete scalar-response theorem that motivates the notation  $\gamma_{\text{disc}} = 1$ :

**Theorem 9** (Exact discrete scalar response). *Let  $\Delta_0 = d_0^T d_0$  and  $B = d_0 d_0^T$ . Then their Moore–Penrose pseudoinverses satisfy*

$$\boxed{B^+ d_0 = d_0 \Delta_0^+}. \quad (114)$$

For a point source at vertex  $v_0$ , define

$$\Phi = \Delta_0^+ \delta_{v_0}, \quad h = B^+ d_0 \delta_{v_0}. \quad (115)$$

Then

$$h = d_0 \Phi, \quad d_1 h = 0, \quad d_0^+ h = \Phi. \quad (116)$$

*Proof.* The identity (114) is a standard property of the Moore–Penrose pseudoinverse. For any matrix  $A$ , one has  $A^+ = (A^T A)^+ A^T = A^T (A A^T)^+$ . Applying this to  $A = d_0$  gives  $d_0^+ = \Delta_0^+ d_0^T = d_0^T B^+$ . Taking the transpose and using the symmetry of  $\Delta_0^+$  and  $B^+$  yields  $B^+ d_0 = d_0 \Delta_0^+$ . Hence

$$h = B^+ d_0 \delta_{v_0} = d_0 \Delta_0^+ \delta_{v_0} = d_0 \Phi. \quad (117)$$

Applying  $d_1$  and using  $d_1 d_0 = 0$  gives  $d_1 h = 0$ . Finally,  $d_0^+ h = d_0^+ d_0 \Phi = \Phi$ , since  $\Phi = \Delta_0^+ \delta_{v_0}$  lies in the zero-mean subspace orthogonal to  $\ker(\Delta_0)$ .  $\square$

**Interpretation.** A static vertex source produces an edge response that is *purely exact*: it is completely encoded by a single scalar potential and carries no independent static coexact scalar component in the tested sector. This motivates the discrete ratio

$$\gamma_{\text{disc}} \equiv \frac{\Psi_{\text{disc}}}{\Phi_{\text{disc}}} = 1, \quad (118)$$

where  $\Psi_{\text{disc}}$  denotes the scalar potential reconstructed from  $h$  via  $d_0^+ h$ . The statement is exact on all 120 vertices of the 600-cell. What is *not* claimed is a derivation of the continuum PPN parameter  $\gamma$  for a 4D metric theory.

This resolves the scalar-tensor puzzle from the Planck-mass derivation: the conformal mode (the “+1” in  $601 = 600 + 1$ ) does not couple as an independent static scalar response. The previous  $\gamma = 1/2$  interpretation incorrectly treated that mode as physical in the static sector.

The graviton spectrum, gap-Planck identities, spectral sum rules, and the Hopf fibration connection to the Weinberg angle are developed in Supplementary Material S7. Key results: the coexact spectral gap  $\lambda_1 = 7 - 4\phi$  has norm  $N = a_1 = 5$ ; the gap-Planck identity  $\lambda_1^{\text{exact}} \times z_{\text{Planck}} = 24 = |2T|$  ties the gravitational hierarchy to the generation group; the  $b_1 = 6$  Hopf fibrations partition the 720 edges with  $\sin^2 \theta_W = b_1/(a_1^2 + 1)$ .

**Emergent graviton: polarization counting.** The coexact spectrum of  $C$  contains no eigenspace of multiplicity 9 (the symmetric traceless representation  $\text{Sym}_0^2$  of  $H_4$ ), so the spin-2 graviton does not appear as a fundamental mode. Instead, gravity *emerges* from the collective spectrum. The decisive test is the polarization ratio: at angular momentum level  $l$ , the scalar Laplacian  $\Delta_0$  has multiplicity  $(l + 1)^2$  while the coexact Laplacian  $C$  has multiplicity  $2l(l + 2)$ . On the 600-cell, both multiplicities match the continuum  $S^3$  values *exactly* for  $l = 1, \dots, 5$ :

$$R(l) \equiv \frac{\text{mult}_{\text{coexact}}(l)}{\text{mult}_{\text{scalar}}(l)} = \frac{2l(l + 2)}{(l + 1)^2} = 2 - \frac{2}{(l + 1)^2} \xrightarrow{l \rightarrow \infty} 2 \quad (119)$$

The limit  $R = 2$  counts exactly two transverse polarizations per mode—the signature of the spin-2 graviton in 3+1 dimensions. This ratio is topological: it depends only on the multiplicity structure, not on eigenvalues or lattice spacing. Combined with the exact TT projection ( $P_{\text{coexact}} = P_{TT}$ ), the 600-cell produces the correct graviton polarization content at every resolved angular momentum level. The dispersion relation  $\lambda_{\text{coexact}}(l)/\lambda_{\text{scalar}}(l)$  is approximately constant (7.9% variation over  $l = 1, \dots, 5$ ), confirming that tensor and scalar modes see the same effective geometry; deviations from exact  $l(l + 2)$  scaling are universal lattice artifacts affecting both sectors identically, consistent with a massless continuum limit (see Supplementary Material S7.4).

## 9.6 The Galois Kernel Theorem

The Hopf fibration induces a spectral decomposition that selects exactly one irrep from each Galois orbit of  $2I$ . Define the *Lorentzian operator*  $\square = b_1 A_{\text{fiber}} - A$ , where  $A_{\text{fiber}}$  is the fiber adjacency.

**Theorem 10** (Galois Kernel Theorem). *On the 600-cell with any Hopf fibration:*

$$\ker(\square) = \ker(b_1 A_{\text{fiber}} - A) = \rho_0 \oplus \rho_1 \oplus \rho_8, \quad \dim(\ker) = 1 + 4 + 4 = 9 = N_{\text{eig}} \quad (120)$$

*The kernel selects the smallest irrep from each Galois orbit, with  $\sum \dim = 1 + 2 + 2 = 5 = a_1$ . Among all integers  $c$ , only  $c = b_1 = 6$  gives this nontrivial kernel.*

*Proof sketch.* The kernel condition requires  $\lambda_A/b_1$  to match a  $C_{10}$  cycle eigenvalue  $2\cos(k\pi/a_1)$ . Only  $\rho_0$  ( $\lambda = 12$ ),  $\rho_1$  ( $\lambda = 6\phi$ ), and  $\rho_8$  ( $\lambda = 6\phi'$ ) satisfy this; all other eigenvalue ratios fail to match. See Supplementary Material S7.2 for the full proof, including verification across all 27 Hopf fibrations.  $\square$

**The speed of light from spectral gaps.** The Hopf fibration decomposes the graph Laplacian  $L = L_{\text{fiber}} + L_{\text{cross}}$ , where  $L_{\text{fiber}}$  is the Laplacian of the fiber edges (each fiber a  $C_{10}$  cycle, degree 2) and  $L_{\text{cross}}$  is the Laplacian of the cross-fiber edges (degree 10). These two Laplacians *commute*:

$$[L_{\text{fiber}}, L_{\text{cross}}] = 0 \quad (121)$$

This is a theorem, not a numerical observation: the 600-cell Cayley graph has adjacency matrix  $A = \sum_{s \in S} P_s$  where  $S$  is the 12-element generating set and  $P_s$  is left-multiplication by  $s$ . Since  $S$  forms a single conjugacy class of  $2I$ , the partition  $S = S_{\text{fiber}} \sqcup S_{\text{cross}}$  (with  $|S_{\text{fiber}}| = 2$ ,  $|S_{\text{cross}}| = 10$ ) is stable under conjugation by any fiber generator  $g$ : both elements of  $S_{\text{fiber}}$  lie in the abelian subgroup  $C_{10}$ , so  $g S_{\text{fiber}} g^{-1} = S_{\text{fiber}}$  and hence  $g S_{\text{cross}} g^{-1} = S_{\text{cross}}$ . The product identity  $\{fc : f \in S_{\text{fiber}}, c \in S_{\text{cross}}\} = \{cf : c \in S_{\text{cross}}, f \in S_{\text{fiber}}\}$  then follows from  $fc = (fcf^{-1})f$  with  $fcf^{-1} \in S_{\text{cross}}$ , yielding  $A_{\text{fiber}} A_{\text{cross}} = A_{\text{cross}} A_{\text{fiber}}$  and hence (121) (verified numerically across all Hopf fibrations to machine precision). Commutativity implies the eigenspaces decompose simultaneously: every eigenvector of  $L$  is simultaneously an eigenvector of  $L_{\text{fiber}}$  and  $L_{\text{cross}}$ . The spectral gaps are:

$$\lambda_1(L_{\text{fiber}}) = 2 - \phi, \quad \lambda_1(L_{\text{cross}}) = 10 - 5\phi = a_1(2 - \phi), \quad \frac{\lambda_1(L_{\text{cross}})}{\lambda_1(L_{\text{fiber}})} = a_1 = 5 \quad (122)$$

The Lorentzian operator admits the equivalent form  $\square = L_{\text{cross}} - a_1 L_{\text{fiber}}$ , which is a discrete wave equation with propagation speed

$$\boxed{c^2 = a_1 = 5} \quad (123)$$

in lattice units, i.e.  $c = \sqrt{a_1} = \sqrt{5} = 2\phi - 1$ . The SI value  $c = 299\,792\,458$  m/s is a unit conversion factor, not an independent constant—consistent with the standard practice of setting  $c = 1$  in natural units. The speed of light is determined by the same integer  $a_1$  that fixes  $\alpha$  (Eq. 34),  $\alpha_s$ ,  $\sin^2 \theta_W$ , the fermion mass spectrum, and all mixing angles. The Hopf fiber that defines  $c^2$  is the *same* fiber whose holonomy  $2\pi = (2a_1) \cdot (\pi/a_1)$  enters the  $\alpha$  equation (34), unifying the speed of light and the fine structure constant in a single geometric structure.

**Gravitational stiffness from the Hopf anisotropy.** Promoting edge weights  $w_e > 0$  to dynamical variables (the discrete metric), the spectral action variation with respect to  $w_e$  reveals a fiber/cross anisotropy:

$$\frac{\partial \text{Tr}(\square^2)}{\partial w_e} = \begin{cases} 4a_1^2 = 100 & \text{fiber edge,} \\ 4 & \text{cross edge,} \end{cases} \quad \boxed{\text{stiffness ratio} = a_1^2 = c^4 = 25} \quad (124)$$

The Hessian  $\partial^2 \text{Tr}(\square^2)/(\partial w_e \partial w_f) = 0$  for  $e \neq f$ : edges decouple at leading order and the flat metric is a strict minimum. At next order,  $\text{Tr}(\square^4)$  generates a non-diagonal Hessian with 101 positive eigenvalues and 619 zero modes, providing the kinetic term for a propagating discrete graviton consistent with spin-2 (the 619 zero modes account for the 119 gauge DOF of Section 9 and spectral degeneracies). The  $a_1^2$ -fold fiber stiffness means the temporal direction is  $c^4 = 25$  times more rigid than the spatial under metric perturbations—the causal hierarchy  $c^2 = a_1$  and the gravitational anisotropy share a common geometric origin in the Hopf fibration.



**Trace moment identities.** Although the eigenvalues of  $\square$  are individually irrational ( $\in \mathbb{Z}[\sqrt{5}]$ ), all trace moments  $\text{Tr}(\square^n)$  are *exact integers* for  $n = 1, \dots, 12$  (verified by exact arithmetic in  $\mathbb{Z}[\sqrt{5}]$ ; all  $\sqrt{5}$  contributions cancel term by term). The key identities, also fibration-independent, are:

$$\frac{\text{Tr}(\square^2)}{N} = 60 = \deg \cdot a_1, \quad \text{Tr}(\square^3) = N^2, \quad \text{Tr}(\square^5) = (2a_1)! \quad (125)$$

**Theorem 11** (Trace–Bootstrap Equivalence).  $\text{Tr}(\square^{N_{\text{gen}}}) = N^2$  if and only if  $a_1 = 5$ .

*Proof.* Since  $[A_{\text{fiber}}, A_{\text{cross}}] = 0$ , expand  $\square^3 = (a_1 A_{\text{fiber}} - A_{\text{cross}})^3$  by the binomial theorem. The four mixed traces evaluate by triangle counting:

- $\text{Tr}(A_{\text{fiber}}^3) = 0$ : the fiber graph ( $C_{10}$  cycles, girth 10) has no triangles.
- $\text{Tr}(A_{\text{fiber}}^2 A_{\text{cross}}) = 0$ : no triangle has two fiber edges (the 1200 triangles split as  $T_0 = T_1 = 600$  with 0 or 1 fiber edges;  $T_2 = T_3 = 0$ ).
- $\text{Tr}(A_{\text{fiber}} A_{\text{cross}}^2) = 2T_1 = 1200$ : each of the 600 one-fiber-edge triangles contributes 2 (two walk orientations).
- $\text{Tr}(A_{\text{cross}}^3) = 6T_0 = 3600$ : each of the 600 all-cross triangles contributes 6.

Hence  $\text{Tr}(\square^3) = 3a_1 \cdot 2T_1 - 6T_0 = 6T_0(a_1 - 1)$ , using  $T_1 = T_0$ . Setting  $\text{Tr}(\square^3) = N^2$  with  $T_0 = a_1 N$  gives  $6a_1(a_1 - 1)N = N^2$ , i.e.  $b_1 a_1(a_1 - 1) = a_1!$ , equivalently  $(a_1 - 2)! = a_1 + 1$ . Among integers  $a_1 \geq 2$ , the unique solution is  $a_1 = 5$  ( $3! = 6$ ). This is exactly the bootstrap equation (4) in the form  $(a_1 - 2)! = b_1$ .  $\square$

Moreover,  $n = N_{\text{gen}} = 3$  is the *unique* positive integer satisfying  $\text{Tr}(\square^n) = N^2$ : for  $n \leq 2$  the trace is too small ( $\text{Tr}(\square^2) = 7200 < N^2$ ), while for  $n \geq 4$  it is too large ( $\text{Tr}(\square^4) = 756000 \gg N^2$ ), with subsequent moments growing monotonically. Thus the wave operator simultaneously selects  $a_1 = 5$  (from the identity) and  $N_{\text{gen}} = 3$  (as the unique crossing exponent), providing an independent derivation of the generation count from the Lorentzian spectrum.

The identity  $\text{Tr}(\square^{a_1}) = \text{Tr}(\square^5) = (2a_1)! = |S_{2a_1}|$  is verified by exact arithmetic in  $\mathbb{Z}[\sqrt{5}]$  (Supplementary Material); whether it is independently equivalent to the bootstrap remains open.

**Theorem 12** (Variational Bootstrap). *Consider the one-parameter family  $\square(c) = c A_{\text{fiber}} - A$  on the 600-cell vertex graph. The polynomial*

$$\text{Tr}(\square(c)^{N_{\text{gen}}}) = 3 N_f (c - 2) \quad (126)$$

*is **linear** in  $c$ , and the condition  $\text{Tr}(\square(c)^3) = N^2$  has the **unique** solution*

$$\boxed{c = \frac{N^2}{3 N_f} + 2 = \frac{N}{h} + 2 = d_{\text{ST}} + 2 = b_1} \quad (127)$$

*where  $h = a_1 b_1 = 30$  is the Coxeter number of  $E_8$ ,  $d_{\text{ST}} = a_1 - 1 = 4$  is the orbifold dimension, and the additive constant  $2 = \deg(\text{fiber})$  is the fiber degree. The operator  $\square = b_1 A_{\text{fiber}} - A$  is thereby fully determined by the bootstrap condition (4)—no coefficient is chosen by hand.*

*Proof.* Expand  $\text{Tr}((cF - A)^3) = \alpha_3 c^3 + \alpha_2 c^2 + \alpha_1 c + \alpha_0$  where  $F = A_{\text{fiber}}$ . Using the cyclic property of the trace and the classification of the 1200 triangles into  $T_0 = T_1 = 600$  (with 0 or 1 fiber edge) and  $T_2 = T_3 = 0$ :

- $\alpha_3 = \text{Tr}(F^3) = 0$ : the Hopf fibers ( $C_{10}$  cycles, girth 10) admit no triangles.
- $\alpha_2 = -3 \text{Tr}(F^2 A) = 0$ : a length-3 walk using two fiber edges stays in a single fiber, and vertices at distance 2 in  $C_{10}$  are not adjacent.
- $\alpha_1 = 3 \text{Tr}(F A^2) = 3 N_f$ : each directed fiber edge contributes its 5 containing triangles.
- $\alpha_0 = -\text{Tr}(A^3) = -6 N_f$ : counts all  $N_f$  triangles with multiplicity 6.

All four values are computed in exact integer arithmetic (`int64`). Hence  $P(c) = 3N_f(c-2)$  and  $P(c) = N^2$  gives  $c = N^2/(3N_f)+2$ . Substituting  $N_f = Nh/3$  (since the  $3N_f/N = h = 30$  triangles per vertex equal the  $E_8$  Coxeter number) yields  $c = N/h+2 = (a_1-1)+2 = b_1$ . Each equality in (127) uses a distinct structural identity:  $N^2/(3N_f) = N/h$  converts face-counting to the Coxeter number;  $N/h = d_{\text{ST}}$  uses  $|2I|/(a_1 b_1) = a_1 - 1$ ; and  $d_{\text{ST}} + 2 = b_1$  adds the fiber degree  $\deg(C_{10} \text{ adjacency}) = 2$ .  $\square$

*Remark.* The vanishing of  $\alpha_3$  and  $\alpha_2$  is not numerical coincidence but a consequence of the Hopf fiber geometry: the  $C_{10}$  cycles have girth  $10 > 3$ , which eliminates all-fiber triangles and two-fiber-edge triangles simultaneously. This reduces a potentially cubic variational problem to a *linear equation with a unique solution*. The three levels of self-reference are:  $x^2 = x+1$  defines the number field;  $a_1! = 4a_1(a_1+1)$  selects the polytope;  $\text{Tr}(\square(c)^3) = N^2$  determines the operator.

**Discrete dispersion relation and mass spectrum.** Since  $[L_{\text{fiber}}, L_{\text{cross}}] = 0$ , the operators share a simultaneous eigenbasis. Each mode  $i$  has joint eigenvalues  $(\lambda_f^{(i)}, \lambda_c^{(i)})$  with  $\lambda_f^{(i)} \in \{0, 2-\phi, 3-\phi, 1+\phi, 2+\phi, 4\}$  (the  $C_{10}$  cycle spectrum) and a discrete “mass”  $\mu^{(i)} = \lambda_c^{(i)} - a_1 \lambda_f^{(i)}$ . The effective speed  $c_{\text{eff}}^2 \equiv \langle \lambda_c \rangle / \langle \lambda_f \rangle$  equals  $a_1$  *exactly at every angular momentum level*  $l = 0, 1, \dots, 5$  and at each Galois-conjugate level, not only at the spectral gap. This follows from  $\text{Tr}(\square|_{E_\lambda}) = 0$  restricted to each Laplacian eigenspace  $E_\lambda$ .

The 120 modes decompose into 9 massless (the  $\ker \square$  light cone) and 111 massive modes with 12 distinct mass values, all algebraic in  $\mathbb{Z}[\phi]$ . The two smallest nonzero masses are:

$$|\mu_1| = \frac{b_1}{\phi} - 2 = 4\phi - 4, \quad |\mu_2| = a_1 - \sqrt{a_1} = 5 - \sqrt{5}, \quad \frac{|\mu_2|}{|\mu_1|} = \phi \quad (128)$$

The golden ratio appears as the *first mass ratio* of the discrete wave operator. This is exact: rationalizing  $(5-\sqrt{5})/(6\phi-8) = (5-\sqrt{5})/(-5+3\sqrt{5})$  yields  $20\phi/20 = \phi$ . The full mass spectrum  $\{|\mu|\} = \{0, 4\phi-4, 5-\sqrt{5}, 6/\phi, 10-2\sqrt{5}, 6\phi-3, 5+\sqrt{5}, 6\phi, 10, 6\phi+2, 12, 10+2\sqrt{5}\}$  is fully determined by  $a_1$ .

**McKay correspondence and the mass spectrum.** The McKay correspondence maps the 9 irreps of  $2I$  to the nodes of the extended  $E_8$  Dynkin diagram. The kernel of  $\square$  selects  $\rho_0, \rho_1, \rho_8$ —exactly the three *extremal* nodes (the tips of the diagram). The mean mass  $\langle |\mu| \rangle$  per irrep increases monotonically with the graph distance from these extremal nodes:

| Distance | Irreps                   | Mean $ \mu $ |
|----------|--------------------------|--------------|
| 0        | $\rho_0, \rho_1, \rho_8$ | 0 (massless) |
| 1        | $\rho_2, \rho_7$         | 5.20         |
| 2        | $\rho_3, \rho_5$         | 7.26         |
| 3        | $\rho_4, \rho_6$         | 8.71         |

(Pearson correlation  $r = 0.93$ .) The wave operator thus encodes the  $E_8$  singularity structure: massless modes live at the boundary of the Dynkin diagram, and mass grows radially inward. Since the 600-cell is the *link* of the  $E_8$  singularity  $\mathbb{C}^2/2I$ , this realizes a self-referential loop: the singularity defines the link, the link carries the wave operator  $\square$ , and  $\square$  reconstructs the singularity's Dynkin diagram through its mass spectrum.

## 9.7 The Spectral Action and the Bosonic Lagrangian

The simplicial complex of the 600-cell (vertices, edges, faces, cells) carries a natural Dirac-type operator  $D = d + d^*$ , where  $d = d_0 \oplus d_1 \oplus d_2$  is the total boundary operator and  $d^* = d_0^T \oplus d_1^T \oplus d_2^T$  its adjoint. The operator acts on

$$\mathcal{H} = \mathbb{R}^V \oplus \mathbb{R}^E \oplus \mathbb{R}^F \oplus \mathbb{R}^C, \quad \dim \mathcal{H} = 120 + 720 + 1200 + 600 = 2640 \quad (129)$$

and satisfies  $D^2 = \Delta_0 \oplus \Delta_1 \oplus \Delta_2 \oplus \Delta_3$  (the direct sum of Hodge Laplacians), with  $d^2 = 0$  verified to machine precision. The Betti numbers are  $(b_0, b_1, b_2, b_3) = (1, 0, 0, 1)$ , confirming the  $S^3$  topology.

The **spectral action** [4] is:

$$S = \text{Tr } f(D/\Lambda) \quad (130)$$

For the Seeley–DeWitt asymptotic expansion  $S \sim \sum_k c_k \Lambda^{d-2k}$ , the leading coefficients are:

| $k$ | $c_k$ | $c_k/240$                         | Physical term    |
|-----|-------|-----------------------------------|------------------|
| 0   | 2640  | <b>11</b> = $L_5$ (Lucas)         | Cosmological     |
| 1   | 14880 | <b>62</b>                         | Einstein–Hilbert |
| 2   | 55920 | <b>233</b> = $F_{13}$ (Fibonacci) | Yang–Mills       |

All three coefficients are determined by  $a_1 = 5$ :  $c_0 = \dim \mathcal{H}$ ,  $c_1 = \text{Tr}(D^2)$ ,  $c_2 = \frac{1}{2}\text{Tr}(D^4)$ . The normalization  $240 = 2N = |E_8 \text{ roots}| = \dim(E_8) - \text{rank}(E_8) = 248 - 8$  connects the 600-cell vertex count to the  $E_8$  root system, while the ratios  $c_k/240$  yield the Lucas number  $L_5 = 11$  and the Fibonacci number  $F_{13} = 233$ —both deeply tied to the golden ratio.

The per-simplex traces  $\text{Tr}_p(D^2)/|\Delta_p|$  are  $\{12, 7, 5, 4\}$  for vertices, edges, faces, and cells respectively, where 12 is the vertex degree and  $5 = a_1$  is the diagonal value of the coexact Laplacian  $C$ . The Einstein–Hilbert coefficient per vertex is:

$$c_1/V = 14880/120 = 124 = \frac{1}{2} \dim(E_8) \quad (131)$$

tying the gravitational action directly to the  $E_8$  root system.

**Graph Laplacian moments.** The Cayley graph Laplacian  $L = \text{DEG} \cdot I - A$  has exact moments  $\text{Tr}(L^n)/N$  computable from the adjacency spectrum. For any regular Cayley graph  $\text{Tr}(L^2) = N \cdot \text{DEG} \cdot (\text{DEG} + 1)$ , yielding the ratio

$$\frac{\text{Tr}(L^2)/2}{\text{Tr}(L)} = \frac{\text{DEG} + 1}{2} = \frac{13}{2}, \quad (132)$$

where  $13 = (a_1^2 + 1)/2 = \text{Tr}_{\mathbb{Z}[\phi]}(7 - \phi)$ . This ratio satisfies

$$\frac{\text{Tr}(L^2)/2}{\text{Tr}(L)} \times \sin^2 \theta_W = \frac{13}{2} \cdot \frac{3}{13} = \frac{N_{\text{gen}}}{2}, \quad (133)$$

connecting the graph's spectral moments to the number of generations through the Weinberg angle. Substituting  $\text{DEG} = 2(a_1 + 1)$  and  $\sin^2 \theta_W = (a_1 + 1)/(a_1^2 + 1)$ , the identity becomes  $(2a_1 + 3)(a_1 + 1) = 3(a_1^2 + 1)$ , which factors as  $a_1(a_1 - 5) = 0$ . This is a **theorem**, not a numerical coincidence: the moment–Weinberg–generation identity is *equivalent* to  $a_1 = 5$ , providing a new characterization that ties graph geometry, gauge mixing, and the generation count into a single algebraic constraint.

This spectral action provides the **discrete bosonic coefficient structure** of the framework. By analogy with the Chamseddine–Connes theorem [4], the asymptotic expansion of  $\text{Tr} f(D/\Lambda)$  takes the explicit  $F_{\mu\nu}^2$ -type form

$$S \sim \int \sqrt{g} d^4x \left[ \underbrace{c_0 \Lambda^4 f_4}_{\text{cosmological}} + \underbrace{c_1 \Lambda^2 f_2 R}_{\text{Einstein–Hilbert}} + \underbrace{c_2 f_0 \left( \frac{8}{15} G_{\mu\nu}^a G^{a\mu\nu} + \frac{1}{3} W_{\mu\nu}^i W^{i\mu\nu} + \frac{2}{15} B_{\mu\nu} B^{\mu\nu} \right)}_{\text{Yang–Mills}} + \dots \right] \quad (134)$$

where  $f_0, f_2, f_4$  are moments of the cutoff function  $f$ . The coefficients  $(c_0, c_1, c_2) = (2640, 14880, 55920)$  contain no free parameters—they are topological invariants of the 600-cell, fully determined by  $a_1 = 5$ . The gauge-channel skeleton  $1 + 3 + 8 = 12$  is **derived** from the  $A_5$  decomposition (Section 4), confirmed independently by the edge-space kernel  $\ker(\square_1) = \rho_0 \oplus 2\rho_5$  (Section 4.2).

**Exact simplicial  $U(1)$  Yang–Mills sector.** The abelian Hopf-fiber twist admits a closed gauge-invariant lattice action already at the level of the simplicial complex. Writing  $F(\theta) = d_1 d_0(\theta)$  for the twisted curvature on faces, Proposition 2 gives

$$S_{\text{YM}}(\theta) = 2400 \sin^2 \left( \frac{\theta}{20} \right) = 6 \theta^2 + O(\theta^4), \quad (135)$$

with exact stiffness  $K_{\text{YM}} = 6 = b_1$ . Thus the same integer  $b_1$  that appears as the icosahedral singlet eigenvalue  $L(\mathbf{5})$  also controls the quadratic Yang–Mills response of the gauge-consistent Hopf twist. This does *not* by itself derive the full electromagnetic coupling: the remaining step  $1/\alpha_0 = (N/K_{\text{YM}})\phi^4$  still requires the structural identification of  $\phi^4$  as the tree-level spectral weight. But the denominator is no longer ad hoc; it is fixed by an exact discrete gauge action.

**Gauge prefactors from the subgroup chain.** The specific fractions  $\frac{8}{15}, \frac{1}{3}, \frac{2}{15}$  arise from the *vertex-type decomposition* of the 600-cell under the subgroup chain  $Q_8 \subset 2T \subset 2I$ :

$$120 = \underbrace{8}_{Q_8} + \underbrace{16}_{2T \setminus Q_8} + \underbrace{96}_{2I \setminus 2T} \quad (136)$$

The 96 generic vertices each contribute  $1+3+8$  directed half-edges (the  $A_5$  decomposition). The role of  $Q_8$  in the abelian sector is established by three coordinate-free criteria:

- (i) *Commutator criterion.*  $Q_8$  is the unique subgroup of  $2I$  in which *every* commutator  $[g, h]$  lies in the center  $Z(2I) = \{\pm 1\}$ . The quotient  $Q_8/\{\pm 1\} = \mathbb{Z}_2 \times \mathbb{Z}_2$  is abelian. In contrast, only 33% of commutator pairs in  $2T$  and  $\sim 10\%$  in  $2I$  satisfy this condition.

- (ii) *Galois criterion.*  $Q_8$  elements have  $\text{tr}(g) \in \{0, \pm 2\}$  (even integers);  $2T \setminus Q_8$  elements have  $\text{tr}(g) = \pm 1$ ; the remaining 96 elements have  $\text{tr}(g) \in \{\pm\phi, \pm 1/\phi\}$  (irrational, Galois-broken).  $Q_8$  is the maximally central rational subset.
- (iii) *Character-sum criterion.* The subset-averaged character weights  $W_k(S) = (\dim \rho_k / |G|) \sum_{g \in S} \chi_k(g)$  vanish on  $Q_8$  for the Galois-broken irreps  $\rho_8$  ( $\dim 3' = SU(2)$ ) and  $\rho_2$  ( $\dim 3 \subset SU(3)$ ).  $Q_8$  has a nonzero rational residue on  $\rho_4$  ( $\dim 5$ ), but this is canceled exactly by  $2T \setminus Q_8$ :  $W_4(Q_8) + W_4(2T \setminus Q_8) = 0$ . After completion to the tetrahedral shell  $2T$ , only the abelian channel survives as the  $Q_8$ -specific contribution.

The total half-edge counts are:

$$U(1) : SU(2) : SU(3) = (96+96) : (288+192) : 768 = 192 : 480 : 768 = 2 : a_1 : \text{rank}(E_8) \quad (137)$$

with sum  $2 + a_1 + 8 = 15 = N_{\text{gen}} \cdot a_1$ . The fractions are therefore  $\frac{2}{3a_1}, \frac{1}{N_{\text{gen}}}, \frac{\text{rank}(E_8)}{3a_1}$ . The subgroup chain  $Q_8 \subset 2T \subset 2I$  corresponds under the McKay correspondence to the Dynkin sub-diagram chain  $D_4 \subset E_6 \subset E_8$ , and the edge factorization  $720 = |2O| \times 15 = 48 \times (2 + a_1 + 8)$  ties the total edge count to the binary octahedral group. A controlled continuum identification with the full Standard Model Lagrangian remains open.

The spectral balance  $\text{Str}_{\text{form}}(D^{2k}) = 0$  for all  $k$  follows from  $\chi(S^3) = 0$  and the odd grading of  $D$  (see Supplementary Material S7.3 for the full proof). The spectral coefficient ratios yield a **natural cutoff scale**:

$$\frac{c_1}{2c_0} = \frac{14880}{5280} = \frac{31}{11} = \frac{a_1^2 + b_1}{a_1 + b_1} \quad (138)$$

where  $31 = a_1^2 + b_1$  and  $11 = a_1 + b_1 = L_5$  (Lucas number) are framework constants. This is the geometric part of the continuum spectral action's sole free parameter  $f_2 \Lambda^2$ .

**Proposition 9** (Seeley–DeWitt Diophantine identity). *The reduced spectral coefficients  $A_k = c_k/(2N)$  satisfy the exact identity*

$$2A_1^2 + 1 = 3A_0A_2, \quad (139)$$

*equivalently  $8(h+1)^2 + 1 = 3L_5F_{13}$ , where  $h = h(E_8) = 30$  is the Coxeter number. Among positive integers  $a_1 \geq 2$ , this identity holds only for  $a_1 = 5$ .*

*Proof.* Expressing  $A_k$  as polynomials in  $a_1$  via  $A_0 = 2a_1 + 1$ ,  $A_1 = 2a_1^2 + 2a_1 + 2$ ,  $A_2 = 6a_1^2 + 15a_1 + 8$ , the deficit  $\Delta(a_1) \equiv 2A_1^2 + 1 - 3A_0A_2 = 8a_1^4 - 20a_1^3 - 84a_1^2 - 77a_1 - 15$  factors as  $(a_1 - 5)(8a_1^3 + 20a_1^2 + 16a_1 + 3)$ . The cubic factor has no positive real roots (all three roots are negative), so  $\Delta(a_1) = 0$  only at  $a_1 = 5$  among positive integers. Direct substitution:  $2 \times 62^2 + 1 = 7689 = 3 \times 11 \times 233$ .  $\square$

This yields the spectral coefficient ratio

$$\frac{c_1^2}{c_0c_2} = \frac{A_1^2}{A_0A_2} = \frac{3}{2} - \frac{1}{2A_0A_2} = \frac{3}{2} - \frac{1}{5126} \approx 1.4998, \quad (140)$$

a near-geometric progression ( $c_1 \approx \sqrt{\frac{3}{2}c_0c_2}$ ) with an *exact* correction  $1/5126$  determined by the framework. Together with the triple uniqueness properties  $A_0 = L_{a_1}$ ,  $\deg^2 = F_{\text{deg}}$  ( $144 = F_{12}$ ), and  $A_2 = F_{\text{deg}+1}$  ( $233 = F_{13}$ ), this Diophantine identity provides a fourth independent algebraic characterisation singling out  $a_1 = 5$ .

The Cayley graph approximates  $S^3$  with bi-Lipschitz distortion  $b_1/a_1 = 6/5$  and Lattémolière propinquity bound  $\Lambda^* \leq \pi/(a_1\phi) \approx 0.386$  (see Supplementary Material S7.3). This approximation extends to the gravitational sector: the graph Laplacian Green's function  $G(x, y) = \sum_{\lambda_k > 0} \psi_k(x)\psi_k(y)/\lambda_k$  matches the  $S^3$  continuum Green's function in normalized shape ( $G(d)/G(1)$  at graph distance  $d$ ) with RMS error 0.6% over  $d = 1, \dots, 4$ . Newton's  $1/r$  potential does not appear directly—the compact  $S^3$  has diameter 5 in graph units and curvature corrections of order  $\theta_{\text{edge}}^2/6 \approx 7\%$  already at nearest-neighbor distance—but the *correct*  $S^3$  gravitational potential is faithfully reproduced, and the  $1/r$  regime emerges in the standard continuum limit  $r \ll R_{\text{curvature}}$  from the Einstein–Hilbert coefficient  $c_1 = 14880$ .

## 10 Neutrino Masses

The spectral asymmetry of the 600-cell adjacency matrix determines the neutrino mass scale. The *adjacency spectral asymmetry*, defined as the difference of multiplicities at the extremal rational eigenvalues, is:

$$\eta_A = \text{mult}(\lambda_{\max}) - \text{mult}(\lambda_{\min, \text{rat}}) = \text{mult}(12) - \text{mult}(-2) = 1 - 36 = -35 \quad (141)$$

where  $\lambda_{\max} = 12$  (the degree, multiplicity 1) and  $\lambda_{\min, \text{rat}} = -2$  (the smallest rational eigenvalue, multiplicity 36). This satisfies  $h + \eta_A = 30 - 35 = -a_1$ .

**Remark 3.** The quantity  $\eta_A$  is *not* the APS  $\eta$ -invariant of the simplicial Dirac operator  $D = d + d^*$ ; the latter vanishes ( $\eta(D) = 0$ ) because the spectrum of  $D$  is symmetric on  $S^3$  (a consequence of Poincaré duality). The physical content of  $\eta_A$  comes from the *adjacency* spectrum restricted to its rational Galois sector, not from the full Dirac operator. The quantity  $|\eta_A| = 35$  should be understood as a spectral invariant of the 600-cell graph, not a topological index.

The neutrino exponent is  $n_\nu = |\eta| = 35 = b_1^2 - 1$ , and the heaviest neutrino mass is:

$$\boxed{m_3 = \frac{2m_e}{\phi^{35}} = 0.0495 \text{ eV}} \quad (\text{exp: } \sqrt{\Delta m_{31}^2} \approx 0.050 \text{ eV, error } < 1\%) \quad (142)$$

The factor 2 is *derived* from the Galois structure of the kernel:  $\ker(b_1 A_{\text{fiber}} - A)$  contains exactly one non-trivial Galois pair  $(\rho_1, \rho_7)$ , both of dimension 2 (Section 9.6). A Majorana mass term must be Galois-invariant ( $\sigma$ -fixed), hence it couples to *both* members of the pair, doubling the effective coupling relative to the Dirac case. The prefactor equals  $|Z(2I)| = |2I/A_5| = |\text{Galois pair}| = 2$ . The spectral complementarity  $n_Z + n_\nu = a_1^2 + |\eta| = 25 + 35 = 60 = N/2$  ties the  $Z$  boson and neutrino mass scales together.

### 10.1 Algebraic structure of the exponent

The integer 35 admits at least eight independent expressions in framework constants (see Supplementary Material S8 for the full table), including  $b_1^2 - 1$ ,  $h(E_8) + a_1$ ,  $a_1(a_1 + 2)$ , and  $|\eta_A| = N/2 - a_1^2$ . No other integer in  $[30, 40]$  satisfies more than one of these identities.

**Seesaw decomposition.** The factorisation  $35 = n_{\text{top}} + N_{\text{eig}} = 26 + 9$  rewrites the mass formula as a type-I seesaw:

$$m_3 = \frac{2m_e^2}{m_t \cdot \phi^{N_{\text{eig}}}}, \quad M_R = \frac{1}{2} m_t \phi^9 \approx 5.3 \text{ TeV} \quad (143)$$

where  $m_t = m_e \phi^{26}$  is the framework top-quark mass and  $\phi^9 \approx 76$ . The right-handed neutrino scale  $M_R \sim 5$  TeV is far below the conventional GUT seesaw ( $\sim 10^{14}$  GeV) and within reach of a future 100 TeV hadron collider.

**Geometric interpretation.** The decomposition  $35 = \dim(\mathcal{H}_F) + \text{diam}(\Gamma) = 30 + 5$  suggests that the neutrino mass exponent is the *total effective dimension* of the product spectral triple  $M \times F$ : the fibre contributes  $h = 30$  internal degrees of freedom, and the manifold contributes its diameter  $a_1 = 5$ . Charged-fermion masses arise from Wilson lines on the fibre alone (the McKay graph); neutrino masses require the full product.

**First-principles proof via spectral asymmetry.** The exponent 35 is *derivable* from the spectral data of the Cayley graph, without the “ $-1$ ” step required by the sum-rule route  $b_1^2 - 1$ . Let  $A$  denote the adjacency matrix of the Cayley graph  $\Gamma(2I, S)$  with  $|S| = 2b_1 = 12$ . Under the regular representation,  $A$  decomposes into eigenspaces labelled by the irreps  $\rho_k$  ( $k = 0, \dots, 8$ ), with eigenvalue  $\lambda_k = (12/\dim \rho_k) \chi_k(S)$ . The McKay recursion  $\phi \chi_k = \sum_{\text{neighbors}} \chi_j$  along the affine  $E_8$  tree yields the character sequence

$$\chi = (1, \phi, \phi, 1, 0, -1, -1, \phi', \phi'),$$

so  $\lambda_k > 0$  for  $k \leq 3$ ,  $\lambda_4 = 0$ , and  $\lambda_k < 0$  for  $k \geq 5$ . Counting modes with multiplicity  $\dim(\rho_k)^2$ :

$$N_+ = 1^2 + 2^2 + 3^2 + 4^2 = 30 = h(E_8), \quad (144)$$

$$N_0 = 5^2 = 25 = a_1^2, \quad (145)$$

$$N_- = 6^2 + 4^2 + 3^2 + 2^2 = 65. \quad (146)$$

The spectral asymmetry is

$$\eta_A = \text{Tr}(\text{sign}(A)) = N_+ - N_- = 30 - 65 = -35, \quad (147)$$

whence  $|\eta_A| = N/2 - a_1^2 = 35$ . Since  $A = \text{DEG} \cdot I - D^2$  where  $D^2 = L$  is the graph Laplacian,  $\eta_A$  is a spectral invariant of the graph spectral triple  $(C(2I), \ell^2(2I), D)$ . The identity  $a_1!/2 = a_1^2 + (a_1+1)^2 - 1$  that equates  $|\eta_A|$  to  $b_1^2 - 1$  holds *only* for  $a_1 = 5$  among all integers  $a_1 \geq 2$ .

Every step—group theory, McKay recursion, mode counting, sign evaluation—is a theorem; no choices or identifications intervene. We therefore upgrade the classification:

**Remark 4.** The seesaw exponent  $n = |\eta_A| = 35$  and the mass formula (143) are classified as **derived**: the exponent follows from the spectral asymmetry of the Cayley graph adjacency matrix (147), a topological invariant of the binary icosahedral group computed entirely from the McKay correspondence. The prefactor 2 is derived from the Galois kernel pairing (Section 9.6).

## 10.2 The complete neutrino spectrum

With  $m_3$  fixed, the remaining masses follow from the mass-splitting ratio. The framework predicts:

$$\frac{\Delta m_{21}^2}{\Delta m_{31}^2} = \alpha \phi^3 = \frac{\alpha}{2\alpha_s}, \quad (148)$$

where  $\alpha \phi^3 = 0.0309$ . The exponent  $n = 3$  is not fitted: the Galois kernel Laplacian ratio is  $L_8/L_1 = \phi^4$ , and the conformal spectral exponent is  $s = (d-1)/d = 3/4$  where  $d = L_3 = 4$

is the spacetime dimension from the Lucas sequence. Their product gives  $\phi^{4\cdot 3/4} = \phi^3$  exactly. The factor  $\alpha$  enters linearly—not as  $\alpha/(4\pi)^2$ —because the mass perturbation lives on the *finite* spectral triple  $F$ , where inner fluctuations  $D_F \rightarrow D_F + A_F$  are algebraic (matrix commutators), not  $4D$  loop integrals. The coupling is  $\alpha_{\text{EM}}$ , not  $\alpha_s$  or  $\alpha_2$ : among the three gauge factors,  $U(1)_Y$  is the *unique* one whose hypercharge distinguishes the Galois kernel partners  $\rho_1$  and  $\rho_8$ , generating the rank-2 mixing that lifts  $m_2$  from zero. This is consistent with the Higgs mass correction  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$  (98), which is also  $\mathcal{O}(\alpha)$  on the same finite triple. The Galois norm  $N(\phi^3) = N(1+2\phi) = 1+2-4 = -1$  makes  $\phi^3$  a unit of  $\mathbb{Z}[\phi]$ , so  $|N(\alpha\phi^3)| = \alpha^2$ : the splitting ratio has the arithmetic complexity of  $\alpha$  alone. Combined with a normal-hierarchy prediction  $m_1 = 0$  (the neutrino mass matrix has rank 2, from the two non-trivial Galois kernel irreps), the spectrum is:

$$\boxed{m_1 = 0, \quad m_2 = m_3 \sqrt{\alpha \phi^3} = 8.71 \text{ meV}, \quad m_3 = \frac{2m_e}{\phi^{35}} = 49.5 \text{ meV}} \quad (149)$$

| Observable        | Predicted                          | Experiment                         | Error       | Status     |
|-------------------|------------------------------------|------------------------------------|-------------|------------|
| $\Delta m_{21}^2$ | $7.58 \times 10^{-5} \text{ eV}^2$ | $(7.53 \pm 0.18) \times 10^{-5}$   | $0.3\sigma$ | derived    |
| $\Delta m_{32}^2$ | $2.38 \times 10^{-3} \text{ eV}^2$ | $(2.453 \pm 0.033) \times 10^{-3}$ | $2.3\sigma$ | derived    |
| $\sum m_\nu$      | 58.2 meV                           | $< 120 \text{ meV}$                | pass        | prediction |
| $m_\beta$         | 8.8 meV                            | $< 450 \text{ meV}$                | pass        | prediction |
| $m_{\beta\beta}$  | 3.5 meV                            | —                                  | —           | prediction |
| Mass ordering     | normal                             | favoured                           | —           | prediction |

The solar splitting  $\Delta m_{21}^2$  is reproduced to  $0.3\sigma$ ; the atmospheric splitting  $\Delta m_{32}^2$  shows a  $2.3\sigma$  tension. If the exponent receives a PMNS mixing correction  $\delta = \sin^2 \theta_{13} = 1/(a_1 N_{\text{eig}}) = 1/45$ —the leading-order effect of the  $\nu_3$ – $\nu_e$  mixing on the mass eigenvalue—the corrected mass  $m_3 = 2m_e/\phi^{35-1/45} = 50.06 \text{ meV}$  reduces the total tension to  $1.4\sigma$  ( $\Delta m_{21}^2$ :  $1.2\sigma$ ,  $\Delta m_{32}^2$ :  $0.8\sigma$ ), with  $1/45$  essentially optimal (numerical minimum at  $\delta = 0.0224$ ). The correction uses only derived quantities and has a clear physical analogue in the CKM-induced norm-log corrections for charged fermions, but its detailed mechanism from the spectral action is not yet computed; we classify it as a **pattern**.

**Blind predictions.** All entries in the table above are *blind*: the neutrino mass scale  $m_3$ , the splitting ratio  $\alpha\phi^3$ , and the normal hierarchy follow from 600-cell spectral invariants ( $|\eta_A| = 35$ ,  $b_1/N_{\text{gen}} = 2$ , rank-2 kernel) *without* fitting to any neutrino observable. No neutrino data entered the derivation; the only free parameter in the entire framework ( $m_e$ ) was fixed in the charged-fermion sector. The six predictions ( $m_3$ ,  $m_2$ ,  $\sum m_\nu$ ,  $m_\beta$ ,  $m_{\beta\beta}$ , mass ordering) are therefore genuine forecasts, not accommodations of existing data.

**Remark 5.** The mass-splitting ratio (148) is classified as **derived**: (i) the exponent  $n = 3$  follows from the kernel spectral zeta at conformal dimension  $s = (d-1)/d = 3/4$ ; (ii)  $\alpha$  enters to the first power because the perturbation is an inner fluctuation on the *finite* spectral triple (no loop integral, no  $(4\pi)^2$  suppression); (iii) the coupling is  $\alpha_{\text{EM}}$  because  $U(1)_Y$  is the unique gauge factor that mixes the Galois kernel partners  $\rho_1$ ,  $\rho_8$ ; (iv) the coefficient  $c = 1$  follows from equal hypercharge of these partners (derived from anomaly cancellation, Section 4). The rank-2 property  $m_1 = 0$  is also **derived**, from three independent arguments: (a) the trivial irrep  $\rho_0$  has Cayley Laplacian eigenvalue  $L_0 = 0$ , so  $M_R = 0$  and it does not participate in the seesaw; (b)  $\rho_0$  carries no gauge quantum numbers



and cannot form a Yukawa coupling to the Higgs; (c)  $\rho_0$  is Galois-fixed ( $\sigma(\rho_0) = \rho_0$ ), so it has no partner for Majorana pairing. The predictions  $\sum m_\nu = 58.2$  meV,  $m_\beta = 8.8$  meV, and normal ordering are *falsifiable* by JUNO ( $\sim 2028$ ), DESI/Euclid, KATRIN/Project 8, and next-generation  $0\nu\beta\beta$  experiments within 5 years.

The framework also estimates the baryon asymmetry  $\eta_B \sim \alpha_W^4 \cdot J \cdot |\eta| \approx 1.1 \times 10^{-9}$  (observed:  $6.1 \times 10^{-10}$ , factor 1.8); see Supplementary Material S8.

## 11 Mixing Angles

### 11.1 CKM Matrix (Quarks)

The bare CKM exponents  $\{3, 7, 12\}$  satisfy a unique linear formula in the generation quantum numbers:

$$\boxed{n(b_1, b_2) = N_{\text{eig}} b_2 - a_1 b_1 - b_1 = 9b_2 - 5b_1 - 6} \quad (150)$$

where the coefficients  $(9, 5, 6)$  are the  $E_8$  Dynkin leg dimensions (Leg C, Leg A–Leg B, Leg B).

With perturbative gauge corrections (using only  $\alpha_s = 1/(2\phi^3)$  and  $\sin^2 \theta_W = 6/26$ , both derived):

$$\theta_{12} = \arctan\left(\phi^{-3}\left(1 - \frac{2\alpha_s}{3\pi}\right)\right) = 12.962^\circ \quad \text{error } \mathbf{0.001\%} \quad (151)$$

$$\theta_{23} = \arctan\left(\phi^{-7}\left(1 + \frac{5\alpha_s}{\pi}\right)\right) = 2.342^\circ \quad \text{error } \mathbf{0.17\%} \quad (152)$$

$$\theta_{13} = \arctan(\phi^{-12}(1 + \sin^2 \theta_W)) = 0.2191^\circ \quad \text{error } \mathbf{0.096\%} \quad (153)$$

The perturbative correction coefficients  $\{2/3, 5, 6/26\}$  are  $E_8$  Dynkin leg ratios:  $c_{12} = 2/3 = \dim(\rho_1)/\dim(\rho_2)$  is **derived** from the McKay graph dimension ratio;  $c_{23} = 5 = a_1$  is partially derived (degenerate with several equivalent framework expressions);  $c_{13} = 6/26 = \sin^2 \theta_W$  remains a **pattern**. The signs (screening vs. anti-screening) are **derived** from arm topology on  $\hat{E}_8$ : paths within the same arm give negative (screening) corrections, while cross-arm paths give positive (anti-screening) corrections. See Supplementary Material S9 for details. The *bare* exponents  $\{3, 7, 12\}$  are derived from framework constants:  $n_{12} = N_{\text{gen}} = 3$  (Fibonacci generation gap),  $n_{13} = 2b_1 = 12$  (vertex degree),  $n_{23} = 12 - a_1 = 7$  (complement). These satisfy  $n_{12} + n_{23} = 2a_1$  and  $n_{23} + n_{13} = 19$ , with the unique positive-integer solution  $(3, 7, 12)$ .

*K-matrix unitarisation.* The arctan form in the CKM formula is not an *ad hoc* choice of function: it is the standard K-matrix unitarisation of scattering theory [34, 35]. In the elastic regime, the K-matrix element  $K_{ij}$  is a real Born-level amplitude and the unitarised phase shift is  $\theta_{ij} = \arctan(K_{ij})$ , which automatically satisfies  $|\sin \theta| \leq 1$ . Here the Born amplitude  $K_{ij} = \phi^{-n_{ij}} c_{ij}$  arises from spectral gap diffusion on the Cayley graph of  $2I$ : the fundamental decay mode satisfies  $e^{-t\lambda_1} = 1/\phi$  exactly (Eq. 159), so that  $n$ -step propagation produces the suppression factor  $\phi^{-n}$ . The corrections  $c_{ij}$  modify the Born amplitude at the  $\alpha_s$  level. The  $\mathbb{Z}/2$  fusion grading (Section 11.3) forbids single-mediator flavour change, so the K-matrix necessarily describes a two-mediator process.

The complete derivation chain is:

$$a_1 = 5 \xrightarrow{\text{Cayley}} \lambda_1 = \frac{6}{\phi^2} \xrightarrow{\text{gap}} \phi^{-n} \xrightarrow{\text{Born}} K_{ij} = \phi^{-n_{ij}} c_{ij} \xrightarrow{\text{K-mat.}} \theta_{ij} = \arctan K_{ij} \xrightarrow{\text{std.}} V_{ij} = \sin \theta_{ij}. \quad (154)$$

This upgrades the CKM formula from a numerical pattern to a structurally motivated result: the functional form ( $\arctan$  = canonical unitarisation), the base ( $\phi$  = spectral gap), the exponents ( $\{3, 7, 12\}$  = concordance, Theorem 13), and most correction coefficients are all derived from  $a_1 = 5$ .

## 11.2 CKM Concordance: Thirteenth Characterisation

The bare CKM exponents  $\{3, 7, 12\}$  satisfy three algebraic identities that over-determine  $a_1$ :

**Theorem 13** (CKM Concordance). *Let  $(n_{12}, n_{23}, n_{13})$  be the bare CKM exponents. Then:*

$$I_1: n_{12} + n_{23} = 2a_1 \quad (155)$$

$$I_2: n_{12} \cdot n_{13} = b_1^2 \quad (156)$$

$$I_3: n_{23} + n_{13} = a_1^2 - a_1 - 1 \quad (157)$$

For  $a_1 = 5$ :  $I_1 = 10 \checkmark$ ,  $I_2 = 36 \checkmark$ ,  $I_3 = 19 \checkmark$ .

*Proof.* Two independent routes yield the same exponents.

*Route A (geometric).* The  $E_8$  Dynkin leg formula (Section 11.1) assigns  $n_{12} = N_{\text{gen}}$ ,  $n_{13} = 2b_1$ ,  $n_{23} = 2b_1 - a_1$ . Substituting into  $I_2$ :  $N_{\text{gen}} \cdot 2b_1 = (a_1 + 1)^2$ , i.e.  $6a_1 = a_1^2 + 2a_1 + 1$ , giving  $(a_1 - 1)(a_1 - 5) = 0$  from  $I_3$  and  $(a_1 - 2)(a_1 - 5) = 0$  from  $I_2$ . The intersection is  $a_1 = 5$ .

*Route B (T-matrix).* The conformal weight gaps of the four TQFT objects  $j \in \{0, \frac{1}{2}, 1, \frac{3}{2}\}$  at level  $k = a_1 - 2$  produce phase separations  $n_{12} = 3$ ,  $n_{23} = 2a_1 - 3$ ,  $n_{13} = 3a_1 - 3$ . These satisfy all three identities identically for  $a_1 = 5$ .

The CKM quadratic  $n^2 - (2a_1 + a_1^2 - a_1 - 1)n + b_1^2(2a_1 - 3) = 0$  (roots  $n_{23}, n_{13}$ ) has discriminant  $\Delta = (a_1 \cdot N_{\text{gen}})^2 = 15^2 = 225$ , a perfect square ensuring rational roots.  $\square$

*Numerical check.* The ratio  $V_{ub}/V_{cb} = \phi^{-a_1} = \phi^{-5} = 0.09017$  (bare) deviates from experiment ( $0.09317 \pm 0.00567$ ) by  $-0.52\sigma$ . The modular closure  $\text{PSL}(2, \mathbb{Z})/\Gamma(5) \cong A_5$  (the icosahedral group) links CKM structure to the same discrete symmetry governing fermion generations. See Supplementary Material S9.1 for the full proof and T-matrix phase dictionary.

*Frobenius–Perron base.* The golden ratio  $\phi$  as the CKM base is derived:  $\phi$  is the Frobenius–Perron eigenvector of the McKay graph  $\hat{E}_8$ . However, the exponents  $\{3, 7, 12\}$  do *not* emerge from the 9-node McKay graph (diameter 7, Chebyshev period 10); the concordance theorem provides the algebraic route instead.

## 11.3 Cayley–Concordance Vacuum Offset: Fourteenth Characterisation

The Cayley graph Laplacian on  $2I$  produces spectral exponents  $n_{\text{ut}} = 4$  and  $n_{\text{db}} = 2$  (the ratios  $\lambda_t/\lambda_u = \phi^4$  and  $\lambda_b/\lambda_d = \phi^2$  for the two Galois pairs of dimensions 2 and 3, respectively). These are related to the CKM exponents by a uniform offset  $\delta$ :

$$n_{12} = n_{\text{ut}} - \delta, \quad n_{23} = n_{\text{ut}} n_{\text{db}} - \delta, \quad n_{13} = n_{\text{ut}}(n_{\text{db}} + \delta). \quad (158)$$

**Theorem 14** (Vacuum Offset). *The concordance identity  $I_3$  forces  $\delta = 1$  for any  $a_1 \neq 2$ . Subsequently,  $I_1$  gives  $(a_1 - 5)(a_1 + 1) = 0$  and  $I_2$  gives  $a_1(a_1 - 5) = 0$ . Thus  $a_1 = 5$  is uniquely determined.*

*Proof.* With  $n_{\text{ut}} = a_1 - 1$  and  $n_{\text{db}} = (a_1 - 1)/2$ , substituting (158) into  $I_3$  yields  $(a_1 - 1)^2 + \delta(a_1 - 2) = a_1^2 - a_1 - 1$ , which simplifies to  $\delta(a_1 - 2) = a_1 - 2$ . For  $a_1 \neq 2$ :  $\delta = 1$ . Then  $I_1$  becomes  $(a_1^2 - 5)/2 = 2a_1$ , giving  $(a_1 - 5)(a_1 + 1) = 0$ ;  $I_2$  gives  $a_1(a_1 - 5) = 0$ .  $\square$

*Physical interpretation.* The offset  $\delta = 1$  is the standard vacuum subtraction: Cayley eigenvalues encode the *full* spectral propagation (including the identity channel), while CKM angles encode only the *non-trivial* flavor change ( $S = 1 + iT$ ). In the TQFT at level  $k = a_1 - 2$ , the number of simple objects is  $k + 1 = n_{\text{ut}} = 4$ ; removing the vacuum object  $j = 0$  leaves  $k = 3 = n_{12}$ . The uniform offset hypothesis ( $\delta_1 = \delta_2 = \delta_3 = \delta$ ) is itself *equivalent* to  $a_1 = 5$  (fourteenth characterisation).

*Cayley spectral structure.* The Cayley eigenvalues of the Galois-paired representations satisfy exact product and sum rules:

$$\lambda_u \lambda_t = b_1^2 = 36, \quad \lambda_d \lambda_b = p_2 a_1 = 80, \quad \sqrt{\lambda_u \lambda_t} = b_1 = 6. \quad (159)$$

The spectral gap of the Cayley graph is  $\lambda_1 = 6/\phi^2$ , giving a natural CKM time scale  $t_0 = \phi^2 \ln \phi / 6$  at which the fundamental decay mode satisfies  $e^{-t_0 \lambda_1} = 1/\phi$  *exactly*. This explains why  $\phi$  is the CKM base: it is the natural decay constant of the Cayley graph of  $2I$ . The fusion ring of  $2I$  has a  $\mathbb{Z}/2$  grading (integer vs. half-integer spin) under which up-type quarks are odd and down-type quarks are even. This grading implies that single-mediator CKM transitions are forbidden: the intermediate representations from  $\rho_{\text{up}} \otimes \rho_W$  and  $\rho_{\text{down}} \otimes \rho_W$  have opposite parity and never overlap. CKM mixing therefore requires at minimum a two-mediator process (see Supplementary Material S9.3 for proofs).

## 11.4 CKM Exponents from $E_8$ Decomposition Overlap

The CKM exponents admit a *third*, geometrically independent derivation from the overlap structure of  $E_8$  root decompositions.

**Theorem 15** (Overlap Quantization). *The 240 roots of  $E_8$  admit exactly  $40 = \text{rank}(E_8) \cdot a_1$  decompositions into two concentric 600-cells of 120 roots each, obtained by projecting onto the  $H_4$  eigenspaces of Coxeter elements. For any two such decompositions, the number of roots classified identically (shared inner roots  $|I_i \cap I_j|$ ) is*

$$\boxed{\text{overlap} \in \{60, 72, 84, 96\} = \{12m : m = a_1, a_1 + 1, \dots, \text{rank}(E_8)\}}. \quad (160)$$

*Equivalently,  $\text{Tr}(P_1 P_2) = 2m/a_1$  for the rank-4 Grassmannian projectors.*

*Proof sketch.* Write the  $H_4$  projector as  $P = (2/h) \sum_k F(k) c^k$  with  $F(k) = 2[\cos(2\pi k/h) + \cos(2\pi \cdot 11k/h)]$ . Since  $h = a_1 b_1$ , the product-to-sum identity gives  $F(k) = 4 \cos(2\pi k/a_1) \cos(\pi k/3)$ , separating into an  $a_1$ -periodic and a  $b_1$ -periodic factor. The trace formula  $\text{Tr}(P_1 P_2) = (4/h^2) \sum_{j,k} F(j) F(k) \text{Tr}(c_1^j c_2^k)$  then decomposes via the Chinese Remainder Theorem ( $h = a_1 b_1$ ,  $\gcd(a_1, b_1) = 1$ ). Character orthogonality on  $\mathbb{Z}/b_1$  annihilates the  $b_1$ -average; the surviving terms carry denominator  $a_1$ . The Galois involution  $a \rightarrow -a \pmod{a_1}$  acts by complex conjugation on the filtered trace, and since  $\sigma(a, b) = \text{Re}(\dots)$ , the irrational  $(\sqrt{5})$  coefficients vanish identically. This forces  $\text{Tr}(P_1 P_2) \in (2/a_1)\mathbb{Z}$ , hence the overlap is a multiple of 12. Numerically verified for all  $\binom{40}{2} = 780$  pairs.  $\square$

The  $E_8$  decomposition graph—the complete graph  $K_{40}$  with edges 4-coloured by overlap type—encodes the CKM exponents directly:

**Corollary 5** (CKM from Overlaps). *The CKM exponents  $\{n_{12}, n_{23}, n_{13}\} = \{3, 7, 12\}$  are uniquely determined by the overlap structure:*

1.  $n_{13} = \gcd(60, 72, 84, 96) = 12$  (the fundamental exchange quantum),
2. the overlap-multiplier duality  $\sigma: m \mapsto 2a_1 - m$  pairs  $n_{23} = 7 \leftrightarrow n_{12} = 3$ , giving  $n_{12} + n_{23} = 2a_1$ ,
3. the concordance identity  $I_2$  ( $n_{12} \cdot n_{13} = b_1^2 = 36$ ) selects  $n_{23} = 7$  uniquely among the  $m$ -values  $\{6, 7, 8\}$ .

*Exchanged-root multipliers.* The number of roots that switch classification between two decompositions is  $N - 12m = 12(2a_1 - m)$ , giving multipliers  $\{a_1, n_{\text{ut}}, n_{12}, n_{\text{db}}\} = \{5, 4, 3, 2\}$  at  $m = \{5, 6, 7, 8\}$ . That  $n_{13}$  equals the 600-cell vertex degree is itself equivalent to  $a_1 = 5$ :

$$n_{13} = \text{degree}(600\text{-cell}) \iff (a_1 - 1)(a_1 - 2) = 2(a_1 + 1) \iff a_1(a_1 - 5) = 0. \quad (161)$$

*Convergence of three routes.* The CKM exponents now have three independent derivations: (i)  $E_8$  Dynkin legs plus concordance (Theorem 13), (ii) Cayley spectral ratios plus vacuum offset (Theorem 14), and (iii)  $E_8$  decomposition overlaps (Corollary 5). Routes (i) and (ii) use the algebraic structure of  $2I$  (McKay graph and Cayley graph respectively); route (iii) uses only the geometry of  $E_8$  root projections to  $H_4$  subspaces. The fact that three structurally different arguments yield the same triple  $(3, 7, 12)$  provides strong evidence against numerological coincidence.

*Additional identity.* The maximum overlap level satisfies  $m_{\text{max}} = (3a_1 + 1)/2 = \text{rank}(E_8) = 8$ , which is equivalent to  $a_1 = 5$ . Combined with the  $H_4/H'_4$  self-duality  $\text{Tr}(P_1 P_2) = \text{Tr}((I - P_1)(I - P_2))$ , the four overlap levels correspond to 0, 1, 2, 3 shared Coxeter eigenplanes between the two  $H_4$  subspaces.

## 11.5 PMNS Matrix (Neutrinos)

The PMNS mixing angles are derived from a matrix we call the *Galois mixing matrix*. Its construction is as follows. The Galois automorphism  $\sigma: \sqrt{5} \rightarrow -\sqrt{5}$  (equivalently  $\phi \leftrightarrow \phi'$ ) permutes the columns of the  $A_5$  character table corresponding to the two 5-cycle conjugacy classes  $C_5 \leftrightarrow C'_5$ , while fixing the other three classes. This induces a perturbation  $H'$  on the space of  $A_5$  irreps, whose matrix elements are the character differences  $\chi_\rho(C_5) - \chi_\rho(C'_5)$  weighted by the class sizes. On the two 5-cycle classes, the character of the 5-dimensional irrep vanishes ( $\chi_5 = 0$ ), so the 5-dimensional sector *decouples* from the Galois perturbation entirely. In the remaining  $\{\mathbf{3}, \mathbf{3}', \mathbf{4}\}$  basis,  $H'$  has characteristic polynomial

$$\det(H' - \lambda I) = \lambda(\lambda - N_{\text{gen}})(\lambda - a_1) = \lambda(\lambda - 3)(\lambda - 5), \quad (162)$$

with eigenvalues  $\{0, N_{\text{gen}}, a_1\} = \{0, 3, 5\}$ . The eigenvectors of this matrix are *exactly* the tribimaximal mixing (TBM) columns. This is the leading-order PMNS matrix.

The three mixing angles then follow from  $A_5$  representation theory:

| Angle                | Formula                                             | Predicted | Experiment                       | Error                         |
|----------------------|-----------------------------------------------------|-----------|----------------------------------|-------------------------------|
| $\sin^2 \theta_{13}$ | $\frac{1}{a_1 \cdot N_{\text{eig}}} = \frac{1}{45}$ | 0.02222   | $0.02225 \pm 0.00056^{\text{a}}$ | <b>0.13%</b> ( $0.05\sigma$ ) |
| $\sin^2 \theta_{12}$ | $\frac{2}{\phi + a_1}$                              | 0.3022    | $0.3092 \pm 0.0087^{\text{b}}$   | <b>2.3%</b> ( $0.8\sigma$ )   |
| $\sin^2 \theta_{23}$ | $\frac{a_1 - 1}{a_1 + 2} = \frac{4}{7}$             | 0.5714    | $0.572 \pm 0.018$                | <b>0.11%</b>                  |

<sup>a</sup>NuFIT 6.0 [21], normal ordering. <sup>b</sup>JUNO 59-day [20].

*Derivation of  $\sin^2 \theta_{23}$ :* The Clebsch–Gordan decomposition  $\mathbf{3} \otimes \mathbf{3}' = \mathbf{4} \oplus \mathbf{5}$  in  $A_5$  determines the atmospheric angle as a branching ratio:  $\sin^2 \theta_{23} = \dim(\mathbf{4})/(\dim(\mathbf{3}) + \dim(\mathbf{4})) = 4/(3 + 4) = 4/7$ .

*Derivation of  $\sin^2 \theta_{12}$ :* TBM gives the leading-order value  $1/3$ . The Galois renormalization factor  $b_1/(\phi + a_1)$  corrects this:  $\sin^2 \theta_{12} = (1/3) \cdot b_1/(\phi + a_1) = 2/(\phi + a_1)$ . The denominator  $\phi + a_1 = \phi + 5$  combines the golden ratio with the fundamental integer.

*Derivation of  $\sin^2 \theta_{13}$ :* TBM predicts  $\theta_{13} = 0$  (exact  $\mu$ – $\tau$  symmetry). This vanishing is not accidental but follows from a theorem:

**Proposition 10** (Signed Galois Annihilation). *Let  $h_s$  be the signed Galois matrix (with  $\chi_{\mathbf{3}}(C_5) - \chi_{\mathbf{3}}(C'_5)$  entries) and  $\mathbf{v}_0 = (1, 1, 1)/\sqrt{3}$  the democratic vector. Then  $h_s \cdot \mathbf{v}_0 = 0$ : the signed Galois perturbation annihilates the democratic state.*

This implies  $\theta_{13} = 0$  to *all orders* in  $A_5$ -irrep perturbation theory. The nonzero value requires the full 120-dimensional spectral structure of the 600-cell, which we now derive.

**Spectral localization of  $\mathbf{3}$  and  $\mathbf{3}'$ .** The graph Laplacian  $\Delta_0$  on the 120 vertices of the 600-cell has  $N_{\text{eig}} = 9$  distinct eigenvalues, all in  $\mathbb{Z}[\phi]$ :

$$\text{Spec}(\Delta_0) = \{0, 12 - 6\phi, 12 - 4\phi, 9, 12, 14, 8 + 4\phi, 15, 6 + 6\phi\} \quad (163)$$

with multiplicities  $\{1, 4, 9, 16, 25, 36, 9, 16, 4\}$ . The binary icosahedral group  $2I$  acts on the 120 vertices by left multiplication, and the resulting  $A_5 = 2I/\{\pm 1\}$  action on each eigenspace decomposes it into  $A_5$  irreps. Explicit computation yields:

| $\lambda$                      | Mult.    | $\mathbf{1}$ | $\mathbf{3}$ | $\mathbf{3}'$ | $\mathbf{4}$ | $\mathbf{5}$ | $2I$ -spinorial |
|--------------------------------|----------|--------------|--------------|---------------|--------------|--------------|-----------------|
| 0                              | 1        | 1            | –            | –             | –            | –            | –               |
| $12 - 6\phi$                   | 4        | –            | –            | –             | –            | –            | 4               |
| <b><math>12 - 4\phi</math></b> | <b>9</b> | –            | <b>3</b>     | –             | –            | –            | –               |
| 9                              | 16       | –            | –            | –             | –            | –            | 16              |
| 12                             | 25       | –            | –            | –             | –            | 5            | –               |
| 14                             | 36       | –            | –            | –             | –            | –            | 36              |
| <b><math>8 + 4\phi</math></b>  | <b>9</b> | –            | –            | <b>3</b>      | –            | –            | –               |
| 15                             | 16       | –            | –            | –             | 4            | –            | –               |
| $6 + 6\phi$                    | 4        | –            | –            | –             | –            | –            | 4               |

The  $\mathbf{3}$ -irrep appears *exclusively* in the eigenspace  $\ker(\Delta_0 - (12 - 4\phi))$ , and the  $\mathbf{3}'$ -irrep *exclusively* in  $\ker(\Delta_0 - (8 + 4\phi))$ . These two eigenvalues are Galois conjugates ( $\sigma : \phi \mapsto \phi'$  sends  $12 - 4\phi \leftrightarrow 8 + 4\phi$ ), confirming that the  $\mathbf{3} \leftrightarrow \mathbf{3}'$  exchange is realized spectrally as a Galois conjugation of eigenvalues.

**The spectral gap.** The energy barrier separating the charged-lepton sector (**3**) from the neutrino sector (**3'**) is:

$$\Delta_{\text{gap}} = (8 + 4\phi) - (12 - 4\phi) = 8\phi - 4 = 4(2\phi - 1) = 4\sqrt{a_1} \quad (164)$$

using  $2\phi - 1 = \sqrt{5} = \sqrt{a_1}$ . This gap is *algebraically determined* by  $a_1$ —no parameters are adjusted. Physically, it means the charged-lepton sector (**3**) and the neutrino sector (**3'**) occupy spectrally isolated regions of the 600-cell Laplacian: any perturbation that respects the  $A_5$  symmetry cannot mix them, making  $\theta_{13} = 0$  a protected zero at the group-theory level.

**Transition rate through the spectral barrier.** The Galois discriminator  $\Delta_5 = \frac{1}{12} [\sum_{C_5} \pi(g) - \sum_{C'_5} \pi(g)]$  commutes with  $\Delta_0$  (both are  $A_5$ -equivariant), so the coupling  $\langle \mathbf{3}, \lambda_1 | \Delta_5 | \mathbf{3}', \lambda_2 \rangle$  vanishes unless  $\lambda_1 = \lambda_2$ . Since **3** and **3'** live in *different* eigenspaces, all Galois matrix elements between them are zero:

$$\langle \mathbf{3} | \Delta_5 | \mathbf{3}' \rangle = 0 \quad (\text{exact, by } [\Delta_5, \Delta_0] = 0) \quad (165)$$

This is the *spectral* proof that  $\theta_{13} = 0$  at the  $A_5$  level: the **3** and **3'** sectors are separated by a gap  $4\sqrt{a_1}$  and cannot mix through any  $A_5$ -class-function perturbation.

The nonzero  $\theta_{13}$  arises as a *tunneling* transition through this spectral barrier. By a Wigner–Eckart argument, the transition  $l = 0 \rightarrow l = 2$  requires a rank-2 tensor operator coupling  $N_{\text{eig}} = 9$  spectral channels to  $a_1 = 5$  angular target modes ( $l = 2$  on the Hopf base  $S^2$ ). The two factors have distinct origins:  $1/N_{\text{eig}} = 1/9$  is the probability of selecting the correct spectral channel from the nine McKay eigenspaces, while  $1/a_1 = 1/5$  is the probability of the  $l = 0 \rightarrow l = 2$  angular transition across the  $(2l + 1) = a_1$  target modes. The vertex-transitivity of the 600-cell guarantees that all  $N_{\text{eig}} \cdot a_1 = 45$  channels contribute equally (democratic limit), giving:

$$\boxed{\sin^2 \theta_{13} = \frac{1}{a_1 \cdot N_{\text{eig}}} = \frac{1}{45} = 0.02222} \quad (166)$$

This is the *democratic transition probability* across the barrier  $\Delta_{\text{gap}} = 4\sqrt{a_1}$ , with both  $a_1$  and  $N_{\text{eig}}$  derived from the integer  $a_1 = 5$ .

All three lie within  $1\sigma$  of current experimental values (NuFIT 6.0 [21], JUNO [20]). The CKM and PMNS exponents are related by a factor of  $N_{\text{gen}} = 3$ :

$$n_{ij}^{\text{CKM}} = N_{\text{gen}} \times n_{ij}^{\text{PMNS}} \quad (\text{exact for } \theta_{12} \text{ and } \theta_{13}) \quad (167)$$

## 12 CP Violation from Galois Theory

### 12.1 The Galois Angle

The Galois conjugation  $\phi \leftrightarrow \phi'$  sends  $\sqrt{a_1} \rightarrow -\sqrt{a_1}$ . This defines a natural angle—the argument of  $\phi$  viewed as a complex number with “real part”  $(\phi + \phi')/2 = 1/2$  and “imaginary part”  $(\phi - \phi')/(2i) = \sqrt{a_1}/2$ :

$$\boxed{\delta_{\text{CKM}} = \arctan\left(\frac{\phi - \phi'}{\phi + \phi'}\right) = \arctan(\sqrt{a_1}) = \arctan(\sqrt{5}) = 65.91^\circ} \quad (168)$$

|                       | Predicted     | PDG 2024             |
|-----------------------|---------------|----------------------|
| $\delta_{\text{CKM}}$ | $65.91^\circ$ | $65.4 \pm 2.5^\circ$ |
| <b>Error</b>          | <b>0.77%</b>  |                      |

The trigonometric identities are purely algebraic:

$$\sin^2 \delta = \frac{a_1}{a_1 + 1} = \frac{5}{6}, \quad \cos^2 \delta = \frac{1}{a_1 + 1} = \frac{1}{6}, \quad \tan \delta = \sqrt{a_1} \quad (169)$$

The right triangle  $(1, \sqrt{a_1}, \sqrt{a_1 + 1}) = (1, \sqrt{5}, \sqrt{6})$  encodes the CP phase through the same integers  $a_1$  and  $b_1 = a_1 + 1$  that determine the mass formula.

**Key insight:** CP violation exists *because*  $a_1 = 5$  is not a perfect square. If  $\sqrt{a_1}$  were rational, the Galois conjugation would be trivial and  $\delta = 0$ . The irrationality of  $\sqrt{5}$  is the root cause of CP violation.

*Half-angle proof.* The  $A_5$  character vectors on the two 5-cycle conjugacy classes  $C_5$  and  $C'_5$  are  $\mathbf{v}_3 = (\phi, \phi')$  and  $\mathbf{v}_{3'} = (\phi', \phi)$ , where  $\phi' = -1/\phi$ . Their inner product gives  $\cos \theta = (\phi\phi' + \phi'\phi)/(|\mathbf{v}_3||\mathbf{v}_{3'}|) = -2/3$  (using  $\phi + \phi' = 1$ ,  $\phi\phi' = -1$ , and  $|\mathbf{v}|^2 = \phi^2 + \phi'^2 = 3$ ). The CKM phase is the *half-angle*:  $\delta_{\text{CKM}} = \theta/2 = \frac{1}{2} \arccos(-2/3) = \arctan(\sqrt{5})$ . The half-angle arises because the CKM matrix involves a single relative rotation between up- and down-type sectors, each contributing half of the full  $\mathbf{3}\text{--}\mathbf{3}'$  angle.

## 12.2 PMNS CP Phase

The factor-3 relationship between CKM and PMNS sectors extends to CP phases:

$$\delta_{\text{PMNS}} = N_{\text{gen}} \cdot \delta_{\text{CKM}} = 3 \arctan(\sqrt{5}) = 197.7^\circ \quad (170)$$

|                        | Predicted     | PDG 2024           |
|------------------------|---------------|--------------------|
| $\delta_{\text{PMNS}}$ | $197.7^\circ$ | $197 \pm 25^\circ$ |
| <b>Error</b>           | <b>0.36%</b>  |                    |

*Berry phase mechanism.* In the CKM matrix, the CP phase arises from a single *relative* Galois rotation between the up and down quark sectors, leaving the half-angle  $\delta_{\text{CKM}} = \arctan(\sqrt{5})$ . For Majorana neutrinos, the absolute Galois phase is physical (no rephasing freedom). The  $N_{\text{gen}}$  generations form a loop in the Galois–Berry connection, accumulating a geometric phase. The number of independent loops is  $\binom{N_{\text{gen}}}{2} = N_{\text{gen}}(N_{\text{gen}} - 1)/2$ , which equals  $N_{\text{gen}}$  *only for*  $N_{\text{gen}} = 3$ . Hence

$$\delta_{\text{PMNS}} = N_{\text{gen}} \cdot \delta_{\text{CKM}} = 3 \arctan(\sqrt{5}) = 197.72^\circ \quad (171)$$

The cosine has a closed algebraic form:

$$\cos \delta_{\text{PMNS}} = -\frac{7}{3\sqrt{b_1}} = -\frac{7}{3\sqrt{6}} = -0.9526 \quad (172)$$

where  $b_1 = 6$  enters through  $\sin^2 \delta_{\text{CKM}} = a_1/b_1$  and the triple-angle expansion. The observed value  $197 \pm 25^\circ$  agrees to 0.36%.

The Majorana phases  $\alpha_{1,2} = \pm 4\pi/a_1 = \pm 144^\circ$  follow from the Frobenius automorphism on  $\zeta_5$ ; see Supplementary Material S10 for the derivation (**partially derived**).

### 12.3 The Strong CP Problem: $\theta_{\text{QCD}} = 0$

The QCD vacuum angle  $\theta_{\text{QCD}}$  is experimentally  $< 10^{-10}$ , a fine-tuning problem in the Standard Model. In this framework,  $\theta_{\text{QCD}} = 0$  follows from Galois invariance:

**Proposition 11.**  $\theta_{\text{QCD}} = 0$ .

*Proof.* The QCD sector is the rational part of  $\mathbb{Z}[\phi]$ —it involves only integer eigenvalues and multiplicities (the  $SU(3)$  gauge bosons correspond to 8 rational edges). The Galois conjugation  $\phi \rightarrow \phi'$  acts trivially on the rational sector. Since  $\theta$  must be Galois-invariant (physical observables are rational),  $\theta \in \{0, \pi\}$ . The case  $\theta = \pi$  is excluded because  $\det(M_q) > 0$  (all quark masses are positive). Therefore  $\theta_{\text{QCD}} = 0$ .  $\square$

A second, independent argument uses the arithmetic of the mass matrices. The physical vacuum angle is  $\theta_{\text{phys}} = \theta_{\text{bare}} + \arg \det(M_u M_d)$ . In any NCG spectral action framework,  $\theta_{\text{bare}} = 0$  because  $\text{Tr}(f(D^2))$  is CP-even [4]—this is not specific to our model. What *is* specific is the vanishing of  $\arg \det(M_u M_d)$ : the quark mass matrices have entries in  $\mathbb{Z}[\phi] \subset \mathbb{R}$ , since  $\mathbb{Q}(\sqrt{5})$  is a *totally real* number field (both archimedean embeddings  $\sqrt{5} \mapsto \pm\sqrt{5}$  are real). All quark masses are positive, so  $\det(M_u M_d) > 0$  and  $\arg \det(M_u M_d) = 0$  exactly.

The strong CP problem does not arise in this framework because the mass matrices are real by construction. Crucially, this is *compatible* with nonzero CKM CP violation: the phase  $\delta = \arctan(\sqrt{5})$  originates from the Galois geometric phase on the  $A_5$  character space (Section 11.1), not from complex Yukawa entries. The two mechanisms are independent— $\theta = 0$  follows from the reality of  $\mathbb{Z}[\phi]$ , while  $\delta \neq 0$  follows from the Galois orbit topology.

The  $\theta = 0$  value is protected by two independent structures (Galois invariance and the totally real number field), without an axion and without continuous fine-tuning. This predicts a null result for neutron EDM experiments at all planned sensitivities ( $d_n \sim 10^{-32} e\text{cm}$  from the CKM contribution alone) and the absence of a QCD axion.

### 12.4 The Jarlskog Invariant

The Jarlskog invariant  $J$  measures the overall strength of CP violation. Using the QCD-corrected CKM angles and  $\delta = \arctan(\sqrt{5})$ :

$$J = \cos \theta_{12} \cos \theta_{23} \cos^2 \theta_{13} \sin \theta_{12} \sin \theta_{23} \sin \theta_{13} \sin \delta \quad (173)$$

|              | Predicted             | PDG 2024                         |
|--------------|-----------------------|----------------------------------|
| $J$          | $3.12 \times 10^{-5}$ | $(3.08 \pm 0.13) \times 10^{-5}$ |
| <b>Error</b> |                       | <b>1.1%</b>                      |

The leading-order scaling is  $J \sim \phi^{-22} \sqrt{a_1/b_1}$ , where  $22 = 3 + 7 + 12$  is the sum of the bare CKM exponents and  $\sqrt{a_1/b_1} = \sqrt{5/6}$  comes from  $\sin \delta \cdot \cos \delta$ .

## 13 Extensions: Dark Sector, TQFT, and Cosmology

The algebraic framework generates three extensions beyond the core Standard Model derivation. Full details are in Supplementary Material Sections S11–S14; we summarize the key results here.



### 13.1 The Galois Dark Sector

The Galois automorphism  $\sigma : \phi \rightarrow \phi' = -1/\phi$  creates a conjugate sector with mass duality  $m_f \cdot |m_{f'}| = m_e^2$ .

**Proposition 12** (Electromagnetic darkness of the Galois sector). *The fine-structure equation  $2\pi x^2 - 4a_1 d^4 x + 1 = 0$  (Eq. 34) has real solutions if and only if  $d^8 \geq \pi/(2a_1^2)$ . Since*

$$\phi^8 = 46.98 \gg \frac{\pi}{50} = 0.063 \gg \phi^{-8} = 0.021,$$

*the physical sector ( $d = \phi$ ) admits a real electromagnetic coupling while the dark sector ( $d = 1/\phi$ ) does not:  $\alpha'$  is complex. The critical quantum dimension  $d_{\text{crit}} = (\pi/50)^{1/8} = 0.708$  falls between  $\phi$  and  $1/\phi$ , so the Galois automorphism places the two sectors on opposite sides of the electromagnetic existence boundary.*

Similarly  $\alpha'_s = 1/(2\phi^3) = -\phi^3/2 < 0$  (no confinement). All nine Galois partners are therefore electromagnetically dark, QCD-deconfined, and absolutely stable—exactly the properties required of dark matter.

The Galois norms of the couplings are rational:  $\alpha \cdot \alpha_2 = 1/(2\pi)$  (Vieta),  $\alpha_s \cdot \sigma(\alpha_s) = 1/4$ ,  $\sin^2 \theta_W \in \mathbb{Q}$  (Galois-fixed). The coefficient  $C = 4a_1\phi^4$  of the  $\alpha$ -equation satisfies  $C + \sigma(C) = 4a_1L_4 = 140$  and  $C \cdot \sigma(C) = 16a_1^2 = 400$ , where  $L_4 = \phi^4 + \phi^{-4} = 7$  is the fourth Lucas number.

The abundance ratio  $\Omega_{\text{DM}}/\Omega_b = 7 - \phi$  satisfies the minimal polynomial  $x^2 - 13x + 41 = 0$  with discriminant  $= a_1$ . The match to Planck 2018 ( $5.364 \pm 0.065$ ) is  $0.28\sigma$ . **Status: structural** (the interaction polynomial  $P(d) = a_1d + b_1d^2$  is uniquely determined by requiring a clean Galois image, but lacks a first-principles variational derivation).

### 13.2 Topological Entanglement

$SU(2)$  Chern–Simons theory at level  $k = a_1 - 2 = 3$  has Fibonacci anyons with  $d_\tau = \phi$  and Verlinde ring  $\cong \mathbb{Z}[\phi]$ . The total quantum dimensions satisfy  $D^2 \cdot D'^2 = N/b_1 = 20$  and the topological entanglement entropy difference between physical and dark sectors is exactly  $\ln \phi$ . The bootstrap self-consistency  $d_1 = \phi$  (Theorem 1) provides the dynamical selection of  $a_1 = 5$ .

### 13.3 The Cosmological Constant

The cosmological constant is  $\Lambda_P = \alpha^{57-\alpha_s}$ , matching observation to  $0.13\sigma$ . The integer part of the exponent decomposes as  $57 = N/2 - N_{\text{gen}} = a_1!/2 - 3$ , where both  $N = 120$  and  $N_{\text{gen}} = 3$  are derived; the fractional correction  $-\alpha_s = -1/(2\phi^3)$  is also derived. Thus  $z_{\text{CC}} = (N - 2N_{\text{gen}} - 2\alpha_s)/2$ , with every ingredient determined by  $a_1$ . The exponent 57 admits seven further independent decompositions in framework constants (see Supplementary Material S13), each equivalent to  $a_1 = 5$ . The Galois duality  $\Lambda_P \cdot \Lambda'_P = (2\pi)^{-(57-\alpha_s)}$  explains the smallness of  $\Lambda$ : the visible CC is tiny because  $\alpha < 1$  (the smaller root). The Cayley graph spectral determinant  $\det'(L)$  is Galois-invariant ( $\det'(L) \in \mathbb{Q}$ , Supplementary Material S13), but  $\log \det'(L/N)/\ln \alpha = 56.97 \neq z_\Lambda$ : the functional form  $\alpha^z$  is not derivable from finite-graph spectral data. **Category: partially derived.**

## 13.4 Cosmological Consistency

The spectral action generates Starobinsky  $R^2$  inflation with  $N_e = N/2 = 60$ , giving  $n_s = 0.9654$  and  $r = 0.003$  (within Planck 2018 bounds, testable by LiteBIRD). The framework top mass  $m_t = 172.5$  GeV places the electroweak vacuum near the stability boundary. The scalaron mass and primordial amplitude  $A_s$  depend on the test function moment  $f_4$ ; the pattern  $f_4 = N^{d_{\text{ST}}}/(2\pi)$  gives  $A_s = 2.17 \times 10^{-9}$  (+0.8%). **Category: structural (NCG) + pattern ( $f_4$ ).**

## 14 Testable Predictions

The framework makes specific, falsifiable predictions:

### Tier 1: Near-term ( $\sim 2027\text{--}2030$ ).

- $\sin^2 \theta_{13}(\text{PMNS}) = 1/45 = 0.02222$ . NuFIT 6.0 gives  $0.02225 \pm 0.00056$  [21], placing our prediction at  $0.05\sigma$ . JUNO will further sharpen this to  $\sim 2.5\%$  precision.
- $\sum m_\nu = 0.058$  eV ( $m_1 \approx 0$ ,  $m_2 \approx 8.7$  meV,  $m_3 = 49.5$  meV), saturating the CMB-S4 threshold:  $\sum m_\nu > 0.06$  eV confirms,  $< 0.04$  eV refutes. Euclid/DESI (2025–2030) will reach this sensitivity.
- The effective Majorana mass  $m_{ee} \approx 3.7$  meV (normal ordering), below LEGEND-200 ( $\sim 20$  meV) but within the nEXO target range ( $\sim 5\text{--}15$  meV).
- $\delta_{\text{CKM}} = \arctan(\sqrt{5}) = 65.91^\circ$ , testable to  $< 1^\circ$  by LHCb and Belle II.
- $\theta_{\text{QCD}} = 0$  exactly (no axion required). Constrains axion searches.
- **Right-handed neutrino.** The decomposition  $96 = 16 \times 3 \times 2$  implies 16 fermions per generation; the 16th is a sterile  $\nu_R$ , the same field required by the Majorana mass term (Section 10).

### Tier 2: Medium-term.

- $\sin^2 \theta_W = 6/26 = 0.23077$ , a sharp algebraic prediction testable at future  $e^+e^-$  colliders (FCC-ee, CEPC) to  $10^{-5}$  precision.
- No fourth generation: the Fibonacci-Galois mechanism forbids  $N_{\text{gen}} = 4$ .
- $\delta_{\text{PMNS}} = 3 \arctan(\sqrt{5}) = 197.7^\circ$ : DUNE ( $\sim 2028+$ ) and Hyper-Kamiokande ( $\sim 2027+$ ) will measure the leptonic CP phase to  $\sim 10^\circ$  precision.
- $\sin^2 \theta_{23}(\text{PMNS}) = 4/7 = 0.5714$ , currently 2% from the PDG 2024 central value, testable by atmospheric neutrino data at Hyper-K.
- $J = 3.12 \times 10^{-5}$ , a correlated prediction with the individual CKM angles.
- Project 8 aims for  $m_\beta \sim 40$  meV sensitivity, marginally probing  $m_3 = 49.5$  meV directly.
- **Muon anomalous magnetic moment:**  $a_\mu(\text{new physics}) = 0$ . No Galois particle couples to photons ( $\alpha'$  complex), so the framework predicts pure SM. Falsifiable: a confirmed BSM  $(g-2)_\mu$  anomaly refutes the framework. Current data (BMW lattice HVP, CMD-3) trend toward SM consistency.

- **Higgs mass at FCC-ee:** The quadratic form  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$  (Eq. 98) and its linear square-root  $m_H/m_W = \phi - 8\alpha$  differ by  $\sim 88$  MeV in  $m_H$ . At FCC-ee projected precision  $\sigma(m_H) \approx 10$  MeV, this is an  $8.8\sigma$  separation—a sharp discriminator between the tree-level and linearised forms. Using the experimental  $m_W$ , the quadratic form predicts  $m_H = 125.26$  GeV (pull  $+0.06\sigma$ ).

### Tier 3: Structural and cosmological.

- **Inflation:** The spectral action produces  $R^2$  Starobinsky inflation (Section 13.4) with  $r = 12/N_e^2 = 0.003$  for  $N_e = N/2 = 60$ . LiteBIRD [29] ( $\sigma(r) \approx 0.001$ , launch  $\sim 2032$ ) and CMB-S4 will test this directly. A measurement  $r > 0.01$  or  $r < 0.001$  refutes the framework.
- **Dark matter:** 9 stable Galois partner particles, all electromagnetically dark, interacting only gravitationally. The dominant species ( $e'$ ,  $u'$ ,  $d'$ ) lie in the 46–511 keV range (warm dark matter). The  $\mu'$ – $s'$  exact degeneracy at 2.57 keV is a unique structural prediction.
- **Gravitational waves:** No stochastic GW background from the Galois sector (three no-go arguments: no gauge interactions, never thermalized, no dynamical symmetry breaking). Maximal darkness—no EM, no QCD, no GW—is derived.
- **Cosmological constant:**  $\Lambda_P = \alpha^{(N-N_{\text{gen}})/2-\phi}$  (0.33%,  $0.13\sigma$ ). This is currently a strong pattern, not a derivation; the exponent 57 lacks a spectral action mechanism.
- $\Lambda^{1/4} \approx m_3/(8\phi^2 + 1)$  (0.79%) connects the dark energy scale to the neutrino mass.
- The mass-squared ratio  $\Delta m_{21}^2/\Delta m_{31}^2 = \alpha\phi^3 = \alpha/(2\alpha_s)$  (Eq. 148; 5% with PDG 2024 central values).
- The graviton spectral gap  $7-4\phi$  predicts a graviton mass  $\sim 10^{-34}$  eV for  $R = R_{\text{Hubble}}$ , far below current bounds.
- The gap-Planck identity  $\lambda_1^{\text{exact}} \cdot 4\phi^2 = 24 = |2T|$  connects gravity to the generation group.

## 15 Discussion and Conclusions

### 15.1 Summary of Predictions

To guard against post-hoc numerical fitting, we distinguish three evidential levels: (i) **Rigorous derivation** ( $a_1 = 5 \rightarrow$  result via established mathematics, no choices): gauge group,  $N_{\text{gen}}$ , the constructive  $(a, b)$  assignment, exact discrete scalar response, the simplicial Yang–Mills stiffness  $K_{\text{YM}} = b_1$ , CKM bare exponents,  $\delta_{\text{CKM}}$ ,  $\theta_{\text{QCD}} = 0$ . (ii) **Algebraically motivated reproduction** (framework constants only, but with a well-motivated identification step): mass corrections ( $\chi^2/\text{dof} = 0.28$ ,  $p = 0.97$ ), coupling constants (all three discrete ingredients of the  $\alpha$  tree-level coefficient derived, with only the KK identification and the  $M_Z$ -scale matching still structural), and continuum bosonic analogies from the discrete spectral action. (iii) **Open problems:** continuum interpretation of the discrete  $\alpha$  coefficient,  $M_Z$  scale, cosmological constant mechanism, nonlinear gravity. Table 5 categorizes every result:

| Quantity                                                                                                                            | Error                      | Status     |
|-------------------------------------------------------------------------------------------------------------------------------------|----------------------------|------------|
| <i>Coupling constants (0 free parameters)</i>                                                                                       |                            |            |
| $\alpha_s = 1/(2\phi^3)$ , $\sin^2 \theta_W = 6/26$                                                                                 | 0.11–0.19%                 | Derived    |
| $\alpha$ (Eq. 34: tree-level coefficient fully discrete; KK identification structural)                                              | 0.0001%                    | Structural |
| <i>Boson masses (+<math>m_e</math>)</i>                                                                                             |                            |            |
| $m_Z$ ( $n=a_1^2$ selected by RG bootstrap, Eq. 96), $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$ , $m_e/m_P = \alpha^{4\phi^2}$           | 0.008–0.24%                | Structural |
| <i>Structural</i>                                                                                                                   |                            |            |
| Gauge group $SU(3) \times SU(2) \times U(1)$ , $N_{\text{gen}} = 3$ (vertex + edge, Sec. 4.2)                                       | Exact                      | Derived    |
| $\ker(\square_1) = 13 = \rho_0 \oplus 2\rho_5$ ; $C _{\text{gauge}} = a_1$ ; $\text{Tr}(\square_1^2)/\text{Tr}(\square_0^2) = 9a_1$ | Exact                      | Derived    |
| Constructive ( $a, b$ ) uniqueness (Thm. 6), McKay sum rules                                                                        | Exact                      | Derived    |
| <i>Fermion masses (0 free params +<math>m_e</math>)</i>                                                                             |                            |            |
| All 9 fermions (with vertex correction)                                                                                             | $\chi^2/\text{dof} = 0.28$ | Derived    |
| Mass corrections: $\alpha = 1 - d_{\text{eff}}/d_{\text{ST}}$ (Eq. 86)                                                              | Exact                      | Derived    |
| <i>Mixing angles and CP violation</i>                                                                                               |                            |            |
| CKM ( $\theta_{12}, \theta_{23}, \theta_{13}$ )                                                                                     | < 0.2%                     | Derived    |
| CKM concordance ( $n_{12} \cdot n_{13} = b_1^2$ etc.)                                                                               | Exact                      | Derived    |
| Cayley–CKM vacuum offset ( $\delta = 1$ , Thm. 14)                                                                                  | Exact                      | Derived    |
| $E_8$ overlap quantization (Thm. 15, 780 pairs)                                                                                     | Exact                      | Derived    |
| PMNS ( $\theta_{12}, \theta_{23}, \theta_{13}$ )                                                                                    | < 2.3%                     | Derived    |
| $\delta_{\text{CKM}} = \arctan(\sqrt{5})$ , $\delta_{\text{PMNS}} = 3\delta_{\text{CKM}}$                                           | < 0.8%                     | Derived    |
| $\theta_{\text{QCD}} = 0$ (Galois), $J = 3.12 \times 10^{-5}$                                                                       | Exact / 1.1%               | Derived    |
| <i>Gravity</i>                                                                                                                      |                            |            |
| Discrete scalar response $\gamma_{\text{disc}} = 1$ (Thm. 9), hierarchy $z_P = 1/\alpha_s + 2$                                      | Exact / structural         | Derived    |
| Spectral coefficients $(c_0, c_1, c_2) = (2640, 14880, 55920)$ , $(A_0, A_1, A_2) = (11, 62, 233)$                                  | Exact                      | Derived    |
| Gauge prefactors $(8/15, 1/3, 2/15)$ from $Q_8 \subset 2T \subset 2I$ vertex decomposition                                          | Structural                 | Structural |
| Emergent graviton: $R(l) \rightarrow 2$ polarizations (Eq. 119)                                                                     | Exact                      | Derived    |
| <i>Neutrinos (+<math>m_e</math>)</i>                                                                                                |                            |            |
| $m_3 = 2m_e/\phi^{35}$ , $\Delta m_{21}^2/\Delta m_{31}^2 = \alpha\phi^3$                                                           | < 5%                       | Derived    |
| <i>Extensions (Sections 13.1–13.4)</i>                                                                                              |                            |            |
| Galois dark sector (9 stable, EM-dark particles)                                                                                    | —                          | Derived    |
| $\Lambda_P = \alpha^{57-\alpha_s}$                                                                                                  | 0.33%                      | Pattern    |
| $\Omega_{\text{DM}}/\Omega_b = 7 - \phi$                                                                                            | 0.41%                      | Pattern    |

Table 5: Summary of all predictions from  $a_1 = 5$  with zero free parameters (all dimensionless ratios algebraic;  $m_e$  is the sole dimensional anchor). “Derived” = deduced via established mathematics; “Pattern” = algebraic formula matching experiment but not uniquely derived. †Structural: three  $d_{\text{eff}}$  regimes and Hölder exponents derived; explicit spectral action normalization partially open (see Section 15.2). ‡The scalar-response theorem is exact on the discrete 600-cell; the continuum PPN interpretation is not claimed. §The vertex-type mechanism ( $Q_8 \subset 2T \subset 2I$ ) is verified on all 120 vertices.  $Q_8$  is the unique maximally central subgroup: all commutators lie in  $Z(2I)$ , and its character weights vanish on the Galois-broken gauge sectors  $\rho_8, \rho_2$  (the residual rational weight on  $\rho_4$  cancels at the tetrahedral shell). The full table with detailed footnotes is in Supplementary Material S15.

## 15.2 Limitations

The principal open problems are:

1. **Mass scale.** The electron is the unique internally selected anchor state: it is the Galois fixed point ( $z_e = 0$ ), the McKay graph origin (trivial irrep  $\rho_0$ ), and the

ground state of the mass ladder ( $n = 0$ ). The physically meaningful dimensionless hierarchy  $m_e/m_P = \alpha^{4\phi^2}$  is derived from the spectral data (0.24% agreement); what remains external is only the conversion from Planck units to laboratory units. This is a genuine no-go: the spectral data are pure numbers, so by dimensional analysis they cannot generate a quantity with units of GeV. But it is a much weaker gap than “ $m_e$  is a free parameter”: the framework determines *all* dimensionless ratios, and the sole residual freedom is a unit convention, not a physical parameter. In particular,  $m_e$  fixes the unit system, not the physics: every prediction in this paper—coupling constants, mass ratios, mixing angles, CP phases—is a pure number derived from  $a_1 = 5$  with zero adjustable parameters. No measurement of  $m_e$  (or any other dimensionful quantity) can falsify or confirm these dimensionless predictions.

2. **Effective action.** The vacuum-selection functional  $S_{\text{eff}} = S_{\text{EW}} + w_c S_{\text{cosmo}}$  is robust and sharply minimized at  $(a_1, n) = (5, 25)$  on the scanned domain. Its *local* form around the minimum is structurally justified: the linear term vanishes identically, the curvature (175) is derived from spectral data, and the quadratic approximation reproduces the exact cost to  $\sim 10\%$  at  $|\delta n| = 1$ . The *global* form need not be specified: the selection is universal within the class of non-negative mismatch functionals with one-loop RG monotonicity ( $L^2$ ,  $L^1$ ,  $L^\infty$ , and Kullback–Leibler norms all select the same vacuum, verified numerically). The sole remaining open problem is the absolute scale conversion (Section 15.6).
3. **Spectral action normalization.** The exact discrete coefficients  $(c_0, c_1, c_2) = (2640, 14880, 55920)$ , the reduced triple  $(11, 62, 233)$ , and the Diophantine identity (139) are established. The gauge prefactors  $(8/15, 1/3, 2/15) = (|Z(2I)| : a_1 : \text{rank}(E_8))/(N_{\text{gen}} \cdot a_1)$  are explained by the vertex-type decomposition under  $Q_8 \subset 2T \subset 2I$  (Eq. 137), with  $Q_8$ ’s abelian role now derived from three coordinate-free criteria (commutator centrality, Galois trace classification, character-weight vanishing on broken sectors). What remains open is a controlled continuum identification with the full bosonic Standard Model action.
4. **Gravity.** The linearized field equation, the exact discrete scalar-response identity  $\gamma_{\text{disc}} = 1$ , and the correct graviton polarization count ( $R \rightarrow 2$ ) are derived on the discrete model; the 4D nonlinear completion and any continuum PPN interpretation remain open.
5. **Cosmological constant.** The formula  $\Lambda_P = \alpha^{57-\alpha_s}$  matches observation to  $0.13\sigma$ . The exponent 57 is fully derived (seven routes); the spectral determinant  $\det'(L)$  is Galois-invariant (Supplementary Material S13) but gives  $\log \det'(L/N)/\ln \alpha = 56.97$ , a near-miss that does not close. The dynamical mechanism selecting the base  $\alpha^z$  remains the framework’s most serious theoretical gap.
6. **Dark matter abundance.** Qualitative properties are derived; the abundance ratio  $\Omega_{\text{DM}}/\Omega_b = 7 - \phi$  requires a reheating temperature not derived from the framework. The spectral action reheating temperature  $T_{\text{reh}} \approx 2.2 \times 10^9$  GeV (Starobinsky preheating) yields  $\Omega_{\text{DM}}/\Omega_b \approx 0.6$  via gravitational freeze-in, an order of magnitude below the observed 5.38; the pattern temperature  $T_{\text{reh}} = M_P/\phi^{11}$  is inconsistent with the spectral value by four orders of magnitude.
7. **Neutrino spectral asymmetry.** The quantity  $|\eta_A| = 35$  is an adjacency spectral invariant, not the APS  $\eta$ -invariant; the operator-theoretic connection remains open.

8. **Holonomy sector dependence.** Berry phase baseline  $1/\phi^4$  is derived, but sector-dependent corrections lack a single-operator origin (three no-go results, Proposition 13).
9. **Mass correction unification.** The three cascade levels are identified as Morrey–Sobolev regimes at effective dimensions  $d_{\text{eff}} = 0, 1, 4$  (Eq. 86). The spectral action provides the  $p = 4$  energy and the Galois anti-symmetric fluctuation generates the corrections. The remaining gaps are: (i)  $d_{\text{eff}} = d_{\text{ST}}$  for non-units rests on a structural argument (spinorial doubling per archimedean place), not a rigorous proof; (ii) the Level 2→Level 3 boundary is discontinuous (+0.163 vs. −0.400), reflecting different  $d_{\text{eff}}$  regimes; (iii) the explicit spectral action computation of the Morrey normalization coefficients has not been carried out.

### 15.3 Statistical Robustness

Three arguments guard against a look-elsewhere effect. *Counting:* the mass formula generates  $\sim 121$  candidates spanning  $\Delta \approx 25$  decades in  $\phi$ ; the probability of all 9 fermion masses matching simultaneously is  $P_9 \lesssim 10^{-12}$ , before imposing the sum-rule constraints that reduce the search space to  $\mathcal{O}(100)$  assignments. *Coherence:* the *same*  $(a, b)$  quantum numbers determine masses, mixing angles, CP phases, and McKay weights; a single mis-assignment destroys multiple predictions simultaneously. *Uniqueness:* Theorem 6 proves the assignment is the unique solution among  $\sim 5.4 \times 10^8$  candidates, with zero continuous parameters. The framework also makes sharp falsifiable predictions:  $a_\mu^{\text{NP}} = 0$ ,  $\theta_{\text{QCD}} = 0$  (no axion), and specific Majorana phases.

### 15.4 Uniqueness Among Regular Polytopes

Among all six regular 4D polytopes, only the 600-cell passes all seven framework tests ( $N_{\text{gen}} = 3$ , gauge group,  $\sin^2 \theta_W$ ,  $\alpha^{-1}$ , mass hierarchy, CKM/PMNS, anomaly cancellation); the other five score 0/7. Even the 120-cell (same algebra,  $a_1 = 5$ ) fails on gauge group (vertex degree 4, no color octet). Full comparison table and failure-mode analysis are in Supplementary Material S16.

### 15.5 Variational Selection

Among all finite subgroups of  $\text{SU}(2)$  (the complete ADE classification),  $2I$  uniquely minimizes the spectral action density  $\rho(G) := \text{Tr}(A_G^2)/|G|$ :  $\rho(2I) = 2/15 = 2/(a_1 N_{\text{gen}})$ , less than half the runner-up  $\rho(2O) = 7/24$ . Combined with the TQFT bootstrap (Theorem 1),  $a_1 = 5$  is the unique self-consistent vacuum *and* the unique action minimum. Full proof and physical interpretation are in Supplementary Material S16.

### 15.6 Vacuum Selection by Effective Action

The RG-bootstrap cost (96) can be extended to include a cosmological sector. Define the effective action

$$S_{\text{eff}}(a_1, n) = S_{\text{EW}}(a_1, n) + w_c S_{\text{cosmo}}(a_1) \quad (174)$$

where  $S_{\text{EW}}$  is the coupling-matching cost (96) and  $S_{\text{cosmo}}(a_1) = (\log \Lambda_P^{\text{pred}} / \log \Lambda_P^{\text{obs}} - 1)^2$  measures the mismatch of the cosmological constant prediction  $\Lambda_P = \alpha^{z_{\text{CC}}}$  with  $z_{\text{CC}} =$

$a_1!/2 - (a_1 - 2)$  against the observed value  $\Lambda_P^{\text{obs}} \approx 10^{-122}$  (Planck units). Three properties hold:

- (i) *Unique global minimum.* The pair  $(a_1, n) = (5, 25)$  minimizes  $S_{\text{eff}}$  over all tested  $(a_1, n)$  with  $a_1 \in \{3, \dots, 10\}$ ,  $n \in \{10, \dots, 40\}$ . The nearest competitor has  $S_{\text{EW}}$  at least  $59\times$  larger; competitors at different  $a_1$  exceed  $10^3 \times S_{\text{EW}}(5, 25)$ . For the adopted observed value of  $\Lambda$ , the cosmological cost is numerically zero at  $a_1 = 5$ , while  $S_{\text{cosmo}}(4) \approx 0.71$  and  $S_{\text{cosmo}}(6)$  is enormously larger.
- (ii) *Weight robustness.* Scanning 131 weight combinations with  $w_1, w_2, w_c$  spanning four decades, the minimum remains at  $(5, 25)$  in 99.2% of cases. The discrete Hessian at the minimum is positive definite with eigenvalue ratio  $\sim 16$  (steeper in  $a_1$  than in  $n$ ).
- (iii) *Exponential dominance.* Interpreting  $\Psi(a_1, n) \propto \exp(-S_{\text{eff}}/S_0)$  as a no-boundary-style wavefunction,  $(5, 25)$  saturates the partition function to machine precision on the scanned domain. The next saddle at  $(5, 26)$  is suppressed by  $\sim 10^{25}$ ; tunneling to any  $a_1 \neq 5$  is suppressed by  $\exp(-10^3)$ .

The cost function (174) uses no external input: the one-loop beta coefficients  $b_i = (41/10, -19/6, -7)$  are themselves derived from the framework's particle content ( $N_{\text{gen}} = 3$  and the gauge group), so the RG running is intrinsic. The exponent  $n_0 = a_1^2 = 25$  is a spectral invariant—the multiplicity of the zero eigenvalue of the adjacency matrix (Table 2)—not an external choice. The local form of the cost around  $n_0$  is structurally constrained:

- (iv) *Structural vanishing of the linear term.* Since  $S_{\text{EW}}$  is a sum of squared relative deviations, each of which vanishes at  $n = n_0$  (where the couplings equal their algebraic values),  $\partial S / \partial n|_{n_0} = 0$  identically. This is a property of any non-negative functional vanishing at a point, not a tuning.

- (v) *Derived curvature.* The quadratic coefficient is

$$\left. \frac{\partial^2 S_{\text{EW}}}{\partial n^2} \right|_{n_0} = \frac{49 \ln^2 \phi}{8\pi^2 \phi^6} + (\text{small } \sin^2 \theta_W \text{ channel}) = 8.1 \times 10^{-3}, \quad (175)$$

where  $49 = b_3^2 = (4N_{\text{gen}}/3 - 11)^2$  is the squared QCD beta coefficient,  $\phi^6 = (2\alpha_s)^{-2}$ ,  $\ln \phi$  is the mass-ladder spacing, and  $\pi$  enters through the Hopf holonomy. The  $\alpha_s$  channel contributes 98.7% of the total curvature; the Weinberg-angle channel contributes 1.3%. Every factor in (175) is a framework constant derived from  $a_1$ . The quadratic approximation  $S \approx \frac{1}{2} c_2 (\delta n)^2$  agrees with the exact one-loop cost to  $\sim 10\%$  at  $|\delta n| = 1$  and  $\sim 30\%$  at  $|\delta n| = 2$ ; beyond that, the exact (logarithmic) form of the RG running becomes relevant.

- (vi) *Electroweak window.* The lattice scale  $m_e \phi^{a_1^2}$  falls in the electroweak range  $[1, 10^4]$  GeV *only* for  $a_1 = 5$  (85.7 GeV). For  $a_1 = 4$ : 0.34 GeV (sub-GeV); for  $a_1 = 6$ :  $1.7 \times 10^5$  GeV (far above). The spectral fixed point and the electroweak window are compatible for a single value of  $a_1$ .

The local cost form is therefore not ad hoc: the quadratic shape, the vanishing linear term, and the curvature all follow from the spectral data. Moreover, the selection

is *universal* within the physically relevant class of functionals: any non-negative mismatch measure  $S \geq 0$  satisfying  $S(n_0) = 0$  and the one-loop RG monotonicity condition  $|\partial g_i / \partial n| > 0$  for  $n \neq n_0$  selects (5, 25) as its unique global minimum. Four representative norms— $L^2$ ,  $L^1$ ,  $L^\infty$ , and Kullback–Leibler—all select the same vacuum (verified numerically on  $n \in [15, 35]$ ). The universality is *not* polynomial (polynomial jet deformations can generate competing minima via negative-cost artifacts), but it *is* physical: no non-negative cost with monotonic RG structure admits a second zero on the scanned domain.

What remains open is the absolute scale conversion—why the lattice spacing is  $1/(m_e \phi^{25})$  rather than, say,  $1/m_P$ —which is the hierarchy problem in discrete form. The 600-cell spectral data are pure numbers in lattice units; no functional of these data can produce a dimensionful quantity without at least one dimensionful input. Given one such input ( $m_e$  or equivalently  $m_P$ ), the spectrum determines all other scales through derived ratios ( $m_e/m_P = \alpha^{4\phi^2}$  to 0.24%,  $m_Z/m_e = \phi^{a_1^2} \cdot R$  to 0.09%). The hierarchy  $m_e/m_P \sim 10^{-22}$  is  $\alpha^{10.5}$ , where both  $1/137$  and  $10.5$  are moderate numbers derived from  $a_1$ ; the large ratio arises from exponentiation, not fine-tuning.

## 15.7 Conclusions

We have shown that a substantial part of Standard Model structure—its gauge group, coupling constants, fermion content, masses, mixing angles, CP structure, and discrete gravitational analogues—can be systematically organized from a single integer  $a_1 = 5$ .

The framework has a *two-layer architecture*. The first layer is a self-referential algebraic seed: the equation  $x^2 = x + 1$  generates the golden ratio  $\phi$ , whose Fibonacci fusion rule defines a topological quantum field theory that selects  $a_1 = 5$  uniquely (Theorem 1). This layer produces  $a_1$ ,  $\phi$ ,  $N_{\text{gen}} = a_1 - 2 = 3$ , and the  $\mathbb{Z}[\phi]$  mass lattice—but no quantitative coupling constants or masses. The second layer is the 600-cell realization: the binary icosahedral group  $2I$  (the unique group of order  $a_1$ !) and its McKay graph ( $\tilde{E}_8$ ) supply the Laplacian spectrum, gauge decomposition, character sums, and Galois structure from which all quantitative predictions follow. The electroweak scale is not postulated: the RG-bootstrap cost function (96) has a sharp, isolated global minimum at  $(a_1, n) = (5, 25)$ , selecting the exponent variationally. The sole irreducible gap between the two layers is the transcendental factor  $2\pi$  in the  $\alpha$  equation, which enters through the topology of the Hopf fiber and cannot be produced from the algebraic discrete data alone.

The key structural results, grouped by theme, are:

1. **Gauge group and generations.** The permutation representation on the icosahedral vertex figure gives  $12 = \mathbf{1} \oplus \mathbf{3} \oplus \mathbf{3}' \oplus \mathbf{5}$ ; Theorem 3 proves  $G = U(1) \times SU(2) \times SU(3)$  is unique. The Generation Theorem gives  $N_{\text{gen}} = 3$  from Galois invariance on  $\mathbb{Z}[\phi]$ .
2. **Coupling constants and gravity.** Two gauge couplings are algebraically derived ( $\alpha_s = 1/(2\phi^3)$  from the  $\mathbb{Z}[\phi]$  factor structure (0.11%),  $\sin^2 \theta_W = 6/26$  from graph combinatorics (0.19%)); the third ( $\alpha$  from the icosahedral Laplacian, 0.0001%) is now better constrained than before: the tree-level coefficient admits the exact discrete form  $1/\alpha_0 = N/(K_{\text{YM}} \lambda_1(L_{\text{fiber}})^2)$ , where  $N = |2I|$  is combinatorial,  $K_{\text{YM}} = b_1$  is the exact simplicial Yang–Mills stiffness, and  $\lambda_1(L_{\text{fiber}}) = 2 - \phi$  is the Hopf-fiber spectral gap. Thus all three factors of the linear coefficient are internally



derived; the remaining structural step is the Kaluza–Klein identification of this discrete tree-level coefficient with the electromagnetic inverse coupling. The  $\alpha$  equation  $\text{Vol}(S^1) \alpha^2 - (N/b_1) \phi^4 \alpha + 1 = 0$  has three coefficients of distinct origin: topological (Hopf holonomy  $2\pi$ ), gauge-theoretic (exact Yang–Mills stiffness  $K_{\text{YM}} = b_1$ ), and spectral (equivalently  $\phi^4 = \lambda_1(L_{\text{fiber}})^{-2}$ ), plus normalization (1). The Vieta product  $\alpha\alpha' = 1/\text{Vol}(S^1)$  is the standard Kaluza–Klein coupling on the Hopf fiber. The hierarchy identity  $m_e/m_P = \alpha^{1/\alpha_s+2}$  connects all three forces. On the bosonic side, the exact discrete spectral-action core is now the coefficient triple  $(2640, 14880, 55920)$  with reduced triple  $(11, 62, 233)$  and identity (139), together with the exact discrete scalar-response theorem  $\gamma_{\text{disc}} = 1$  and the exact abelian Yang–Mills response  $S_{\text{YM}}(\theta) = 2400 \sin^2(\theta/20)$ ; the gauge prefactors  $(8/15, 1/3, 2/15)$  are now explained structurally by the vertex-type decomposition  $120 = 8(Q_8) + 16(2T) + 96$  under the subgroup chain  $Q_8 \subset 2T \subset 2I$  (Eq. 137); what remains open is the continuum identification.

3. **Fermion masses and uniqueness.** The Fibonacci mass formula  $m_f = m_e \phi^{5a+6b+\delta_1+\delta_2}$  with zero-parameter corrections (norm-log for quarks, Galois conjugate for leptons, Verlinde radiative corrections with  $c_{\text{eff}} = \sqrt{2/a_1} (1 - b_1 \alpha^2)$  from the  $S$ -matrix prefactor and fusion matrix Frobenius norm) achieves  $\chi^2 = 2.2/8$  dof ( $p = 0.97$ , RMS pull  $0.53\sigma$ ) across all 9 fermions. The  $(a, b)$  assignment is no longer only a bounded-search observation: Theorem 6 gives a constructive derivation from McKay main-chain weights, minimal local edge lifts in  $\mathbb{Z}[\phi]$ , branch identification from affine- $E_8$  geometry and chirality, and the prime-sector Galois pairing  $z_b = \phi \sigma(z_t)$ . The older conditions (C1)–(C2) survive as arithmetic corollaries and as a computational cross-check inside the bounded domain. The McKay mass correspondence forces the main-chain weights to  $[N_{\text{gen}}, F(3), b_1, 0, a_1]$  uniquely (Theorem 16). The three correction regimes (logarithmic, power-law, constant) are unified by the Morrey–Sobolev embedding at effective dimension  $d_{\text{eff}} = 0, 1, d_{\text{ST}}$ , with Hölder exponent  $\alpha = 1 - d_{\text{eff}}/d_{\text{ST}}$  (Eq. 86). The Higgs-to- $W$  mass ratio  $m_H^2/m_W^2 = \phi^2 - 16\alpha\phi$  follows from  $L(\mathbf{3}')/L(\mathbf{3}) = \phi^2$  with a derived unit coefficient (Proposition 5).
4. **Mixing angles and CP violation.** CKM bare exponents  $(3, 7, 12) = (N_{\text{gen}}, \text{degree} - a_1, \text{degree})$  with  $E_8$  leg-ratio corrections ( $c_{12} = 2/3$  derived from McKay graph, signs from arm topology); PMNS angles from the Galois mixing matrix with eigenvalues  $\{0, 3, 5\}$  and TBM eigenvectors;  $\theta_{13} = 0$  exact at the  $A_5$  level (spectral gap  $4\sqrt{a_1}$ ), with  $\sin^2 \theta_{13} = 1/45$  as the democratic tunneling rate. The three bare CKM exponents satisfy algebraic identities whose intersection singles out  $a_1 = 5$  (Theorem 13, thirteenth characterisation). The Cayley spectral exponents  $(n_{\text{ut}}, n_{\text{db}}) = (4, 2)$  are connected to the CKM exponents by a uniform vacuum offset  $\delta = 1$  forced by the concordance identities (Theorem 14, fourteenth characterisation). CP phases  $\delta_{\text{CKM}} = \arctan(\sqrt{5})$  and  $\delta_{\text{PMNS}} = 3\delta_{\text{CKM}}$ ;  $\theta_{\text{QCD}} = 0$  by Galois invariance.
5. **Galois dark sector and cosmology.** The Galois conjugation generates 9 stable, electromagnetically dark particles ( $\alpha'$  complex,  $\alpha'_s < 0$ ) with mass duality  $m_f |m_{f'}| = m_e^2$ . The cosmological constant  $\Lambda_P = \alpha^{57-\alpha_s}$  matches observation to  $0.13\sigma$ , with seven mutually consistent decompositions of 57 (see Supplementary Material S13), each connecting 57 to independent structural data of the 600-cell.
6. **McKay–Cayley sum rules.** The Cayley-graph Laplacian eigenvalues satisfy  $\sum a_i = 12 = \dim(G)$  and  $\sum b_i = 8 = \text{rank}(E_8)$ , and the  $2I$  character table is

recovered exactly from  $\hat{E}_8$  eigenvectors.

7. **Galois kernel (Theorem 10).** The Lorentzian operator  $\square = b_1 A_{\text{fiber}} - A$  on any Hopf fibration has  $\ker(\square) = \rho_0 \oplus \rho_1 \oplus \rho_8$  with  $\dim = 9 = N_{\text{eig}}$ . The coefficient  $b_1$  is the *unique* nontrivial integer giving a nonzero kernel (see Supplementary Material S7.2), and the kernel selects the smallest irrep from each Galois orbit, with dimension sum  $1 + 2 + 2 = 5 = a_1$ . The kernel is stable across all 27 Hopf fibrations.

With  $\sim 38$  quantities constrained from framework data and the Galois structure of  $\mathbb{Z}[\phi]$  simultaneously organizing CP violation, the strong CP problem, the generation count, dark matter darkness, and the cosmological constant scale, the framework represents a tightly constrained mathematical structure. The effective vacuum-selection principle is no longer locally ad hoc: its local form is structurally induced (vanishing linear term, positive curvature (175) dominated by the  $\alpha_s$  channel, all coefficients derived from  $a_1$ ), and the selection is universal within the class of non-negative mismatch functionals with one-loop RG monotonicity.

The electron is the unique internally selected anchor state—the Galois fixed point of the mass lattice, the trivial irrep on the McKay graph, and the ground state of the mass operator. In Planck units, its mass is a derived pure number:  $m_e/m_P = \alpha^{4\phi^2}$  (0.24%). The physically meaningful dimensionless hierarchy is therefore explained; what remains external is only the conversion from Planck units to laboratory units. Whether even that residual convention can be eliminated—for example, by coupling the discrete geometry to a gravitational path integral that generates a Planck mass dynamically—remains open.

## 15.8 Related Work

The framework shares conceptual ground with the Chamseddine–Connes–Marcolli spectral action programme [4, 14] (same dynamical principle, but finite geometry *determined* by  $a_1 = 5$  rather than chosen), the Boyle–Farnsworth algebraic axioms [25] (for which the 600-cell provides the missing selection principle), and the Furey–Hughes division-algebra triality [22] (both exploit  $D_4$  branching for  $N_{\text{gen}} = 3$ ). Detailed comparisons with these and with Todorov–Drenska [26] and Singh [23] are in Supplementary Material S16.

## Acknowledgments

We acknowledge the foundational work on spectral geometry by Alain Connes, Ali Chamseddine, and Matilde Marcolli [4, 14], the geometric optimality results of Cohn and Kumar [1, 2], and the 600-cell GUT framework developed by Christopher C. O’Neill [5, 6]. Experimental data from CODATA 2022 [9] and the Particle Data Group [10].

## Data and Code Availability

All numerical results reported in this paper can be reproduced using the self-contained Python scripts available at <https://github.com/razvananghelina/One-Integer-Three-Generation>. The repository contains five verification scripts covering: coupling constants, 600-cell Laplacian spectrum and irrep localization, fermion masses and mixing angles, Berry phase computation, and spectral action properties. Each script requires only NumPy and SciPy.

# References

- [1] H. Cohn and A. Kumar, “Universally optimal distribution of points on spheres,” *J. Amer. Math. Soc.* **20**, 99–148 (2007).
- [2] H. Cohn and A. Kumar, “Optimality and uniqueness of the Leech lattice among lattices,” *Ann. of Math.* **170**, 1003–1050 (2009).
- [3] H. Cohn, A. Kumar, S. D. Miller, D. Radchenko, and M. Viazovska, “Universal optimality of the  $E_8$  and Leech lattices and interpolation formulas,” *Ann. of Math.* **196**, 983–1082 (2022).
- [4] A. H. Chamseddine and A. Connes, “The Spectral Action Principle,” *Commun. Math. Phys.* **186**, 731–750 (1997).
- [5] C. C. O’Neill, “The 600-Cell Grand Unified Theory: A Geometric and Group-Theoretic Model of the Standard Model and Beyond,” preprint (2025).
- [6] C. C. O’Neill, “Isomorphism of a Modified Standard Model Particle Assignment with the Binary Octahedral Group (24-cell),” preprint (2023).
- [7] J. H. Conway and N. J. A. Sloane, “Quaternionic lattices and the icosians,” in *Sphere Packings, Lattices and Groups*, 3rd ed., Springer, 1998.
- [8] V. Elser and N. J. A. Sloane, “A highly symmetric four-dimensional quasicrystal,” *J. Phys. A* **20**, 6161–6168 (1987).
- [9] E. Tiesinga et al., “CODATA recommended values of the fundamental physical constants: 2022,” *Rev. Mod. Phys.* (2024).
- [10] R. L. Workman et al. (Particle Data Group), “Review of Particle Physics,” *Prog. Theor. Exp. Phys.* **2024**, 083C01 (2024).
- [11] T. Regge, “General relativity without coordinates,” *Nuovo Cimento* **19**, 558–571 (1961).
- [12] A. G. Lisi, “An Exceptionally Simple Theory of Everything,” arXiv:0711.0770 [hep-th] (2007).
- [13] C. Furey, “Standard Model physics from an algebra?” Ph.D. thesis, University of Waterloo (2015). arXiv:1611.09182 [hep-th].
- [14] A. H. Chamseddine, A. Connes, and M. Marcolli, “Gravity and the Standard Model with neutrino mixing,” *Adv. Theor. Math. Phys.* **11**, 991–1089 (2007).
- [15] M. Garny, M. Sandora, and M. S. Sloth, “Planckian interacting massive particles as dark matter,” *Phys. Rev. Lett.* **116**, 101302 (2016).
- [16] Y. Ema, K. Nakayama, and Y. Tang, “Production of purely gravitational dark matter: the case of fermions and vectors,” *JHEP* **07**, 060 (2019).
- [17] J. Distler and S. Garibaldi, “There is no ‘Theory of Everything’ inside  $E_8$ ,” *Commun. Math. Phys.* **298**, 419–436 (2010).

- [18] ATLAS Collaboration, “Measurement of the Higgs boson mass with  $H \rightarrow \gamma\gamma$  and  $H \rightarrow ZZ^* \rightarrow 4\ell$  events using  $pp$  collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector,” arXiv:2308.04775 [hep-ex] (2023).
- [19] CMS Collaboration, “Measurement of the Higgs boson mass and width using the four-lepton final state in proton-proton collisions at  $\sqrt{s} = 13$  TeV,” arXiv:2409.13663 [hep-ex] (2024).
- [20] JUNO Collaboration, “Measurement of the solar neutrino oscillation parameters with the first data of JUNO,” arXiv:2511.14593 [hep-ex] (2025).
- [21] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, and T. Schwetz, “NuFIT 6.0,” arXiv:2410.05380 [hep-ph] (2024).
- [22] N. Furey and M. J. Hughes, “Three Generations and a Trio of Trialities,” *Phys. Lett. B* (2025). arXiv:2409.17948 [hep-th].
- [23] T. P. Singh, “Mass ratios of charged leptons and quarks from the exceptional Jordan algebra,” arXiv:2508.10131 [hep-ph] (2025).
- [24] F. Latrémolière, “The quantum Gromov–Hausdorff propinquity,” *Trans. Amer. Math. Soc.* **368** (2016) 365–411. arXiv:1506.04341 [math.OA].
- [25] L. Boyle and S. Farnsworth, “The standard model, the Pati–Salam model, and ‘Jordan geometry’,” *New J. Phys.* **22**, 073023 (2020). arXiv:1910.11888 [hep-th].
- [26] I. Todorov and S. Drenska, “Octonions, exceptional Jordan algebra and the role of the group  $F_4$  in particle physics,” *Adv. Appl. Clifford Algebras* **28**, 82 (2018). arXiv:1805.06739 [hep-th].
- [27] A. A. Starobinsky, “A new type of isotropic cosmological models without singularity,” *Phys. Lett. B* **91**, 99–102 (1980).
- [28] Y. Akrami et al. (Planck Collaboration), “Planck 2018 results. X. Constraints on inflation,” *Astron. Astrophys.* **641**, A10 (2020). arXiv:1807.06211 [astro-ph.CO].
- [29] E. Allys et al. (LiteBIRD Collaboration), “Probing cosmic inflation with the LiteBIRD cosmic microwave background polarization survey,” *Prog. Theor. Exp. Phys.* **2023**, 042F01 (2023). arXiv:2202.02773 [astro-ph.IM].
- [30] A. Kitaev and J. Preskill, “Topological entanglement entropy,” *Phys. Rev. Lett.* **96**, 110404 (2006). arXiv:hep-th/0510092.
- [31] M. Levin and X.-G. Wen, “Detecting topological order in a ground state wave function,” *Phys. Rev. Lett.* **96**, 110405 (2006). arXiv:cond-mat/0510613.
- [32] Y. Koide, “New viewpoint in lepton mass matrix — a prediction  $m_\tau = 1776.97$  MeV,” *Phys. Rev. Lett.* **47**, 1241 (1981); “Fermion–boson two-body model of quarks and leptons and Cabibbo mixing,” *Lett. Nuovo Cim.* **34**, 201 (1982).
- [33] G. Degrandi, S. Di Vita, J. Elias-Miró, J. R. Espinosa, G. F. Giudice, G. Isidori, and A. Strumia, “Higgs mass and vacuum stability in the Standard Model at NNLO,” *JHEP* **1208**, 098 (2012), arXiv:1205.6497.

- [34] G. F. Chew and S. Mandelstam, “Theory of the low-energy pion–pion interaction,” *Phys. Rev.* **119**, 467 (1960). See also S. U. Chung, J. Brose, R. Hackmann, E. Klempt, S. Spanier, and C. Strassburger, “Partial wave analysis in K-matrix formalism,” *Ann. Phys. (Leipzig)* **4**, 404 (1995).
- [35] R. H. Dalitz and S. F. Tuan, “The phenomenological representation of K-nucleon scattering and reaction amplitudes,” *Ann. Phys.* **10**, 307 (1960).

## A Key Propositions

We collect the geometric propositions referenced in the main text. Full proofs and extended discussion are in Supplementary Material S17.

**Proposition 13** (Static geometry no-go). *The sector-dependent holonomy corrections cannot arise from the static 600-cell geometry: (i) edge weights are isotropic ( $w_R = d_R/12$ ); (ii) Hopf and gauge decompositions are independent ( $\chi^2 = 0$ ); (iii) holonomy is face-transitive (Berry phase =  $1/\phi^4$  universally).*

**Proposition 14** (McKay–CG Yukawa structure). *The finite Dirac operator  $D_F$  on the McKay graph of  $2I$  satisfies: (i) all 16 CG maps have multiplicity 1; (ii)  $\|Y_e\|_F = 1/\sqrt{2}$  universally; (iii)  $\text{Tr}(D_F^2) = 8 = \text{rank}(E_8)$ ; (iv) spectral gap  $4\sqrt{a_1}$  between Galois-conjugate Cayley eigenvalues.*

**Proposition 15** (McKay–Cayley spectral sum rules). *The Cayley-graph Laplacian eigenvalues  $\lambda_i = a_i + b_i\phi$  satisfy:*

$$\sum_{i=1}^9 a_i = 12 = \dim(G), \quad (176)$$

$$\sum_{i=1}^9 b_i = 8 = \text{rank}(E_8). \quad (177)$$

*The  $2I$  character table is recovered exactly from the  $\hat{E}_8$  eigenvectors.*

**Proposition 16** (Galois anti-symmetric Cayley eigenvalue). *For each irrep  $\rho_i$  of  $2I$ , define  $G_i = (\lambda_i(10a) - \lambda_i(5a))/2$ . Then  $G_i = 0$  for WHITE/ $A_5$  irreps;  $G_i \neq 0$  for BLACK/spin irreps; and the BLACK values are  $G(\rho_2) = -6\phi$ ,  $G(\rho_4) = -3$ ,  $G(\rho_6) = +2$ ,  $G(\rho_8) = +6/\phi$ . In particular,  $|G(\rho_4)| = 3 = N_{\text{gen}}$ .*

**Theorem 16** (McKay mass correspondence). *On the McKay graph  $\hat{E}_8$  of  $2I$ , assigning fermion mass exponents as weighted graph distances from  $\rho_1$ , the main-chain weights are uniquely forced:*

$$\rho_1 \xrightarrow{N_{\text{gen}}} \rho_2 \xrightarrow{F(3)} \rho_4 \xrightarrow{b_1} \rho_7 \xrightarrow{0} \rho_8 \xrightarrow{a_1} \rho_9 \quad (178)$$

*i.e.  $[w_1, \dots, w_5] = [N_{\text{gen}}, F(3), b_1, 0, a_1]$  in all solutions.*

**Corollary 6** (Complementarity). *Three structures encode complementary aspects of fermion masses: (i)  $D_F$  (selection rules), (ii) McKay distance (hierarchy), (iii)  $\mathbb{Z}[\phi]$  lattice (absolute scale).*

**Fermionic Lagrangian.** The product spectral triple has  $\mathcal{H} = \mathcal{H}_M \otimes \mathcal{H}_F$  ( $\dim = 79,200$ ),  $D = D_M \otimes \mathbf{1}_{30} + \gamma_{\text{form}} \otimes D_F$ , and  $\gamma = \gamma_{\text{form}} \otimes \gamma_F$ , satisfying  $\{\gamma, D\} = 0$ . Physical masses are **Wilson lines** on the McKay tree:

$$m_f/m_e = \phi^{n_f}, \quad n_f = 5a + 6b. \quad (179)$$

The full fermionic Lagrangian, NCG comparison, and open problems are in Supplementary Material S17.