

Statistics with Spa OWS

Lecture 14

Julia Schroeder

Julia.schroeder@imperial.ac.uk

Outline

- Repeatability
- Pitfalls

Repeatability

- How consistent something is within a group, compared to the whole sample

Repeatability

- How consistent something is within a group, compared to the whole sample
- Sounds familiar?

Repeatability

- How consistent something is within a group, compared to the whole sample
- Sounds familiar?

$$R = \frac{\sigma_A^2}{\sigma_T^2}$$

Repeatability

- How consistent something is within a group, compared to the whole sample
- Sounds familiar?
- It's the ratio of among-group variance over the total variance

$$R = \frac{\sigma_A^2}{\sigma_T^2}$$

Repeatability

- How consistent something is within a group, compared to the whole sample
- Sounds familiar?
- It's the ratio of among-group variance over the total variance

$$R = \frac{\sigma_A^2}{\sigma_A^2 + \sigma_W^2}$$

- It's got lots of interesting uses in biology.

Observer repeatability

- Measuring tarsus consistently is not easy
- Some people measure it 3 times and take the mean
- Are observers consistent in their measures?



Individual behaviour - personality

- Do birds always behave the same way?
- Different from others?

Ecology

- If you calculate diversity indices, are the one in one region more alike than those between regions?

Ecology

- If you calculate diversity indices, are the one in one region more alike than those between regions?
- Is earthworm abundance consistent between day in the same plots?

Repeatability calculated from ANOVA output

$$R = \frac{\sigma_A^2}{\sigma_A^2 + \sigma_W^2}$$

$$\sigma_A^2 = \frac{MS_A - MS_W}{n_0}$$

$$\sigma_W^2 = MS_W$$

Repeatability calculated from ANOVA output

$$R = \frac{\sigma_A^2}{\sigma_A^2 + \sigma_W^2}$$

$$\sigma_A^2 = \frac{MS_A - MS_W}{n_0}$$

$$\sigma_W^2 = MS_W$$

Repeatability calculated from ANOVA output

- N_0 not sample size
- Among-group variance difficult to compute (we've had trouble with SS before, remember!)

Repeatability calculated from ANOVA output

- N_0 not sample size
- Among-group variance difficult to compute (we've had trouble with SS before, remember!)

- Balanced dataset

$a_i = 1, 1, 2, 1, 1, 1$

$b_i = 3, 3, 4, 3, 3, 3$

$c_i = 5, 5, 4, 5, 5, 1$

Repeatability calculated from ANOVA output

- N_0 not sample size
 - Among-group variance difficult to compute (we've had trouble with SS before, remember!)
-
- Unbalanced dataset
 - Heterogeneous dataset

$a_i = 1, 1, 2, 1, 1, 1$

$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7$

$c_i = 5, 5$

Repeatability calculated from ANOVA output

- N_0 not sample size
 - Among-group variance difficult to compute (we've had trouble with SS before, remember!)
-
- Unbalanced dataset
 - Heterogeneous dataset
- $a_i = 1, 1, 2, 1, 1, 1$
- $b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7$
- $c_i = 5, 5$
- In ecology we have those often. In exact sciences less so (medicine, any planned experiments)

Repeatability calculated from ANOVA output

- N_0 not sample size
- Among-group variance difficult to compute (we've had trouble with SS before, remember!)

$$n_0 = \left[\frac{1}{a-1} \left[\sum_{i=1}^a n_i - \left(\frac{\sum_{i=1}^a n_i^2}{\sum_{i=1}^a n_i} \right) \right] \right]$$

a = number of groups

n_i = sample size in each group i

Repeatability calculated from ANOVA output

$$n_0 = \left[\frac{1}{a-1} \left[\sum_{i=1}^a n_i - \left(\frac{\sum_{i=1}^a n_i^2}{\sum_{i=1}^a n_i} \right) \right] \right]$$

a = number of groups

n_i = sample size in each group i

$a_i = 1, 1, 2, 1, 1, 1$

$b_i = 3, 3, 4, 3, 3, 3$

$c_i = 5, 5, 4, 5, 5, 1$

$a_i = 1, 1, 2, 1, 1, 1$

$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7$

$c_i = 5, 5$

Repeatability calculated from ANOVA output

$$n_0 = \left[\frac{1}{a-1} \left[\sum_{i=1}^a n_i - \left(\frac{\sum_{i=1}^a n_i^2}{\sum_{i=1}^a n_i} \right) \right] \right]$$

a = number of groups

n_i = sample size in each group i

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3 \quad n = 6$$

$$c_i = 5, 5, 4, 5, 5, 1 \quad n = 6$$

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7 \quad n = 11$$

$$c_i = 5, 5 \quad n = 2$$

$$n_0 = \frac{(3 * 6) - \left(\frac{3 * 36}{3 * 6} \right)}{2}$$

Repeatability calculated from ANOVA output

$$n_0 = \left[\frac{1}{a-1} \left[\sum_{i=1}^a n_i - \left(\frac{\sum_{i=1}^a n_i^2}{\sum_{i=1}^a n_i} \right) \right] \right]$$

a = number of groups

n_i = sample size in each group i

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3 \quad n = 6$$

$$c_i = 5, 5, 4, 5, 5, 1 \quad n = 6$$

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7 \quad n = 11$$

$$c_i = 5, 5 \quad n = 2$$

$$n_0 = \frac{(3 * 6) - \left(\frac{3 * 36}{3 * 6} \right)}{2}$$
$$n_0 = 6$$

Repeatability calculated from ANOVA output

$$n_0 = \left[\frac{1}{a-1} \left[\sum_{i=1}^a n_i - \left(\frac{\sum_{i=1}^a n_i^2}{\sum_{i=1}^a n_i} \right) \right] \right]$$

a = number of groups

n_i = sample size in each group i

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3 \quad n = 6$$

$$c_i = 5, 5, 4, 5, 5, 1 \quad n = 6$$

$$n_0 = \frac{(3 * 6) - \left(\frac{3 * 36}{3 * 6} \right)}{2}$$

$$n_0 = 6$$

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7 \quad n = 11$$

$$c_i = 5, 5 \quad n = 2$$

$$n_0 = \frac{(6 + 11 + 2) - \left(\frac{36 + 121 + 4}{6 + 11 + 2} \right)}{2}$$

$$n_0 = 5.27$$

Repeatability calculated from ANOVA output

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3 \quad n = 6$$

$$c_i = 5, 5, 4, 5, 5, 1 \quad n = 6$$

$$n_0 = 6$$

$$a_i = 1, 1, 2, 1, 1, 1 \quad n = 6$$

$$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7 \quad n = 11$$

$$c_i = 5, 5 \quad n = 2$$

$$n_0 = 5.27$$

Repeatability calculated from ANOVA output

$a_i = 1, 1, 2, 1, 1, 1$ $n = 6$

$b_i = 3, 3, 4, 3, 3, 3$ $n = 6$

$c_i = 5, 5, 4, 5, 5, 1$ $n = 6$

$$n_0 = 6$$

$$R = \frac{\sigma_A^2}{\sigma_A^2 + \sigma_W^2}$$

$a_i = 1, 1, 2, 1, 1, 1$ $n = 6$

$b_i = 3, 3, 4, 3, 3, 3, 4, 2, 5, 5, 7$ $n = 11$

$c_i = 5, 5$ $n = 2$

$$n_0 = 5.27$$

$$\sigma_A^2 = \frac{MS_A - MS_W}{n_0}$$

$$\sigma_W^2 = MS_W$$

Learning aim

- Repeatability is intra-class correlation coefficient
- Ratio of how much variance is explained by groups
- N_0 is horrible

Do it NOW!

- HO 14
- Calculate the repeatability of body mass within individual birds.
- –Calculate the between–observer repeatability of your study group (A or B) of both wing, and tarsus! Explain what you did, and why. Justify!
- Report in writing, and on blackboard. Discuss!