

Statistics with Spa OWS

Lecture 10

Julia Schroeder

Julia.schroeder@imperial.ac.uk

Outline

- Hypothesis testing in linear models
- Interpretation (a bit)
- Standardizing
- Reporting

Hypothesis testing in linear models

- What do we actually test?



$$y_i = b_0 + b_1 x_i + \varepsilon_i$$

y i
5 1
3 2
6 3
10 4
4 5



$b_0 = ?$

$b_1 = ?$

x	i	ε	i
3	1	?	1
1	2	?	2
4	3	?	3
8	4	?	4
2	5	?	5



We tested whether the SS of a straight line with slope 0 ($b_1 = 0$) was different from the line we guesstimated, with a different slope



We tested whether the SS of a straight line with slope 0 ($b_1 = 0$) was different from the line we guesstimated, with a different slope

- Test if slope (b_1) estimate is different from 0
- T-test!

Hypothesis testing:

```
Call:
lm(formula = Mass ~ Tarsus, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.83246    0.98195     5.94 3.48e-09 ***
Tarsus        1.18466    0.05295    22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
F-statistic: 500.6 on 1 and 1642 DF,  p-value: < 2.2e-16
```

Tests null hypotheses that
 $b_0 = 0$
 $b_1 = 0$


Inference

Tests null hypotheses that

$$b_0 = 0$$

$$b_1 = 0$$

What does it mean if the null hypothesis is rejected?



Inference


Tests null hypotheses that

$$b_0 = 0$$

$$b_1 = 0$$

What does it mean if the null hypothesis is rejected?

Usually not much for b_0



Inference

```
Call:
lm(formula = Mass ~ Tarsus, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.83246    0.98195    5.94 3.48e-09 ***
Tarsus        1.18466    0.05295   22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
F-statistic: 500.6 on 1 and 1642 DF,  p-value: < 2.2e-16
```

Tests null hypotheses that
 $b_0 = 0$, rejected!
 $b_1 = 0$, rejected!

Inference

```
Call:
lm(formula = Mass ~ Tarsus, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.83246    0.98195    5.94 3.48e-09 ***
Tarsus       1.18466    0.05295   22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
F-statistic: 500.6 on 1 and 1642 DF,  p-value: < 2.2e-16
```

Tests null hypotheses that
 $b_0 = 0$, rejected!
 $b_1 = 0$, rejected!

The intercept (where Tarsus
= 0) is 5.8. A bird with no
tarsus would weigh 5.8g.

Inference

```
Call:
lm(formula = Mass ~ Tarsus, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.83246    0.98195    5.94 3.48e-09 ***
Tarsus       1.18466    0.05295   22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
F-statistic: 500.6 on 1 and 1642 DF,  p-value: < 2.2e-16
```

Tests null hypotheses that
 $b_0 = 0$, rejected!
 $b_1 = 0$, rejected!

The intercept (where Tarsus = 0) is 5.8. A bird with no tarsus would weigh 5.8g.

The slope is 5.8
For each mm longer tarsus, a sparrow weights 1.18g more.

Inference

Tests null hypotheses that


$$b_0 = 0$$

$$b_1 = 0$$

What does it mean if the null hypothesis is rejected?


Usually not much for b_0

b_1 is more interesting biologically speaking




Why standardize data?


- z-scores:
- Normal distributed
- Mean of 0
- Sd of 1




Why standardize data?




Why standardize data?




Why standardize data?




Why standardize data?



Why standardize data?



Why standardize data?



(black: mm, red: z-standardized)

Call:
lm(formula = Mass ~ zTarsus, data = d2)

Residuals:

Min	1Q	Median	3Q	Max
-7.7271	-1.2202	-0.1302	1.1592	7.5036

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	27.77895	0.04539	611.94	<2e-16 ***
zTarsus	1.01596	0.04541	22.37	<2e-16 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared: 0.2336, Adjusted R-squared: 0.2332
F-statistic: 500.6 on 1 and 1642 DF, p-value: < 2.2e-16

Why standardize data?

Intercept becomes
meaningful!

It's the mean of y!



(black: mm, red: z-standardized)

```
Call:
lm(formula = Mass ~ zTarsus, data = d2)
```

Residuals:

Min	1Q	Median	3Q	Max
-7.7271	-1.2202	-0.1302	1.1592	7.5036

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	27.77895	0.04539	611.94	<2e-16 ***
zTarsus	1.01596	0.04541	22.37	<2e-16 ***


Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared: 0.2336, Adjusted R-squared: 0.2332
F-statistic: 500.6 on 1 and 1642 DF, p-value: < 2.2e-16

Why standardize data?

Intercept becomes
meaningful!

It's the mean of y!



(black: mm, red: z-standardized)

```
Call:
lm(formula = Mass ~ zTarsus, data = d2)
```

Residuals:

Min	1Q	Median	3Q	Max
-7.7271	-1.2202	-0.1302	1.1592	7.5036

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	27.77895	0.04539	611.94	<2e-16 ***
zTarsus	1.01596	0.04541	22.37	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared: 0.2336, Adjusted R-squared: 0.2332
F-statistic: 500.6 on 1 and 1642 DF, p-value: < 2.2e-16

Why standardize data?

- Also, units. What if we'd measured tarsus in cm?
- `d2$Tarsuscm<-d2$Tarsus*10`

Why standardize data?

- Also, units. What if we'd measured tarsus in cm?
- `d2$Tarsuscm<-d2$Tarsus*10`
- `Model4<-lm(Mass~Tarsuscm,data=d2)`

Why standardize data?

- Also, units. What if we'd measured tarsus in cm?
- `d2$Tarsuscm<-d2$Tarsus*10`
- `Model4<-lm(Mass~Tarsuscm,data=d2)`

```
Call:
lm(formula = Mass ~ Tarsuscm, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.832455   0.981952   5.94 3.48e-09 ***
Tarsuscm     0.118466   0.005295  22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
```


Why standardize data?

- Also, units. What if we'd measured tarsus in cm?
- `d2$Tarsuscm<-d2$Tarsus*10`
- `Model4<-lm(Mass~Tarsuscm,data=d2)`

```
Call:
lm(formula = Mass ~ zTarsus, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 27.77895    0.04539   611.94  <2e-16 ***
zTarsus      1.01596    0.04541    22.37  <2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
F-statistic: 500.6 on 1 and 1642 DF,  p-value: < 2.2e-16
```

```
Call:
lm(formula = Mass ~ Tarsuscm, data = d2)

Residuals:
    Min       1Q   Median       3Q      Max
-7.7271 -1.2202 -0.1302  1.1592  7.5036

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  5.832455    0.981952     5.94 3.48e-09 ***
Tarsuscm      0.118466    0.005295    22.37 < 2e-16 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.841 on 1642 degrees of freedom
Multiple R-squared:  0.2336,    Adjusted R-squared:  0.2332
```

Interpretation of statistics – take home

- Always consider units!
 - Standardize to make intercept meaningful
 - T-tests test for null hypothesis that parameter estimates equal –
-
- Always think of the biological meaning in units!

How to report?

Methods:

To test whether heavier birds also had longer tarsi, I used a linear model, where body mass (g) was the response variable, and tarsus length (mm) the explanatory variable.

How to report?

Methods:

To test whether heavier birds also had longer tarsi, I used a linear model, where body mass (g) was the response variable, and tarsus length (mm) the explanatory variable. I z-standardized tarsus length to a mean of 0 and SD of one, so that the intercept could be interpreted as the mean of body mass.

How to report?

Methods:

To test whether heavier birds also had longer tarsi, I used a linear model, where body mass (g) was the response variable, and tarsus length (mm) the explanatory variable. I z-standardized tarsus length to a mean of 0 and SD of one, so that the intercept could be interpreted as the mean of body mass. Following the analysis, I used visual inspection of residual plots to assess that the assumption that the residuals follow a normal distribution was not violated.

How to report?

Methods:

To test whether heavier birds also had longer tarsi, I used a linear model, where body mass (g) was the response variable, and tarsus length (mm) the explanatory variable. I z-standardized tarsus length to a mean of 0 and SD of one, so that the intercept could be interpreted as the mean of body mass. Following the analysis, I used visual inspection of residual plots to assess that the assumption that the residuals follow a normal distribution was not violated. I report results as statistically significant if p equals or is smaller than 0.05. I used R version 3.3 (R Core Team 2015) for all analysis and plotting.

How to report?

Results:

I used data from 1646 observations.

How to report?

Results:

I used data from 1646 observations.
The sparrows weighed on average
27.78 g (SD 2.10, range: 18.60–36.20).

How to report?

Results:

I used data from 1646 observations.
The sparrows weighed on average
27.78 g (SD 2.10, range: 18.60–36.20).
The tarsi of the sparrows were on
average 18.5mm long (SD 0.86, range:
15.00–21.10).

How to report?

Results:

I used data from 1646 observations.
The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20).
The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1).

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

Table 1: Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

- Clear headers!
- Headers of columns center aligned!

Table 1: Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report?

- Clear headers!
- Headers of columns center aligned!
- Always report all variables in the model. Don't be selective!
- Right align first column!

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

Table 1: Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

- Clear headers!
- Headers of columns center aligned!
- Always report all variables in the model. Don't be selective!
- Right align first column!
- Left align numbers!
- Round to two digits (or what else is biologically reasonable)!
- Always report parameter estimates (b)

Table 1: Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

- Clear headers!
- Headers of columns center aligned!
- Always report all variables in the model. Don't be selective!
- Right align first column!
- Left align numbers!
- Round to two digits (or what else is biologically reasonable)!
- Always report parameter estimates (b)
- Report meaningful p-values (no e-16 bs or similar)!

Table 1: Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report?

Results:

I used data from 1646 observations. The sparrows weighed on average 27.78 g (SD 2.10, range: 18.60–36.20). The tarsi of the sparrows were on average 18.5mm long (SD 0.86, range: 15.00–21.10). I found a positive, statistically significant association between mass and tarsus (Table 1). An increase in 1SD Tarsus length meant an increase of 1.02g in body mass.

- Clear headers!
- Headers of columns center aligned!
- Always report all variables in the model. Don't be selective!
- Right align first column!
- Left align numbers!
- Round to two digits (or what else is biologically reasonable)!
- Always report parameter estimates (b)
- Report meaningful p-values (no e-16 bs or similar)!
- Table legend goes on top of tables (below figures)
- Table legend should be self-explanatory without referring to text!

Table 1. Results from a linear model explaining body mass of Lundy sparrows with tarsus length. Tarsus length was z-standardized. N = 1646.

Variable	<i>b</i>	<i>SE</i>	<i>t</i>	<i>p</i>
Intercept	27.78	0.05	611.94	<0.001
Tarsus	1.02	0.05	22.37	<0.001

How to report – take home

- Methods:
- Always describe *all analyses you present in results. Be precise and specific. Describe what's the response variable. Say what is the explanatory variable, and say WHY you used those. Give the units. Say when you standardize, and WHY.
- Justify, justify, justify.

How to report – take home

- Results:
- Start with describing the dataset.
- Sample size, mean, range, missing values ect.
- Explain results of each analysis
- Make nice tables and think before copy/pasting values from R!
- Legends need be self-explanatory!

HO 10– DO IT NOW!

- Do HO 10! Do the excersises on HO 10!
- Run diagnostics for a model with sex as explanatory variable. Interpret the plots.
- Run a linear model, where you test the hypothesis that sparrows with bigger bills can eat more. The prediction is that the larger the bill, the heavier the sparrow.
- Write a short (1A4) report on methods and results on this last model.
- Before you go into the linear model, you should first describe your data, say how many sparrows, how many females and males, **whether there is a difference in your response between the sexes. If that difference is meaningful, you should test the sexes separately.** Write this section as you would write it for a scientific article.