

The impact of temperature and absolute humidity on the coronavirus disease 2019 (COVID-19) outbreak - evidence from China

Peng Shi^{1#}, Yinqiao Dong^{2#}, Huanchang Yan^{3#}, Xiaoyang Li^{1#}, Chenkai Zhao^{1#}, Wei Liu¹, Miao He¹,
Shixing Tang³, Shuhua Xi^{1*}

¹ Department of Environmental Health, School of Public Health, China Medical University, Shenyang,
China

² Department of Occupational Health, School of Public Health, China Medical University, Shenyang,
China

³ Department of Epidemiology, School of Public Health, Southern Medical University, Guangzhou,
China

[#] Peng Shi, Yinqiao Dong, Huanchang Yan, Xiaoyang Li, and Chenkai Zhao contributed equally to this
work.

* Correspondence to: Prof Shuhua Xi, Department of Environmental Health, School of Public
Health, China Medical University, No. 77 Puhe Road, Shenyang North New Area, Shenyang,
Liaoning Province, China. Email address: shxi@cmu.edu.cn

21 **What is already known on this topic**

- 22 ● Many infectious diseases present an environmental pattern in their incidence.
- 23 ● Environmental factors, such as climate and weather condition, could drive the space and time
- 24 correlations of infectious diseases, including influenza.
- 25 ● Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can be transmitted through
- 26 aerosols, large droplets, or direct contact with secretions (or fomites) as influenza virus can.
- 27 ● Little is known about environmental pattern in COVID-19 incidence.

28 **What this study adds**

- 29 ● The significant association between COVID-19 daily incidence and temperature was
- 30 confirmed, using 3 methods, based on the data on COVID-19 and weather from 31
- 31 provincial-level regions in mainland China.
- 32 ● Environmental factors were considered on the basis of SEIR model, and a modified
- 33 susceptible-exposed-infectious-recovered (M-SEIR) model was developed.
- 34 ● Simulations of the COVID-19 outbreak in Wuhan presented similar effects of temperature on
- 35 incidence as the incidence decrease with the increase of temperature.

36 **ABSTRACT**

37 **OBJECTIVE**

38 To investigate the impact of temperature and absolute humidity on the coronavirus disease 2019
39 (COVID-19) outbreak.

40 **DESIGN**

41 Ecological study.

42 **SETTING**

43 31 provincial-level regions in mainland China.

44 **MAIN OUTCOME MEASURES**

45 Data on COVID-19 incidence and climate between Jan 20 and Feb 29, 2020.

46 **RESULTS**

47 The number of new confirm COVID-19 cases in mainland China peaked on Feb 1, 2020. COVID-19
48 daily incidence were lowest at -10 °C and highest at 10 °C, while the maximum incidence was
49 observed at the absolute humidity of approximately 7 g/m³. COVID-19 incidence changed with
50 temperature as daily incidence decreased when the temperature rose. No significant association
51 between COVID-19 incidence and absolute humidity was observed in distributed lag nonlinear models.
52 Additionally, A modified susceptible-exposed-infectious-recovered (M-SEIR) model confirmed that
53 transmission rate decreased with the increase of temperature, leading to further decrease of infection
54 rate and outbreak scale.

55 **CONCLUSION**

56 Temperature is an environmental driver of the COVID-19 outbreak in China. Lower and higher
57 temperatures might be positive to decrease the COVID-19 incidence. M-SEIR models help to better
58 evaluate environmental and social impacts on COVID-19.

59

60 **Keywords:** COVID-19, Temperature, Humidity, Dynamic transmission model.

61

62 INTRODUCTION

63 In December 2019, an outbreak of novel coronavirus pneumonia occurred in Wuhan, Hubei Province,
64 China, and then were declared as an international public health emergency by the World Health
65 Organization (WHO) on January 30 2020. The disease was officially named as coronavirus disease
66 2019 (COVID-19) and the newly emerged virus was named as SARS-CoV-2 in February 2020.¹

67 Previous studies on early cases showed that the disease severity of COVID-19 with a 2.3%
68 case-fatality rate,² is much lower than Middle East Respiratory Syndrome (MERS) and Severe Acute
69 Respiratory Syndrome (SARS).³ However, as Li et al. reported,⁴ the number of COVID-19 cases
70 doubled every 7.4 days between December 2019 and January 2020, indicating COVID-19 might be
71 more infectious than SARS and MERS. In March 2020, the outbreak of COVID-19 was declared as a
72 global pandemic for the coronavirus rapidly expanded throughout China and to 116 other countries and
73 territories worldwide.

74 Many infectious diseases present an environmental pattern in their incidence. A few studies on
75 environmental issues, such as climate and weather condition, indicated that environmental factor could
76 drive the space and time correlations of infectious diseases.⁵⁻⁷ Based on analysis on climate predictors,
77 James D et al. found that humidity and temperature are optimal indicators in predicting influenza
78 epidemics in tropical regions.⁸ Temperate regions of the Northern and Southern Hemispheres are
79 characterized by highly synchronized annual influenza circulations during their winter months
80 respectively.^{5 7 8} In the United States, an epidemiological study indicated that lower specific humidity
81 is related to the occurrence of pandemic influenza, which is consistent with earlier finding in laboratory
82 experiments.⁹ Absolute humidity, the actual mass of water vapor, is identified as a main cause of

83 seasonal influenza epidemics.¹⁰ The influenza presents significant seasonal fluctuation in temperate
 84 monsoon climate regions as the absolute humidity varies greatly in summer and winter, which could
 85 help the multiplication of virus.

86 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can be transmitted through
 87 aerosols, large droplets, or direct contact with secretions (or fomites) as influenza virus can.¹¹ However,
 88 the environmental pattern remains to be elucidated in COVID-19 incidence. Based on dynamical
 89 equations, susceptible-exposed-infectious-recovered (SEIR) modeling has been developed and used to
 90 estimate key epidemic parameter to better characterize mechanism for the epidemic dynamics.¹²⁻¹⁴
 91 Therefore, we explored the association between daily incidence and climate conditions using locally
 92 weighted regression and smoothing scatterplot (LOESS) and distributed lag nonlinear models (DLNMs)
 93 based on the data on COVID-19 and weather from 31 provincial-level regions in mainland China,
 94 between Jan 20 and Feb 29, 2020. Furthermore, we took account of environmental factors on the basis
 95 of SEIR model, and developed a modified susceptible-exposed-infectious-recovered (M-SEIR) model
 96 to characterize the climate impacts on epidemic dynamics.

97

98 **METHODS**

99 **Study data**

100 Data on COVID-19, including the number of new confirmed and probable cases were obtained from
 101 the China National Health Commission (CNHC) using the CoV2019 package¹⁵
 102 (<http://www.nhc.gov.cn/>). COVID-19 data were collected among all of the 31 provincial-level regions
 103 in mainland China and Wuhan city, between Jan 20 and Feb 29, 2020. COVID-19 emerged in Wuhan

city at the end of 2019 and rapidly spread across mainland China. Thus, population dynamic factors, including birth rate and death rate, were not considered here. Finally, daily incidences among the 31 provincial-level regions and Wuhan city were calculated by dividing the number of new confirmed cases by the population size at the end of 2018 respectively, and was reported per 100,000 population.

Daily temperatures (T) and relative humidity (RH) of 344 cities of the corresponding period were collected from the meteorological authority in mainland China. Means of temperatures and absolute humidity were further calculated for every provincial-level region. The Clausius-Clapeyron relation equation was used to calculate absolute humidity (AH) as following:

$$AH = \frac{6.112 \times e^{\frac{17.67T}{T+243.5}} \times RH \times 2.1674}{273.15 + T}$$

Data on climate conditions and population were retrieved from official reports previously released in mainland China. Therefore, the ethical review was not required.

Statistical analysis

Trends of climate factors and daily COVID-19 incidence indicators, including the incidence and the common logarithm of numbers of newly confirmed cases (lgN), were analyzed with locally weighted regression and smoothing scatterplot (LOESS) in 31 provincial-level regions in mainland China from Jan 20 to Feb 29, 2020.

Developed on the definition of a cross-basis, DLNMs were used to infer the exposure-lag-response associations between climate factors and daily confirmed cases of COVID-19. DLNMs were constructed for mainland China outside of Hubei Province, Hubei Province outside of Wuhan city, and Wuhan city respectively. To induce the redundant analysis, temperature and absolute

124 humidity of mainland China were represented by data on the capital, Beijing. Additionally, temperature
125 and absolute humidity means of the sites in Hubei Province outside of Wuhan, were calculated as a
126 representative of Hubei Province data.

127 To better understand the potential environmental driver of COVID-19, we took account of
128 environmental factors on the basis of SEIR model and constructed the M-SEIR model to simulate the
129 COVID-19 outbreak dynamic in Wuhan after travel restriction was put into force. Further sensitivity
130 analysis was performed for quantitative risk assessment to evaluate the relationships between
131 environmental parameter and COVID-19 incidence.

132 The equations of M-SEIR model were given in the following:

$$\begin{aligned}\frac{dS(t)}{dt} &= \frac{-\beta_t S(t)I(t)}{N} \\ \frac{dE(t)}{dt} &= \frac{\beta_t S(t)I(t)}{N} - \sigma E(t) \\ \frac{dI(t)}{dt} &= \sigma E(t) - \gamma I(t) \\ \frac{dR(t)}{dt} &= \gamma I(t) \\ \beta_t &= \beta_1(1 + \beta_2 AH + \beta_3 T)\end{aligned}$$

133 where $S(t)$, $E(t)$, $I(t)$, and $R(t)$ were the number of susceptible, exposed, infectious, and removed
134 individuals at time t ; $\frac{1}{\sigma}$ and $\frac{1}{\gamma}$ were the mean latent and infectious period; β_t was a time dependent
135 rate of infectious contact; β_1, β_2 and β_3 were constant coefficients.

136 The simulations of COVID-19 dynamic and sensitivity analysis were conducted by using the
137 system dynamic section in AnyLogic software (version 8.5.2). The specific depict of parameter values

138 in modified model and basic model details were included in **Supplementary Table 1**.

139

140 **RESULTS**

141 80,981 cases of COVID-19 (cases of decrease in accounting not removed) was confirmed in 31
142 provincial-level regions in mainland China, between Jan 20 and Feb 29, 2020. Out of 80,981 cases,
143 68,034 (84.01%) were diagnosed in Hubei Province. Daily number of new confirmed cases and daily
144 incidence in mainland China were presented in **Figure 1** and **Supplementary Table 1**. Daily number of
145 cases peaked on Feb 12 and then it decreased, due to the adjustment in the diagnostic criteria of Hubei
146 Province. And the number of cases and the incidence in China (outside of Hubei Province) have begun
147 to decline early in Feb.

148 From Jan 20 to Feb 29, 2020, temperature and absolute humidity varied in 31 provincial-level
149 regions in mainland China (**Figure 2**). The highest temperature (26 °C) and absolute humidity (19.45
150 g/m³) were observed in Hainan Province and the lowest temperature (-22 °C) and absolute humidity
151 (0.54 g/m³) were observed in Jilin Province, which resulted from the geographical location. COVID-19
152 daily incidence indicators (daily incidence and lgN) increased as the absolute humidity rose and
153 declined slightly when absolute humidity reached approximately 7 g/m³ (**Figure 3**). Analysis for Hubei
154 Province (outside of Wuhan) and Wuhan showed highly similar results (**Supplementary Figure 1 and**
155 **Supplementary Figure 2**). Differences lay in the fact that cases clinically diagnosed without nucleic
156 acid testing had been counted as confirmed cases in Hubei Province since Feb 12, which might increase
157 potential bias in the model.

158 Associations between temperature and COVID-19 relative risk (RR) in mainland China (outside

of Hubei Province), Hubei Province (outside of Wuhan) and Wuhan were presented as three-dimensional plots in **Figure 4**, compared with a reference value of 0 °C. The plots showed significant effect on COVID-19 incidence of temperature. In mainland China (outside of Hubei Province), the highest RR (1.71, 95% CI: 1.28-2.27) was observed at a cold temperature (-6 °C), suggesting the COVID-19 incidence were most likely to increase at -6 °C. The RR of 0.59 (95% CI: 0.44-0.78) at 6 °C rose to 1.06 (95% CI: 0.96-1.18) when temperature dropped to -6 °C. However, no statistical significance was found in lag-specific relative risk at lag 2 to lag 4, suggesting no delayed effect at any temperature. For example, the relative risk maintained at lag 2-4, as lag-specific RR was 1.14 (95% CI: 0.90-1.44) at lag 2 and 1.03 (95% CI: 0.86-1.33) at lag 4 when temperature was -6 °C. In Hubei Province (outside of Wuhan), RR was significantly higher at 8°C (RR 1.22, 95% CI: 1.07-1.38) and 10 °C (RR 1.92, 95% CI: 1.21-3.03) in lag 0. Conversely, lag-specific RR ranged from lag 0 to lag 7 at 8-10 °C, suggesting positive delayed effect on decreasing COVID-19 incidence during the condition. In Wuhan city, the highest RR 1.04 (95% CI: 0.92-1.17) without significance was observed at approximately 9 °C. However, the incidence was more likely to decrease with immediate and delayed effect at a lower or higher temperature than 9 °C. For example, RR was in a range of 0.64 (95% CI: 0.46-0.87) to 0.88 (95% CI: 0.73-0.99) at lag 0 to 5 days when the temperature was 4 °C and similar results were observed when the temperature was 16 °C.

Overall pictures of the effect of absolute humidity on incidence in mainland China (outside of Hubei Province), Hubei Province (outside of Wuhan) and Wuhan were presented in **Figure 4**, showing 3-D graphs of COVID-19 relative risk (RR) along absolute humidity and lags compared with a reference value of 7.5 g/m³. The plots showed inconsistent effect of absolute humidity on COVID-19 incidence. In mainland China, immediate effect on COVID-19 incidence was strongest at absolute

181 humidity of 4 g/m³ (RR: 1.13, 95%CI:1.02-1.27), indicating COVID-19 incidence was more likely to
 182 increase during the condition. When absolute humidity rose to 5 g/m³, values of lag-specific RR were
 183 in range of 0.60 (95% CI: 0.36-0.99) to 0.62 (95% CI: 0.41-0.93) at lag 3 to lag 5 (**Supplementary**
 184 **Figure 3**), indicating a strong delayed effect on COVID-19 incidence at absolute humidity of 5 g/m³. In
 185 Hubei Province, immediate effect on reducing COVID-19 incidence was observed when absolute
 186 humidity ranged from 4.5 g/m³ (RR 0.40, 95% CI: 0.19-0.84) to 5.5g/m³ (RR 0.65, 95% CI: 0.44-0.96)
 187 (**Supplementary Figure 4**). However, no significant difference was observed in absolute humidity in
 188 Wuhan city (**Supplementary Figure 5**).

189 Considering the environmental impacts, we constructed the M-SEIR model to simulate the
 190 dynamic of COVID-19 by using the system dynamic section in AnyLogic software. SEIR dynamic
 191 transmission model compartmentalized the population into four states including susceptible, exposed,
 192 infected, and recovered, and further analyzed the relationships and interconnection using stock and set
 193 parameters, flows and table function (**Figure 5A; Supplemental video 1**). We set the initial values of
 194 the parameter and incorporated the temperature index in Wuhan city from Jan 20 and Feb 29, 2020,
 195 into the modified SEIR model. **Supplemental table 3** presented the comparison of modified SEIR
 196 model in our study and classic SEIR models in similar studies. The four curves were stratified by types
 197 of state, and showed a similar pattern: the population size increased early in epidemic and then
 198 decreased as the period ends (e.g., due to recovery). As the M-SEIR model predicted, the number of
 199 infections would peak around Mar 5, reaching the inflection point, and the COVID-19 outbreak in
 200 Wuhan would be expected to end by late April (**Figure 5B; Supplemental video 1**). Furthermore, a
 201 sensitivity analysis on the transmission rate adjusted by temperature indicated high stability of our
 202 M-SEIR model (**Figure 5C; Supplemental video 2**). We set the transmission rate from 0 to 1 with a

203 step of to 0.1, and conducted the simulations to reduce the bias involving in the model, parameters, and
204 functional relationships. Finally, we found that the transmission rate decreased with the increase of
205 temperature, leading to the decrease of infection rate and outbreak size.

206

207 **DISCUSSION**

208 We inferred that the number of new confirm COVID-19 cases in mainland China peaked on Feb 1,
209 2020. COVID-19 daily incidence were lowest at -10 °C and highest at 10 °C, while the maximum
210 incidence was observed at the absolute humidity of approximately 7 g/m³. We found significant
211 association between temperature and COVID-19 daily incidence due to the immediate and delayed
212 effect observed using DLNMs. As predicted in M-SEIR model, the COVID-19 outbreak would peak
213 around March 5, 2020 and end in late April in Wuhan. Additionally, we found that transmission rate
214 decreased with the increase of temperature, leading to further decrease of infection rate and outbreak
215 size. Therefore, temperature drive the space and time correlations of COVID-19, and it can be used as
216 an optimal predictor.

217 In this study, we inferred the significant association between temperature and COVID-19 daily
218 incidence using LOESS, DLNMs and M-SEIR model, suggesting that temperature plays an important
219 role in the outbreak of COVID-19 and can be used in predicting the potential spread of COVID-19.
220 Lower and higher temperatures may be positive to decrease the COVID-19 incidence, which help to
221 shed new light on the environmental drivers of COVID-19 in China. Our results are in line with the
222 findings in SARS. Based on data on SARS and climate in 4 cities, Tan et al. found that temperature is a
223 powerful indicator for SARS-CoV transmission, in which the risk of increased daily incidence differed

224 between the effects of high and low temperatures.¹⁶ Additionally, Lowen's laboratory work evidenced
225 that temperature affect the virus spread of aerosol using a guinea pig model.¹⁷ However, the
226 temperature DLNM in Hubei Province, showed different patterns from those in mainland China and
227 Wuhan city, as COVID-19 relative risk rose at a moderate temperature.

228 In our analysis, we failed to observe a significant relationship between absolute humidity and
229 COVID-19 incidence based on the data of mainland China. However, absolute humidity has been
230 reported as a strong correlation with influenza epidemic, due to the seasonal pattern that influences the
231 multiplications and spread of influenza.^{9 18 19} In another study on MERS, caused a lethality of more
232 than 35%, confirmed that the activity of MERS-CoV in droplet or aerosol, decreases significantly as
233 absolute humidity increases though the mechanism is not yet clear.²⁰ The difference between our study
234 and previous finding may result from the fact that absolute humidity remained stable in a region during
235 a very limited period. Additionally, rapid and strong actions taken by the government could biased our
236 study. Despite of the negative consequence in our study, further studies on absolute humidity are
237 required to perform.

238 Combination of infectious disease dynamics model and environmental patterns is required to
239 better explain the relationship between environmental factors and infection.²¹ Dynamic transmission
240 model was usually performed to predict the genesis and development trend of infectious diseases as
241 well as to evaluate the effect of intervention but few dynamic transmission models included
242 environmental factors for the increasing uncertainty. However, to reveal the dynamic of an infectious
243 disease, it would be much better to take account of environmental impact on the basis of dynamic
244 transmission model.^{22 23}

Environmental factors, characterized by lag effects and threshold effects, can target at two objects, host and virus, during infectious disease outbreak. On one hand, human activity patterns and immunity can be influenced by environmental factors. But the effect caused by environmental condition was limited during the COVID-19 outbreak, due to the absence of extreme weather and specific immunity for a newly emerging virus. On the other hand, environmental impacts on the SARS-CoV-2 are more significant than the host population because the transmission and virulence of the virus varies in different conditions. Finally, environmental impacts on transmission of virus should be characterized in the dynamic model, because infectiousness estimated in the traditional dynamic model is actually a confounding effect with environmental effect. It is necessary to take account of environmental issues on the basis of dynamic transmission model so that the impacts could be isolated and qualified. A dynamic model is not only compatible with the infectious disease transmission mode for virus itself, but also can be well coupled with surveillance data on environmental issues.²⁴ Consequently, we constructed a M-SEIR model to correct the potential deviation of temperature to simulate the dynamic epidemic of COVID-19. The M-SEIR model predicted that the outbreak would reach its peak reach an inflection point around March 5, 2020, which is consistent with the actual situation based on data released by the NHC.²⁵⁻²⁹ And it is expected that the COVID-19 outbreak in Wuhan would end in late April. In addition, we conducted a sensitivity analysis on the temperature-adjusted transmission rate. Finally, we found transmission rate decreased with the increase of temperature, leading to further decrease of infection rate and epidemic size.

Our analysis is subject to limitations. First, the COVID-19 dynamics are determined by multiple factors, including virus, climate, socio-economic development, population mobility, population immunity, and urbanization. However, not all those factors were considered in this study. Second, the

parameters of M-SEIR models were optimized, based on the previous analysis which might be biased by the lack of official data and the adjustment of diagnostic criteria in the outbreak. Third, it's an ecological analysis in very short period so that we cannot avoid the bias caused by other ecological factors changed over time.

Conclusions and public health implications

Temperature is an environmental driver of the COVID-19 outbreak in China. Lower and higher temperatures might be positive to decrease the COVID-19 incidence. As predicted in M-SEIR model, the COVID-19 outbreak would peak around March 5, 2020 and end in late April in Wuhan. Modified-SEIR models help to better evaluate and identify national and international prevention and intervention targeted COVID-19. The COVID-19 outbreak would not last for a long period of time with the increase of temperature, but the scale of the outbreak would be influenced by the measures taken among countries.

Contributions: All authors contributed to the study concept and design. PS, HY, and XL wrote the paper. PS and YD collected the data. PS, YD and CZ analyzed the data. PS, HY, ST, WL, MH, and SX reviewed and revised the manuscript before submission. All authors approved the final submitted version.

Funding: The work was supported by the National Key Research and Development Program of China to Prof Shuhua Xi (2018YFC1801204). The funders had no role in design and conduct of the study;

287 collection, management, analysis, and interpretation of the data; preparation, review, and approval of
288 the manuscript; or the decision to submit the manuscript for publication.

289 **Competing interests:** All authors have completed the ICMJE uniform disclosure form at
290 http://www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and
291 declare: no support from any organization for the submitted work other than those described above; no
292 financial relationships with any organizations that might have an interest in the submitted work in the
293 previous three years; no other relationships or activities that could appear to have influenced the
294 submitted work.

295 **Transparency declaration:** The manuscript's guarantor affirms that the manuscript is an honest,
296 accurate, and transparent account of the study being reported; that no important aspects of the study
297 have been omitted; and that any discrepancies from the study have been explained.

298 **Data sharing:** No additional data available.

299

300 **References**

301 1. Zi Yue Zu M, Meng Di Jiang M, Peng Peng Xu M, Wen Chen M, Qian Qian Ni P, Guang Ming Lu M,
302 et al. Coronavirus Disease 2019 (COVID-19): A Perspective from China.
303 doi:10.1148/radiol.2020200490

304 2. She J, Jiang J, Ye L, Hu L, Bai C, Song Y. 2019 novel coronavirus of pneumonia in Wuhan, China:
305 emerging attack and management strategies. *Clin Transl Med* 2020;9(1):19.
306 doi:10.1186/s40169-020-00271-z

- 307 3. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019
308 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for
309 Disease Control and Prevention. *JAMA* 2020. doi:10.1001/jama.2020.2648
- 310 4. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China,
311 of Novel Coronavirus-Infected Pneumonia. *The New England journal of medicine* 2020.
312 doi:10.1056/NEJMoa2001316
- 313 5. Bedford T, Riley S, Barr IG, Broor S, Chadha M, Cox NJ, et al. Global circulation patterns of seasonal
314 influenza viruses vary with antigenic drift, *Nature*.2015. doi:10.1038/nature14460
- 315 6. Lemaitre J, Pasetto D, Perez-Saez J, Sciarra C, Wamala JF, Rinaldo A. Rainfall as a driver of
316 epidemic cholera: Comparative model assessments of the effect of intra-seasonal precipitation events.
317 *Acta Trop*. 2019;190:235–243. doi:10.1016/j.actatropica.2018.11.013
- 318 7. Sooryanarain H, Elankumaran S. Environmental role in influenza virus outbreaks. *Annu Rev Anim*
319 *Biosci* 2015;3:347-73. doi:10.1146/annurev-animal-022114-111017
- 320 8. Tamerius JD, Shaman J, Alonso WJ, Bloom-Feshbach K, Uejio CK, Comrie A, et al. Environmental
321 predictors of seasonal influenza epidemics across temperate and tropical climates. *PLoS Pathog*
322 2013;9(3):e1003194. doi:10.1371/journal.ppat.1003194
- 323 9. Dalziel BD, Kissler S, Gog JR, Viboud C, Bjornstad ON, Metcalf C, et al. Urbanization and humidity
324 shape the intensity of influenza epidemics in U.S. cities. *Science* 2018;362(6410):75-79.
325 doi:10.1126/science.aat6030
- 326 10. Thai PQ, Choisy M, Duong TN, Thiem VD, Yen NT, Hien NT, et al. Seasonality of absolute humidity

327 explains seasonality of influenza-like illness in Vietnam. *Epidemics* 2015;13:65-73.
328 doi:10.1016/j.epidem.2015.06.002

329 11. Li Y, Huang X, Yu IT, Wong TW, Qian H. Role of air distribution in SARS transmission during the
330 largest nosocomial outbreak in Hong Kong. *Indoor Air* 2005;15(2):83-95.
331 doi:10.1111/j.1600-0668.2004.00317.x

332 12. Chanprasopchai P, Pongsumpun P, Tang IM. Effect of Rainfall for the Dynamical Transmission
333 Model of the Dengue Disease in Thailand. *Comput Math Methods Med* 2017;2017:2541862.
334 doi:10.1155/2017/2541862

335 13. Liu T, Zhu G, He J, Song T, Zhang M, Lin H, et al. Early rigorous control interventions can largely
336 reduce dengue outbreak magnitude: experience from Chaozhou, China. *BMC Public Health*
337 2017;18(1):90. doi:10.1186/s12889-017-4616-x

338 14. Niakan Kalhori SR, Ghazisaeedi M, Azizi R, Naserpour A. Studying the influence of mass media and
339 environmental factors on influenza virus transmission in the US Midwest. *Public Health* 2019;170:17-22.
340 doi:10.1016/j.puhe.2019.02.006

341 15. Tianzhi Wu, Xijin Ge, Guangchuang Yu, Erqiang Hu. Open-source analytics tools for studying the
342 COVID-19 coronavirus outbreak. *medRxiv*. doi:10.1101/2020.02.25.20027433

343 16. Tan J. An initial investigation of the association between the SARS outbreak and weather: with the
344 view of the environmental temperature and its variation. *Journal of Epidemiology & Community Health*
345 2005;59(3):186-192. doi:10.1136/jech.2004.020180

346 17. Lowen AC, Mubareka S, Steel J, Palese P. Influenza virus transmission is dependent on relative

347 humidity and temperature. *PLoS Pathog* 2007;3(10):1470-6. doi:10.1371/journal.ppat.0030151

348 18. Shaman J, Goldstein E, Lipsitch M. Absolute humidity and pandemic versus epidemic influenza. *Am*

349 *J Epidemiol* 2011;173(2):127-35. doi:10.1093/aje/kwq347

350 19. Shoji M, Katayama K, Sano K. Absolute humidity as a deterministic factor affecting seasonal

351 influenza epidemics in Japan. *Tohoku J Exp Med* 2011;224(4):251-6. doi:10.1620/tjem.224.251

352 20. van Doremalen N, Bushmaker T, Munster VJ. Stability of Middle East respiratory syndrome

353 coronavirus (MERS-CoV) under different environmental conditions. *Euro Surveill* 2013;18(38).

354 doi:10.2807/1560-7917.es2013.18.38.20590

355 21. He D, Ionides EL, King AA. Plug-and-play inference for disease dynamics: measles in large and

356 small populations as a case study. *J R Soc Interface* 2010;7(43):271-83. doi:10.1098/rsif.2009.0151

357 22. Martinez PP, King AA, Yunus M, Faruque AS, Pascual M. Differential and enhanced response to

358 climate forcing in diarrheal disease due to rotavirus across a megacity of the developing world. *Proc Natl*

359 *Acad Sci U S A* 2016;113(15):4092-7. doi:10.1073/pnas.1518977113

360 23. Bakker KM, Martinez-Bakker ME, Helm B, Stevenson TJ. Digital epidemiology reveals global

361 childhood disease seasonality and the effects of immunization. *Proc Natl Acad Sci U S A*

362 2016;113(24):6689-94. doi:10.1073/pnas.1523941113

363 24. Pitzer VE, Viboud C, Alonso WJ, Wilcox T, Metcalf CJ, Steiner CA, et al. Environmental drivers of

364 the spatiotemporal dynamics of respiratory syncytial virus in the United States. *PLoS Pathog*

365 2015;11(1):e1004591. doi:10.1371/journal.ppat.1004591

366 25. Tang Z, Li X, Li H. Prediction of New Coronavirus Infection Based on a Modified SEIR Model.

367 *medRxiv* 2020:2020.03.03.20030858. doi:10.1101/2020.03.03.20030858

368 26. Shi P, Cao S, Feng P. SEIR Transmission dynamics model of 2019 nCoV coronavirus with
369 considering the weak infectious ability and changes in latency duration. *medRxiv*
370 2020:2020.02.16.20023655. doi:10.1101/2020.02.16.20023655

371 27. Pan J, Yao Y, Liu Z, Li M, Wang Y, Dong W, et al. Effectiveness of control strategies for Coronavirus
372 Disease 2019: a SEIR dynamic modeling study. *medRxiv* 2020:2020.02.19.20025387.
373 doi:10.1101/2020.02.19.20025387

374 28. Hong N, He J, Ma Y, Jiang H, Han L, Su L, et al. Evaluating the secondary transmission pattern and
375 epidemic prediction of the COVID-19 in metropolitan areas of China. *medRxiv*
376 2020:2020.03.06.20032177. doi:10.1101/2020.03.06.20032177

377 29. Zhang L, Wan K, Chen J, Lu C, Dong L, Wu Z. When will the battle against novel coronavirus end in
378 Wuhan: a SEIR modeling analysis. *medRxiv* 2020:2020.02.16.20023804.
379 doi:10.1101/2020.02.16.20023804

380

381 **Figure Legends**

382 **Figure 1** Daily number of new confirmed cases of COVID-19 in mainland China between Jan 20 and
383 Feb 29, 2020.

384 **Figure 2** Between Jan 20 and Feb 29, 2020, temperature values (left columns) and absolute humidity
385 values (right columns) in 31 provincial-level regions in mainland China.

386 **Figure 3** COVID-19 daily incidence indicators (daily incidence and $\lg N$) and the expected values
387 based on the temperature and absolute humidity in mainland China (outside of Hubei Province) from
388 Jan 20 to Feb 29, respectively. The black line represents the expected value of a daily incidence and
389 $\lg N$ based upon a LOESS regression for all days of available estimates. LOESS, locally weighted
390 regression and smoothing scatterplots.

391 **Figure 4** 3-D plot of RR of COVID-19 along climate factors (temperature and absolute humidity) and
392 lags in mainland China (outside of Hubei Province), Hubei Province (outside of Wuhan), and Wuhan
393 city.

394 **Figure 5 COVID-19 dynamic trends and sensitivity analysis using M-SEIR model in Wuhan. (A)**

395 The over-all structure of M-SEIR model constructed by using the system dynamic section in AnyLogic
396 software. **(B)** The snapshot represents the different population proportion of susceptible, exposed,

397 infected, and recovered states under the specific time-point and forecasts the trend of the COVID-19

398 outbreak in Wuhan city. **(C)** Sensitivity analysis under different temperature scenarios in Wuhan city.

399 As the temperature-corrected transmission index rises, the peak of the curve increased under different

400 times gradually. M-SEIR model, modified susceptible-exposed-infectious-recovered model; TI, the

401 temperature-corrected transmission index (i.e. The transmission rate for susceptible to exposed, β_t).

Figures




Figure 1 Daily number of new confirmed cases of COVID-19 in mainland China between Jan 20 and Feb 29, 2020.




Figure 2 Between Jan 20 and Feb 29, 2020, temperature values (left columns) and absolute humidity values (right columns) in 31 provincial-level regions in mainland China.




Figure 3 COVID-19 daily incidence indicators (daily incidence and lgN) and the expected values based on the temperature and absolute humidity in mainland China (outside of Hubei Province) from Jan 20 to Feb 29, respectively. The black line represents the expected value of a daily incidence and lgN based upon a LOESS regression for all days of available estimates. LOESS, locally weighted regression and smoothing scatterplots.




Figure 4 3-D plot of RR of COVID-19 along climate factors (temperature and absolute humidity) and lags in mainland China (outside of Hubei Province), Hubei Province (outside of Wuhan), and Wuhan city.




Figure 5 COVID-19 dynamic trends and sensitivity analysis using M-SEIR model in Wuhan. (A)

The over-all structure of M-SEIR model constructed by using the system dynamic section in AnyLogic software. **(B)** The snapshot represents the different population proportion of susceptible, exposed,

infected, and recovered states under the specific time-point and forecasts the trend of the COVID-19

outbreak in Wuhan city. (C) Sensitivity analysis under different temperature scenarios in Wuhan city.

As the temperature-corrected transmission index rises, the peak of the curve increased under different

times gradually. M-SEIR model, modified susceptible-exposed-infectious-recovered model; TI, the

temperature-corrected transmission index (i.e. The transmission rate for susceptible to exposed, β_t).