

characterize many carnivorous non-mammalian synapsids²⁴. The molariform teeth at the back of the dentition of *Repenomamus* are small with blunt crowns; they probably played a minor role in food processing. Although mammals are considered definitive chewers within amniotes²⁵, the dental morphology and large pieces of prey in the stomach of *Repenomamus* suggest that chewing as a derived feature in mammals was probably not present in *Repenomamus*.

It is not easy to assess whether *Repenomamus* was a predator or scavenger. Scavengers are relatively rare among mammals—among extant carnivorous mammals, only two species of hyenas are habitual scavengers^{12,26}. Compared to their hunting cousins, these hyenas have smaller second upper incisors and less jaw muscle leverage, which probably reflect their inability to capture and handle live prey. In contrast, the enlarged incisors and strong jaw muscles of *Repenomamus* are well shaped for catching prey, favouring it as a predator rather than a scavenger.

For fossil mammals, body size is one of the most important factors influencing life history strategy²⁷. Early mammals or their close relatives, such as morganocodontids and kuehneotheriids in the Late Triassic to Early Jurassic periods, were small and considered to be nocturnal insectivores^{2,3}; the same is true of most later Mesozoic mammals²⁸ (Fig. 4). The reason for the very small size of Mesozoic mammals is uncertain⁵, but it has often been hypothesized that well-established larger (and presumably diurnal) reptilian carnivores and herbivores, particularly dinosaurs, prevented mammals from invading those niches²⁹. *Repenomamus* extend significantly the upper limit of body size of Mesozoic mammals (Fig. 4) and are actually larger than several small dinosaurs, particularly dromaeosaurid dinosaurs, from the same fauna¹¹. Larger animals can live longer and move faster, but they also need a larger food supply and broader home range³⁰. Judging from their body size, *R. giganticus* could feed on larger prey and forage a wider area for food. These large Mesozoic mammals were probably carnivores that competed with dinosaurs for food and territory. □

Received 29 May; accepted 8 October 2004; doi:10.1038/nature03102.

- Bakker, R. T. Dinosaur physiology and the origin of mammals. *Evolution* **25**, 636–658 (1971).
- Hopson, J. A. Endothermy, small size and the origin of mammalian reproduction. *Am. Nat.* **107**, 446–452 (1973).
- Jerison, H. J. *Evolution of the Brain and Intelligence* (Academic, New York, 1973).
- Crompton, A. W., Taylor, C. R. & Jagger, J. A. Evolution of homeothermy in mammals. *Nature* **272**, 333–336 (1978).
- Lillegraven, J. A. in *Mesozoic Mammals: The First Two-thirds of Mammalian History* (eds Lillegraven, J. A., Kielan-Jaworowska, Z. & Clemens, W. A.) 1–6 (Univ. California Press, Berkeley, 1979).
- Li, J.-L., Wang, Y., Wang, Y.-Q. & Li, C.-K. A new family of primitive mammal from the Mesozoic of western Liaoning, China [in Chinese]. *Chin. Sci. Bull.* **45**, 2545–2549 (2000).
- Wang, Y.-Q., Hu, Y.-M., Meng, J. & Li, C.-K. An ossified Meckel's cartilage in two Cretaceous mammals and origin of the mammalian middle ear. *Science* **294**, 357–361 (2001).
- Zhou, Z.-H., Barrett, P. M. & Hilton, J. An exceptionally preserved Lower Cretaceous ecosystem. *Nature* **421**, 807–814 (2003).
- Li, C.-K., Wang, Y.-Q., Hu, Y.-M. & Meng, J. A new species of *Gobiconodon* from the Jehol Biota and its implication to the age of the fauna. *Chin. Sci. Bull.* **48**, 177–182 (2003).
- Wang, S.-S., Wang, Y.-Q., Hu, H.-G. & Li, H.-M. The existing time of Sihetun vertebrate in western Liaoning, China—Evidence from U-Pb dating of zircon [in Chinese with English abstract]. *Chin. Sci. Bull.* **46**, 779–782 (2001).
- Xu, X. & Wang, X.-L. A new dromaeosaur (Dinosauria: Theropoda) from the Early Cretaceous Yixian Formation of western Liaoning. *Vert. Palaeontol.* **42**, 111–119 (2004).
- Nowak, R. M. *Walker's Mammals of the World* 6th edn (Johns Hopkins Univ. Press, Baltimore, 1999).
- Silva, M. & Downing, J. A. *CRC Handbook of Mammalian Body Mass* (CRC Press, Boca Raton, 1995).
- Jenkins, F. A. Jr Limb posture and locomotion in the Virginia opossum (*Didelphis marsupialis*) and in other non-cursorial mammals. *J. Zool.* **165**, 303–315 (1971).
- Fischer, M. S., Schilling, N., Schmidt, M., Dieter Haarmann, D. & Witte, H. Basic limb kinematics of small therian mammals. *J. Exp. Biol.* **205**, 1315–1338 (2002).
- Wilson, R. W. Late Cretaceous (Fox Hills) multituberculates from the Red Owl Local Fauna of western South Dakota. *Dakoterra* **3**, 118–132 (1987).
- Clemens, W. A., Wilson, G. P. & Molnar, R. E. An enigmatic (synapsid?) tooth from the Early Cretaceous of New South Wales, Australia. *J. Vert. Paleontol.* **23**, 232–237 (2003).
- Jenkins, F. A. Jr & Schaff, C. R. The Early Cretaceous mammal *Gobiconodon* (Mammalia, Triconodontata) from the Cloverly Formation in Montana. *J. Vert. Paleontol.* **8**, 1–24 (1988).
- Rougier, G. W. *Vincelestes neuquenianus Bonaparte (Mammalia, Theria), un Primitivo Mammifero del Cretacico Inferior de la Cuenca Neuquina* PhD Thesis, Univ. Nacional de Buenos Aires, Buenos Aires (1993).
- Alexander, R. McN., Jayes, A. S., Maloij, G. M. O. & Wathuta, E. M. Allometry of limb bones of mammals from shrews (*Sorex*) to elephant (*Loxodonta*). *J. Zool.* **189**, 305–314 (1979).
- Van Valkenburgh, B. in *Body Size in Mammalian Paleobiology: Estimation and Biological Implication*

- (eds Damuth, J. & MacFadden, B. J.) 181–205 (Cambridge Univ. Press, Cambridge, 1990).
- Coombs, W. P. Jr Juvenile specimens of the ornithischian dinosaur *Psittacosaurus*. *Palaeontology* **25**, 89–107 (1982).
- Carbone, C., Mace, G. M., Roberts, S. C. & Macdonald, D. W. Energetic constraints on the diet of terrestrial carnivores. *Nature* **402**, 286–288 (1999).
- Van Valkenburgh, B. & Jenkins, I. Evolutionary patterns in the history of Permo-Triassic and Cenozoic synapsid predators. *Paleontol. Soc. Pap.* **8**, 267–288 (2002).
- Reilly, S. M., McBrayer, L. D. & White, T. D. Prey processing in amniotes: biomechanical and behavioral patterns of food reduction. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **128**, 397–415 (2001).
- Van Valkenburgh, B., Sacco, T. & Wang, X.-M. Pack hunting in Miocene Borophagine dogs: Evidence from craniodental morphology and body size. *Bull. Am. Mus. Nat. Hist.* **279**, 147–162 (2004).
- Damuth, J. & MacFadden, B. J. in *Body Size in Mammalian Paleobiology: Estimation and Biological Implication* (eds Damuth, J. & MacFadden, B. J.) 1–10 (Cambridge Univ. Press, Cambridge, 1990).
- Lillegraven, J. A., Kielan-Jaworowska, Z. & Clemens, W. A. (eds) *Mesozoic Mammals: The First Two-thirds of Mammalian History* (Univ. California Press, Berkeley, 1979).
- Crompton, A. W. in *Comparative Physiology: Primitive Mammals* (eds Schmidt-Nielsen, K., Bolis, L. & Taylor, C. R.) 1–12 (Cambridge Univ. Press, Cambridge, 1980).
- Eisenberg, J. F. in *Body Size in Mammalian Paleobiology: Estimation and Biological Implication* (eds Damuth, J. & MacFadden, B. J.) 25–38 (Cambridge Univ. Press, Cambridge, 1990).

Supplementary Information accompanies the paper on www.nature.com/nature.

Acknowledgements We thank M.-M. Chang, Z.-H. Zhou, X.-L. Wang, X. Xu, F.-C. Zhang, Y. Wang, F. Jin and J.-Y. Zhang for help coordinating the research and fieldwork; X. Xu, X.-L. Wang, F.-C. Zhang, Z.-H. Zhou, and M. Norell for discussions on the research subject, and S.-H. Xie, S.-J. Li and A. Davidson for specimen preparation. This work was supported by funding from the Chinese Ministry of Science and Technology, the National Natural Science Foundation of China and the Chinese Academy of Sciences. Y.H. is also supported by a fellowship from the American Museum of Natural History, through the City University of New York.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to Y. H. (yhu@amnh.org).

The simplicity of metazoan cell lineages

Ricardo B. R. Azevedo¹, Rolf Lohaus^{1,2}, Volker Braun², Markus Gumbel², Muralikrishna Umamaheshwar¹, Paul-Michael Agapow³, Wouter Houthoofd⁴, Ute Platzer², Gaëtan Borgonie⁴, Hans-Peter Meinzer² & Armand M. Leroi³

¹Department of Biology and Biochemistry, University of Houston, Houston, Texas 77204-5001, USA

²Division of Medical and Biological Informatics, German Cancer Research Center, D-69120, Heidelberg, Germany

³Department of Biology, Imperial College, Silwood Park, Ascot SL5 7PY, UK

⁴Department of Biology, Ghent University, B-9000 Ghent, Belgium

Developmental processes are thought to be highly complex, but there have been few attempts to measure and compare such complexity across different groups of organisms^{1–5}. Here we introduce a measure of biological complexity based on the similarity between developmental and computer programs^{6–9}. We define the algorithmic complexity of a cell lineage as the length of the shortest description of the lineage based on its constituent sublineages^{9–13}. We then use this measure to estimate the complexity of the embryonic lineages of four metazoan species from two different phyla. We find that these cell lineages are significantly simpler than would be expected by chance. Furthermore, evolutionary simulations show that the complexity of the embryonic lineages surveyed is near that of the simplest lineages evolvable, assuming strong developmental constraints on the spatial positions of cells and stabilizing selection on cell number. We propose that selection for decreased complexity has played a major role in moulding metazoan cell lineages.

Biological systems are obviously complex in both structure and