



POLITECNICO DI MILANO
SCHOOL OF INDUSTRIAL AND INFORMATION ENGINEERING
ACADEMIC YEAR 2024-2025

Real and Functional Analysis

Professor: Gianmaria Verzini

Last updated: December 20, 2024

**This document is intended for educational purposes only.
These are unreviewed notes and may contain errors.**

Made by Roberto Benatuil Valera

Contents

1	Set Theory	3
1.1	Basic notions	3
1.2	Families of subsets	4
1.3	Characteristic functions	6
1.4	Equivalence relations and Quotient sets	6
2	Measure Theory	8
2.1	Measure spaces	8
2.2	Generation of a σ -algebra	9
2.3	Borel sets	10
2.4	Measures	11
2.4.1	Properties of measures	12
2.5	Sets of measure zero, negligible sets, complete measures	15
2.6	Towards the Lebesgue measure	16
2.6.1	Carathéodory's criterion	18
2.7	Lebesgue measure	20
2.8	Measurable functions	25
2.8.1	Operations on measurable functions	27
2.8.2	Properties holding almost everywhere	29
2.8.3	Simple functions	30
2.9	Lebesgue integral	31
2.9.1	Integral of non-negative simple functions	31
2.9.2	Integral of non-negative measurable functions	33

Chapter 1

Set Theory

1.1 Basic notions

Definition 1.1.1. Let X, Y be sets. We say:

- X, Y are **equipotent** if there exists a bijection $f : X \rightarrow Y$.
- X has a **cardinality greater or equal** to Y if there exists an surjection $f : X \rightarrow Y$.
- X is **finite** if it is equipotent to $\{1, 2, \dots, n\}$ for some $n \in \mathbb{N}$. X is infinite otherwise.

Remark: X is infinite \iff it is equipotent to a proper subset of itself.

E.g.: The set of natural numbers \mathbb{N} is infinite. In fact, the set of even natural numbers $E = \{2, 4, 6, \dots\} \subset \mathbb{N}$ is equipotent to \mathbb{N} , as we can define the bijection $f : \mathbb{N} \rightarrow E$ as $f(n) = 2n$.

Definition 1.1.2. Let X be an infinite set. We say X is **countable** if it is equipotent to \mathbb{N} . X is **uncountable** otherwise, in which case it is **more than countable**.

Definition 1.1.3. X has the **cardinality of the continuum** if it is equipotent to $[0, 1] \subset \mathbb{R}$. Any such set is uncountable.

E.g.: We have that:

- $\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ are countable.
- $\mathbb{R}, \mathbb{R}^n, (0, 1), [0, 1]$ are uncountable.
- Countable union of countable sets is countable.

1.2 Families of subsets

Let X be a set. The “Power set” of X is the set of all subsets of X , denoted by $\mathcal{P}(X)$.

$$\mathcal{P}(X) = \{E : E \subseteq X\}$$

Note that $\mathcal{P}(X)$ has always a cardinality greater than X . For example, if $X = \mathbb{N}$, then $\mathcal{P}(X)$ has the cardinality of the continuum.

Definition 1.2.1. Let X be a set. A **family of subsets** of X is a set E such that:

$$E \subseteq \mathcal{P}(X)$$

We usually denote $E = \{E_i\}_{i \in I}$, where I is an index set.

Definition 1.2.2. Let $E = \{E_i\}_{i \in I}$ be a family of subsets of X . We define:

- The **union** of E as:

$$\bigcup_{i \in I} E_i = \{x \in X : x \in E_i \text{ for some } i \in I\}$$

- The **intersection** of E as:

$$\bigcap_{i \in I} E_i = \{x \in X : x \in E_i \text{ for all } i \in I\}$$

Definition 1.2.3. Let $E = \{E_i\}_{i \in I}$ be a family of subsets of X . We say F is **pairwise disjoint** if:

$$E_i \cap E_j = \emptyset \quad \forall i, j \in I, i \neq j$$

Definition 1.2.4. We say that the family $E = \{E_i\}_{i \in I}$ of subsets of X is a **covering** of X if:

$$X = \bigcup_{i \in I} E_i$$

Any subfamily of E , $E' = \{E_i\}_{i \in I'}$ is a **subcovering** of X if it is a covering of X itself.

E.g.: Let $X = \mathbb{R}$. We define:

$$\mathcal{T} = \{E \subset X : E \text{ is open}\}$$

We say that \mathcal{T} is the standard topology of X . More generally, this can be done in

“metric spaces” (X, d) .

Properties of \mathcal{T} (open sets):

- $\emptyset, X \in \mathcal{T}$.
- Finite intersection of elements in \mathcal{T} is in \mathcal{T} .
- Arbitrary union of elements in \mathcal{T} is in \mathcal{T} .

We can also define **sequences of sets**. Let X be a set. A sequence of sets in X is a family of sets $\{E_n\}_{n \in \mathbb{N}}$.

Definition 1.2.5. Let X be a set. A sequence of sets $\{E_n\}_{n \in \mathbb{N}}$ is said to be:

- **Increasing** if:

$$E_n \subseteq E_{n+1} \quad \forall n \in \mathbb{N}$$

It is denoted by $\{E_n\} \uparrow$.

- **Decreasing** if:

$$E_n \supseteq E_{n+1} \quad \forall n \in \mathbb{N}$$

It is denoted by $\{E_n\} \downarrow$.

Let now $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{P}(X)$ be a sequence of sets in X :

Definition 1.2.6. We define the following:

- The **limit superior** of $\{E_n\}$ as:

$$\limsup_{n \rightarrow \infty} E_n = \bigcap_{n \in \mathbb{N}} \bigcup_{k \geq n} E_k$$

- The **limit inferior** of $\{E_n\}$ as:

$$\liminf_{n \rightarrow \infty} E_n = \bigcup_{n \in \mathbb{N}} \bigcap_{k \geq n} E_k$$

- If the limit superior and limit inferior are equal, we say that

$$\lim_{n \rightarrow \infty} E_n = \limsup_{n \rightarrow \infty} E_n = \liminf_{n \rightarrow \infty} E_n$$

Exercise: Let X be a set and $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{P}(X)$ be a sequence of sets in X . Prove that:

$$(i) \quad \{E_n\} \uparrow \Rightarrow \lim_{n \rightarrow \infty} E_n = \bigcup_{n \in \mathbb{N}} E_n \quad (ii) \quad \{E_n\} \downarrow \Rightarrow \lim_{n \rightarrow \infty} E_n = \bigcap_{n \in \mathbb{N}} E_n$$

1.3 Characteristic functions

Definition 1.3.1. Let X be a set and $E \subseteq X$. The **characteristic function** of E is the function $\mathbb{1}_E : X \rightarrow \{0, 1\}$ defined as:

$$\mathbb{1}_E(x) = \begin{cases} 1 & \text{if } x \in E \\ 0 & \text{if } x \notin E \end{cases}$$

This function is also called the **indicator function** of E .

Remark: Let $E, F \subseteq X$. We have that:

- $\mathbb{1}_{E \cap F} = \mathbb{1}_E \cdot \mathbb{1}_F$.
- $\mathbb{1}_{E \cup F} = \mathbb{1}_E + \mathbb{1}_F - \mathbb{1}_{E \cap F}$.
- $\mathbb{1}_{E^c} = 1 - \mathbb{1}_E$.
- $\mathbb{1}_{\limsup_{n \rightarrow \infty} E_n} = \limsup_{n \rightarrow \infty} \mathbb{1}_{E_n}$.
- $\mathbb{1}_{\liminf_{n \rightarrow \infty} E_n} = \liminf_{n \rightarrow \infty} \mathbb{1}_{E_n}$.

1.4 Equivalence relations and Quotient sets

Definition 1.4.1. A relation R on a set X is a subset of $X \times X$. For any $x, y \in X$, we say that x is related to y if $(x, y) \in R$. We denote this as xRy .

Definition 1.4.2. A relation R on a set X is an **equivalence relation** if it satisfies:

- **Reflexivity:**

$$xRx \quad \forall x \in X$$

- **Symmetry:**

$$xRy \Rightarrow yRx \quad \forall x, y \in X$$

- **Transitivity:**

$$xRy, yRz \Rightarrow xRz \quad \forall x, y, z \in X$$

Every equivalence relation on X induces a partition of X . We define the **equivalence class** of $x \in X$ as:

$$[x] = \{y \in X : xRy\}$$

The set of all equivalence classes is called the **quotient set** of X by R , denoted by X/R .

$$X/R = \{[x] : x \in X\}$$

E.g.: Let $X = \mathbb{Z} \times \mathbb{Z}_0$ such that $\mathbb{Z}_0 = \mathbb{Z} \setminus \{0\}$. We define the relation R on X as:

$$(a, b)R(c, d) \iff ad = bc$$

We can prove that R is an equivalence relation. The equivalence classes are:

$$[(a, b)] = \{(c, d) \in X : ad = bc\}$$

Notice that:

$$[(a, b)] = \{(a, b), (2a, 2b), (3a, 3b), \dots\}$$

If we denote a class $[(a, b)]$ as $[a/b]$, then we have that:

$$X/R = \{[a/b] : a, b \in \mathbb{Z}_0\} = \mathbb{Q}$$

Chapter 2

Measure Theory

2.1 Measure spaces

Definition 2.1.1. Let X be a non-empty set. A family $\mathcal{M} \subseteq \mathcal{P}(X)$ is a **σ -algebra** if:

- (i) $\emptyset \in \mathcal{M}$.
- (ii) $E \in \mathcal{M} \implies E^c \in \mathcal{M}$.
- (iii) $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M} \implies \bigcup_{n \in \mathbb{N}} E_n \in \mathcal{M}$.

If instead of (iii) we have that $E_1, E_2 \in \mathcal{M} \implies E_1 \cup E_2 \in \mathcal{M}$, then \mathcal{M} is called an **algebra**.

Remark: If \mathcal{M} is a σ -algebra, then we say that (X, \mathcal{M}) is a **measurable space**. Any set $E \in \mathcal{M}$ is called a **measurable set**.

E.g.: Let $X \neq \emptyset$. Then:

- $\mathcal{P}(X)$ is a σ -algebra.
- $\{\emptyset, X\}$ is a σ -algebra.
- $\{\emptyset, E, E^c, X\}$ is a σ -algebra for any $E \subseteq X$.
- $X = \mathbb{R}$, $\mathcal{M} = \{E \subseteq \mathbb{R} : E \text{ is open}\}$ is NOT a σ -algebra.

Properties 2.1.1. Let (X, \mathcal{M}) be a measurable space. Then:

- (i) $X = \emptyset^c \in \mathcal{M}$
- (ii) \mathcal{M} is also an algebra. Indeed, if $\{E_1, E_2\} \subseteq \mathcal{M}$, $E_n = \emptyset \forall n \geq 3$, then $E_1 \cup E_2 = \bigcup_{n \in \mathbb{N}} E_n \in \mathcal{M}$.
- (iii) $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M} \implies \bigcap_n E_n \in \mathcal{M}$.
- (iv) $E, F \in \mathcal{M} \implies E \setminus F \in \mathcal{M}$
- (v) $\Omega \subseteq X$. Then, the **restriction** of \mathcal{M} to Ω is:

$$\mathcal{M}|_{\Omega} := \{F \subseteq \Omega : F = E \cap \Omega, E \in \mathcal{M}\}$$

Then, $(\Omega, \mathcal{M}|_{\Omega})$ is a measurable space.

2.2 Generation of a σ -algebra

Theorem 2.2.1. Take any family $\mathcal{A} \subseteq \mathcal{P}(X)$. Then, it is well-defined the σ -algebra generated by \mathcal{A} , denoted by $\sigma_0(\mathcal{A})$, as the smallest σ -algebra containing \mathcal{A} . It is characterized by:

- (i) $\sigma_0(\mathcal{A})$ is a σ -algebra.
- (ii) $\mathcal{A} \subseteq \sigma_0(\mathcal{A})$.
- (iii) If \mathcal{M} is a σ -algebra and $\mathcal{A} \subseteq \mathcal{M}$, then $\sigma_0(\mathcal{A}) \subseteq \mathcal{M}$.

Sketch of proof. Define $V = \{\mathcal{M} \subseteq \mathcal{P}(X) : \mathcal{M} \text{ is } \sigma\text{-algebra, } \mathcal{A} \subseteq \mathcal{M}\}$. Notice that $V \neq \emptyset$ because $\mathcal{P}(X) \in V$. Then, define:

$$\sigma_0(\mathcal{A}) = \bigcap_{\mathcal{M} \in V} \mathcal{M}$$

Then, $\sigma_0(\mathcal{A})$ is a σ -algebra as it satisfies the properties of a σ -algebra, denoted in definition 2.1.1. ■

Remark: This is relevant. Often, to check that a σ -algebra has certain properties, it is enough to check the property on a set of generators.

2.3 Borel sets

Take (X, d) as a metric space, so that open sets are defined. Recall that:

$$\mathcal{T} = \{E \subseteq X : E \text{ is open}\}$$

Definition 2.3.1. The σ -algebra generated by \mathcal{T} is called the **Borel σ -algebra** of X , denoted by:

$$\mathcal{B}(X) := \sigma_0(\mathcal{T})$$

Any set $E \in \mathcal{B}(X)$ is a **Borel set**.

Remark: The following are Borel sets:

- Open sets
- Closed sets
- Countable intersections of open sets (G_δ -sets)
- Countable unions of closed sets (F_σ -sets)

We will deal with:

$$X = \mathbb{R} = (-\infty, \infty)$$

but also:

$$X = \overline{\mathbb{R}} = [-\infty, \infty] = \mathbb{R} \cup \{-\infty, \infty\}$$

Let us define the arithmetic operations on $\overline{\mathbb{R}}$. Let $a \in \mathbb{R}$:

- $a \pm \infty = \pm\infty$
- $a > 0 : a \cdot \pm\infty = \pm\infty$
- $a < 0 : a \cdot \pm\infty = \mp\infty$
- $a = 0 : 0 \cdot \pm\infty = 0$
- $\infty - \infty, \infty/\infty, 0/0$ are not defined.

Also, the open intervals in $\overline{\mathbb{R}}$ are the following:

- (a, b) , with $a, b \in \mathbb{R}$
- $[-\infty, b)$
- $(a, \infty]$

Remark: We have that:

$$\begin{aligned}\mathcal{B}(\mathbb{R}) &:= \sigma_0(\{\text{open sets}\}) \\ &= \sigma_0(\{(a, b) : a < b\}) \\ &= \sigma_0(\{[a, b] : a < b\}) \\ &= \sigma_0(\{(a, \infty) : a \in \mathbb{R}\})\end{aligned}$$

$$\begin{aligned}\mathcal{B}(\overline{\mathbb{R}}) &:= \sigma_0(\{\text{open sets}\}) \\ &= \sigma_0(\{(a, \infty] : a \in \mathbb{R}\})\end{aligned}$$

$$\mathcal{B}(\mathbb{R}^N) = \sigma_0(\{\text{open rectangles}\})$$

2.4 Measures

Let (X, \mathcal{M}) be a measurable space.

Definition 2.4.1. A function $\mu : \mathcal{M} \rightarrow [0, \infty]$ is a (positive) **measure** on \mathcal{M} if:

- (i) $\mu(\emptyset) = 0$
- (ii) $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M}, \text{ disjoint} \implies \mu(\bigcup_{n \in \mathbb{N}} E_n) = \sum_{n \in \mathbb{N}} \mu(E_n)$

Note: To avoid nonsenses, we always assume that $\exists E \in \mathcal{M} \text{ s.t. } \mu(E) < \infty$

Terminology: Let X, \mathcal{M}, μ defined as above:

- (X, \mathcal{M}, μ) is a **measure space**.
- If $\mu(X) = 1$, then (X, \mathcal{M}, μ) is a **probability space** and μ is a **probability measure**.

Definition 2.4.2. A measure μ is:

1. **Finite** if $\mu(X) < \infty$
2. **σ -finite** if $\exists \{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M} \text{ s.t.}$

$$\mu(E_n) < \infty \quad \forall n \in \mathbb{N} \quad \wedge \quad X = \bigcup_{n \in \mathbb{N}} E_n$$

E.g.: Some examples of measures are:

1. (Trivial measure): For any (X, \mathcal{M}) , define μ as $\mu(E) = 0 \quad \forall E \in \mathcal{M}$
2. (Counting measure): For any (X, \mathcal{M}) , typically $\mathcal{M} = \mathcal{P}(X)$, define $\mu_{\#}$ as:

$$\mu_{\#}(E) = \begin{cases} \#\{\text{elements of } E\} & E \text{ finite} \\ \infty & \text{otherwise} \end{cases}$$

3. (Dirac measure): For any (X, \mathcal{M}) , pick $x_0 \in X$. Then, define δ_{x_0} as:

$$\delta_{x_0}(E) = \begin{cases} 1 & x_0 \in E \\ 0 & x_0 \notin E \end{cases}$$

2.4.1 Properties of measures

Theorem 2.4.1 (Basic properties). *Let (X, \mathcal{M}, μ) be a measure space. Then:*

- (i) μ is finitely additive: $E, F \in \mathcal{M}, E \cap F = \emptyset \implies \mu(E \cup F) = \mu(E) + \mu(F)$
- (ii) (Monotonicity): $E, F \in \mathcal{M}, E \subseteq F \implies \mu(E) \leq \mu(F)$
- (iii) (Excision property): $E, F \in \mathcal{M}, E \subseteq F, \mu(E) < \infty \implies \mu(F \setminus E) = \mu(F) - \mu(E)$

Proof. The proof is straightforward:

- (i) Let $E, F \in \mathcal{M}, E \cap F = \emptyset$. Then:

$$\mu(E \cup F) = \mu(E) + \mu(F)$$

Proof. Obvious, using $E_n = \emptyset$ for $n \geq 3$. ■

- (ii) Let $E, F \in \mathcal{M}, E \subseteq F$. Then:

$$\mu(E) \leq \mu(F)$$

Proof. Let $F = E \cup (F \setminus E)$. Then:

$$\mu(F) = \mu(E) + \mu(F \setminus E) \geq \mu(E)$$
■

- (iii) Let $E, F \in \mathcal{M}, E \subseteq F, \mu(E) < \infty$. Then:

$$\mu(F \setminus E) = \mu(F) - \mu(E)$$

Proof. Let $F = E \cup (F \setminus E)$. Then:

$$\mu(F) = \mu(E) + \mu(F \setminus E)$$

This concludes the proof. ■

Theorem 2.4.2 (Continuity among monotone sequences). *Let (X, \mathcal{M}, μ) be a measure space. Let $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M}$ be a sequence of measurable sets. Then:*

(i) *If $\{E_n\} \uparrow$, $E := \lim_n E_n = \bigcup_n E_n$, then:*

$$\mu(E) = \lim_n \mu(E_n)$$

(ii) *If $\{E_n\} \downarrow$, $E := \lim_n E_n = \bigcap_n E_n$, and $\mu(E_1) < \infty$, then:*

$$\mu(E) = \lim_n \mu(E_n)$$

Proof. The proof goes as follows:

(i) If $\mu(E_n) = \infty$ for some n , then the proof is trivial. Otherwise, let $F_1 = E_1$ and $F_n = E_n \setminus E_{n-1}$ for $n \geq 2$. Then, we can check that:

- $F_n \in \mathcal{M}, \forall n \in \mathbb{N}$
- $\{F_n\}$ is a disjoint sequence.
- $E_n = \bigcup_{k=1}^n F_k$
- Because of the disjointness, we have that:

$$\mu(E_n) = \sum_{k=1}^n \mu(F_k)$$

Then, we have that:

$$\begin{aligned} \mu(E) &= \mu\left(\lim_n E_n\right) = \mu\left(\bigcup_{n=1}^{\infty} E_n\right) = \\ &= \mu\left(\bigcup_{n=1}^{\infty} F_n\right) = \sum_{n=1}^{\infty} \mu(F_n) = \\ &= \sum_{n=1}^{\infty} (\mu(E_n) - \mu(E_{n-1})) = \lim_n \mu(E_n) \end{aligned}$$

(ii) Define $G_n = E_1 \setminus E_n$. Then, check that:

- $G_n \in \mathcal{M}, \forall n \in \mathbb{N}$
- $\{G_n\} \uparrow$

By the previous result, we have that:

$$\mu \left(\bigcup_{n=1}^{\infty} G_n \right) = \lim_n \mu(G_n)$$

Then, on the right-hand side:

$$\begin{aligned} \lim_n \mu(G_n) &= \lim_n \mu(E_1 \setminus E_n) = \\ &= \mu(E_1) - \lim_n \mu(E_n) \end{aligned}$$

On the left-hand side:

$$\begin{aligned} \mu \left(\bigcup_{n=1}^{\infty} G_n \right) &= \mu \left(\bigcup_{n=1}^{\infty} (E_1 \setminus E_n) \right) = \\ &= \mu \left(E_1 \setminus \bigcap_{n=1}^{\infty} E_n \right) = \\ &= \mu(E_1) - \mu(E) \end{aligned}$$

Finally, we have that:

$$\mu(E_1) - \mu(E) = \mu(E_1) - \lim_n \mu(E_n)$$

And because $\mu(E_1) < \infty$, we have that:

$$\mu(E) = \lim_n \mu(E_n)$$

■

Remark: In (ii), the condition $\mu(E_1) < \infty$ is essential. Consider $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu_{\#})$. Then, define the following sequence:

$$E_n = \{n, n+1, n+2, \dots\}$$

Note that $E_n \subseteq E_{n-1}$. Also, note that for any $n \in \mathbb{N}$, we have that:

$$\mu_{\#}(E_n) = \infty$$

Then, we have that:

$$\mu_{\#} \left(\bigcap_n E_n \right) = \mu_{\#}(\emptyset) = 0$$

But:

$$\lim_n \mu_{\#}(E_n) = \infty$$

This shows that the condition $\mu(E_1) < \infty$ is essential.

Theorem 2.4.3 (σ -subadditivity). *Let (X, \mathcal{M}, μ) be a measure space. Let $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{M}$ be a sequence of measurable sets. Then:*

$$\mu \left(\bigcup_n E_n \right) \leq \sum_n \mu(E_n)$$

Proof. Let $F_1 = E_1$ and $F_n = E_n \setminus (\bigcup_{k=1}^{n-1} E_k)$ for $n \geq 2$. Then, we have that:

- $F_n \in \mathcal{M}, \forall n \in \mathbb{N}$
- $F_n \subseteq E_n, \forall n \in \mathbb{N}$
- $\{F_n\}$ is a disjoint sequence.
- $\bigcup_n E_n = \bigcup_n F_n$

Then, we have that:

$$\begin{aligned} \mu \left(\bigcup_n E_n \right) &= \mu \left(\bigcup_n F_n \right) = \\ &= \sum_n \mu(F_n) \leq \sum_n \mu(E_n) \end{aligned}$$

■

2.5 Sets of measure zero, negligible sets, complete measures

Definition 2.5.1. Let (X, \mathcal{M}, μ) be a measure space. Then:

1. A set $E \in \mathcal{M}$ is a **set of measure zero** if $\mu(E) = 0$.
2. A set $F \in X$ (not necessarily measurable) is a **negligible set** if $\exists E \in \mathcal{M}$ s.t. $F \subseteq E$ and E is a set of measure zero.

Definition 2.5.2. Let (X, \mathcal{M}, μ) be a measure space. Then, we say that μ is a **complete measure** (alternatively, that (X, \mathcal{M}, μ) is a **complete measure space**) all negligible sets are measurable.

Remark (Completion of a measure space): A measure space (X, \mathcal{M}, μ) may not be complete. However, we can define the following:

$$\overline{\mathcal{M}} = \{E \subseteq X : \exists F_1, F_2 \in \mathcal{M} : F_1 \subseteq E \subseteq F_2, \mu(F_2 \setminus F_1) = 0\}$$

One can show that $\overline{\mathcal{M}}$ is a σ -algebra, and that $\mathcal{M} \subseteq \overline{\mathcal{M}}$. Moreover, if E, F_1, F_2 are as above, define:

$$\overline{\mu}(E) = \mu(F_1) = \mu(F_2)$$

One can check that $(X, \overline{\mathcal{M}}, \overline{\mu})$ is a complete measure space.

2.6 Towards the Lebesgue measure

We would like to define a measure λ with $X = \mathbb{R}$ (or $X = \mathbb{R}^N$) s.t. $\forall a < b$:

- $\lambda((a, b)) = b - a$ (**length of the interval**)
- $\forall E, \lambda(E + x) = \lambda(E)$ (**translation invariance**)

In principle, we would like to define it in $\mathcal{P}(\mathbb{R})$. Such a measure should satisfy $\lambda(\{a\}) = 0$.

Theorem 2.6.1 (Ulam). *The only measure on $\mathcal{P}(\mathbb{R})$ that satisfies $\lambda(\{a\}) = 0 \forall a \in \mathbb{R}$ is the trivial measure.*

Therefore, we need to choose an $\mathcal{M} \subsetneq \mathcal{P}(\mathbb{R})$. We can construct one as follows:

- Starting family with a “measure”, e.g., $\mathcal{T} = \{(a, b) : a < b\}$ and $f((a, b)) = b - a$.
- Construct an “outer measure” μ^* on $\mathcal{P}(\mathbb{R})$.
- Restrict μ^* to a well-chosen σ -algebra $\mathcal{M} \subsetneq \mathcal{P}(\mathbb{R})$.

Definition 2.6.1. Let X be a set. An **outer measure** μ^* on X is a function

$$\mu^* : \mathcal{P}(X) \rightarrow [0, \infty]$$

such that:

1. $\mu^*(\emptyset) = 0$
2. (Monotonicity) $E \subseteq F \subseteq X \implies \mu^*(E) \leq \mu^*(F)$
3. (σ -subadditivity) $\{E_n\}_{n \in \mathbb{N}} \subseteq \mathcal{P}(X) \implies \mu^*\left(\bigcup_{n \in \mathbb{N}} E_n\right) \leq \sum_{n \in \mathbb{N}} \mu^*(E_n)$

Remark: Any measure μ is an outer measure. However, the converse is not true.

Proposition 2.6.2. Let $\mathcal{E} \subseteq \mathcal{P}(X)$, $f : \mathcal{E} \rightarrow [0, \infty]$. Assume that $\emptyset, X \in \mathcal{E}$, $f(\emptyset) = 0$. Then, $\forall E \subseteq X$ define:

$$\mu^*(E) = \inf \left\{ \sum_{n=1}^{\infty} f(A_n) : A_n \in \mathcal{E}, E \subseteq \bigcup_n A_n \right\}$$

Then, μ^* is an outer measure.

Proof. The proof is omitted. ■

Remark: In this generality, if $E \in \mathcal{E}$, then $f(E)$ and $\mu^*(E)$ may not be equal. We can only guarantee that $\mu^*(E) \leq f(E)$.

E.g.: There are some important examples:

- $X = \mathbb{R}$, $\mathcal{E} = \{(a, b) : a, b \in \overline{\mathbb{R}}, a \leq b\}$

$$f((a, b)) = \text{length}((a, b)) = b - a$$

- $X = \mathbb{R}^N$, $\mathcal{E} = \{(a_1, b_1) \times \dots \times (a_N, b_N) : a_i, b_i \in \overline{\mathbb{R}}, a_i \leq b_i\}$

$$f(\underline{a}, \underline{b}) = \text{volume}(\underline{a}, \underline{b}) = \prod_{i=1}^N (b_i - a_i)$$

In both cases, the outer measure μ^* is called the **Lebesgue outer measure**. We will denote it by λ^* (or λ_N^* in the second case). Note that in this case, $\lambda^*(E) = f(E)$ for any $E \in \mathcal{E}$.

Remark: Any μ measure on $\mathcal{P}(X)$ is an outer measure. However, the converse is not true. In particular, $\exists A, B \subseteq \mathbb{R}$ s.t. $A \cap B = \emptyset$ and $\lambda^*(A \cup B) < \lambda^*(A) + \lambda^*(B)$.

2.6.1 Carathéodory's criterion

Definition 2.6.2 (Carathéodory's condition). Let μ^* be an outer measure on $\mathcal{P}(X)$. A set $E \subseteq X$ is μ^* -**measurable** if $\forall A \subseteq X$:

$$\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Lemma 2.6.3 (Equivalence of Carathéodory's condition). E is μ^* -measurable $\iff \forall A \subseteq X, \mu^*(A) < \infty$:

$$\mu^*(A) \geq \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Proof. The proof is as follows:

(\Rightarrow) : Trivial

(\Leftarrow) : Let $A \subseteq X$, such that $\mu^*(A) < \infty$ and:

$$\mu^*(A) \geq \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Then, notice that $\{A \cap E, A \cap E^c\}$ is a covering of A . By subadditivity:

$$\mu^*(A) \leq \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Therefore, we have that:

$$\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \cap E^c)$$

Meaning that E is μ^* -measurable. This concludes the proof. ■

Theorem 2.6.4 (Carathéodory). *Let μ^* be an outer measure on $\mathcal{P}(X)$. The family:*

$$\mathcal{M} = \{E \subseteq X : E \text{ is } \mu^*\text{-measurable}\}$$

is a σ -algebra, and μ^ restricted to \mathcal{M} (denoted $\mu = \mu^*|_{\mathcal{M}}$) is a complete measure.*

Remark: (X, \mathcal{M}, μ) as in the above theorem is sometimes called the “abstract Lebesgue measure space”. We will only prove the completeness of μ .

Lemma 2.6.5. *Let (X, \mathcal{M}, μ) be the measure space as in Carathéodory’s theorem. Then, any $N \subseteq X$ s.t. $\mu^*(N) = 0$ is μ -measurable, i.e., $N \in \mathcal{M}$, and $\mu(N) = 0$.*

Proof. We have to show that N satisfies Carathéodory’s condition, or equivalently, that it satisfies the lemma 2.6.3. Let $A \subseteq X$ be arbitrary. Then, notice that:

$$\mu^*(A \cap N) \leq \mu^*(N) = 0$$

Also, notice that:

$$\mu^*(A \cap N^c) \leq \mu^*(A)$$

Therefore, we have that:

$$\mu^*(A \cap N) + \mu^*(A \cap N^c) \leq 0 + \mu^*(A) = \mu^*(A)$$

By lemma 2.6.3, we have that N is μ^* -measurable. By Carathéodory’s theorem, we have that N is μ -measurable. Finally, we have that $\mu(N) = \mu^*(N) = 0$. ■

Corollary 2.6.5.1. *μ as in Carathéodory’s theorem is a complete measure.*

Proof. Let $N \subseteq E$, and $\mu(E) = 0$ ($E \in \mathcal{M}$). Then, we have that:

$$\mu(E) = 0 \implies \mu^*(E) = 0$$

By monotonicity, we have that:

$$\mu^*(N) \leq \mu^*(E) = 0$$

Then, $\mu(N) = \mu^*(N) = 0$, thus $N \in \mathcal{M}$. This concludes the proof. ■

2.7 Lebesgue measure

Definition 2.7.1. Let $\mathcal{E} = \{(a, b) : a, b \in \overline{\mathbb{R}}, a \leq b\}$. Define:

$$\lambda^*((a, b)) = b - a$$

Then, λ^* is the **Lebesgue outer measure** on \mathbb{R} .

Theorem 2.7.1. Let λ^* be the Lebesgue outer measure on $\mathcal{E} = \{(a, b) : a, b \in \overline{\mathbb{R}}, a \leq b\}$. Then, by Carathéodory's theorem, the family:

$$\mathcal{L}(\mathbb{R}) = \{E \subseteq \mathbb{R} : E \text{ is } \lambda^*\text{-measurable}\}$$

is a σ -algebra, called the **Lebesgue σ -algebra**, and λ^* restricted to $\mathcal{L}(\mathbb{R})$ (denoted $\lambda = \lambda^*|_{\mathcal{L}(\mathbb{R})}$) is a complete measure, called the **Lebesgue measure**.

Proof. The proof is omitted. ■

Remark: The measure space $(\mathbb{R}, \mathcal{L}(\mathbb{R}), \lambda)$ is called the **Lebesgue measure space**.

Proposition 2.7.2. Let λ be the Lebesgue measure on \mathbb{R} . Then:

- (i) $a \in \mathbb{R} \implies \{a\} \in \mathcal{L}(\mathbb{R})$ and $\lambda(\{a\}) = 0$
- (ii) $E \subset \mathbb{R}$ at most countable $\implies E \in \mathcal{L}(\mathbb{R})$ and $\lambda(E) = 0$

Proof. The proof is as follows:

- (i) Let $a \in \mathbb{R}$. Then, we have that, for any $\varepsilon > 0$:

$$E_1 = (a - \varepsilon, a + \varepsilon), \quad E_2 = E_3 = \dots = \emptyset$$

is a covering of $\{a\}$. Then, by definition of λ^* :

$$0 \leq \lambda^*(\{a\}) \leq \sum_{n=1}^{\infty} \lambda(E_n) = 2\varepsilon$$

As ε is arbitrary, we have that $\lambda^*(\{a\}) = 0$. By Lemma 2.6.5, we then have that $\{a\} \in \mathcal{L}(\mathbb{R})$ and $\lambda(\{a\}) = 0$.

(ii) Let $E \subseteq \mathbb{R}$ be at most countable. Then, we have that:

$$E = \bigcup_{a \in E} \{a\}$$

Because $\{a\} \in \mathcal{L}(\mathbb{R})$ and $\lambda(\{a\}) = 0$, we have that $E \in \mathcal{L}(\mathbb{R})$ and:

$$\lambda(E) = \lambda\left(\bigcup_{a \in E} \{a\}\right) = \sum_{a \in E} \lambda(\{a\}) = 0$$

■

Remark: We can point out 2 important consequences of the above proposition:

1. Arguing as in (i), we can show that the Lebesgue measure is translation invariant. That is, $\forall E \in \mathcal{L}(\mathbb{R}), \forall x \in \mathbb{R}$:

$$\lambda(E + x) = \lambda(E)$$

2. In particular, since \mathbb{Q} is countable, we have that $\mathbb{Q} \in \mathcal{L}(\mathbb{R})$ and $\lambda(\mathbb{Q}) = 0$. In the measure sense, \mathbb{Q} has very few elements with respect to \mathbb{R} . On the other hand, \mathbb{Q} is dense in \mathbb{R} . In the topology sense, \mathbb{Q} has a lot of points.

Proposition 2.7.3. *We have that: $\mathcal{B}(\mathbb{R}) \subset \mathcal{L}(\mathbb{R})$*

Proof. Since $\mathcal{B}(\mathbb{R}) = \sigma_0(\{(a, \infty) : a \in \mathbb{R}\})$, if we show that $(a, \infty) \in \mathcal{L}(\mathbb{R}), \forall a \in \mathbb{R}$, then the prop. follows.

Take $A \subset \mathbb{R}$, s.t. $\lambda^*(A) < \infty$. Then, we must show that:

$$\lambda^*(A) \geq \lambda^*(A \cap (a, \infty)) + \lambda^*(A \cap (-\infty, a])$$

Moreover, by a previous remark, one can assume that $a \notin A$. Then, take any countable covering of A by open intervals:

$$A \subseteq \bigcup_n I_n$$

Then, let us define $A_{left} = A \cap (-\infty, a]$ and $I_{n,left} = I_n \cap (-\infty, a]$. Then, we notice that $\{I_{n,left}\}$ is a covering of A_{left} .

In the same way, we define $A_{right} = A \cap (a, \infty)$ and $I_{n,right} = I_n \cap (a, \infty)$. Then, we notice that $\{I_{n,right}\}$ is a covering of A_{right} .

Then, we have that:

$$\lambda^*(A_{left}) \leq \sum_n \lambda^*(I_{n,left})$$

$$\lambda^*(A_{right}) \leq \sum_n \lambda^*(I_{n,right})$$

Summing both inequalities, we have that:

$$\begin{aligned} \lambda^*(A_{left}) + \lambda^*(A_{right}) &\leq \sum_n \lambda^*(I_{n,left}) + \sum_n \lambda^*(I_{n,right}) \\ &= \sum_n \lambda^*(I_n) \end{aligned}$$

Taking the infimum over all countable coverings of A , we have that:

$$\lambda^*(A_{left}) + \lambda^*(A_{right}) \leq \lambda^*(A)$$

■

Remark: In particular, we have that $\forall (a, b) \subset \mathbb{R}$:

$$(a, b) \in \mathcal{L}(\mathbb{R}), \quad \lambda((a, b)) = b - a$$

We know that:

$$\mathcal{B}(\mathbb{R}) \subset \mathcal{L}(\mathbb{R}) \subset \mathcal{P}(\mathbb{R})$$

Are these inclusions strict? We already know that $\mathcal{L}(\mathbb{R}) \neq \mathcal{P}(\mathbb{R})$, by Ulam's theorem. In particular, $\exists E \subset \mathbb{R}$ not Lebesgue measurable (**Vitali sets**). More precisely:

$$\forall E \in \mathcal{L}(\mathbb{R}), \text{ s.t } \lambda(E) > 0, \exists V \subset E, \text{ s.t } V \notin \mathcal{L}(\mathbb{R})$$

The relation between $\mathcal{B}(\mathbb{R})$ and $\mathcal{L}(\mathbb{R})$ is more subtle. It is clarified by the following proposition:

Proposition 2.7.4 (Regularity of the Lebesgue measure). *Let $E \in \mathbb{R}$. Then, the following are equivalent:*

(i) $E \in \mathcal{B}(\mathbb{R})$

(ii) $\forall \varepsilon > 0, \exists A \subset \mathbb{R}$ open set s.t.

$$E \subset A \quad \text{and} \quad \lambda^*(A \setminus E) < \varepsilon$$

(iii) $\forall \varepsilon > 0, \exists G \subset \mathbb{R}$ of class G_δ s.t.

$$E \subset G \quad \text{and} \quad \lambda^*(G \setminus E) = 0$$

(iv) $\forall \varepsilon > 0, \exists C \subset \mathbb{R}$ closed set s.t.

$$C \subset E \quad \text{and} \quad \lambda^*(E \setminus C) < \varepsilon$$

(v) $\forall \varepsilon > 0, \exists F \subset \mathbb{R}$ of class F_σ s.t.

$$F \subset E \quad \text{and} \quad \lambda^*(E \setminus F) = 0$$

We get as a consequence the following:

Corollary 2.7.4.1. $\forall E \in \mathcal{L}(\mathbb{R}), \exists F, G \in \mathcal{B}(\mathbb{R})$ s.t. $F \subset E \subset G$ and

$$\lambda(G \setminus F) = \lambda(G \setminus E) + \lambda(E \setminus F) = 0$$

In other words:

$$\mathcal{L}(\mathbb{R}) = \overline{\mathcal{B}(\mathbb{R})}$$

(But $\mathcal{L}(\mathbb{R}) \neq \mathcal{B}(\mathbb{R})$).

Proof. (Regularity of the Lebesgue measure). The proof goes as follows:

(i) \Rightarrow (ii) :

Let $E \in \mathcal{B}(\mathbb{R})$. Note that, since $A \in \mathcal{L}(\mathbb{R})$ for all A open, we have that:

$$\lambda^*(A \setminus E) = \lambda(A \setminus E)$$

By definition of λ^* , we have that $\forall \varepsilon > 0, \exists \{I_n\}_{n \in \mathbb{N}}$ s.t.

$$E \subset \bigcup_n I_n \quad \text{and} \quad \sum_n \lambda(I_n) < \lambda^*(E) + \varepsilon$$

Then, set $A = \bigcup_n I_n$. We have that A is open, $E \subset A$ and:

$$\begin{aligned} \lambda(A) &\leq \sum_n \lambda(I_n) < \lambda(E) + \varepsilon \\ \implies \lambda(A \setminus E) &= \lambda(A) - \lambda(E) < \varepsilon \end{aligned}$$

(ii) \Rightarrow (iii) :

Assume $\forall \varepsilon > 0$, $\exists A_\varepsilon$ open s.t. $E \subset A_\varepsilon$ and $\lambda(A_\varepsilon \setminus E) < \varepsilon$. Then, set $\varepsilon = 1/n$, $n \geq 1$ (for ease of notation, $A_n = A_{1/n}$) and define:

$$G = \bigcap_n A_n$$

Then, G is a G_δ set, $E \subset G$ and:

$$0 \leq \lambda^*(G \setminus E) \leq \lambda^*(A_n \setminus E) < 1/n$$

Taking the limit, we have that $\lambda(G \setminus E) = 0$.

(iii) \Rightarrow (i) :

We know that $E \subset G$, $G \in \mathcal{L}(\mathbb{R})$ with $\lambda(G \setminus E) = 0$. Then, we have that:

$$E = G \setminus (G \setminus E) \in \mathcal{L}(\mathbb{R})$$

since $G \in \mathcal{L}(\mathbb{R})$ and $G \setminus E \in \mathcal{L}(\mathbb{R})$. The last is because it is a negligible set and λ is complete. ■

E.g. (Cantor set): Let $T_0 = [0, 1]$. Then, construct T_{n+1} from T_n (recursively) by removing the inner third part of every interval in T_n :

$$\begin{aligned} T_0 &= [0, 1], \\ T_1 &= [0, 1/3] \cup [2/3, 1], \\ T_2 &= [0, 1/9] \cup [2/9, 1/3] \cup [2/3, 7/9] \cup [8/9, 1], \dots \end{aligned}$$

Then, define the **Cantor set** as:

$$C = \bigcap_n T_n$$

It can be proven that:

- C has the cardinality of \mathbb{R}
- $\lambda(C) = 0$
- C is compact
- C is nowhere dense (has no interior points), i.e., $\text{int}(C) = \emptyset$
- $\exists E \subset C$ s.t. $E \in \mathcal{L}(\mathbb{R})$ but $E \notin \mathcal{B}(\mathbb{R})$

2.8 Measurable functions

Definition 2.8.1. Given $f : X \rightarrow Y$, it is well-defined the **preimage** (or counterimage) of f as:

$$f^{-1} : \mathcal{P}(Y) \rightarrow \mathcal{P}(X), \quad f^{-1}(E) = \{x \in X : f(x) \in E\}$$

Remark: Preimages work well with set operations:

- $f^{-1}(E \cup F) = f^{-1}(E) \cup f^{-1}(F)$
- $f^{-1}(E \cap F) = f^{-1}(E) \cap f^{-1}(F)$
- $f^{-1}(E \setminus F) = f^{-1}(E) \setminus f^{-1}(F)$

Note this is not true for images.

Definition 2.8.2. Let (X, \mathcal{M}) and (Y, \mathcal{N}) be measurable spaces. A function $f : X \rightarrow Y$ is **measurable** if $\forall E \in \mathcal{N}$, we have that $f^{-1}(E) \in \mathcal{M}$. We also say that f is **$(\mathcal{M}, \mathcal{N})$ -measurable**.

Proposition 2.8.1. Let (X, \mathcal{M}) and (Y, \mathcal{N}) be measurable spaces, and $\rho \subset \mathcal{N}$ s.t. $\mathcal{N} = \sigma_0(\rho)$. Then, $f : X \rightarrow Y$ is measurable $\iff \forall E \in \rho$, we have that $f^{-1}(E) \in \mathcal{M}$.

Proof. The proofs goes as follows:

(\Rightarrow) : Trivial

(\Leftarrow) : Define $\Sigma := \{E \subset Y : f^{-1}(E) \in \mathcal{M}\}$. We have:

- $\rho \subset \Sigma$ as a consequence of $(\forall E \in \rho, f^{-1}(E) \in \mathcal{M})$

- Σ is a σ -algebra (check as an exercise)

Then, we have that $\mathcal{N} = \sigma_0(\rho) \subset \Sigma$. Therefore, f is measurable. ■

Definition 2.8.3. Suppose that $\mathcal{M} \supseteq \mathcal{B}(X)$ and $\mathcal{N} = \mathcal{B}(Y)$. We say that $f : X \rightarrow Y$ is:

- **Borel measurable** if f is $(\mathcal{B}(X), \mathcal{B}(Y))$ -measurable.
- **Lebesgue measurable** if it is $(\mathcal{M}, \mathcal{B}(Y))$ -measurable.

Remark: If $f : X \rightarrow Y$ is Borel measurable, then it is Lebesgue measurable. The converse is not true.

Also, we never deal with $\mathcal{L}(Y)$.

Corollary 2.8.1.1. f is Borel measurable $\iff f^{-1}(E) \in \mathcal{B}(X), \forall E \in Y$ open.
Also, f is Lebesgue measurable $\iff f^{-1}(E) \in \mathcal{M}, \forall E \in Y$ open.

Proof. It follows from the previous proposition, since $\mathcal{B}(Y) = \sigma_0(\{E \subset Y : E \text{ open}\})$. ■

Definition 2.8.4. We say that f is **continuous** $\iff f^{-1}(E) \subset X$ is open $\forall E \subset Y$ open.

Proposition 2.8.2. If $f : X \rightarrow Y$ is continuous, then f is Borel measurable (and thus Lebesgue measurable).

Proof. Let $E \subset Y$ be open. By continuity of f , we have that $f^{-1}(E)$ is open. Then $f^{-1}(E) \in \mathcal{B}(X)$, and thus f is Borel measurable.

Note that the proposition is false when $\mathcal{N} \supsetneq \mathcal{B}(Y)$. ■

2.8.1 Operations on measurable functions

Proposition 2.8.3. *Let $f : X \rightarrow Y$ be Lebesgue measurable, and $g : Y \rightarrow Z$ be continuous. Then:*

$$g \circ f : X \rightarrow Z \text{ is Lebesgue measurable}$$

Corollary 2.8.3.1. *Let $f : X \rightarrow Y$ be Lebesgue measurable. Then:*

- $f^+(x) = \max\{f(x), 0\}$ is Lebesgue measurable
- $f^-(x) = \max\{-f(x), 0\}$ is Lebesgue measurable
- $|f(x)| = f^+(x) + f^-(x)$ is Lebesgue measurable

Proof. Let f be Lebesgue measurable, and $g : \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Then, take $E \subset Z$ open. We have that:

$$(g \circ f)^{-1}(E) = f^{-1}(g^{-1}(E))$$

Since g is continuous, $g^{-1}(E)$ is open. Then, $f^{-1}(g^{-1}(E)) \in \mathcal{M}$ ■

Proposition 2.8.4. *Let $f, g : X \rightarrow \mathbb{R}$ be Lebesgue measurable, and $\Phi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous. Then, $h(x) = \Phi(f(x), g(x))$ is Lebesgue measurable.*

Proof. Let $h(x) = \Phi(f(x), g(x)) = (\Phi \circ \Psi)(x)$, where $\Psi : X \rightarrow \mathbb{R}^2$ is defined as

$$\Psi(x) = (f(x), g(x))$$

Then, we should prove that Ψ is Lebesgue measurable for applying the previous proposition. For this, we have to show that $\forall (a, b) \times (c, d) \subset \mathbb{R}^2$, we have that:

$$\Psi^{-1}((a, b) \times (c, d)) = \{x \in X : f(x) \in (a, b), g(x) \in (c, d)\} \in \mathcal{M}$$

This can be done using the fact that f and g are Lebesgue measurable. ■

Corollary 2.8.4.1. *Let $f, g : X \rightarrow \mathbb{R}$ be Lebesgue measurable. Then:*

- $f + g$ is Lebesgue measurable
- $f \cdot g$ is Lebesgue measurable

Proposition 2.8.5. *Let (X, \mathcal{M}) be a measurable space (with $\mathcal{M} \supseteq \mathcal{B}(X)$), and $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of Lebesgue measurable functions $f_n : X \rightarrow \mathbb{R}$. Then, the following functions are Lebesgue measurable:*

1. $\sup_n f_n$
2. $\inf_n f_n$
3. $\limsup_n f_n$
4. $\liminf_n f_n$

In particular, if $\lim_n f_n$ exists, then it is Lebesgue measurable.

Proof. The proof goes as follows:

1. Since $\mathcal{B}(\mathbb{R}) = \sigma_0(\{(a, \infty) : a \in \mathbb{R}\})$, it is enough to show that $\forall a \in \mathbb{R}$, we have that:

$$(\sup_n f_n)^{-1}((a, \infty)) = \{x \in X : \sup_n f_n(x) > a\} \in \mathcal{M}$$

This can be done by using the fact that f_n is Lebesgue measurable. Indeed, we have that:

$$\begin{aligned} \{x \in X : \sup_n f_n(x) > a\} &= \bigcup_n \{x \in X : f_n(x) > a\} \\ &= \bigcup_n f_n^{-1}((a, \infty)) \in \mathcal{M} \end{aligned}$$

because $f_n^{-1}((a, \infty)) \in \mathcal{M}$ for all n .

2. The proof is analogous to the previous case, taking that:

$$\inf_n f_n = -\sup_n (-f_n)$$

3. We have that:

$$\limsup_n f_n = \inf_n \sup_{k \geq n} f_k$$

4. We have that:

$$\liminf_n f_n = \sup_n \inf_{k \geq n} f_k$$

■

2.8.2 Properties holding almost everywhere

Definition 2.8.5. Let (X, \mathcal{M}, μ) be a complete measure space. We say that a property $P(x)$ holds μ -almost everywhere (a.e) if:

$$\mu(\{x \in X : P(x) \text{ is false}\}) = 0$$

In other words, $P(x)$ holds μ -almost everywhere if it holds everywhere except for a set of measure zero.

E.g.: Let $f(x) = x^2$. Is it true that $f(x) > 0$ a.e.?

We have that $\{x : x^2 \leq 0\} = \{0\}$

- In $(\mathbb{R}, \mathcal{L}(\mathbb{R}), \lambda)$, the property is true a.e., since $\lambda(\{0\}) = 0$
- In $(\mathbb{R}, \mathcal{P}(\mathbb{R}), \mu_{\#})$ (counting measure), the property is false a.e., since $\mu_{\#}(\{0\}) = 1$

Proposition 2.8.6. Let (X, \mathcal{M}, μ) be a measure space:

1. $f : X \rightarrow \overline{\mathbb{R}}$ s.t. $f = g$ a.e, with g measurable $\implies f$ is measurable
2. $\{f_n\}_{n \in \mathbb{N}}$ a sequence of measurable functions s.t. $f_n \rightarrow f$ a.e., then f is measurable.

2.8.3 Simple functions

Definition 2.8.6. Let (X, \mathcal{M}) be a measurable space. A function $s : X \rightarrow \overline{\mathbb{R}}$ is measurable and **simple** if s is measurable and $s(X)$ is a finite set:

$$s(X) = \{a_1, a_2, \dots, a_k\}$$

where $a_i \in \overline{\mathbb{R}} \forall i$, with $a_i \neq a_j$ for $i \neq j$. Then, s can be written as:

$$s(x) = \sum_{i=1}^k a_i \cdot \chi_{A_i}(x)$$

where $A_i = s^{-1}(\{a_i\})$, $A_i \cap A_j = \emptyset$ for $i \neq j$, $\bigcup_{i=1}^k A_i = X$ and $A_i \in \mathcal{M}$, $\forall i$.

Particular case:

If $X = \mathbb{R}$ (or $(a, b) \subset \mathbb{R}$) and A_i is an interval $\forall i$, then s is called a **step function**.

On the other hand, $\chi_{\mathbb{Q}}$ is a simple function, but not a step function:

$$\chi_{\mathbb{Q}}(x) = \begin{cases} 1 & x \in \mathbb{Q} \\ 0 & x \notin \mathbb{Q} \end{cases}$$

Remark: One may define simple functions without measurability requirements.

Goal:

Approximate any measurable function $f : X \rightarrow \overline{\mathbb{R}}$ with (measurable and) simple functions.

Theorem 2.8.7 (Simple approximation theorem (SAT)). *Take (X, \mathcal{M}) measurable space and $f : X \rightarrow [0, \infty]$, measurable. Then $\exists \{s_n\}_{n \in \mathbb{N}}$ a sequence of measurable, simple functions s.t. $s_1 \leq s_2 \leq \dots \leq f$ pointwise (i.e., $\forall x \in X$) and:*

$$\lim_{n \rightarrow \infty} s_n(x) = f(x) \quad \forall x \in X$$

Moreover, if f is bounded, the convergence is uniform:

$$\lim_{n \rightarrow \infty} \sup_{x \in X} |s_n(x) - f(x)| = 0$$

Proof. In case f is bounded, say $0 \leq f < 1$.

For any $n \geq 1$, divide $[0, 1)$ into 2^n intervals of length 2^{-n} , and define:

$$A_n^{(i)} = \{x \in X : \frac{i}{2^n} \leq f(x) < \frac{i+1}{2^n}\}$$

and:

$$s_n(x) = \sum_{i=0}^{2^n-1} \frac{i}{2^n} \cdot \chi_{A_n^{(i)}}(x)$$

One can show that such a sequence has the required properties ■

2.9 Lebesgue integral

- First, we will define the integral for non-negative simple (measurable) functions.
- Second, we will extend the definition to non-negative measurable functions.
- Finally, we will define the integral for general measurable functions.

2.9.1 Integral of non-negative simple functions

Definition 2.9.1. Let $s : X \rightarrow [0, \infty]$ be a measurable and simple function:

$$s = \sum_{i=1}^k a_i \cdot \chi_{A_i}$$

where $a_i \geq 0$ and $A_i \in \mathcal{M}$. Let $E \in \mathcal{M}$. Then, we define the **(Lebesgue) integral** of s over E as:

$$\int_E s d\mu = \sum_{i=1}^k a_i \cdot \mu(A_i \cap E)$$

Remark: There are some remarks:

1. $s : [a, b] \rightarrow [0, \infty)$, $\mu, \mu = \lambda$ (Lebesgue measure)
Then, $\int_{[a,b]} s d\mu = \text{area under the graph of } s \text{ in } [a, b]$
2. We are already using $0 \cdot \infty = 0$ in the definition. In particular,

$$a_i \cdot \mu(A_i \cap E) = \begin{cases} 0 & a_i = 0 \\ \infty & a_i > 0 \end{cases}$$

if $\mu(A_i \cap E) = \infty$.

3. $D \in \mathcal{M}$, then χ_D is a simple function, and:

$$\int_E \chi_D d\mu = \mu(D \cap E)$$

4. More generally, s simple and measurable, $E \in \mathcal{M}$, then:

$$\int_E s d\mu = \int_X s \cdot \chi_E d\mu$$

Properties 2.9.1 (Basic properties). Let $N, E, F \in \mathcal{M}$, $s_1, s_2 : X \rightarrow [0, \infty)$ simple and measurable functions. Then:

(i) If $\mu(N) = 0$, then:

$$\int_N s_1 d\mu = 0$$

(ii) If $0 \leq c \leq \infty$, then:

$$\int_E c \cdot s_1 d\mu = c \cdot \int_E s_1 d\mu$$

(iii) $\int_E (s_1 + s_2) d\mu = \int_E s_1 d\mu + \int_E s_2 d\mu$

(iv) If $s_1 \leq s_2$, then:

$$\int_E s_1 d\mu \leq \int_E s_2 d\mu$$

(v) if $E \subset F$, then:

$$\int_E s_1 d\mu \leq \int_F s_1 d\mu$$

The properties (ii) and (iii) are called **linearity** of the integral. The properties (iv) and (v) are called **monotonicity** of the integral.

Proposition 2.9.1. *Let $s : X \rightarrow [0, \infty)$ be a simple measurable function. Then, the function:*

$$\phi(E) := \int_E s \, d\mu : \mathcal{M} \rightarrow [0, \infty]$$

is a measure on (X, \mathcal{M}) .

Proof. Let $s = \sum_{i=1}^k a_i \cdot \chi_{A_i}$, $0 \leq a_i \leq \infty$. We have to show that:

1. $\phi : \mathcal{M} \rightarrow [0, \infty]$?: Yes, since $s \geq 0$, $\phi(E) \geq 0$, $\forall E \in \mathcal{M}$.
2. $\phi(\emptyset) = 0$?: Yes, since $\int_{\emptyset} s \, d\mu = 0$, as $\mu(\emptyset) = 0$.
3. σ -additivity?: Let $\{E_n\}_{n \in \mathbb{N}}$ be a sequence of pairwise disjoint sets in \mathcal{M} , and $E = \bigcup_n E_n$. Then, we have that:

$$\begin{aligned} \phi(E) &= \int_E s \, d\mu = \int_X s \cdot \chi_E \, d\mu = \sum_{i=1}^k a_i \cdot \mu(A_i \cap E) \\ &= \sum_{i=1}^k a_i \cdot \mu\left(\bigcup_n A_i \cap E_n\right) \end{aligned}$$

Since μ is σ -additive, we have that:

$$\begin{aligned} &= \sum_{i=1}^k a_i \sum_n \mu(A_i \cap E_n) \\ &= \sum_n \sum_{i=1}^k a_i \cdot \mu(A_i \cap E_n) \\ &= \sum_n \int_{E_n} s \, d\mu = \sum_n \phi(E_n) \end{aligned}$$

■

2.9.2 Integral of non-negative measurable functions

Definition 2.9.2. Let $f : X \rightarrow [0, \infty]$ be a measurable function, $E \in \mathcal{M}$. Then, we define the **(Lebesgue) integral** of f over E as:

$$\int_E f \, d\mu = \sup \left\{ \int_E s \, d\mu : s \text{ simple, measurable and } 0 \leq s \leq f \right\}$$

Remark: There are some remarks:

1. If f is simple, then the definition coincides with the previous one.
2. $(\mathbb{N}, \mathcal{P}(\mathbb{N}), \mu_{\#})$. Then $f : \mathbb{N} \rightarrow [0, \infty]$ is a sequence. Indeed, if we name $f_n = f(n)$, then:

$$\int_{\mathbb{N}} f d\mu_{\#} = \sum_n f_n$$

3. All the basic properties of the integral for simple functions above hold for this new definition.