

Brave Care
6924 NE Sandy Blvd
Portland, OR 97213
5039637963

Patient Name: BERTLER, RACHEL SHEHER
Patient DOB: 11/03/1988
Accession#: 7968517
Ord Phys: ROBERTS, CHELSEA NPI

Gender: F

Patient ID: 195596354

Lab Account#: 97213002 (Quest - Seattle)
Lab Accession#: OW173145J
Comment:

Collected: 05/13/20 11:00

Received: 05/14/20 02:27
Reported: 05/15/20 14:50

Test Name	Result	Unit	Range	Reference	Status	Date/Tm	Lab
DHEA SULFATE	187	mcg/dL	23 - 266		Final	05/15/20 14:50	EN
HEMOGLOBIN A1c	5.0	% of total Hgb	<5.7		Final	05/15/20 14:50	NW

For the purpose of screening for the presence of diabetes:

<5.7% Consistent with the absence of diabetes
5.7-6.4% Consistent with increased risk for diabetes (prediabetes)
> or =6.5% Consistent with diabetes

This assay result is consistent with a decreased risk of diabetes.

Currently, no consensus exists regarding use of hemoglobin A1c for diagnosis of diabetes in children.

According to American Diabetes Association (ADA) guidelines, hemoglobin A1c <7.0% represents optimal control in non-pregnant diabetic patients. Different metrics may apply to specific patient populations. Standards of Medical Care in Diabetes(ADA).

INSULIN	---	The following results were previously reported on a prior report with lab accession OW187879J.---	3.2	uIU/mL	Final	05/15/20 11:44	NW
---------	-----	---	-----	--------	-------	----------------	----

Reference Range < or = 19.6

Risk:
Optimal < or = 19.6
Moderate NA
High >19.6

Adult cardiovascular event risk category cut points (optimal, moderate, high) are based on Quest Diagnostics population data from 12/2011.

This insulin assay shows strong cross-reactivity for some insulin analogs (lispro, aspart, and glargine) and much lower cross-reactivity with others (detemir, glulisine).