

# Math Template

Rohan Mukherjee

May 24, 2024

As per the usual, let  $H_1 = \begin{pmatrix} 1 \end{pmatrix}$  and define the  $n$ th hadamard matrix of size  $2^n \times 2^n$  recursively by:

$$H_n = \begin{pmatrix} H_{n-1} & H_{n-1} \\ H_{n-1} & -H_{n-1} \end{pmatrix}$$

Decompose the vector  $x \in \{-1, 1\}^{2^n}$  as  $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$  where  $x_1, x_2 \in \{-1, 1\}^{2^{n-1}}$ . Then, we can write:

$$H_n x = \begin{pmatrix} H_{n-1} & H_{n-1} \\ H_{n-1} & -H_{n-1} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} H_{n-1}x_1 + H_{n-1}x_2 \\ H_{n-1}x_1 - H_{n-1}x_2 \end{pmatrix}$$

If the max coordinate in  $H_{n-1}x_1$  has the same sign as the max coordinate in  $H_{n-1}x_2$ , then the max coordinate in  $H_n x$  is just  $|H_{n-1}x_1 + H_{n-1}x_2|$ . On the other hand, if they have differing signs, the max coordinate is just  $|H_{n-1}x_1 - H_{n-1}x_2|$ . Since  $|H_{n-1}x_1| \approx \beta(H_{n-1})$ , we have that that  $|H_n x| \approx 2\beta(H_{n-1})$ . Normalizing  $H_n$  will give that  $|\frac{1}{\sqrt{2^n}} H_n x| \approx \frac{2}{\sqrt{2}} \beta\left(\frac{1}{\sqrt{2^{n-1}}} H_{n-1}\right)$ . This would say that  $\beta(H_n) \approx \sqrt{2^n}$  up to constants.

We write

$$2^8 = 2^4 \cdot 7 + 2^7 + 2^4$$

$$2^{16} = 2^7 \cdot 7 + 2^{11} \cdot 7 + 2^5 \cdot 5^3 \cdot 7 + 2^9 \cdot 5 \cdot 7 + 2^8 \cdot 3 \cdot 5 + 2^9 + 2^5$$