
What is left of the axon response when the controls have the same behaviour
square cue 0-3 s | spout + reward
square 3-10 s | spout + reward
square cue 0-3 s | no spout
square 3-10 s | no spout
not testable (too few behaviour matches)
rewarded - spontaneous lick bout
hit - miss trial
not testable (too few behaviour matches)
flash
moving bar
expanding loom
receding loom
expanding - receding
not testable (too few behaviour matches)
4/4 sess, 1 animal, 94% matched
p shift = 0.040 | p benchmark = 0.040
4/4 sess, 1 animal, 92% matched
p shift = 0.30 | p benchmark = 0.65
3/3 sess, 2 animals, 78% matched
p shift = 0.45 | p benchmark = 0.64
n=0/3 sessions
3/4 sess, 1 animal, 81% matched
p shift = 0.88 | p benchmark = 0.65
n=0/4 sessions
4/5 sess, 2 animals, 65% matched
p shift = 0.003 | p benchmark = 0.030
5/5 sess, 3 animals, 97% matched
p shift = 0.040 | p benchmark = 0.21
4/4 sess, 2 animals, 97% matched
p shift = 0.040 | p benchmark = 0.035
3/4 sess, 2 animals, 100% matched
p shift = 0.053 | p benchmark = 0.025
n=0/3 sessions
−0.050 −0.025 0.000 0.025 0.050 0.075 0.100 0.125 0.150
event-window axon dF/F minus control windows
naive: event - all controls
regression, no matching
matched (filled = unflagged, hollow = flagged)
behaviour-only benchmark