
What is left of the axon response when the controls have the same behaviour
square cue 0-3 s | spout + reward
square 3-10 s | spout + reward
square cue 0-3 s | no spout
square 3-10 s | no spout
rewarded - spontaneous lick bout
hit - miss trial
flash
moving bar
expanding loom
receding loom
expanding - receding
not testable (too few behaviour matches)
not testable (too few behaviour matches)
not testable (too few behaviour matches)
5/5 sess, 1 animal, 94% matched
p shift = 0.090 | p benchmark = 0.100
5/5 sess, 1 animal, 92% matched
p shift = 0.99 | p benchmark = 0.95
3/3 sess, 2 animals, 78% matched
p shift = 0.57 | p benchmark = 0.71
n=0/3 sessions
4/5 sess, 1 animal, 81% matched
p shift = 0.23 | p benchmark = 0.42
n=0/5 sessions
4/5 sess, 2 animals, 65% matched
p shift = 0.010 | p benchmark = 0.040
5/5 sess, 3 animals, 98% matched
p shift = 0.72 | p benchmark = 0.83
3/4 sess, 2 animals, 98% matched
p shift = 0.056 | p benchmark = 0.040
3/4 sess, 2 animals, 100% matched
p shift = 0.030 | p benchmark = 0.015
n=0/3 sessions
−0.050 −0.025 0.000 0.025 0.050 0.075 0.100 0.125
event-window axon dF/F minus control windows
naive: event - all controls
regression, no matching
matched (filled = unflagged, hollow = flagged)
behaviour-only benchmark